Pacemaker extrusion from innocuous direct impact to the chest wall

A 59-year-old man presented with giddiness, shortness of breath and lethargy in 2011. ECG showed complete heart block, which necessitated temporary pacing. Echocardiography revealed no regional wall motion abnormality and normal systolic ejection fraction. His cardiovascular risk factors were type 2 diabetes mellitus and hypertension. Subsequently, a dual-chamber pacemaker was implanted with direct puncture of the left subclavian vein for placement of two 6.1 F silicone-coated leads (figure 1). The pacing parameters were acceptable. About

Figure 1 The pacemaker pulse generator protruding through the eroded skin pocket.

19 months post implantation, his 2-year-old grandson bumped into him. The toddler's head hit his left upper chest wall. There was pain and swelling of the left pulse generator (PG) pocket site. Two days later, he noticed protrusion of the PG edge through the eroded skin. Inflammatory blood markers were not raised. There was no evidence of pus at the pocket site. The affected area was thoroughly cleaned and debrided. The PG was wiped with povidone iodine solution before being placed back into a deeper subcutaneous pocket. He completed 1 week course of cloxacillin. His wound healed nicely and remained afebrile during the following month clinic review. This case highlights the occurrence of PG extrusion from an innocuous direct impact force to the chest wall, possibly affected by the superficially placed PG in the thin subcutaneous skin layer. The exposed PG could still be salvaged by meticulous debridement and cleaning with antiseptic and a course of prophylactic antibiotic. Surveillance should be continued for occult or late-onset cardiovascular implantable electronic device infection.

Kuan Leong Yew,¹ Anderson Steven,¹ Farah Razali²

¹Cardiology Department, Heart Center Sarawak General Hospital, Kota Samarahan, Sarawak, Malaysia

²Sarawak General Hospital, Kuching, Sarawak, Malaysia

Correspondence to Dr Kuan Leong Yew, Cardiology Department, Heart Center Sarawak General Hospital, Kota Samarahan, Sarawak 94300, Malaysia; yewkuanleong@yahoo.com

Collaborators Siong Hee Lim

Contributors KLY wrote the manuscript. FR and AS gave suggestion on improvement and proofread the manuscript.

Competing interests None.

Patient consent Obtained.

Provenance and peer review Not commissioned; internally peer reviewed.

To cite Yew KL, Steven A, Razali F. *Heart Asia* Published Online First: [please include Day Month Year] doi:10.1136/heartasia-2013-010355

Heart Asia 2013;0:100. doi:10.1136/heartasia-2013-010355