Development of Novel 1,2,3,4-Tetrahydroquinoline Scaffolds as Potent NF- κ B Inhibitors and Cytotoxic Agents

Hyeju Jo,[†] Minhoo Choi,[†] Arepalli Sateesh Kumar,[†] Yeongeun Jung,[†] Sangeun Kim,[†] Jieun Yun,[‡] Jong-Soon Kang,[‡] Youngsoo Kim,[†] Sang-bae Han,[†] Jae-Kyung Jung,[†] Jungsook Cho,^{||} Kiho Lee,[§] Jae-Hwan Kwak,[⊥] and Heesoon Lee^{*,†}

[†]Department of Pharmacy, Chungbuk National University, Chungbuk 362-763, Republic of Korea

[‡]Korea Research Institute of Bioscience and Biotechnology, Ochang 363-883, Republic of Korea

[§]College of Pharmacy, Korea University, Sejong 339-700, Republic of Korea

^{||}College of Pharmacy, Dongguk University, Goyang 410-773, Republic of Korea

[⊥]College of Pharmacy, Kyungsoo University, Busan 608-736, Republic of Korea

S Supporting Information

ABSTRACT: 1,2,3,4-Tetrahydroquinolines have been identified as the most potent inhibitors of LPS-induced NF- κ B transcriptional activity. To discover new molecules of this class with excellent activities, we designed and synthesized a series of novel derivatives of 1,2,3,4-tetrahydroquinolines (**4a–g**, **5a–h**, **6a–h**, and **7a–h**) and bioevaluated their *in vitro* activity against human cancer cell lines (NCI-H23, ACHN, MDA-MB-231, PC-3, NUGC-3, and HCT 15). Among all synthesized scaffolds, **6g** exhibited the most potent inhibition (53 times that of a reference compound) of LPS-induced NF- κ B transcriptional activity and the most potent cytotoxicity against all evaluated human cancer cell lines.

Inhibition of NF- κ B activity	IC ₅₀ (μM)
Human Cancer Cell lines	GI ₅₀ (μM)
NCI-H23	0.292±0.0111
ACHN	0.526±0.178
MDA-MB-231	0.288±0.0992
PC-3	0.729±0.131
NUGC-3	0.754±0.129
HCT 15	0.797±0.173

KEYWORDS: 1,2,3,4-Tetrahydroquinolines, NF- κ B inactivation, *in vitro* cytotoxicity, human cancer cell lines

NF- κ B is a lymphoid-specific protein that binds to the enhancer of kappa light chain in the nucleus of B cells; NF- κ B was discovered by Sen and Baltimore.¹ NF- κ B is involved in the regulation of many immune and inflammatory responses, cellular growth, and apoptosis.^{2,3} At present, NF- κ B and its signaling is one of the most exciting and extensively studied research fields since NF- κ B dysregulation is associated with many diseases such as cancer, AIDS, asthma, arthritis, diabetes, and inflammatory bowel disease.^{4–6} Several natural and synthetic compounds, including some drugs, have been tested for their potential to inhibit NF- κ B, but very few of them are suitable for anticancer therapy.^{7,8} Therefore, it has been suggested that the development of novel NF- κ B inhibitors with antitumor and anti-inflammatory activities is most important. Our group has been involved in the development of novel potent NF- κ B inhibitors^{9–11} with anticancer activity.

Besides, quinolines and tetrahydroquinolines are important ubiquitous structural motifs in biologically active natural products and pharmacologically relevant therapeutic agents.^{12–16} However, Khan et al.¹⁷ demonstrated that tetrahydroquinolines can be used as NF- κ B inhibitors as well as anti-HIV agents, anti-Parkinson's diseases, etc. María José Abad et al.¹⁸ also tested quinoline-based compounds as modulators of HIV transcription through NF- κ B and Sp1

inhibition. These distinct inhibitory activities have encouraged us to prepare such core motifs to test our hypothesis that their derivatives would act as most potent NF- κ B inhibitors and have anticancer activity. We designed and synthesized different novel derivatives of 1,2,3,4-tetrahydroquinoline-2-carboxylic acid *N*-(substituted)phenyl amide and tested them as potential NF- κ B inhibitors and also evaluated their cytotoxicities against six human cancer cell lines (NCI-H23, ACHN, MDA-MB-231, PC-3, NUGC-3, and HCT-15). Based on our previous reports,^{9–11} we wish to maintain amide functionality to the core motifs, and the effect of the synthesized molecules on NF- κ B transcriptional activity was measured by using a reported procedure.¹⁹ *In vitro* cytotoxicity assay was performed using the number of cells measured indirectly by the sulforhodamine B method according to the National Cancer Institute (USA) protocol.²⁰

We commenced our synthesis from commercially available quinoline-2-carboxylic acid, which underwent amidation reaction with various substituted aromatic amines in the presence of the coupling reagent 1,1'-carbonyldiimidazole

Received: January 6, 2016

Accepted: February 16, 2016

Published: February 16, 2016

Table 1. continued

No.	substituents (R group)	% inhibition at 100 μ M	IC ₅₀ ^c (μ M)	No.	% inhibition at 100 μ M	IC ₅₀ ^c (μ M)	No.	% inhibition at 100 μ M	IC ₅₀ ^c (μ M)
5d	octyl	22		6d	63	27 $\pm$ 0.71	7d	58	51 $\pm$ 21
5e	phenyl	>100	1.4 $\pm$ 0.071	6e	44		7e	66	29 $\pm$ 0.71
5f	2-chlorophenyl	78	20 $\pm$ 0.71	6f	82	0.90 $\pm$ 0.071	7f	90	23 $\pm$ 0.0
5g	3-chlorophenyl	58	78 $\pm$ 2.1	6g	90	0.70 $\pm$ 0.071	7g	68	27 $\pm$ 0.71
5h	4-chlorophenyl	82	20 $\pm$ 2.8	6h	92	2.7 $\pm$ 0.42	7h	92	20 $\pm$ 2.1

^cIC₅₀ values are means of the concentration (μ M) exhibiting 50% inhibition of LPS-induced NF- κ B transcriptional activity.

Figure 1. Efficacy of 4a–g, 5a–h, 6a–h, and 7a–h analogues in inhibition of NF- κ B transcriptional activities.

prepared to study the effect of the core 1,2,3,4-tetrahydroquinoline-2-carboxamide motif. We also examined the electronic influence of different groups at R₁, R₂, R₃, and R₄, including electron-deficient (–CF₃) and electron-rich groups (–OCH₃ and –OH). Next, we introduced various acyl and aroyl groups at the 1-position of tetrahydroquinoline-2-carboxamides, which resulted in 5a–h, 6a–h, and 7a–h analogues. Sizes of different aliphatic chains (–R) and substituents at the aromatic ring were then examined. The inhibitory effects of these compounds (4a–g, 5a–h, 6a–h, and 7a–h) depended on the nature of the substitution at the R-position and the substituents (R₁, R₂, R₃, and R₄) on the N-phenyl ring of tetrahydroquinoline-2-carboxamides. For instance, with R = an alkyl group (methyl, ethyl, propyl, or octyl), we did not notice any significant inhibitory effects, but with R = phenyl. The following compounds exhibited excellent inhibitory effects on LPS-induced NF- κ B transcriptional activity (Figure 2): 5e (IC₅₀: 1.4 $\pm$ 0.071 μ M, about 26 and 37 times more potent than PDTC and KL-1156, respectively), 6f (IC₅₀: 0.90 $\pm$ 0.071 μ M, about 41 and 58 times more potent), 6g (IC₅₀: 0.70 $\pm$ 0.071 μ M, about 53 and 75 times), and 6h (IC₅₀: 2.7 $\pm$ 0.42 μ M, about 13 and 19 times). We also tested other substitutions at the R-position and different substituents at the R₁, R₂, R₃, and R₄ positions of the 1,2,3,4-tetrahydroquinoline core motif (Table 1). To explain the trend toward NF- κ B inhibition interestingly, the electron-withdrawing group –CF₃ at the R₁ position resulted in outstanding inhibitory effects on LPS-induced NF- κ B transcriptional activity in comparison with any electron-releasing group (for example, –OH). We also noticed that –I and +M effect groups (such as –Cl) at the second and third positions of the aryl group of R in combination with –CF₃ at the R₁ position also showed excellent inhibition of NF- κ B transcriptional activity (6g, IC₅₀: 0.70 $\pm$ 0.071 μ M; 6f, IC₅₀: 0.90 $\pm$ 0.071 μ M). Overall, most of the synthesized compounds (4d, 4e, 4g, 5f, 5h, 6d, 7a–c, and 7e–h) were more potent inhibitors of NF- κ B transcriptional activity than PDTC and KL-1156 (Figure 2). We next studied the *in vitro* cytotoxicity of the synthesized compounds and initially evaluated quinoline-2-carboxamide derivatives (3a–g) with the human lung cancer cell line (NCI-H23) and doxorubicin (ADR) as a reference compound.

Figure 2. *In vitro* efficacy of 4a–g, 5a–h, 6a–h, and 7a–h analogues in inhibiting growth of human cancer cell lines.

Table 2. Cytotoxicity of 3a–g Series against Lung Cancer (NCI-H23) Cell Lines

No.	R ₁	R ₂	R ₃	R ₄	GI ₅₀ (μM) ^d
ADR					0.101 ± 0.00602
3a	H	H	H	H	108 ± 8.74
3b	OH	H	H	H	87.5 ± 9.77
3c	H	H	OH	H	102 ± 11.4
3d	H	H	OCH ₃	H	7.65 ± 1.93
3e	CF ₃	H	H	H	>90
3f	H	H	CF ₃	H	>90
3g	H	CF ₃	H	CF ₃	1.44 ± 0.463

^dGI₅₀ values are taken as a mean from three experiments and correspond to the agent's concentration causing a 50% decrease in net cell growth.

As shown in Table 2, electron-withdrawing substituents such as –CF₃ at the R₂ and R₄ positions of the phenyl ring (3g, GI₅₀ 1.44 ± 0.463 μM) resulted in the most potent cytotoxicity. The series 4a–g, 5a–h, 6a–h, and 7a–h were also evaluated for *in vitro* cytotoxicity against six human cancer cells: NCI-H23, ACHN (renal), MDA-MB-231 (breast), PC-3 (prostate), NUGC-3 (gastric), and HCT15 (colon) (Table 3). Any substitution on the phenyl ring was not beneficial, and only 4b (GI₅₀ 2.23 ± 0.455 μM) exhibited better cytotoxic activities against all tested cell lines than other analogues of the 4a–g series (Table 3). To further confirm that the tetrahydroquinoline motif is beneficial for cytotoxicity, we executed acylation reaction with triethyl amine in tetrahydrofuran with 4b, 4e, and 4g; introduction of electron-rich or electron-withdrawing substituents at the R₁ position afforded 5a–h, 6a–h, and 7a–h analogues. As expected, these analogues had improved cytotoxicity against all tested cell lines (Figure 2), suggesting that substitutions at the R₁ position and the first position of the tetrahydroquinoline motif are most important (Table 3). Compound 5e exhibited the highest cytotoxicity (Table 3) against all evaluated cell lines (NCI-H23, GI₅₀ 3.49 ± 0.999 μM; NUGC-3, GI₅₀ 3.78 ± 0.618 μM; HCT 15, GI₅₀ 3.83 ± 0.994 μM). The importance of an electron-withdrawing group was also confirmed by 6g and 6h derivatives, which have –CF₃ and –Cl on both phenyl rings and exhibited outstanding cytotoxicity (6g, GI₅₀ 0.292 ± 0.111–0.797 ± 0.173 μM; 6h, GI₅₀ 0.307 ± 0.0941–0.839 ± 0.0610 μM) against all tested cell lines (Figure 2, Table 3). The –CF₃ group at the R₂ and R₄ positions of the phenyl ring in 7g also resulted in potent cytotoxicity against all tested cell lines (0.420–1.19 μM; Table 3). Compound 7h also exhibited potent cytotoxicity against lung (NCI-H23 (GI₅₀ 0.889 ± 0.102 μM) and gastric (NUGC-3, GI₅₀ 1.66 ± 0.406 μM) cancer cell lines and moderate cytotoxicity against the other four cell lines (Table 3).

In summary, we identified a new class of 1,2,3,4-tetrahydroquinolines as the most potent NF-κB inhibitors and potent cytotoxic agents against human cancer cell lines.

The first round of screening starting from the initial lead scaffold 4a led to the discovery of 6f, 6g, and 6h analogues. These new lead analogues inhibited LPS-induced NF-κB transcriptional activity 41, 53, and 13 times more potently, respectively, than the reference compound PDTC. Analogues 6f, 6g, and 6h have also shown the most potent *in vitro* cytotoxicity against all evaluated human cancer cell lines. Thus, 6f, 6g, 6h, and related analogues provide new chemical tools for development of pathway-selective NF-κB inhibitors with

Table 3. Cytotoxicity against NCI-H23, ACHN, MDA-MB-231, PC-3 NUGC-3, and HCT-15 Cancer Cell Lines

No.	R	R ₁	R ₂	R ₃	R ₄	NCI-H23	ACHN	MDA-MB-231	PC-3	NUGC-3	HCT 15
ADR						0.0726 ± 0.00940	0.0995 ± 0.00848	0.112 ± 0.0132	0.0923 ± 0.0103	0.125 ± 0.0104	0.0854 ± 0.00329
4a	H	H	H	H	H	97.1 ± 15.3	72.9 ± 11.4	108 ± 17.8	110 ± 15.6	59.0 ± 9.88	109 ± 15.0
4b	OH	H	H	H	H	2.23 ± 0.455	3.99 ± 1.33	5.26 ± 1.77	4.31 ± 1.34	2.68 ± 0.824	5.61 ± 1.31
4c	H	H	H	OH	H	>100	>100	>100	>100	>100	>100
4d	H	H	H	OCH ₃	H	55.7 ± 9.80	38.0 ± 11.7	51.4 ± 11.2	56.9 ± 17.6	51.9 ± 11.0	50.2 ± 8.10
4e	CF ₃	H	H	H	H	56.8 ± 11.4	57.6 ± 11.7	45.5 ± 13.1	53.3 ± 9.65	39.3 ± 8.58	40.4 ± 4.68
4f	H	H	H	CF ₃	H	25.8 ± 4.12	25.0 ± 8.17	24.2 ± 5.10	20.3 ± 7.35	18.6 ± 5.76	17.6 ± 3.45
4g	H	H	CF ₃	H	CF ₃	8.41 ± 1.01	8.19 ± 3.78	8.27 ± 2.74	8.12 ± 5.09	8.76 ± 1.36	8.70 ± 3.02

Table 3. continued

No.	R	R ₁	R ₂	R ₃	R ₄	GI ₅₀ (μM) ^d					
						NCI-H23	ACHN	MDA-MB-231	PC-3	NUGC-3	HCT 15
5a	methyl	OH	H	H	H	65.7 ± 8.84	50.7 ± 6.32	89.6 ± 9.29	70.7 ± 14.6	49.7 ± 10.9	54.5 ± 7.60
5b	ethyl	OH	H	H	H	>90	>90	>90	>90	>90	>90
5c	propyl	OH	H	H	H	43.1 ± 6.24	50.6 ± 11.2	38.0 ± 3.55	72.5 ± 9.36	35.4 ± 7.10	58.3 ± 9.89
5d	octyl	OH	H	H	H	55.8 ± 8.08	65.0 ± 6.13	54.3 ± 10.0	45.2 ± 7.03	31.1 ± 5.10	38.4 ± 6.27
5e	phenyl	OH	H	H	H	3.49 ± 0.999	4.26 ± 1.21	4.21 ± 0.892	5.69 ± 1.21	3.78 ± 0.618	3.83 ± 0.994
5f	2-chlorophenyl	OH	H	H	H	46.9 ± 2.99	44.2 ± 9.55	47.4 ± 5.04	60.0 ± 14.3	36.8 ± 6.41	44.9 ± 8.04
5g	3-chlorophenyl	OH	H	H	H	11.6 ± 2.75	17.2 ± 3.54	10.6 ± 1.60	23.3 ± 4.39	11.0 ± 2.34	23.1 ± 5.28
5h	4-chlorophenyl	OH	H	H	H	19.2 ± 3.77	19.6 ± 5.06	19.6 ± 3.54	25.1 ± 4.62	13.4 ± 2.20	20.2 ± 2.97
No.	R	R ₁	R ₂	R ₃	R ₄	NCI-H23	ACHN	MDA-MB-231	PC-3	NUGC-3	HCT 15
6a	methyl	CF ₃	H	H	H	>80	>80	>80	>80	>80	>80
6b	ethyl	CF ₃	H	H	H	>80	>80	>80	>80	>80	>80
6c	propyl	CF ₃	H	H	H	>70	>70	>70	>70	>70	>70
6d	octyl	CF ₃	H	H	H	4.22 ± 1.26	8.33 ± 3.09	3.35 ± 0.482	4.32 ± 1.04	6.18 ± 1.30	7.70 ± 1.11
6e	phenyl	CF ₃	H	H	H	6.34 ± 2.52	8.09 ± 1.68	5.29 ± 0.846	5.56 ± 0.990	8.12 ± 1.44	6.50 ± 1.14
6f	2-chlorophenyl	CF ₃	H	H	H	2.15 ± 0.680	3.90 ± 0.946	2.53 ± 0.625	5.14 ± 1.02	2.73 ± 0.554	4.22 ± 0.303
6g	3-chlorophenyl	CF ₃	H	H	H	0.292 ± 0.111	0.526 ± 0.178	0.288 ± 0.0992	0.729 ± 0.131	0.754 ± 0.129	0.797 ± 0.173
6h	4-chlorophenyl	CF ₃	H	H	H	0.307 ± 0.0941	0.824 ± 0.0708	0.533 ± 0.0824	0.786 ± 0.0996	0.551 ± 0.126	0.839 ± 0.0610
No.	R	R ₁	R ₂	R ₃	R ₄	NCI-H23	ACHN	MDA-MB-231	PC-3	NUGC-3	HCT 15
7a	methyl	H	CF ₃	H	CF ₃	>70	>70	>70	>70	>70	>70
7b	ethyl	H	CF ₃	H	CF ₃	7.82 ± 1.62	7.72 ± 2.40	7.28 ± 0.965	8.54 ± 1.99	7.22 ± 1.35	7.62 ± 0.902
7c	propyl	H	CF ₃	H	CF ₃	15.0 ± 2.67	13.9 ± 4.03	15.2 ± 2.44	18.6 ± 2.87	14.4 ± 3.49	21.9 ± 3.11
7d	octyl	H	CF ₃	H	CF ₃	3.23 ± 0.397	4.52 ± 1.13	7.53 ± 1.69	7.43 ± 1.89	2.23 ± 0.579	4.97 ± 1.09
7e	phenyl	H	CF ₃	H	CF ₃	2.13 ± 0.638	2.92 ± 0.640	2.43 ± 0.632	3.50 ± 0.630	1.68 ± 0.238	3.54 ± 1.20
7f	2-chlorophenyl	H	CF ₃	H	CF ₃	2.08 ± 0.482	4.70 ± 0.733	5.25 ± 1.08	8.04 ± 2.55	1.93 ± 0.404	4.05 ± 0.820
7g	3-chlorophenyl	H	CF ₃	H	CF ₃	0.420 ± 0.0607	0.813 ± 0.271	1.19 ± 0.313	1.05 ± 0.254	0.560 ± 0.189	0.625 ± 0.0674
7h	4-chlorophenyl	H	CF ₃	H	CF ₃	0.889 ± 0.102	2.57 ± 0.981	2.00 ± 0.683	4.23 ± 1.67	1.66 ± 0.406	2.22 ± 0.216

^d GI_{50} values are taken as a mean from three experiments and correspond to the agent's concentration causing a 50% decrease in net cell growth.

anticancer activity. Work on the enhancement of potency and pharmacological profiles of these probe molecules are underway.

■ ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acsmedchemlett.6b00004.

Synthetic procedures, characterization of final products, biological assay protocols, and data and pharmacology profiles (PDF)

■ AUTHOR INFORMATION

Corresponding Author

*Tel: +82 43 261 2811. Fax: +82 43 268 2732. E-mail: medchem@chungbuk.ac.kr.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

This research was supported by Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education, Science and Technology (NRF-2013R1A1A2009381), and Medical Research Center Program (2008-0062275).

■ ABBREVIATIONS

LPS, lipopolysaccharide; NF- κ B, nuclear factor kappa-light-chain-enhancer of activated B cells

■ REFERENCES

- (1) Sen, R.; Baltimore, D. Multiple nuclear factors interact with the immunoglobulin enhancer sequences. *Cell* **1986**, *46*, 705–716.
- (2) Hayden, M. S.; Ghosh, S. NF- κ B, the first quarter-century: Remarkable progress and outstanding questions. *Genes Dev.* **2012**, *26*, 203–234.
- (3) Siebenlist, U.; Franzoso, G.; Brown, K. Structure, regulation and function of NF- κ B. *Annu. Rev. Cell Biol.* **1994**, *10*, 405–455.
- (4) Kumar, A.; Takada, Y.; Boriek, A. M.; Aggarwal, B. B. Nuclear factor- κ B: Its role in health and disease. *J. Mol. Med.* **2004**, *82*, 434–448.
- (5) Arepalli, S. K.; Choi, M.; Jung, J. K.; Lee, H. Novel NF- κ B inhibitors: A patent review (2011–2014). *Expert Opin. Ther. Pat.* **2015**, *25*, 319–334.
- (6) Kwak, J. H.; Jung, J. K.; Lee, H. Nuclear factor-kappa B inhibitors: A patent review (2006 - 2010). *Expert Opin. Ther. Pat.* **2011**, *21*, 1897–1910.
- (7) Srinivasan, B.; Johnson, T. E.; Lad, R.; Xing, C. Structure - Activity relationship studies of chalcone leading to 3-hydroxy-4,3',4',5'-tetramethoxychalcone and its analogues as potent nuclear factor κ B inhibitors and their anticancer activities. *J. Med. Chem.* **2009**, *52*, 7228–7235.
- (8) Chen, H.; Yang, Z.; Ding, C.; Chu, L.; Zhang, Y.; Terry, K.; Liu, H.; Shen, Q.; Zhou, J. Discovery of O-alkylamino-tethered niclosamide derivatives as potent and orally bioavailable anticancer agents. *ACS Med. Chem. Lett.* **2013**, *4*, 180–185.
- (9) Choi, M.; Hwang, Y. S.; Kumar, A. S.; Jo, H.; Jeong, Y.; Oh, Y.; Lee, J.; Yun, J.; Kim, Y.; Han, S. B.; Jung, J. K.; Cho, J.; Lee, H. Design and synthesis of 3,4-dihydro-2H-benzo[h]chromene derivatives as potential NF- κ B inhibitors. *Bioorg. Med. Chem. Lett.* **2014**, *24*, 2404–2407.
- (10) Choi, M.; Jo, H.; Park, H. J.; Sateesh Kumar, A.; Lee, J.; Yun, J.; Kim, Y.; Han, S. B.; Jung, J. K.; Cho, J.; Lee, K.; Kwak, J. H.; Lee, H.

Design, synthesis, and biological evaluation of benzofuran- and 2,3-dihydrobenzofuran-2-carboxylic acid N-(substituted)phenylamide derivatives as anticancer agents and inhibitors of NF- κ B. *Bioorg. Med. Chem. Lett.* **2015**, *25*, 2545–2549.

(11) Kwak, J. H.; Kim, Y.; Park, H.; Jang, J. Y.; Lee, K. K.; Yi, W.; Kwak, J. A.; Park, S. G.; Kim, H.; Lee, K.; Kang, J. S.; Han, S. B.; Hwang, B. Y.; Hong, J. T.; Jung, J. K.; Kim, Y.; Cho, J.; Lee, H. Structure-activity relationship of indoline-2-carboxylic acid N-(substituted)phenylamide derivatives. *Bioorg. Med. Chem. Lett.* **2010**, *20*, 4620–4623.

(12) Sridharan, V.; Suryavanshi, P. A.; Menéndez, J. C. Advances in the chemistry of tetrahydroquinolines. *Chem. Rev.* **2011**, *111*, 7157–7259.

(13) Katritzky, A. R.; Rachwal, S.; Rachwal, B. Recent progress in the synthesis of 1,2,3,4-tetrahydroquinolines. *Tetrahedron* **1996**, *52*, 15031.

(14) Asolkar, K. N.; Schröder, D.; Heckman, R.; Lang, S.; Wagner-Döbler, I.; Laatsch, H. Helquinoline, a new tetrahydroquinoline antibiotic from *Janibacter limosus* Hel 1. *J. Antibiot.* **2004**, *57*, 17–23.

(15) Pitta, M. G. R.; Pitta, M. G. R.; Rêgo, M. J. B. M.; Galdino, S. L. The evolution of drugs on schistosoma treatment: Looking to the past to improve the future. *Mini-Rev. Med. Chem.* **2013**, *13*, 493–508.

(16) Leeson, P. D.; Carling, R. W.; Moore, K. W.; Moseley, A. M.; Smith, J. D.; Stevenson, G.; Chan, T.; Baker, R.; Foster, A. C.; Grimwood, S.; Kemp, J. A.; Marshall, G. R.; Hoogsteen, K. 4-Amido-2-carboxytetrahydroquinolines. Structure-activity relationships for antagonism at the glycine site of the NMDA receptor. *J. Med. Chem.* **1992**, *35*, 1954–1968.

(17) Khan, P. M.; Correa, R. G.; Divlianska, D. B.; Peddibhotla, S.; Sessions, E. H.; Magnuson, G.; Brown, B.; Suyama, E.; Yuan, H.; Mangravit-Novo, A.; Vicchiarelli, M.; Su, Y.; Vasile, S.; Smith, L. H.; Diaz, P. W.; Reed, J. C.; Roth, G. P. 2011. Identification of inhibitors of NOD1-induced nuclear factor- κ B activation. *ACS Med. Chem. Lett.* **2011**, *2*, 780–785.

(18) Bedoya, L. M.; Abad, M. J.; Calonge, E.; Saavedra, L. A.; Gutierrez, C. M.; Kouznetsov, V. V.; Alami, J.; Bermejo, P. 2010. Quinoline-based compounds as modulators of HIV transcription through NF- κ B and Sp1 inhibition. *Antiviral Res.* **2010**, *87*, 338–344.

(19) Moon, K. Y.; Hahn, B. S.; Lee, J.; Kim, Y. S. A cell-based assay system for monitoring NF- κ B activity in human HaCaT transfectant cells. *Anal. Biochem.* **2001**, *292*, 17–21.

(20) Greten, F. R.; Eckmann, L.; Greten, T. F.; Park, J. M.; Li, Z. W.; Egan, L. J.; Kagnoff, M. F.; Karin, M. IKK β links inflammation and tumorigenesis in a mouse model of colitis-associated cancer. *Cell* **2004**, *118*, 285–296.

(21) Nurmi, A.; et al. Pyrrolidine dithiocarbamate inhibits translocation of nuclear factor kappa-B in neurons and protects against brain ischaemia with a wide therapeutic time window. *J. Neurochem.* **2004**, *91*, 755–765.

(22) Hayakawa, M.; Miyashita, H.; Sakamoto, I.; Kitagawa, M.; Tanaka, H.; Yasuda, H.; Karin, M.; Kikugawa, K. Evidence that reactive oxygen species do not mediate NF- κ B activation. *EMBO J.* **2003**, *22*, 3356–3366.

(23) Liu, S. F.; Ye, X.; Malik, A. B. In Vivo Inhibition of Nuclear Factor- κ B Activation Prevents Inducible Nitric Oxide Synthase Expression and Systemic Hypotension in a Rat Model of Septic Shock. *J. Immunol.* **1997**, *159*, 3976–3983.

(24) Ziegler-Heitbrock, H. W. L.; Sternsdorf, T.; Liese, J.; Belohradsky, B.; Weber, C.; Wedel, A.; Schreck, R.; Bäuerle, P.; Ströbel, M. Pyrrolidine dithiocarbamate inhibits NF- κ B mobilization and TNF production in human monocytes. *J. Immunol.* **1993**, *151*, 6986–6993.

(25) Schreck, R.; Meier, B.; Männel, D. N.; Dröge, W.; Bäuerle, P. A. Dithiocarbamates as potent inhibitors of nuclear factor κ B activation in intact cells. *J. Exp. Med.* **1992**, *175*, 1181–1194.

(26) Kim, B. H.; Reddy, A. M.; Lee, K. H.; Chung, E. Y.; Cho, S. M.; Lee, H.; Min, K. R.; Kim, Y. Inhibitory mechanism of chroman compound on LPS-induced nitric oxide production and nuclear factor- κ B activation. *Biochem. Biophys. Res. Commun.* **2004**, *325*, 223–228.