CASE REPORT

Glossopharyngeal neuralgia associated with cardiac syncope

Laura Burfield, Faheem Ahmad, Jacqueline Adams

Queen Elizabeth University
Hospital, Glasgow, UK

Correspondence to

Dr Laura Burfield,
lauracburfield@gmail.com

Accepted 10 April 2016

SUMMARY

Glossopharyngeal neuralgia is a rare pain syndrome presenting with paroxysms of pain in the region of the glossopharyngeal nerve. Even more uncommon is the association between glossopharyngeal neuralgia and cardiac syncope. In these patients, the cardiovascular consequences may include bradycardia, hypotension and cardiac arrest. We describe the case of a 40-year-old patient who presented with this rare association of glossopharyngeal neuralgia and syncope. Multiple pauses including one lasting 14 s were noted on ambulatory ECG monitoring. In this case, the patient declined pharmacological treatment with carbamazepine or with permanent pacing and so far has been in remission from symptoms for 3 months.

BACKGROUND

Glossopharyngeal neuralgia (GN) is a rare facial pain syndrome characterised by paroxysms of severe pain in the distribution of the glossopharyngeal nerve. Even rarer is the association between GN and syncope, when it may then be termed vagoglossopharyngeal neuralgia.¹ In these patients, GN may be associated with episodes of bradycardia, severe hypotension and even cardiac arrest leading to syncope.² While GN is commonly misdiagnosed as the much more common trigeminal neuralgia, this is an important diagnosis not to miss, as the potential cardiac consequences may be life-threatening.

We present a case of a patient with GN associated with cardiac syncope, presenting with collapse and significant nocturnal pauses on ECG monitoring.

CASE PRESENTATION

A 40-year-old man initially presented to his general practitioner, with symptoms of throat pain associated with light-headedness and palpitations. The throat pain had been occurring intermittently over the preceding 3 months and was described as 'shooting, rippling or pulsing' deep in the right side of his throat, worse on swallowing. This pain never occurred on the left side. While on holiday 9 months previously, he had experienced several episodes of light-headedness associated with the same pain and described several brief episodes of collapse. He was admitted twice while on holiday and reported being told that he was dehydrated and his blood pressure was low. On his return, he was admitted to hospital elsewhere, with throat pain and associated collapse. The only abnormality found at that time was sinus bradycardia.

On discharge, he was advised to attend his general practitioner to be referred for 24 h ECG monitoring.

He had a medical history of generalised anxiety disorder and post viral fatigue syndrome. There was no family history of note.

His general and neurological examination, including cranial nerve examination, was normal.

INVESTIGATIONS

Chest radiograph, echocardiogram, full blood count, routine biochemistry and a short synacthen test were all normal.

Twenty-four hour ECG recording reported sinus rhythm with a pause of 14 s at 05:22 am along with multiple other shorter nocturnal pauses (figure 1). On questioning, it was revealed that the patient had been awake at this time and experiencing throat pain, but had not collapsed. Following this, he underwent a tilt table test, during which he experienced multiple paroxysms of throat pain that allowed a clear relationship between paroxysms of pain and episodes of bradycardia to be demonstrated (figures 2 and 3).

A diagnosis of GN with cardiac syncope was made and the patient was referred to the ear, nose and throat physicians to exclude a structural lesion. A CT scan of his neck did not reveal an abnormality.

DIFFERENTIAL DIAGNOSIS

The differential diagnosis included trigeminal neuralgia, superior laryngeal neuralgia (affecting the superior laryngeal branch of the Xth cranial nerve) and naevus intermedius neuralgia (affecting the somatic sensory branch of the VIIth cranial nerve).¹ GN occurs with a frequency of about 1% that of trigeminal neuralgia but has very similar pain characteristics as well as adjacent locations of pain and similar triggers.³ GN occurs more frequently on the left, while trigeminal neuralgia occurs more commonly on the right.⁴ Bilateral involvement is found more commonly in trigeminal neuralgia.¹

Superior laryngeal neuralgia is a paroxysmal pain syndrome characterised by paroxysms of pain usually localised to one side of the larynx. Identified trigger factors include swallowing, straining the voice, singing and turning the head.⁵ Application of topical anaesthesia to the larynx or blockade of the superior laryngeal nerve should stop the pain and can be used as a diagnostic test.¹ More rarely, GN may need to be distinguished from nervus intermedius neuralgia, which produces intermittent stabbing pain in the ear. Nervus

CrossMark

To cite: Burfield L,
Ahmad F, Adams J. *BMJ*
Case Rep Published online:
[please include Day Month
Year] doi:10.1136/bcr-2015-
214104

Figure 1 Twenty-four hour ECG recording showing a pause of 14 s during an episode of throat pain.

intermedius neuralgia is always unilateral and usually occurs spontaneously, although it may be triggered by stimulation of the ear canal, or swallowing or talking.⁶ The medical management is identical to that of GN.¹

OUTCOME AND FOLLOW-UP

In this case, the patient declined pharmacological treatment with carbamazepine or insertion of a permanent pacemaker. He remains in remission and has had no further throat pain and no episodes of collapse for the past 3 months.

DISCUSSION

Paroxysms of pain in the region of the glossopharyngeal nerve were first described in 1910 by Weisenberg⁷ in a 35-year-old patient with a right cerebellopontine angle tumour. The International Headache Society (IHS) has described GN as 'a severe transient stabbing pain experienced in the ear, base of tongue, tonsillar fossa or beneath the angle of the jaw'.⁸ The onset of GN is typically sudden. Episodes of pain last from several seconds to minutes, with intervals between the paroxysms lasting from minutes to hours. The intervals between clusters are irregular but can range from days to years with patients being asymptomatic between episodes.¹ Patients often have difficulty in identifying triggers of pain, most likely as it involves the deep structures of the mouth, pharynx and ear. Most frequently, swallowing triggers the paroxysms of pain, although chewing, yawning and speaking have also been implicated.⁹

Most cases of GN are idiopathic—most likely these are caused by compression of the nerve by the posteroinferior cerebellar artery as the nerve enters or exits the brainstem.¹⁰ Secondary GN can occur due to compression of the glossopharyngeal nerve by structural lesions including cerebellopontine angle tumours, laryngeal and nasopharyngeal tumours, cranial base tumours, oropharynx and tongue tumours, and by intracranial vascular compression parapharyngeal abscesses, trauma, dental extractions, Paget's disease, direct carotid puncture and an elongated styloid process (Eagle's syndrome), among other causes.^{1 7 9 11–13}

As there is no specific diagnostic test for GN, the diagnosis is made on clinical grounds. It has been recommended that all patients with GN be referred to an ear, nose and throat physician to investigate a potential structural cause.¹ Imaging of the brain stem to identify tumours, or possible vascular compression, is recommended. Furthermore, a high resolution CT scan of the neck can identify an elongated styloid process.¹

In 1942, Riley *et al*¹⁴ first reported the association of GN with brief episodes of bradycardia, hypotension and asystole causing cardiac syncope. The association of GN with cardiac syncope is rare. In a Mayo clinic review of 217 patients with GN, in 1981, only four were identified as having cardiac syncope.¹¹

The mechanism by which GN leads to cardiac syncope is not yet fully elucidated. Usually, there appears to be a latency period between the symptoms of neuralgia and the first syncopal

Figure 2 Twelve-lead ECG recorded during a paroxysm of throat pain.

Figure 3 Twelve-lead ECG recorded immediately after paroxysm of throat pain resolved.

event.¹⁵ Very rarely, GN has presented with syncope and without the associated pain syndrome.¹⁶ Most proposed mechanisms are based on the close anatomical relationship of the glossopharyngeal and vagus nerves in the medulla oblongata and the possibility of the formation of a vagoglossopharyngeal reflex arc.^{17 18} It is proposed that the afferent nerve impulses travelling within the glossopharyngeal nerve may reach the nucleus of the tractus solitarius in the midbrain and, through collaterals, reach the dorsal nucleus of the vagus nerve. Additionally, the fibres of the carotid sinus nerve, also known as Hering's nerve, join the glossopharyngeal nerve and terminate in the dorsal nucleus of the vagus nerve, providing a route by which glossopharyngeal afferent impulses may produce vagal nerve stimulation.^{17 18} Finally, Gardner *et al*¹⁹ proposed that there may be anomalous communication between the nucleus of the tractus solitarius and nucleus ambiguus mediated by the development of synapses on the fibres of cranial nerves IX and X.

Therapeutic options described for GN may be pharmacological or surgical. For idiopathic GN, the first-line treatment is usually pharmacological and similar to the management of trigeminal neuralgia. Anticonvulsants including carbamazepine, gabapentin, lamotrigine, phenytoin, oxcarbazepine and pregabalin, have all been used in the treatment of GN with the most evidence being available for carbamazepine.^{9 20 21} Indeed, treatment with carbamazepine can lead to resolution of pain from neuralgia as well as resolution of cardiac symptoms.¹⁵

In patients who are unresponsive to pharmacological interventions, surgical options may be considered. The surgical procedure with the highest rate of success for patient with idiopathic GN is microvascular decompression. In a recent review, Rey-Dios and Cohen-Gadol²² reported the outcomes of 454 patients from 14 different series undergoing microvascular decompression of cranial nerves IX and X. The mean follow-up duration was 4.9 years. The rate of resolution of pain for patients undergoing microvascular decompression was 84.7% with recurrence in 7% of patients. Transient cranial nerve X dysfunction defined as dysphagia, hoarseness or both, occurred in 13.2% of patients treated with microvascular decompression and a permanent cranial nerve deficit occurred following 5.5% procedures. In contrast, the long-term rate of pain control for 157 patients who underwent rhizotomy of the upper cranial nerve rootlets of cranial nerve X was higher at 87.3%, however,

the rate of transient and permanent cranial nerve X deficits was increased at 25% and 19.1%, respectively. Overall, rhizotomy appears to have higher rates of pain control at the expense of high rates of postoperative cranial nerve X dysfunction.²²

Gamma Knife radiosurgery has also been reported to be effective in the treatment of GN, particularly for an elderly patient group who may not be fit for conventional surgery and may benefit from a minimally invasive approach. O'Connor and Bidiwala²³ presented a case series in which 13 of the 15 (87%) patients treated with radiosurgery achieved significant pain relief without any reported adverse effects.

Pacemakers can prevent bradycardia during paroxysms of pain. In the treatment of GN associated with syncopal episodes, pacing has been used with varying degrees of success. Johnston and Redding¹⁵ reported a case of a 53-year-old patient in whom pain and syncope completely resolved following the insertion of a permanent pacemaker and treatment with carbamazepine. Furthermore, in the case presented by Korkes *et al*²⁴ where the patient had syncope associated with swallowing, since the insertion of a permanent pacemaker, the patient has been free of syncopal episodes (35 months of follow-up). Similarly, Hie and Ruitei²⁵ described a case of a 60-year-old patient treated with a permanent dual-chamber pacemaker, who had complete resolution of syncopal episodes, although she did continue to experience episodic neuralgia. In contrast, the patient presented in Barbash *et al*² failed to respond to the insertion of a temporary intravenous pacemaker. Likewise, the patient described by Roa and Krupin²⁶ continued to be hypotensive despite having a capturing transvenous pacemaker. It is important to note that, while a pacemaker may prevent syncopal episodes, it will have no therapeutic effect on the pain syndrome, and this is hence why some authors feel that, where possible, a surgical option should be pursued where pharmacological therapy has proved ineffective.

In summary, GN is an unusual pain syndrome, important not only as it has the potential to cause extreme distress among sufferers, but also as it has the potential to cause significant cardiovascular compromise including severe hypotension, syncope and cardiac arrest. While trigeminal neuralgia is a far more common pain syndrome, we urge clinicians to bear in mind the potential diagnosis of GN in those presenting with paroxysmal pain in the region of the tongue, pharynx or ear and to be aware of the potential relationship with syncope.

Learning points

- ▶ Glossopharyngeal neuralgia is a rare pain syndrome commonly misdiagnosed as trigeminal neuralgia.
- ▶ Consider glossopharyngeal neuralgia in patients presenting with paroxysmal throat pain and be aware of the potential cardiovascular consequences: bradycardia, severe hypotension and even cardiac arrest.
- ▶ Therapeutic options include pharmacological modalities to target the pain syndrome and device therapy to alleviate bradycardia.

Competing interests None declared.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Blumenfeld A, Nikolskaya G. Glossopharyngeal neuralgia. *Curr Pain Headache Rep* 2013;17:343.
- 2 Barbash GI, Keren G, Korczyn AD, et al. Mechanisms of syncope in glossopharyngeal neuralgia. *Electroencephalogr Clin Neurophysiol* 1986;63:231–5.
- 3 Kazemi B, Akbarzadeh F. Syncopal storm caused by glossopharyngeal neuralgia. *Am J Emerg Med* 2012;30:2101.e5–7.
- 4 Katusic S, Williams DB, Beard CM, et al. Incidence and clinical features of glossopharyngeal neuralgia, Rochester, Minnesota, 1945–1984. *Neuroepidemiology* 1991;10:266–75.
- 5 Baugh RF, Baugh A, Bunge F. Superior laryngeal nerve syndrome and the evaluation of anterior neck pain. *Am J Otolaryngol* 2012;33:481–3.
- 6 Tubbs RS, Steck DT, Mortazavi MM, et al. The nervus intermedius: a review of its anatomy, function, pathology and role in neurosurgery. *World Neurosurg* 2013;79:763–7.
- 7 Weisenberg TH. Cerebello-pontine tumor diagnosed for six years as tic douloureux: the symptoms of irritation of the 9th and 12th cranial nerves. *JAMA* 1910;54:1600–4.
- 8 Headache Classification Subcommittee of the International Headache Society. The international classification of headache disorders. *Cephalgia* 2004;24:1–160.
- 9 Elias J, Kuniyoshi R, Carloni WWH, et al. Glossopharyngeal neuralgia associated with cardiac syncope. *Arq Bras Cardiol* 2002;78:510–19.
- 10 Alafaci C, Granata F, Cutugno M, et al. Glossopharyngeal neuralgia caused by a complex neurovascular conflict: case report and review of the literature. *Surg Neurol Int* 2015;6:19.
- 11 Rushton JG, Stevens JC, Miller RH. Glossopharyngeal (vagoglossopharyngeal) neuralgia: a study of 217 cases. *Arch Neurol* 1981;38:201–5.
- 12 Koratkar H, Parashar V, Koratkar S. A review of neuropathic pain conditions affecting teeth. *Gen Dent* 2010;58:436–41.
- 13 Soh KB. The glossopharyngeal nerve, glossopharyngeal neuralgia and the Eagle's syndrome—current concepts and management. *Singapore Med J* 1999;40:659–65.
- 14 Riley HA, Greenan WJ, Wortis H, et al. Glossopharyngeal neuralgia initiating or associated with cardiac arrest. *Trans Am Neurol Assoc* 1942;68:28–9.
- 15 Johnston RT, Redding VJ. Glossopharyngeal neuralgia associated with cardiac syncope: long term treatment with permanent pacing and carbamazepine. *Br Heart J* 1990;64:403–5.
- 16 Reddy K, Hobson DE, Gomori A, et al. Painless glossopharyngeal “neuralgia” with syncope: a case report and literature review. *Neurosurgery* 1987;21:916–19.
- 17 Ferrante L, Artico M, Nardacci B, et al. Glossopharyngeal neuralgia with cardiac syncope. *Neurosurgery* 1995;36:58–63.
- 18 Aguiar PH, Tella OJr, Pereira CU, et al. Chiari type I presenting as left glossopharyngeal neuralgia with cardiac syncope. *Neurosurg rev* 2002;25:99–102.
- 19 Gardner WJ. Concerning the mechanism of trigeminal neuralgia and hemifacial spasm. *J Neurosurg* 1962;19:947–58.
- 20 De Simone R, Ranieri A, Bilo L, et al. Cranial neuralgias: from physiopathology to pharmacological treatment. *Neurol Sci* 2008;29:S69–78.
- 21 Titlic M, Jukic I, Tonkic A, et al. Use of lamotrigine in glossopharyngeal neuralgia: a case report. *Headache* 2006;46:167–9.
- 22 Rey-Dios R, Cohen-Gadol AA. Current neurosurgical management of glossopharyngeal neuralgia and technical nuances for microvascular decompression surgery. *Neurosurg focus* 2013;34:E8.
- 23 O'Connor JK, Bidiwala S. Effectiveness and safety of Gamma Knife radiosurgery for glossopharyngeal neuralgia. *Proc (Bayl Univ Med Cent)* 2013;26:262–4.
- 24 Korkes H, de Oliveira EM, Brollo L, et al. Cardiac syncope induced by glossopharyngeal “neuralgia”: a rare presentation. *Arq Bras Cardiol* 2006;87:e189–91.
- 25 Hie CH, Ruiters AJH. Glossopharyngeal neuralgia with syncope: a case report. *Neth Heart J* 2002;10:150–3.
- 26 Roa NC, Krupin BR. Glossopharyngeal neuralgia with syncope-anaesthetic considerations. *Anesthesiology* 1981;54:426–8.

Copyright 2016 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow