Published in final edited form as:

Angew Chem Int Ed Engl. 2013 June 3; 52(23): 6068–6071. doi:10.1002/anie.201301264.

Direct Synthesis of β -*N*-Glycosides by the Reductive Glycosylation of Azides with Protected and Native Carbohydrate Donors**

Dr. Jianbin Zheng,

Department of Chemistry, Yale University, 225 Prospect Street, New Haven, CT 06520-8107 (USA). Shanghai Key Laboratory of New Drug Design, School of Pharmacy, East China University of Science and Technology (P.R. China)

Dr. Kaveri Balan Urkalan, and

Department of Chemistry, Yale University, 225 Prospect Street, New Haven, CT 06520-8107 (USA)

Dr. Seth B. Herzon

Department of Chemistry, Yale University, 225 Prospect Street, New Haven, CT 06520-8107 (USA). Homepage : <http://www.chem.yale.edu/faculty/herzon.html>

Seth B. Herzon: seth.herzon@yale.edu

Keywords

aspartylation; glycosides; glycosylation; Wittig reactions

The β -*N*-glycoside linkage is embedded within structurally diverse natural products such as the anthraquinone antibiotics (e.g. mycorrhodin),^[1] indigo glycosides (e.g. akashin C),^[2] and the ansamycin antibiotics (e.g. ansacarbamitocin A; Scheme 1).^[3, 4] This linkage is also found in a large number of glycopeptides, which exhibit a broad spectrum of biological functions;^[5] erythropoietin (EPO) is a well-known example.

Reported methods for the synthesis of *N*-linked glycoside bonds include the functionalization of glycosyl azides^[6] and the condensation of ammonia,^[7,8] *N,O*-dialkylhydroxylamines,^[9] and acyl hydrazides^[10] with reducing sugars. We describe herein a simple and complementary method that proceeds by mild thermolysis of alkyl and aryl azides in the presence of reducing sugars and a tertiary phosphine; it is essentially an aza-Wittig reaction in which a carbohydrate is used as a latent carbonyl group [Eq. (1)]. Despite the simplicity of this approach, to our knowledge, a single report describing the

** Financial support from the National Institute of General Medical Sciences (R01GM090000), the Searle Scholars Program, Shanghai Committee of Science and Technology (11DZ2260600, Shanghai Municipal Education Commission Fellowship to J.Z.), and Yale University is gratefully acknowledged. We thank Dr. Seann Mulcahy for preliminary experiments and Dr. Le Li for helpful discussions.

Correspondence to: Seth B. Herzon, seth.herzon@yale.edu.

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.201301264>.

condensation of a polyfluorinated iminophosphorane and lactose^[11] stands as the only direct precedent for this reaction.

(1)

The condensation of ammonia and *N,O*-dialkyl hydroxyl-amines with unactivated carbohydrates is well-developed, and this method has found great utility in the synthesis of glycoproteins^[12] and natural product glycoconjugate libraries.^[9a–c] However, the reaction of ammonia with a reducing sugar requires long reaction times and a large excess of ammonia to drive the process to completion, thus rendering this process unsuitable for the synthesis of *N*-glycosides incorporating precious amine fragments. The condensation of *N,O*-dialkyl hydroxylamines with carbohydrates is more efficient, but necessarily provides an unnatural (neoglycoside) linkage. The method we report herein is characterized by short reaction times, a small (0.5 equiv) excess of reagent, and high selectivity for the β -*N*-glycoside product.

The reaction between an azide, phosphine, and carbohydrate to form an *N*-glycoside and phosphine oxide may proceed by several pathways. Regardless, the overall transformation is formally a Staudinger reduction–aza-Wittig sequence and we therefore began by evaluating the ability of various phosphines to promote the coupling of benzyl azide (**1a**) and *O*-allyl-*N,N*-dimethyl-D-pyrrolosamine (**2a**, Table 1).^[13] Products derived from the protected 2,6-dideoxy-glycoside **2a** can be readily purified by flash-column chromatography, and exhibit well-resolved resonances in their ¹H NMR spectra, which facilitated the characterization of the products and optimization of the reaction conditions. Heating a mixture of **1a** (1.5 equiv) and **2a** in the presence of triphenylphosphine (1.5 equiv) in tetrahydrofuran as the solvent resulted in the formation of the *N*-glycoside **3a** in 24% yield and with a > 15:1 selectivity for the β anomer (¹H NMR spectroscopic analysis; $J_{\text{H1-H2ax}} = 10.5$ Hz). The reaction did not proceed at lower temperatures (24 °C). When the more reactive dimethylphenylphosphine was employed, the product **3a** was isolated in 56 % yield (entry 2). Reasoning that protic acids might promote the pyranose–hydroxylaldehyde isomerization, a variety of acidic additives were evaluated. The addition of *para*-toluenesulfonic acid (5 mol %) increased the yield of the *N*-glycoside **3a** to 90 % (entry 3). Trimethylphosphine was also effective as the reductant (88 % yield, entry 4). The product was obtained in lower yield when *N,N*-dimethylformamide was employed as the solvent (43 %, entry 5).

This reductive *N*-glycosylation reaction has proven to be relatively general in nature (Table 2). As shown in entries 1–4, primary alkyl azides coupled with **2a** in high yields (72–90 %). Phenyl azide (**1e**) also formed the corresponding *N*-phenyl-*N*-glycoside **3e** in 83 % yield (entry 5), thus establishing that this method is not limited to alkyl azide substrates. Ethyl azidoacetate (**1f**) and ω -azidolysine methyl ester (**1g**) both coupled with **2a** in high yields (77 and 78%, entries 6 and 7, respectively). Benzyl azide (**1a**) also coupled efficiently with *O*-allyl-L-oleandrose (**2b**, 88%; entry 8). 2,3,5-Tribenzyl-D-ribose (**2c**) underwent

conjugation with *para*-methoxybenzyl azide (**1b**) in 73% yield (entry 9). *N*-Acetyl- D - glucosamine (**2d**) and the fully deprotected carbohydrates D-glucose (**2 e**) and maltose (**2 f**) were shown to couple with benzyl and phenyl azide in yields of 72–82% (entries 10–13). The product **3i**, derived from ribose (entry 9), was formed with a β : α selectivity of 3:2; in all other cases, the β anomer was obtained with > 15:1 selectivity (¹H NMR spectroscopic analysis).

The results in Table 2 demonstrate that β -linked *N*-glycosides are readily formed from simple azides and reducing sugars. It was of interest to determine if the method is also amenable to the synthesis of *N*-linked glycopeptides and, in particular, to the formation of the β -Asn-*N*-acetylglucosamine linkage, which is abundant in natural glycoproteins.^[5] Toward this end, we envisioned a sequence comprising the *N*-glycosylation of an ammonia equivalent, followed by aspartylation.^[12, 14] After much experimentation, we found that heating trimethylsilylmethyl azide (**1h**)^[15] and *N*-acetyl-D-glucosamine (**2d**) with dimethylphenylphosphine resulted in the formation of the β -linked *N*-glycoside **3 n** in 75 % yield as a 30:1 mixture of β and α anomers (Scheme 2). Benzoylation of **3n** (benzoyl chloride) afforded the benzamide **4a** in 85% yield. Finally, the trimethylsilylmethyl substituent was cleaved by treatment with ceric ammonium nitrate in acetonitrile (90%).^[16] The benzamide **5a** was obtained without detectable erosion of the anomeric site (¹H NMR spectroscopic analysis). This sequence provides a complementary and highly practical alternative to the reaction of carbohydrates with ammonia, as primary *N*-glycosides are notoriously difficult to purify and are prone to epimerization at the anomeric position.

This approach was readily adapted toward amino acid coupling reactions by employing the corresponding amino acid mixed anhydride in the acylation step. A selection of glycosylated amino acids and peptides prepared in this way are shown in Table 3. As a consequence of their high polarities, the intermediates in these sequences were employed without purification, although they may be isolated by reverse-phase chromatography, if desired. This approach led to the anomerically pure glucosamine derivatives **5b** and **5c** in high overall yields (76 % and 75% for **5b** and **5c**, respectively). The Asn-linked tripeptide was also obtained in excellent yield and with complete β selectivity (**5d**, 70 %). Finally, we discovered that the method is amenable to the direct introduction of disaccharides (**5e**, 61 %), which suggests its application in more complex settings.

Although glycosyl azides have been widely used for the synthesis of *N*-glycosides,^[6] the reductive condensation of azides with unactivated sugars does not appear to have been developed. We have disclosed herein a simple and straightforward method for the stereocontrolled synthesis of β -linked *N*-glycosides by employing alkyl and aryl azides as the sources of nitrogen atoms. The method is compatible with protected and deprotected carbohydrate donors, and does not require activation of the anomeric position. A simple, high-yielding procedure for the synthesis of β -Asn-linked glycosides has been developed, and this method may be of great utility in the synthesis of complex glycopeptides. Efforts to fully delineate the scope of the azide and carbohydrate components, as well as applications in natural products synthesis, are underway.

Experimental Section

Representative experimental procedure (Table 2, entry 1): Benzyl azide (**1a**; 67.0 mg, 500 μ mol, 1 equiv) and THF (300 μ L) were added in sequence to an oven-dried 4 mL vial equipped with a teflon-lined cap and magnetic stir bar. Dimethylphenylphosphine (70.0 mg, 500 μ mol, 1.00 equiv) was added dropwise by syringe over 5 min. The reaction vessel was sealed under an atmosphere of argon and the mixture stirred for 30 min at 24°C. A solution of **2a** (161 mg, 750 μ mol, 1.50 equiv) in THF (100 μ L) and *para*-toluenesulfonic acid (5.0 mg, 25.0 μ mol, 0.05 equiv) were then added in sequence. The vial was sealed under argon and then placed in an oil bath that had been preheated to 55°C. The reaction mixture was stirred and heated for 12 h at 55°C before cooling the product mixture to 23°C. The cooled product mixture was concentrated to dryness and the residue obtained was purified by flash-column chromatography (deactivated with 1% triethylamine/5 % acetone in hexanes; eluting with 5% acetone in hexanes) to provide the *N*-glycoside **3a** as a clear, colorless oil (two runs: 139 mg and 133 mg, average yield = 90%).

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

References

1. a) Takeda U, Okada T, Takagi M, Gomi S, Itoh J, Sezaki M, Ito M, Miyadoh S, Shomura T. *J Antibiot.* 1988; 41:417. [PubMed: 3372348] b) Takeda U, Okada T, Takagi M, Gomi S, Itoh J, Sezaki M, Ito M, Miyadoh S, Shomura T. *J Antibiot.* 1988; 41:425. [PubMed: 3372349]
2. a) Maskey RP, Grün-Wollny I, Fiebig HH, Laatsch H. *Angew Chem.* 2002; 114:623. *Angew. Chem. Int. Ed.* 2002, 41, 597; b) Maskey RP, Grün-Wollny I, Laatsch H. *Nat Prod Res.* 2005; 19:137. [PubMed: 15715257]
3. Snipes CE, Duebelbeis DO, Olson M, Hahn DR, Dent WH, Gilbert JR, Werk TL, Davis GE, Lee-Lu R, Graupner PR. *J Nat Prod.* 2007; 70:1578. [PubMed: 17892263]
4. For additional examples, see Omura S, Iwai Y, Hirano A, Nakagawa A, Awaya J, Tsuchiya H, Takahashi Y, Masuma R. *J Antibiot.* 1977; 30:275. [PubMed: 863788] Nettleton DE, Doyle TW, Krishnan B, Matsumoto GK, Clardy J. *Tetrahedron Lett.* 1985; 26:4011. Bonjouklian R, Smitka TA, Doolin LE, Molloy RM, Debono M, Shaffer SA, Moore RE, Stewart JB, Patterson GML. *Tetrahedron.* 1991; 47:7739. Matsunaga S, Fusetani N, Kato Y, Hirota H. *J Am Chem Soc.* 1991; 113:9690.
5. For selected reviews, see Spiro RG. *Glycobiology.* 2002; 12:43R. Gamblin DP, Scanlan EM, Davis BG. *Chem Rev.* 2009; 109:131. [PubMed: 19093879] Kan C, Danishefsky SJ. *Tetrahedron.* 2009; 65:9047. [PubMed: 20161377] Larkin A, Imperiali B. *Biochemistry.* 2011; 50:4411. [PubMed: 21506607]
6. For selected examples, see Mizuno M, Muramoto I, Kobayashi K, Yaginuma H, Inazu T. *Synthesis.* 1999:162. He Y, Hinklin RJ, Chang J, Kiessling LL. *Org Lett.* 2004; 6:4479. [PubMed: 15548055] Doores KJ, Mimura Y, Dwek RA, Rudd PM, Elliott T, Davis BG. *Chem Commun.* 2006:1401. for a recent review, see Györgydeák Z, Thiem J. *Adv Carbohydr Chem Biochem.* 2006; 60:103. [PubMed: 16750443]

7. a) Likhoshesterov LM, Novikova OS, Derevitskaja VA, Kochetkov NK. Carbohydr Res. 1986; 146:C1. b) Urge L, Kollat E, Hollosi M, Laczko I, Wroblewski K, Thuring J, Otvos L Jr. Tetrahedron Lett. 1991; 32:3445.
8.
The condensation of allyl- and benzylamine (10 equiv, 4 days reaction time) with unactivated carbohydrates has been reported: Cipolla L, Lay L, Nicotra F, Pangrazio C, Panza L. Tetrahedron. 1995; 51:4679.
9. a) Langenhan JM, Peters NR, Guzei IA, Hoffmann FM, Thorson JS. Proc Natl Acad Sci USA. 2005; 102:12305. [PubMed: 16105948] b) Ahmed A, Peters NR, Fitzgerald MK, Watson JA Jr, Hoffmann FM, Thorson JS. J Am Chem Soc. 2006; 128:14224. [PubMed: 17076473] c) Peltier-Pain P, Timmons SC, Grandemange A, Benoit E, Thorson JS. ChemMedChem. 2011; 6:1347. [PubMed: 21714096] for reviews, see d) Griffith BR, Langenhan JM, Thorson JS. Curr Opin Biotechnol. 2005; 16:622. [PubMed: 16226456] e) Langenhan JM, Griffith BR, Thorson JS. J Nat Prod. 2005; 68:1696. [PubMed: 16309329]
10. Peluso, Sp; Imperiali, B. Tetrahedron Lett. 2001; 42:2085.
11. El Ghoul M, Escoula B, Rico I, Lattes A. J Fluorine Chem. 1992; 59:107.
12.
For selected examples, see Wang P, Zhu J, Yuan Y, Danishefsky SJ. J Am Chem Soc. 2009; 131:16669. [PubMed: 19886622] Nagorny P, Sane N, Fasching B, Aussedat B, Danishefsky SJ. Angew Chem. 2012; 124:999. Wang P, Li X, Zhu J, Chen J, Yuan Y, Wu X, Danishefsky SJ. J Am Chem Soc. 2011; 133:1597. [PubMed: 21207981]
13. Gholap SL, Woo CM, Ravikumar PC, Herzon SB. Org Lett. 2009; 11:4322. [PubMed: 19719089]
14. Anisfeld ST, Lansbury PT. J Org Chem. 1990; 55:5560.
15. Nishiyama K, Tanaka N. J Chem Soc Chem Commun. 1983:1322.
16. Palomo C, Aizpurua JM, Legido M, Mielgo A, Galarza R. Chem Eur J. 1997; 3:1432.

Scheme 1.

Representative natural products that contain the β -N-glycoside linkage.

Scheme 2.

Synthesis of the *N*-benzoyl-*N*-glycoside **5a**. TMS = trimethylsilyl, Bz = benzoyl, THF = tetrahydrofuran, CAN = ceric ammonium nitrate.

Table 1Optimization of the reductive glycosylation.^[a]

Entry	Phosphine	Additive	Solvent	Yield [%] (β/α) ^[b]
1	PPh_3	–	THF	24 (> 15 : 1)
2	$\text{P}(\text{CH}_3)_2\text{Ph}$	–	THF	56 (> 15 : 1)
3	$\text{P}(\text{CH}_3)_2\text{Ph}$	PTSA (5 mol %)	THF	90 (> 15 : 1)
4	$\text{P}(\text{CH}_3)_3$	PTSA (5 mol %)	THF	88 (> 15 : 1)
5	$\text{P}(\text{CH}_3)_2\text{Ph}$	PTSA (5 mol %)	DMF	43 (> 15 : 1)

^[a] Conditions : Benzyl azide (**1a**, 1.50 equiv), **2a** (1 equiv), phosphine (1.50 equiv), 55°C , 12 h.^[b] Yield of isolated product after purification by flash-column chromatography.THF = tetrahydrofuran, PTSA = *p*-toluenesulfonic acid, DMF = *N,N*-dimethylformamide, PMB = *para*-methoxybenzyl, CBz = benzyloxycarbonyl.

Table 2Scope of the reductive glycosylation reaction.^[a]

Entry	Azide	Carbohydrate donor	Yield [%] ^[b]
1	BnN ₃ 1 a		90 3 a
2	PMBN ₃ 1 b		72 3 b
3	<i>p</i> -(CH ₃)C ₆ H ₄ N ₃ 1 c		78 3 c
4	 1 d		84 3 d
5	PhN ₃ 1 e		83 3 e
6	 1 f	2 a	77 3 f
7	 1 g		78 3 g
8	BnN ₃ 1 a		88 3 h
9	PMBN ₃ 1 b		73 3 i 3 : 2 β : α
10 ^[c]	BnN ₃ 1 a		76 3 j
11 ^[c]	BnN ₃ 1 a		82 3 k
12 ^[c]	PhN ₃ 1 e	2 e	78 3 l

Entry	Azide	Carbohydrate donor	Yield [%] ^[b]
13 ^[c]	BnN ₃ 1 a	2 f	72 3 m

^[a] Conditions : Azide (1 equiv), PhP(CH₃)₂ (1.00 equiv), carbohydrate (1.50 equiv), PTSA (5 mol %), THF, 55 °C, 12 h. All reactions were conducted on a 0.5 mmol scale.

^[b] Yield of isolated product after purification by flash-column chromatography. β : α selectivity > 15 : 1, unless otherwise noted.

^[c] Reaction conducted with 1 equiv carbohydrate, 1.50 equiv azide, and 1.50 equiv PhP(CH₃)₂, THF, 55 °C, 12 h.

PMB = *p*-methoxybenzyl ; CBz = benzyloxycarbonyl.

Table 3Glycosylated amino acids and peptides prepared by the sequence outlined in Scheme 2.^[a]

1. **1h**, P(CH₃)₂Ph
 2. R'-CO₂H, ClCO₂tBu, TEA
 3. CAN, CH₃CN / H₂O

Entry	Product	Yield [%] ^[b]
1	<p>5b</p>	76
2	<p>5c</p>	75
3	<p>5d</p>	70
4	<p>5e</p>	61

^[a] Conditions : Step 1: **1h** (1.50 equiv), PhP(CH₃)₂ (1.50 equiv), **2d** or **2f** (1 equiv), THF, 55 °C. Step 2 : ClCO₂tBu (1.20 equiv), amino acid (1.10 equiv), TEA (1.20 equiv), THF, -40 → 24 °C. Step 3 : CAN (5.00 equiv), CH₃CN/H₂O (30 : 1, v/v), MW, 90 °C, 10 min.

^[b] Yield of isolated product after purification by flash-column chromatography. β : α selectivity > 15 : 1.

TEA = triethylamine, MW = microwave, Bn = benzyl, Fmoc = 9-fluorenylmethoxycarbonyl.