Published in final edited form as:

J Mol Cell Cardiol. 2016 April ; 93: 162–174. doi:10.1016/j.yjmcc.2015.11.010.

Integrins and Integrin-related Proteins in Cardiac Fibrosis

Chao Chen, Ruixia Li, Robert S. Ross¹, and Ana Maria Manso¹

Department of Medicine, Cardiology, UCSD School of Medicine, La Jolla, CA, 92093-0613 USA, and Veterans Administration San Diego Healthcare System, San Diego, CA, 92161 USA

Abstract

Cardiac fibrosis is one of the major components of the healing mechanism following any injury of the heart and as such may contribute to both systolic and diastolic dysfunction in a range of pathophysiologic conditions. Canonically, it can occur as part of the remodeling process that occurs following myocardial infarction or that follows as a response to pressure overload. Integrins are cell surface receptors which act in both cellular adhesion and signaling. Most importantly, in the context of the continuously contracting myocardium, they are recognized as mechanotransducers. They have been implicated in the development of fibrosis in several organs, including the heart. This review will focus on the involvement of integrins and integrin-related proteins, in cardiac fibrosis, outlining the roles of these proteins in the fibrotic responses in specific cardiac pathologies, discuss some of the common end effectors (Angiotensin II, transforming growth factor beta 1 and mechanical stress) through which integrins function and finally discuss how manipulation of this set of proteins may lead to new treatments which could prove useful to alter the deleterious effects of cardiac fibrosis.

Abstract

¹Address correspondence to: Ana Maria Manso or Robert S. Ross, Biomedical Research Facility 2, Room 2A-17, 9500 Gilman Drive #0613-C, La Jolla, CA 92093-0613, Phone: (858) 246-0624; Fax: (858) 822-3027, amanso@ucsd.edu or rross@ucsd.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Disclosures of all authors: None

Keywords

Fibrosis; integrins; myocardium; transforming growth factor beta; angiotensin; mechanical stress

1. Introduction

Heart failure is a major cause of morbidity and mortality in the western world with limited numbers of therapeutics that impact the primary disease process [1]. Both heart failure with reduced ejection fraction (HFrEF) and heart failure with preserved ejection fraction (HFpEF) display a range of physiological and morphological changes, including fibrosis of the myocardium. Fibrosis is the excessive deposition of extracellular matrix (ECM) proteins into tissues, leading to scar formation, disruption of normal tissue architecture and potentially to organ failure [2]. In particular, cardiac fibrosis is one of the major components of the healing mechanism following any injury of the heart and as such may contribute to both systolic and diastolic dysfunction in a range of pathophysiologic conditions [3]. Canonically, it can occur as part of the remodeling process that occurs following myocardial infarction (MI) or that follows as a response to pressure overload (PO). It can even be currently tracked non-invasively in man, through use of gadolinium-based magnetic resonance imaging [4].

Multiple organs in the body can be affected by fibrosis, and a large effort has been focused on studying the fibrotic response that occurs in diseases of lung, kidney, liver and skin. Although the triggering events which lead to the fibrotic disorders in these non-cardiac organs could be quite different, the fundamental processes that drive fibrosis are likely to be common in most tissues throughout the body, including the heart. Importantly, disorders that lead to fibrosis can share the complex interplay between inflammatory, epithelial, myofibroblast and ECM responses [5–7]. Myofibroblasts are a main cell type that fuel fibrosis and have combined characteristics of fibroblasts and smooth muscle cells. As such they have a contractile phenotype and contribute significantly to the formation of scarring by secreting ECM components [8]. The origin of myofibroblasts in different organs has been intensely studied, with potential sources suggested to include resident fibroblasts, circulating progenitors (fibrocytes) and also potentially cells that arise by the process of epithelial – mesenchymal transition (EMT) [9, 10].

Integrins are cell surface receptors which act in both cellular adhesion and signaling. Most importantly, in the context of the continuously contracting myocardium, they are recognized as mechanotransducers [11–13]. They have been implicated in the development of fibrosis in several organs, including the heart [14]. In addition to their direct effects on cellular proliferation, migration and survival, mediated by their binding to ECM proteins, integrins can potentiate signals from soluble growth factors such as transforming growth factor β 1 (TGF β), and act as a receptors for matricellular proteins [15]. All of these properties allow integrins and proteins which interact with them, to play essential roles in the fibrotic process.

This review will focus on the involvement of integrins and integrin-related proteins, in cardiac fibrosis. While fibrosis can occur as a component of a wide variety of myocardial diseases, this review will focus on examples that occur following MI and as a result of

hemodynamic overload, aging and diabetes. Here we will provide information on how integrins are affected and involved in cardiac fibrosis. In the first part we will introduce basic information about integrins and integrin-related proteins. Then, we will outline the roles of these proteins in the fibrotic responses of specific cardiac pathologies, discuss some of the common end effectors through which integrins function and finally discuss how manipulation of this set of proteins may lead to new treatments which could prove useful to alter the deleterious effects of cardiac fibrosis.

2. Integrins and integrin-related proteins

There have been several recent excellent reviews written about the structure, function, expression and extensive animal modeling studies of integrins and integrin-related proteins. These include ones relevant to the heart, by our own group and others [11, 12]. Given this, here we will only provide a brief background on these proteins prior to our discussion of their role in the fibrotic process.

2.1 Integrins

Integrins are transmembrane receptors which act as bridges for cell-ECM connections and in some instances, cell-cell interactions. Thus one of their prime functions is to couple the ECM outside cells, to the cytoskeleton inside the cell. Integrin receptors are obligate heterodimers, composed of two different chains, termed the α and β subunits. In mammals there are 18 α and 8 β subunits, which combine to make up 24 different integrin combinations [16]. The integrin subunits can vary from 90 – 160 kDa and generally consist of a large extracellular domain, a single transmembrane spanning domain, and a short cytoplasmic tail [17]. The cytoplasmic domain of many of the β subunits is highly homologous, while the α subunit sequences are significantly more diverse. It is through the cytoplasmic tail, dominantly of β subunits, that the integrins bind both cytoskeletal linkers and also signal. (Figure 1)

Cell attachment to the ECM is a basic requirement to build a multicellular organism. Integrins are part of the cell adhesion complexes, which along with many cytoplasmic structural and signaling proteins, such as talin (Tln), vinculin (Vcl), paxillin (Pax), focal adhesion kinase (FAK), α -actin and integrin-linked kinase (ILK), serve to link two networks across the plasma membrane: the ECM and the intracellular actin filamentous system [18]. Thus integrins and their associated proteins, are not simply hooks in a cellular meshwork, but provide the cell with critical inputs about the nature of its surroundings. Although integrins do not possess their own enzymatic activity, they are potent bidirectional signaling receptors, playing an important role in cell signaling [19, 20]. When triggered by ligands, integrins can influence a host of downstream biochemical pathways in the cell interior, a process commonly termed **outside-in signaling**. This type of signaling may allow sensing of both chemical composition and mechanical status of the ECM outside the cell. Then, depending on the integrin's regulatory impact, the cell can experience growth, proliferation, division, differentiation or other means of remodeling. In addition, the control of integrin function occurs via regulatory signals that originate within the cell cytoplasm and are then transmitted to the external ligand-binding domain of the receptor. This concept is known as

inside-out signaling. It can increase both binding of integrin to ligand (ECM) and lead to clustering of multiple integrins in close spacing within the cell membrane.

The variety of integrin receptors expressed on a particular cell type can be unique. Further, expression of integrins may not only be restricted to a particular cell type, but can vary depending on developmental stage or pathological state. In addition, functional complexity of integrins also occurs since a single integrin receptor can bind to one or several ligands, and in addition, a single ligand can be bound by several integrin heterodimers. For example, in the cardiac myocyte (CM), the integrin heterodimers most highly expressed are $\alpha 1\beta 1$, $\alpha 5\beta 1$, and $\alpha 7\beta 1$, which are predominantly collagen, fibronectin, and laminin-binding receptors, respectively. In addition, $\alpha 6$, $\alpha 9$, and $\alpha 10$ are also detected in myocytes. $\beta 1$ is the dominant CM β integrin subunit, but $\beta 3$ and $\beta 5$ subunit function have also been studied [21–23]. In contrast, cardiac fibroblasts (CFs) express $\alpha 1\beta 1$, $\alpha 2\beta 1$, $\alpha 3\beta 1$, $\alpha 4\beta 1$, $\alpha 5\beta 1$, $\alpha 6\beta 1$, as well as $\alpha \nu\beta 1$, $\alpha \nu\beta 3$ and $\alpha \nu\beta 5$. These integrin pairs play critical roles in cardiac remodeling via ECM-integrin interaction. Endothelial cells (EC) express $\alpha 1\beta 1$, $\alpha 2\beta 1$, $\alpha 5\beta 1$ and $\alpha \nu\beta 3$ integrins. As will be relevant to later discussion, $\beta 2$ integrins are confined to leukocytes, being expressed on cells such as macrophages and neutrophils.

Matrix metalloproteinases (MMPs), a large family of calcium-dependent zinc-containing endopeptidases can mediate degradation of ECM components during various physiological and pathological processes. Some ECM components, through interaction with integrin receptor and modulation of downstream signaling, are capable of regulating expression and activity of several MMPs. One example is that $\alpha 4\beta 1$, $\alpha 5\beta 1$ as well as $\alpha \nu\beta 3$ integrins can mediate expression and activity of MMPs and their effector responses, in different cellular systems. Damsky's group found that when FN is bound to $\alpha 4\beta 1$ and $\alpha 5\beta 1$ integrins, MMP expression in rabbit synovial fibroblasts is regulated [24]. This occurred when the fibroblasts were cultured on surfaces coated with the central RGD-containing region of FN (120FN) as well as when the cells were incubated with anti- $\alpha 5\beta 1$ or anti- $\alpha 4$ antibodies. Brooks and colleagues later found that MMP-2 was localized in a proteolytically-active form on the surface of invasive cells when they bound directly to $\alpha \nu\beta 3$ integrin[25].

Although integrins can alter the expression and the localization of MMPs, MMPs can also cleave integrins and affect the integrin signaling pathway. For example, MMP-2 impairs $\beta 1$ integrin-mediated survival signals produced by activation of focal adhesion kinase (FAK) to protect β –adrenergic receptor-stimulated apoptosis of adult CMs. Overexpression of $\beta 1$ integrins also inhibited apoptosis induced by purified, active MMP-2, in adult CMs [26]. Recently, MMP-9 was found to cause shedding of the $\beta 2$ integrin subunit (CD18) from macrophages, which suggests that MMP-9 can play an important role in $\beta 2$ integrin post-translational modification [27]. Further, other studies found that MMP-2 is up-regulated in invasive colorectal tumors and with this, caused shedding of $\beta 1$ integrin followed by subsequent integrin degradation. This lead to decreased adhesion and enhanced cell motility [28]. MMP mediated-shedding of integrins is a posttranslational modification which can alter integrin activation state, as well as integrin receptor internalization and degradation.

2.2 Integrin-Related Proteins

Integrins do not possess their own enzymatic or actin-binding activity. Therefore, various adaptor proteins that bind to the cytoplasmic tails of α and β subunits are required to mediate structural or scaffolding properties, and produce catalytic activity (i.e. outside-in signaling), or activate integrins to effect ECM binding (inside-out signaling). Since some of these proteins crucial for integrin function have not been investigated in the fibrotic process, we will only introduce a few of these proteins below.

Talin (Tln) is a ubiquitously expressed, cytosolic protein that is found in high concentrations in focal adhesions (FA). It is a large protein (270kDa) which links integrins to the actin cytoskeleton either directly or indirectly, by interacting with vinculin (below) and α -actinin [29]. Tln has been shown to be an essential protein for force generation and mechanotransduction [30]. In addition to its structural role, Tln is essential for integrin activation [31]. In vertebrates, there are 2 Tln isoforms, Tln1 and Tln2 [29].

Integrin Linked Kinase (ILK) is a 51 kDa serine/threonine protein kinase that interacts with $\beta 1$ and $\beta 3$ integrin cytoplasmic domains, linking integrins to the actin cytoskeleton and mediating integrin signaling in diverse types of cells. ILK is suggested to elicit its biologic activity through its role as a scaffolding protein, binding proteins such as PINCH and Parvin, as opposed to a role as a kinase protein [32].

Focal adhesion kinase (FAK) is a 125kDa ubiquitously expressed non-receptor tyrosine kinase that plays a major role in integrin-mediated signal transduction. FAK can be activated by either ECM (through integrins) or by growth factors, regulating multiple signaling pathway outputs [33]. The C-terminal domain of FAK promotes its colocalization with integrins through its association with integrin-associated proteins such as Pax and Tln [34]. In vivo and in vitro studies showed that FAK is rapidly activated by mechanical stress [35–37]. Proline-rich tyrosine kinase-2 (Pyk2) is a tyrosine kinase related to FAK that shares a similar domain structure and has common phosphorylation sites [38].

Kindlins are 78kDa cytosolic proteins that directly interact with the cytoplasmic tail of β integrin and are required for the correct assembly of FAs. They act as crucial co-activators of integrins. There are 3 subtype kindlins: kindlin-1, -2, and -3, that have varied expression patterns. Kindlin-1 and kindlin-2 are widely expressed in murine and human tissues, while kindlin-3 is restricted to hematopoietic tissues. There are 4 kindlin-binding proteins: ILK, migfilin, $\beta 1$ integrin, and $\beta 3$ integrin [39].

3. Integrins and Integrin-related proteins in cardiac fibrosis

Cardiac fibrosis, characterized by the excessive production and deposition of scar tissue, is often a result of chronic conditions such as hypertension and diabetes, or acute conditions such as MI or myocarditis. It also occurs with normal aging. There are different types of fibrosis in heart.

With death of CMs as occurs after MI, the regions of the heart from which the myocytes are lost undergoes fibrotic change termed “replacement fibrosis.” With stress related remodeling

of cardiac chambers as can occur with PO states (e.g. with hypertension or valvular stenosis,) or in the remote regions following MI, “reactive fibrosis” occurs in an interstitial or perivascular pattern. All of these types of fibrosis may lead to systolic and / or diastolic decrements in function, and further, may even predispose to arrhythmias given the loss of myocyte-myocyte connections, and potential influence of non-myocytes (e.g. fibroblasts or other cells) as they interface with myocytes [40].

Importantly, a cell death mechanism termed anoikis must be considered in the context of how integrins may be related to fibrosis. Anoikis is defined as programmed cell death induced by the loss of cell-matrix interactions [41, 42]. Thus integrins are central to this process. Anoikis may play a physiological role by regulating cell homeostasis in developing or mature tissues. However, anoikis can also be involved in pathological processes. While anoikis has been principally linked to a role in cancer, it may also serve important roles in the cardiovascular system. For instance it likely is a means to produce CM cell loss in the context of multiple myocardial stressors [43]. Lorell and colleagues found that the localization of $\beta 1$ integrin was increased on the cell surface of CMs, and in the ECM surrounding CMs, during the early stage of heart failure, as perhaps integrins were shed from the cells [44]. Loss of CMs occurred soon after left ventricular pressure overload, and with this anoikis was indicated to contribute to the progression towards heart failure. Since pressure loading produced abnormal myocyte-ECM anchorage, and subsequent anoikis mediated-cell death, replacement fibrosis could ensue. Similar results could occur with other cardiovascular pathologies. Having the above as background, we will now focus on the behavior and role that integrins and associated proteins play in the fibrotic process in some key cardiac pathologies.

3.1 Cardiac fibrosis after myocardial infarction

Cardiac remodeling and particularly fibrosis, both in the post-MI infarcted and non-infarcted regions, is recognized to be a major determinant of subsequent development of impaired ventricular function [45, 46]. After MI, both replacement and interstitial fibrosis occur and are determined by the extent of myofibroblast proliferation, and also by its associated collagen deposition. Wound healing after MI entails a complex cascade of events that involves interplay between several different cell types, including inflammatory cells, endothelial cells (EC), fibroblasts and myofibroblasts [47]. Infarct healing can be divided into three distinct, but overlapping phases in man: the inflammatory phase (from infarct onset to 2 days post-MI), the proliferative phase (from 3 – 4 days to 2 weeks after infarction), and the maturation phase (from 2 weeks after MI, onward) – see below [47]. In response to post-MI cell death, intracellular contents are released from the necrotic cells and activate innate immune mechanisms, initiating an intense, but transient inflammatory response. As the infarct zone is cleared of dead cells, the inflammatory response is suppressed and infiltration of CF and EC into the infarct zone occurs. This marks the beginning of the proliferation phase. During this phase, CFs undergo dramatic phenotypic and functional changes showing high proliferative and migratory activity, acquisition of the myofibroblast phenotype with α -smooth muscle actin expression (α -SMA), and augmented matrix synthetic capacity (collagens I and III), all of which is important to maintain the structural integrity of the infarct area. TGF β signaling, mechanical stress and specialized

matrix proteins such as the ED-A isoform of cellular fibronectin (ED-A FN), are some of the main promoters of CF to myofibroblast transdifferentiation [48]. ED-A is one type of fibronectin subunit that is expressed commonly during wound healing and in areas of fibrotic change. Each FN subunit is formed from a series of repeating, homologous modules, and contains binding sites for cell surface receptors such as integrins, as well as other ECM components. FN polymorphisms follow from the alternative splicing of the type III segments and these variants are termed ED-A, ED-B, and IIICS [49]. Other factors that play important roles during this phase are angiotensin II (Ang II) and the matricellular proteins thrombospondin-1 (TSP-1), osteopontin (OPN), periostin and tenascin-C [48].

As the infarcted zone is filled with matrix, cellular proliferation is inhibited and the maturation phase begins. As the infarct matures, matrix cross-linking results in the formation of a dense, collagen-based scar. Collagen deposition also occurs in the un-infarcted, remote myocardial region, predominantly in the interstitium, where these deposits clearly contribute to ventricular stiffness, arrhythmogenesis and dysfunction [50]. Neurohumoral factors, such as Ang II and TGF β 1, are preferentially expressed at the infarct border. There they traverse the common interstitial space and enhance collagen deposition at sites distant from the MI, contributing to the remote interstitial fibrosis. Some studies also suggest that mechanical alteration in the infarcted myocardium leads to persistent myofibroblast expansion and excessive ECM production, due to the failure of the apoptotic mechanism of these cells, favoring remote zone fibrosis [51].

Undoubtedly, since integrins serve at the interface between cells and ECM, they are involved in the different healing / remodeling phases after MI. Initial work has been performed to evaluate the changes in several α integrin subunits after MI in the rat [52]. In hearts without infarction, no expression of α 1 integrin, moderate expression of the α 3 integrin and only slight expression of the α 5 subunit were observed in myocardium. In the first week after MI, the α 1 subunit, collagen and fibronectin were increased only in the peri-infarcted area, while the α 5 subunit was increased both in peri-infarcted and non-infarcted areas. At day 42, the expression of the α 1 subunit and collagen were still increased, although the α 5 subunit and fibronectin were decreased. The expression of the α 3 subunit was not altered throughout the experimental period. Thus α 1 integrin remained increased during a significant duration after MI, in the peri-infarcted and non-infarcted areas; α 5 integrin expression levels increased during the healing process and decreased during the remodeling process. This data suggests that different α integrin subunits may play varied roles during the MI process.

β integrins may also play an important coordinating role in ECM synthesis and remodeling in the heart, as was demonstrated previously in skin and lung injury [53–55]. Liu's group studied the role of β integrins after MI using rat and mouse model systems [56]. Both β 1 and β 3 integrins had low expression basally, but increased at the infarct zone by day 3 post-MI, peaking at day 7 post-MI, then gradually declined thereafter and returned towards baseline. In line with the cell-type specific expression of β 1 integrin isoforms, the β 1A isoform was found primarily expressed on CF and inflammatory cells, while the β 1D isoform was expressed on CMs. β 3 integrin was detected principally on ECs and smooth muscle cells in the peri-infarct vessels [56]. The temporal expression and spatial localization of these various integrin subunits suggests they play an important role in healing and remodeling

processes after MI. In other studies, Singh's group used $\beta 1$ integrin heterozygous knockout (hKO) mice and studied them one month after MI [57]. MI increased $\beta 1$ expression in both KO and control groups, with the increase mainly observed in the peri-infarct and non-infarcted areas. However, the increase was lower in hKO, that had one $\beta 1$ integrin allele deleted, and the hKO group displayed worsened cardiac function than the WT controls, after MI. Increased apoptosis and cardiac fibrosis in the hKO was also found after MI. Together this data suggests that $\beta 1$ integrins are crucial in post-MI remodeling [57].

ILK expression was also found to be increased following MI, in the infarct area [58]. Forced increase of ILK expression, using gene therapy, has been shown to improve cardiac remodeling in rats after MI. Recombinant adenoviral-ILK was delivered during coronary ligation induced MI, in the rat. At the 4-week follow-up study, ILK treated animals showed improved cardiac function, along with decreased: a) infarct size, b) interstitial fibrosis and c) CM apoptosis, as compared to the control group [59, 60]. Additionally, ILK- overexpression in mesenchymal stem cells injected into porcine peri-infarct myocardium 7 days post-MI, preserved cardiac function and myocardial perfusion, reduced fibrosis, increased CM proliferation, and enhanced angiogenesis [61]. These studies suggest ILK overexpression may preserve cardiac function and reduce cardiac fibrosis post-MI.

Although activated myofibroblasts are the main effector cells in the fibrotic heart, monocytes / macrophages, lymphocytes, and mast cells also contribute to the fibrotic response. The mechanisms how these immune cells work on healing and remodeling post-MI has been recently extensively reviewed and will not be discussed here [62–67]. Yet, different immune cells express varied integrins. In leukocytes, $\beta 2$ as well as $\alpha 4\beta 1$ and $\alpha 4\beta 7$ (LPAM-1) integrins play roles in cellular recruitment and in inflammatory disorders. The $\beta 2$ -integrins consist of a common β subunit (CD18) that associates with 4 different α subunits. They comprise the $\alpha L\beta 2$ -integrin (lymphocyte function-associated antigen 1 (LFA-1); CD11a/CD18) and the $\alpha M\beta 2$ -integrin (macrophage-1 antigen (Mac-1) also designated complement receptor 3 (CR3); CD11b/CD18), which are the most crucial $\beta 2$ -integrins for leukocyte recruitment, as well as the $\alpha X\beta 2$ (CD11c/CD18; p150,95; CR4) and the $\alpha D\beta 2$ (CD11d/CD18) integrins [68]. LFA-1 is expressed by neutrophils, monocytes and lymphocytes, whereas Mac-1 is found mainly on neutrophils and monocytes, and VLA-4 is expressed on monocytes and T lymphocytes [68–71]. $\alpha X\beta 2$ is present on macrophages and dendritic cells [72, 73], and $\alpha D\beta 2$ is expressed on monocytes/macrophages, especially foam cells, which are macrophages found in atherosclerotic lesions [74, 75].

Since immune cells are an essential part of the healing and remodeling process, and different immune cells express varied integrin receptors, anti-integrin therapeutic approaches could be promising to reduce infarction size and adverse post-MI remodeling. Both antibody neutralization studies (anti-Mo1, anti-CD11b) [76] and genetic loss-of-function models [77], documented the crucial role of $\beta 2$ integrins, in recruitment of neutrophils into the infarcted myocardium, and in modifying how adherent neutrophils transmigrate into the infarct through interactions that may involve endothelial transmembrane proteins, including platelet endothelial cell adhesion molecule (PECAM)-1, Intercellular Adhesion Molecule 1(ICAM-1), vascular-endothelial (VE)-cadherin and members of the junctional adhesion molecule family. Several independent studies in animal models have suggested marked

reduction in infarct size upon administration of anti-CD11/CD18 antibodies [76, 78, 79]. Unfortunately, three small clinical trials showed no effects of early anti-integrin approaches in reducing infarct size in human patients with MI [79–81].

Administration of the anti- $\alpha 4$ integrin antibody (natalizumab) was found to block macrophage trafficking to the heart in a Simian immunodeficiency virus (SIV) infection model of AIDS. With this, it decreased macrophage numbers in cardiac tissues, and decreased fibrosis in the SIV-infected group [82]. These data demonstrate a role for macrophages in the development of cardiac inflammation and fibrosis, and suggest that blocking $\alpha 4$ integrin, with subsequent reduction of monocyte/ macrophage traffic to the heart, can alleviate SIV/HIV and potentially be useful as a therapy with other viral-associated myocarditis and associated fibrosis.

As even further evidence of integrin-linked immune responses to cardiac fibrotic disease, are studies with CD11b/CD18 integrin knockout (KO) mice which showed reduced Ang II induced atrial fibrosis and atrial fibrillation, likely by inhibition of neutrophil infiltration during this process. In summary, modification of integrin expression / function on inflammatory cells might be a target that can reduce cardiac fibrosis.

3.2 Pressure overload and fibrosis

Fibrosis also underlies the remodeling response which occurs in the mammalian heart as it responds to hemodynamic loading provoked by PO. Arterial hypertension induces long term structural changes in the myocardium that together may act as risk factors for the evolution from a compensated state, towards heart failure. These responses include cardiac hypertrophy and cardiac fibrosis. Remodeling secondary to hypertension is attributed to mechanical stress, and to the hypertrophic, pro-inflammatory and pro-fibrotic effects of vasoactive factors such as Ang II, catecholamines, aldosterone, endothelin and TGF β 1, which are released by the cells within the stressed myocardium. In this process, fibrosis is found in spatially distinct locations - interstitial fibrosis that occurs around CMs, and perivascular fibrosis, which is detected in the vicinity of large coronary vessels. With development of fibrotic remodeling of the cardiac interstitium, increased stiffness, arrhythmogenesis and diastolic dysfunction can result [83]. Fibroblasts respond to alterations in mechanical loading by enhancing their matrix-synthetic capacity and as discussed above, by transdifferentiating into myofibroblasts. Matricellular proteins also contribute to the fibrosis following PO [15].

Recent work has begun to show that integrins and their downstream kinase FAK, are affected and involved in PO mediated cardiac fibrosis [84]. For example, Burgess, et. al. studied rats subjected to treadmill exercise or hypertension. In these models, $\beta 1$ integrin expression was elevated in CFs, while $\alpha 1$ and $\alpha 2$ integrin levels both decreased, but $\alpha 5$ integrin increased, in the exercise group and decreased in the hypertensive one [85]. These results show that CFs respond to changing environments in pathophysiological conditions by modulating integrin expression. FAK also was found to play a role in the activation of CFs in response to cyclic stretch and in the development of cardiac fibrosis provoked by PO [86]. FAK silencing by small-interfering RNA (siRNA) attenuated fibrosis, collagen content, and activity of matrix metalloproteinase-2 (MMP-2) in the overloaded left ventricle (LV) [87].

Another study showed that CF $\beta 3$ integrin is critical for ECM accumulation during PO hypertrophy in mouse. $\beta 3$ KO mice displayed attenuated cardiac fibrosis compared to WT controls. CFs from $\beta 3$ KO mice exhibited a significant reduction in cell-matrix adhesion, cell spreading, proliferation and migration. In addition, the activation of platelet-derived growth factor (PDGF) receptor associated tyrosine kinase, and the non-receptor tyrosine kinase Pyk2, were impaired in $\beta 3$ null cells following PDGF stimulation. These results indicate that $\beta 3$ integrin-mediated Pyk2 signaling in CF plays a critical role in PO -induced cardiac fibrosis [88].

3.3 Cardiac fibrosis in aging

Fibrosis is also a hallmark of cardiovascular aging [89]. Development of heart disease has high prevalence in the elderly and has multifactorial causes that can summate in heart failure. Accumulation of collagen with resultant fibrosis is considered as a major contributor to reduced ventricular compliance, potentially contributing to HFpEF [90]. The increase in fibrosis not only affects the mechanical properties of the heart but has also been proposed to affect electrical properties, which may lead to arrhythmias and sudden cardiac death. Since integrins and their related protein partners mechanically couple the ECM with the intracellular cytoskeleton, it is proposed that these connections can be a component of aging related cardiac fibrosis [91].

With numerous changes occurring in the ECM of aging heart, it is reasonable to speculate that age-associated changes in integrin expression are a component of this process. Aged mice (20 months) express higher levels of collagen and fibronectin than middle aged (12 months) and younger mice (2 months). Consistent with the temporal increase in the fibronectin and collagen content within the heart, are parallel increases in the expression of integrin subunits which can bind these ECM proteins, i.e., $\alpha 1$ and $\alpha 5$ integrins. In contrast, the $\beta 1$ integrin levels in old mouse hearts is significantly less than that detected in middle aged and young mice [92]. Age-associated decreases in $\beta 1$ integrin was also observed in the myocardium of Wistar-Kyoto rats [93]. Likewise, adult CFs express lower levels of $\beta 1$ integrins when compared to neonatal CF [94] and aged myocytes exhibit decreased levels of $\beta 1$, $\alpha 3$, and $\alpha 3\beta 1$ integrins, when compared to younger adult myocytes [95]. While little work has been performed investigating how loss of integrin subunits might affect cardiac aging (e.g. using various KO mouse models) it is interesting to note that reduction of $\beta 1$ integrin in CMs also induced cardiac fibrosis [96], which suggests that maintenance of normal $\beta 1$ integrin levels in CMs may be required to prevent cardiac fibrosis in the aged myocardium.

3.4 Cardiac fibrosis in diabetic cardiomyopathy

Fibrosis is a frequent complication of diabetes mellitus in many organs and tissues but the mechanism of how diabetes impacts the formation of fibrotic lesions is not well defined [97]. Diabetic cardiomyopathy is characterized by the production of a disorganized, fibrotic matrix in the absence of coronary atherosclerosis and associated ischemic damage, or hypertension. Myocardial fibrosis and collagen deposition are the primary structural changes observed in diabetic cardiomyopathy [98–101]. Cardiac fibroblasts may play a key role in the in the elaboration of fibrosis in the diabetic heart with increased proliferation causing

remodeling of collagen [102], and also by induction of a switch from a fibroblast phenotype, to that of a myofibroblast, which is both profibrotic and highly contractile [103, 104].

In a recent paper, in which streptozotocin (STZ)-treated Sprague-Dawley rats were used as a diabetes model, Talior-Volodarsky and collages found that $\alpha 11$ integrin transcript and protein expression were increased in isolated CFs from STZ rats vs. controls. Knock-down of $\alpha 11$ integrin expression with siRNA, in human fibroblasts plated on methylglyoxal-treated collagen (a glycated collagen present in the diabetic heart) blocked the increase in transcript levels of TGF β 2 (a critical end-effector in the fibrotic process, see below) and the increased protein expression of α -SMA. This result suggests that $\alpha 11$ integrin and TGF β 2 could mediate myofibroblast differentiation in CFs in the diabetic heart, and therefore this integrin subunit may contribute to fibrosis associated with diabetic cardiomyopathy [105].

4. End effectors of the fibrotic process

A complex interaction among a network of growth factors, cytokines and hormones are responsible for initiating and maintaining fibrotic responses in vivo. In particular, Ang II and TGF β 1 appear to work together to induce activation of resident fibroblasts, promote persistence of myofibroblasts and induce expression of a variety of ECM components, in the myocardium [106]. Recently, it has become increasingly appreciated that in addition to these factors, the cellular microenvironment also plays a critical role in cardiac fibrosis. Here we will discuss how these common effectors lead to fibrosis in most tissues, including the heart. They have been shown to be regulated by or modulated through integrins and thus are particularly relevant to the current topic.

4.1 Integrin and integrin-related proteins in Ang II induced cardiac fibrosis

Ang II plays a critical role in cardiac remodeling. This peptide promotes CM hypertrophy and interstitial fibrotic changes associated with PO hypertrophy, post-MI remodeling, and congestive heart failure. Importantly, Ang II is both expressed and activated by macrophages and myofibroblasts during remodeling after MI [107]. In CM and CF, Ang II induces the expression of TGF β 1 (see below) through the angiotensin type I receptor [108], and, in vivo, TGF β is required for Ang II to cause both cardiac hypertrophy and fibrosis [109]. This data supports the hypothesis that Ang II is upstream of TGF β 1 in driving cardiac fibrosis. In this regard, angiotensin receptor blockers such as losartan, are effective in reducing cardiac fibrosis in both animals and humans [110, 111].

Ang II can affect integrin expression and at the same time, integrins can regulate the expression of the angiotensinogen gene. Ang II treatment of CFs in vitro has been shown to increase the expression of $\alpha v\beta 3$ and $\alpha 8\beta 1$ integrins [112–114]. A 14 day infusion of Ang II into rats increased the expression of $\alpha 8\beta 1$ in the myocardium, where it was localized in vascular smooth muscle cells (VSMC) and myofibroblasts [115]. Carver et al., showed that $\alpha 8\beta 1$ integrin plays a role in collagen contraction by CFs [116]. Though remaining controversial, some data show that deletion of the $\alpha 8$ integrin gene does not protect mice from myocardial fibrosis in a model of desoxycorticosterone-acetate (DOCA)-salt hypertension [117]. In addition, $\beta 1$ integrin has been found to regulate expression of angiotensinogen in CFs following mechanical stretch. This effect is mediated by activation

of Rac1 and inhibition of RhoA, intracellular kinases involved in cytoskeletal organization and contraction [118]. Further, Ang II activation can provoke a stabilized scaffold of integrin-related proteins such as FAK and Tln in cells, including ones in the cardiovascular system, producing more adherence to the ECM, thus influencing the fibrotic process [119].

Ang II also induces cardiovascular injury in part, by activating an inflammatory response. In two recent papers, macrophages and neutrophils have been shown to play an important role in cardiac fibrosis following Ang II infusion. In mice treated with Ang II, Tenascin-C was unregulated, accelerating macrophage migration and synthesis of proinflammatory and profibrotic cytokines in the myocardium, ultimately resulting in a significant fibrotic response [120]. Tenascin-C is a matricellular protein not detected in normal adult heart, but expressed in several heart diseases closely associated with inflammation. It binds to multiple cell surface receptors, including several integrins [121]. In one study, Shimojo et. al. demonstrated that $\alpha v \beta 3$ is the functional receptor in macrophages for Tenascin-C, mediating migration and the inflammatory response [120]. In another study, Friedrichs et. al. demonstrated that Ang II-induced infiltration of neutrophils into atrial tissue was mediated by $\alpha M \beta 2$ integrin. In turn, this caused fibrosis which promoted the initiation and propagation of atrial fibrillation [122].

Ang II is also a potent inducer of OPN expression in different cell types such as smooth muscle cells and fibroblasts [123]. OPN is a glycoprotein that acts as a cytokine and also as a matricellular protein when immobilized to the matrix. It is expressed by many immune cells such as macrophages, and is markedly upregulated in response to tissue injury [124]. OPN has two types of cell surface receptors, CD44 and integrins ($\alpha v \beta 1$, $\alpha v \beta 3$, $\alpha v \beta 5$, $\alpha v \beta 6$, $\alpha 5 \beta 1$, as well as $\alpha 4$ and $\alpha 9$ subunits) [125, 126], and is markedly upregulated after MI and in models of cardiac hypertrophy and fibrosis due to PO and Ang II infusion [127–129]. In a model of Ang II-induced fibrosis, OPN KO mice had markedly reduced fibrosis compare to WT mice [128]. OPN may thus promote matrix deposition by enhancing macrophage infiltration or by modulating fibroblast function. It has been shown to mediate proliferation in CFs stimulated with Ang II [130]. Whether interactions with specific integrins or CD44 mediate the profibrotic actions of OPN needs further investigation.

4.2 Integrins and integrin related proteins in TGF β 1 activation, Epithelial to Mesenchymal Transition (EMT) and fibrosis

TGF β 1 has been reported to be a key mediator of CF activation and is recognized as a major factor in the development of fibrosis in multiple organs and in a variety of cardiac pathologies [131]. In cardiac fibrosis and remodeling, TGF β 1 has been shown to be initially derived from immune cells and then subsequently to be produced by myofibroblasts [132]. Inhibition of TGF β 1 prevented late cardiac remodeling in a mouse MI model [133]. In PO induced cardiac remodeling, antibodies against TGF β reduced myocardial fibrosis without affecting resultant hypertrophy or cardiac function [134]. TGF β 1 appears to be also implicated in the pathogenesis of age-related fibrotic cardiomyopathy that can potentially be a component of HFpEF. TGF β 1 heterozygous KO mice displayed an improved lifespan that was associated with reduced myocardial fibrosis and improved left ventricular compliance when mice were examined at 24 months of age [135].

There are three isoforms of TGF β , and all are synthesized as precursor proteins that are processed by proteolytic cleavage in the endoplasmic reticulum and assemble as a non-covalent complex of a disulfide-linked homodimer of the mature cytokine (a short C-terminal fragment) and a disulfide-linked homodimer of a large amino terminal fragment termed the latency associated peptide (LAP) (Figure 2). The LAP homodimer prevents the mature TGF β from binding to its receptors and therefore inhibits further downstream signaling from TGF β . This complex termed the small latent complex, SLC, is further modified and linked to another family of proteins called latent TGF β binding proteins (LTBP) to form the large latent complex (LLC). (Figure 2) Upon secretion LTBP is chemically cross-linked to the ECM, allowing it to store and tether TGF β in a latent form in the extracellular space [136, 137]. Latent TGF β 1 is converted into its active form by various mechanisms. Proteolytic cleavage of LLC and liberation of active TGF β 1 can be performed by bone morphogenetic protein 1, several MMPs, plasmin, elastase, thrombin, and cathepsin. Independent of proteolytic cleavage, interaction between LAP and TSP-1 also promote latent TGF β 1 activation [138].

Integrins have also been shown to bind, and in some cases activate, latent TGF β 1. α v integrins (α v β 1, α v β 3, α v β 5, α v β 6, and α v β 8), integrins α 5 β 1, α 8 β 1 and the platelet integrin α IIb β 3, can bind latent TGF β 1 [139]. Integrins α v β 3, α v β 5, α v β 6 and α v β 8 have been shown to bind the RGD sequence in the LAP of TGF β 1 and TGF β 3 [140–144].

Two main models are proposed to explain how integrins contribute to latent TGF β 1 activation. In the first, integrins α v β 8 and α v β 3 are suggested to simultaneously bind the latent TGF β 1 complex and MMPs. This allows the latent TGF β 1 complex and proteases to be positioned in close proximity and facilitates the enzymatic cleavage and release of active TGF β 1. Alternatively, it is suggested that integrins α v β 3, α v β 5, α v β 6 and α v β 8 change the conformation of the latent TGF β 1 complex by transmitting cell traction forces (Figure 2). This second mechanism requires the association of the latent complex with ECM, and is independent from proteolysis [138]. Knowing that different integrins use varied mechanisms to activate latent TGF β depending on the cell type and organ, may help us to develop new therapeutic strategies directed specifically at the organ and cell type planned for the targeted therapy.

4.2.1 TGF β 1 activation by epithelial cells (α v β 6)—One of the best characterized integrin-dependent mechanisms of TGF β activation has been studied in epithelial cells which contain the α v β 6 integrin receptor. Although this integrin is largely restricted to epithelial cells which are not expressed in heart, these studies provide a foundation to understand how integrins interface with TGF β activation and thus are relevant to our discussion. Specifically, several studies support the role of α v β 6 integrin in TGF β 1 activation and link it with pulmonary, liver, kidney and skin fibrosis. In healthy lungs, the α v β 6 integrin is minimally expressed in alveolar epithelial cells. Yet after lung injury, it is rapidly upregulated [145]. Similarly, α v β 6 integrin expression is low in kidney and liver, but marked induction is observed in epithelial cells in several renal and hepatic diseases associated with inflammation and fibrosis [146–148]. Genetic ablation of β 6 integrin expression or treatment with anti- α v β 6 antibodies or α v β 6 inhibitors (EMD527040) have

been shown to attenuate the accumulation of myofibroblasts and the interstitial deposition of collagen in models of lung, kidney and liver fibrosis [146, 148–150].

The $\alpha v\beta 6$ integrin can bind directly to the LAP of TGF β 1 and TGF β 3. Cells expressing $\alpha v\beta 6$ can activate TGF β 1 in vitro, a process which can be completely inhibited by anti- $\beta 6$ integrin blocking antibodies [151]. Activation of TGF β 1 in these cells can also be inhibited by the use of actin polymerization and Rho inhibitors, which indicates the important role of force generation by the actin cytoskeleton in this process [142, 152]. This was confirmed by Shi and colleagues, in studies where they solved the crystal structure of the small latent complex of TGF β and demonstrated that mechanical force generated by the contractile actomyosin cytoskeleton and transmitted by integrins, is a common mechanism to activate TGF β [153]. (Figure 2)

In addition to integrins themselves, some integrin-related proteins have also been linked to TGF β 1 activation in epithelial cells and therefore in fibrosis. Depletion of kindlin-2 by siRNA ameliorated renal tubulointerstitial fibrosis after unilateral ureteral obstruction. In vitro studies showed that kindlin-2 knockdown in tubular epithelial cells suppressed the activation of TGF β downstream signaling [154].

4.2.2 TGF β 1 activation by mesenchymal cells (fibroblasts, mesangial and hepatic stellate cells) ($\alpha v\beta 8$, $\alpha v\beta 3$, $\alpha v\beta 5$)—Myofibroblasts are derived from mesenchymal cells and as discussed above, one of the main cell types involved in the fibrotic response in the myocardium. They are a major source of ECM proteins, and also can liberate and activate TGF β 1. Several studies have shown that integrins in mesenchymal cells are involved in the development of fibrosis in different organs and act via activation of TGF β 1. Though some of the following discussion centers on varied organs, the overall discussion has relevance to cardiac fibrosis, as parallel studies may evolve from future studies in heart.

$\alpha v\beta 8$ integrin is increased in the airway fibroblasts of chronic obstructive pulmonary disease patients [155]. Conditional deletion of the $\beta 8$ integrin subunit in fibroblasts inhibited airway fibrosis in IL-1 β and ovalbumin-induced airway fibrosis, in mice [156]. Similar to $\alpha v\beta 6$, $\alpha v\beta 8$ also binds and activates TGF β 1, yet while TGF β 1 activation by $\alpha v\beta 6$ is resistant to protease inhibitors, metalloprotease inhibitors abolish $\alpha v\beta 8$ -mediated TGF β 1 activation [141]. MT1-MMP (MMP14) has been shown to be the MMP involved in this process [141]. $\alpha v\beta 8$ activates TGF β 1 by binding the RGD sequence in the LAP and presenting the latent complex to cell-surface MMPs, which in turn cleave and degrade LAP and then release the active TGF β 1 form [141].

In mesangial cells, modified pericytes in the kidney, $\alpha v\beta 8$ appear to have the opposite effect. Its binding to the latent TGF β 1 complex maintained TGF β 1 in an inactive state [157]. This highlights how the same integrin can have different effects depending in the cell type and/or organ in which it is expressed.

Hepatic stellate cells (HSC), the local pericyte population in the liver, are the major source of myofibroblasts in liver fibrosis. $\alpha v\beta 3$ integrin is expressed in human and rat HSCs, and its inhibition reduced proliferation and increase apoptosis of these cells in vitro [158]. Though

still somewhat controversial, the use of cilengitide (a pentapeptide containing the RGD sequence and that acts as an inhibitor for $\alpha v\beta 3$ and $\alpha v\beta 5$ function) increased the collagen deposition in two models of liver fibrosis [159].

$\alpha v\beta 3$ and $\alpha v\beta 5$ are also found to be upregulated in skin fibroblasts from patients with scleroderma. Both of these integrins have been shown to be involved in activation of latent TGF β 1 in primary cultures of these cells. Further, treatment of these cells with antibodies against $\alpha v\beta 3$ and $\alpha v\beta 5$ reduced the expression of collagen I [140, 144, 160, 161].

As we discussed above, αv integrin-mediated activation of latent TGF β 1 is a key event which promotes fibrosis in many organs. In a paper by Henderson et al., αv integrin subunit expression was reduced in myofibroblasts, leading to marked reduction in fibrosis in liver, lung and kidney, identifying myofibroblast αv -containing integrins as components of a core pathway widely shared by pathological fibrosis in multiple organs [162]. In another recent paper from the same group [163], they developed a highly specific small molecule inhibitor of $\alpha v\beta 1$ integrin (c8), which is highly expressed in active fibroblasts, and they obtained the same level of protection against fibrosis in liver and lung, as the one they reported in the mice with conditional deletion of all αv integrins from activated fibroblasts [162], suggesting that $\alpha v\beta 1$ is the major αv integrin responsible for that effect.

Further, FAK expression and activity were upregulated in fibrotic foci in lungs from pulmonary fibrosis patients, while siRNA-mediated down-regulation or inhibition (PF-562,271) of FAK prevented bleomycin-induced lung fibrosis in mice. They also proved that FAK deficiency impaired myofibroblast differentiation [164]. FAK has been shown to be a key mediator of TGF β signaling in lung and skin fibroblasts [165, 166].

The heart does not possess epithelial cells. Therefore, it is suggested that mesenchymal cells there may contribute to mechanical activation of latent TGF β 1. The mesenchymal integrins $\alpha v\beta 5$ and $\alpha v\beta 3$ are expressed in normal heart, and become upregulated during heart fibrosis [167] and after TGF β 1 and Ang II treatment of CFs [113]. Recent work identified integrins $\alpha v\beta 3$ and $\alpha v\beta 5$ as activators of latent TGF β 1 in human CF and showed their role in regulating CF to myofibroblast differentiation, through TGF β 1 activation [167]. Balasubramanian et. al., demonstrated that $\beta 3$ integrin KO mice show a substantially reduced accumulation of interstitial fibronectin and collagen after PO, compared to control WT mice [88]. These results are compatible with the hypothesis that loss of $\beta 3$ integrin expression, likely related specifically to $\alpha v\beta 3$, reduced activation of TGF β 1. Proof of this hypothesis requires further investigation.

$\alpha v\beta 1$ integrin is expressed in CFs but there is no information about how this integrin behaves in the heart undergoing remodeling, or in CFs stimulated with fibrogenic stimuli such as TGF β 1 or Ang II. Still, given recent data in other organs that implicates the important role of this integrin in organ fibrosis, work with relevant cardiac disease models should be explored to investigate the role of $\alpha v\beta 1$ in cardiac fibrosis.

4.2.3 Integrins and the epithelial-mesenchymal transition (EMT)—As mentioned above, the origin of myofibroblasts has been intensely studied, with potential sources

including resident fibroblasts, circulating progenitors, and also the possible derivation during the process of epithelial – mesenchymal transition (EMT) [10, 168]. EMT involves the conversion of differentiated epithelial cells into fibroblasts and myofibroblasts. This process occurs normally during embryonic development, but in aberrant wound repair processes, EMT may also lead to fibrosis. EMT has been associated with fibrosis in several organs such as kidney and lung. TGF β 1 is one of the main factors triggering EMT.

In lung, this process has shown to be regulated by the epithelial integrin α 3 β 1. Mice with epithelial cell specific deletion of α 3 integrin showed a reduction in myofibroblast number and collagen I deposition in the lungs, following bleomycin injury (a known inducer of fibrosis) compared to control treated mice [169]. In addition, the integrin related proteins ILK, PINCH and FAK have shown to be involved in EMT and kidney fibrosis [170–173]. Although earlier reports supported the contribution of epithelial cells to the myofibroblast population in kidney fibrosis, more recent genetic tracing studies failed to confirm these observations [174, 175]. Still, inhibition of EMT has been shown to be protective against kidney and lung fibrosis. Recent studies suggest that the pro-fibrogenic consequences of EMT activation are not mediated by simply inducing myofibroblast proliferation, but rather also occurs through impaired regenerative potential and cell-cell communication of epithelial cells [176, 177].

Endothelial–mesenchymal transition (EndMT) is a complex biological process distinct from, but similar to EMT, whereby ECs lose their specific markers and acquire a mesenchymal or myofibroblastic phenotype. With this they express mesenchymal cell products such as α -SMA and type I collagen. EndMT was originally observed during heart development, but recent studies have suggested that it also plays a role in pathological settings such as cancer and fibrosis. Several papers have shown that cardiac fibrosis is associated with the emergence of fibroblasts originating from ECs, and that TGF β 1 is the main factor that induces this process, suggesting that EndMT may well be relevant not only in development, but also in cardiac fibrotic disease [178–180]. Unfortunately, no studies about the regulation of this process by integrins or integrin related proteins in cardiac fibrosis has been performed to date. Still recent papers using genetic tracing techniques demonstrate that the contribution of EndMT to the fibroblast/ myofibroblast population after TAC and MI is negligible [181, 182], yet the authors did not rule out the possibility that this process could contribute to cardiac fibrosis without contributing to the myofibroblast population, as happens in kidney fibrosis [176, 177].

4.3 Integrins in mechanical stress and cardiac fibrosis

Beside biochemical factors, such as Ang II and TGF β 1 mentioned above, cues produced via mechanical strain and ECM stiffness, also play important roles in regulating myofibroblast differentiation and therefore fibrosis. Specifically, lung fibroblasts cultured on a stiff matrix have higher rates of proliferation and α -SMA expression, but decreased apoptosis compared with those grown on a compliant matrix [183]. Similarly, hepatic stellate cells become progressively activated with increasing levels of mechanical tension in the ECM [184]. On the other hand, decreasing substrate stiffness on which myofibroblasts are cultured, has been

shown to reverse myofibroblast differentiation and promote the quiescent fibroblast phenotype, in vitro [185].

Cardiac cells reside within one of the most mechanically dynamic environments of the body, and with each contraction they experience cyclic strain (relative deformation) and stress (force per unit area). During pathological conditions such as occur with MI or PO, those parameters change, affecting cardiac cell behavior. Several studies show that CFs respond in a varied way to stimulation by different magnitudes of strain, particularly as regards differentiation of myofibroblasts [186]. Furthermore, as mentioned above for other types of organs, several studies have shown that myofibroblast differentiation from CFs is regulated by ECM stiffness [187, 188]. Indeed, the relative high stiffness of injured and fibrotic areas within the heart may promote myofibroblast differentiation, which could prolong the existence of fibrosis via increase secretion of TGF β [189].

Integrins are mechanosensor adhesion proteins that transduce mechanical signals between the cells and their microenvironment, stimulating cellular responses such as cell growth and differentiation. Early studies in CFs by MacKenna et al demonstrated that integrins act as mechanotransducers in these cells, activating downstream signaling in response to mechanical stretch [190]. As mentioned above, β 1 integrins regulate the angiotensinogen gene in CFs, through p38 MAPK signaling, in response to mechanical stretch [118]. This effect is mediated by Rac1 and RhoA kinases. These two cases are examples of integrin outside-in signaling. In addition, as also mentioned above, integrins can participate in inside-out signaling. In this case, intracellular forces generated by actin fibers, can be transmitted through integrins to the ECM to alter its organization. A recent paper demonstrates that β 1 integrin binding plays a role in the constant traction force generation in response to varying stiffness for cells grown on cardiac ECM [191]. One relevant integrin inside-out signaling case relevant for fibrosis, is the activation of TGF β 1. TGF β 1 is secreted into the ECM in an inactive form, mechanical force generated by the contractile actomyosin cytoskeleton and transmitted by α v integrins, can change the conformation of the latent TGF β 1 complex, releasing its active form [153]. The contribution of mechanobiology and mechanosensing to the development of fibrosis are just beginning to emerge and their role in the myocardium requires significantly more study.

5. Integrins and FAK as therapeutics targets against fibrosis. Could manipulation of integrin expression / function be used therapeutically to modify cardiac fibrosis?

Because TGF β has been shown to have a crucial role in promoting the fibrotic response, initial approaches to fight fibrosis in different organs have focused on inhibition of this cytokine. Many of these studies have been hampered due to the multifunctional nature of TGF β . Due to the anti-proliferative role of TGF β on epithelial cells, its inhibition could have carcinogenic effects, which is particularly relevant for liver fibrosis [192]. Due to the critical role of TGF β as an immunosuppressant, generalized blockade of TGF β activity may lead to excessive autoimmunity and inflammation [193, 194]. Therefore, some anti-fibrotic strategies that aim to diminish TGF β signaling or prevent TGF β activation have gained much

more relevance in the fight against fibrosis. As above, the latent TGF β complex may be activated by proteolytic mechanisms or by traction forces mediated by integrins (mainly $\alpha\text{v}\beta 6$, $\alpha\text{v}\beta 3$ and $\alpha\text{v}\beta 5$). In this regard, an $\alpha\text{v}\beta 6$ -specific monoclonal antibody is undergoing phase II evaluation to treat idiopathic pulmonary fibrosis [195]. Small molecules inhibitors such as EMD527040, a non-peptide $\alpha\text{v}\beta 6$ integrin antagonist, have also shown to have therapeutic potential in animal models of liver fibrosis [150]. Also, a small molecule inhibitor for all αv integrins (CWHM12), has been shown to be effective in the prevention of liver and lung fibrosis in a mouse model [162]. Recently, a small molecule (c8) that inhibits $\alpha\text{v}\beta 1$ has been also found to be very effective against pulmonary and liver fibrosis [163]. Therefore, these two compounds could have a great potential to be broadly useful for treatment of diseases associated with excessive tissue fibrosis. Future studies will show the relevance of this work in other organs to treatment of cardiac fibrosis.

Although the myocardium does not have epithelial cells, inhibition of mesenchymal integrins in CFs ($\alpha\text{v}\beta 3$, $\alpha\text{v}\beta 5$, and $\alpha\text{v}\beta 1$), could be potential targets to suppress cardiac fibrosis through the inhibition of TGF β activation and fibroblast - myofibroblast differentiation. Therefore, the use of small molecules inhibitors such as CWHM12 or c8, should be evaluated in models of cardiac fibrosis. Still, one must be aware that generalized inhibition of $\alpha\text{v}\beta 3$ and $\alpha\text{v}\beta 5$ integrins could affect function of these integrins in ECs, thus altering angiogenesis that can be required for appropriate post-MI remodeling. Another important issue that has to be considered in post-MI remodeling is the timing of the treatment administration after MI. As it is broadly accepted, treatments that decrease collagen production should be avoided early after MI because they could jeopardize the formation of the mature scar and lead to ventricular rupture, therefore we posit that use of integrin inhibitors that target TGF β activation should be avoided during early phases of remodeling when they could affect collagen deposition and scar formation.

A variety of in vitro and in vivo studies with FAK inhibitors, suggests that FAK plays a key role in the development of fibrotic disorders in lung, skin and liver, and thus it appears to be an attractive target for anti-fibrotic therapy [196]. In recent years, several orally bioavailable ATP-competitive FAK inhibitors have been developed and have entered in clinical trials as anti-tumor treatments [197, 198]. PF-562,271 was one of the first clinically available FAK inhibitors. It inhibits FAK phosphorylation and in studies with animals has been shown to prevent bleomycin-induced lung fibrosis [164]. Another, FAK inhibitor PF-573,228 has been shown to be effective against skin fibrosis [199]. To date, no clinical studies in humans have evaluated the effects of FAK inhibitors in fibrotic diseases.

In the heart, FAK has been associated with CF proliferation, migration, in response to growth factors such as PDGF [84]. FAK has also been associated with cardiac fibrosis in in vivo models of MI and PO [87, 200]. Therefore, FAK inhibitors could be another potential therapeutic tool against cardiac fibrosis, although, as before, the timing of their use in MI patients may be difficult to predict, as importantly they should be limited to use in later phases of remodeling, when the scar is completely mature. Further studies in animal models of cardiac fibrosis with FAK inhibitors will be necessary to evaluate the potential use of these treatments.

6. Conclusion and future directions

A great amount of effort has been focused on the study of the varied mechanisms and factors that regulate fibrosis in different organs. Integrins have been shown to be one of the main common regulatory factors of this process. (Figure 3) They are able to regulate the main cytokine involved in myofibroblast trans-differentiation, TGF β 1, and are also well known as transducers of mechanical stress. It is apparent that increased integrin expression in specific cell types, may be linked to fibrosis. This has made them an attractive anti-fibrotic therapeutic target. Recently, several integrin inhibitors have been developed and preliminary animal studies have shown great potential for their use against fibrosis in multiple organs.

The study of integrins in cardiac fibrosis has not been as extensively investigated as it has in other organs. Indeed, some of the cardiac-oriented studies have simply been descriptive or observational in nature. Additional studies using genetically modified animals (knockouts, transgenics, etc.) with specific attention to sub-types of cardiac cells evaluated, as well as studies with specific integrin inhibitors, are necessary to better understand the role of integrins in the development of cardiac fibrosis. Since development of cardiac fibrosis is a factor in many cardiac pathologies that can lead to frank heart failure, development of new treatments focused on modulating integrin and integrin-related protein function may offer novel treatments that can favorably alter the cardiac remodeling process. Much new work is required for this purpose.

Acknowledgements

R. S. Ross (RSR) was supported by NIH grant HL115933, HL103566, HL066941 and HL127806.

Abbreviations

α-SMA	α -smooth muscle actin
Ang II	Angiotensin II
CFs	Cardiac fibroblasts
CM	Cardiac myocyte
ED-A FN	ED-A fibronectin
EC	Endothelial cells
EndMT	Endothelial–mesenchymal transition
EMT	Epithelial – mesenchymal transition
ECM	Extracellular matrix
FAK	Focal adhesion kinase
HFpEF	Heart failure with preserved ejection fraction
HFrfEF	Heart failure with reduced ejection fraction
HSC	Hepatic stellate cells

ILK	Integrin-linked kinase
LLC	Large Latent Complex
LAP	Latency Associated Peptide
LTBP	Latent TGF β Binding Proteins
LV	Left Ventricle
LFA-1	Lymphocyte function-associated antigen
Mac-1	Macrophage-1 antigen
MMP	Metalloproteinase
MI	Myocardial infarction
OPN	Osteopontin
Pax	Paxillin
PECAM-1	Platelet Endothelial Cell Adhesion Molecule
PDGF	Platelet-derived growth factor
PO	Pressure overload
Pyk2	Proline-rich tyrosine kinase-2
SIV	Simian immunodeficiency virus
SLC	Small latent complex
STZ	Streptozotocin
VSMC	Vascular smooth muscle cells
Tln	Talin
TSP-1	Thrombospondin-1
TGFβ1	Transforming growth factor beta 1
Vcl	Vinculin

References

1. Miner EC, Miller WL. A look between the cardiomyocytes: the extracellular matrix in heart failure. *Mayo Clin Proc.* 2006; 81:71–76. [PubMed: 16438481]
2. Nanthakumar CB, Hatley RJ, Lemma S, Gaudie J, Marshall RP, Macdonald SJ. Dissecting fibrosis: therapeutic insights from the small-molecule toolbox. *Nat Rev Drug Discov.* 2015
3. Weber KT, Sun Y, Bhattacharya SK, Ahokas RA, Gerling IC. Myofibroblast-mediated mechanisms of pathological remodelling of the heart. *Nat Rev Cardiol.* 2013; 10:15–26. [PubMed: 23207731]
4. Ambale-Venkatesh B, Lima JA. Cardiac MRI: a central prognostic tool in myocardial fibrosis. *Nat Rev Cardiol.* 2015; 12:18–29. [PubMed: 25348690]
5. Galli SJ, Borregaard N, Wynn TA. Phenotypic and functional plasticity of cells of innate immunity: macrophages, mast cells and neutrophils. *Nat Immunol.* 2011; 12:1035–1044. [PubMed: 22012443]
6. Henderson NC, Iredale JP. Liver fibrosis: cellular mechanisms of progression and resolution. *Clin Sci (Lond).* 2007; 112:265–280. [PubMed: 17261089]

7. Varga J, Abraham D. Systemic sclerosis: a prototypic multisystem fibrotic disorder. *J Clin Invest*. 2007; 117:557–567. [PubMed: 17332883]
8. van den Borne SW, Diez J, Blankesteyn WM, Verjans J, Hofstra L, Narula J. Myocardial remodeling after infarction: the role of myofibroblasts. *Nat Rev Cardiol*. 2010; 7:30–37. [PubMed: 19949426]
9. Derynck R, Muthusamy BP, Saeteurn KY. Signaling pathway cooperation in TGF-beta-induced epithelial-mesenchymal transition. *Curr Opin Cell Biol*. 2014; 31:56–66. [PubMed: 25240174]
10. Barriere G, Fici P, Gallerani G, Fabbri F, Rigaud M. Epithelial Mesenchymal Transition: a double-edged sword. *Clin Transl Med*. 2015; 4:14. [PubMed: 25932287]
11. Iwamoto DV, Calderwood DA. Regulation of integrin-mediated adhesions. *Curr Opin Cell Biol*. 2015; 36:41–47. [PubMed: 26189062]
12. Israeli-Rosenberg S, Manso AM, Okada H, Ross RS. Integrins and integrin-associated proteins in the cardiac myocyte. *Circ Res*. 2014; 114:572–586. [PubMed: 24481847]
13. Hynes RO. Integrins: bidirectional, allosteric signaling machines. *Cell*. 2002; 110:673–687. [PubMed: 12297042]
14. Agarwal SK. Integrins and cadherins as therapeutic targets in fibrosis. *Front Pharmacol*. 2014; 5:131. [PubMed: 24917820]
15. Frangogiannis NG. Matricellular proteins in cardiac adaptation and disease. *Physiol Rev*. 2012; 92:635–688. [PubMed: 22535894]
16. Humphries JD, Byron A, Humphries MJ. Integrin ligands at a glance. *J Cell Sci*. 2006; 119:3901–3903. [PubMed: 16988024]
17. Nermut MV, Green NM, Eason P, Yamada SS, Yamada KM. Electron microscopy and structural model of human fibronectin receptor. *The EMBO journal*. 1988; 7:4093–4099. [PubMed: 2977331]
18. Wang N, Tytell JD, Ingber DE. Mechanotransduction at a distance: mechanically coupling the extracellular matrix with the nucleus. *Nat Rev Mol Cell Biol*. 2009; 10:75–82. [PubMed: 19197334]
19. Schwartz MA, Ginsberg MH. Networks and crosstalk: integrin signalling spreads. *Nat Cell Biol*. 2002; 4:E65–E68. [PubMed: 11944032]
20. Clark EA, Brugge JS. Integrins and signal transduction pathways: the road taken. *Science*. 1995; 268:233–239. [PubMed: 7716514]
21. Suryakumar G, Kasiganesan H, Balasubramanian S, Kuppuswamy D. Lack of beta3 integrin signaling contributes to calpain-mediated myocardial cell loss in pressure-overloaded myocardium. *J Cardiovasc Pharmacol*. 2010; 55:567–573. [PubMed: 20224428]
22. Johnston RK, Balasubramanian S, Kasiganesan H, Baicu CF, Zile MR, Kuppuswamy D. Beta3 integrin-mediated ubiquitination activates survival signaling during myocardial hypertrophy. *FASEB J*. 2009; 23:2759–2771. [PubMed: 19364763]
23. Le Gat L, Bonnel S, Gogat K, Brizard M, Van Den Berghe L, Kobetz A, et al. Prominent beta-5 gene expression in the cardiovascular system and in the cartilaginous primordia of the skeleton during mouse development. *Cell Commun Adhes*. 2001; 8:99–112. [PubMed: 11936189]
24. Huhtala P, Humphries MJ, McCarthy JB, Tremble PM, Werb Z, Damsky CH. Cooperative signaling by alpha 5 beta 1 and alpha 4 beta 1 integrins regulates metalloproteinase gene expression in fibroblasts adhering to fibronectin. *J Cell Biol*. 1995; 129:867–879. [PubMed: 7537277]
25. Brooks PC, Stromblad S, Sanders LC, von Schalscha TL, Aimes RT, Stetler-Stevenson WG, et al. Localization of matrix metalloproteinase MMP-2 to the surface of invasive cells by interaction with integrin alpha v beta 3. *Cell*. 1996; 85:683–693. [PubMed: 8646777]
26. Menon B, Singh M, Ross RS, Johnson JN, Singh K. beta-Adrenergic receptor-stimulated apoptosis in adult cardiac myocytes involves MMP-2-mediated disruption of beta1 integrin signaling and mitochondrial pathway. *Am J Physiol Cell Physiol*. 2006; 290:C254–C261. [PubMed: 16148033]
27. Vaisar T, Kassim SY, Gomez IG, Green PS, Hargarten S, Gough PJ, et al. MMP-9 sheds the beta2 integrin subunit (CD18) from macrophages. *Mol Cell Proteomics*. 2009; 8:1044–1060. [PubMed: 19116209]

28. Kryczka J, Stasiak M, Dziki L, Mik M, Dziki A, Cierniewski C. Matrix metalloproteinase-2 cleavage of the beta1 integrin ectodomain facilitates colon cancer cell motility. *J Biol Chem.* 2012; 287:36556–36566. [PubMed: 22898815]
29. Critchley DR. Biochemical and structural properties of the integrin-associated cytoskeletal protein talin. *Annu Rev Biophys.* 2009; 38:235–254. [PubMed: 19416068]
30. Wang L, Tassioulas I, Park-Min KH, Reid AC, Gil-Henn H, Schlessinger J, et al. 'Tuning' of type I interferon-induced Jak-STAT1 signaling by calcium-dependent kinases in macrophages. *Nat Immunol.* 2008; 9:186–193. [PubMed: 18084294]
31. Beckerle MC, Burridge K, DeMartino GN, Croall DE. Colocalization of calcium-dependent protease II and one of its substrates at sites of cell adhesion. *Cell.* 1987; 51:569–577. [PubMed: 2824061]
32. Zhang Y, Chen K, Tu Y, Velyvis A, Yang Y, Qin J, et al. Assembly of the PINCH-ILK-CH-ILKBP complex precedes and is essential for localization of each component to cell-matrix adhesion sites. *J Cell Sci.* 2002; 115:4777–4786. [PubMed: 12432066]
33. Mitra SK, Hanson DA, Schlaepfer DD. Focal adhesion kinase: in command and control of cell motility. *Nat Rev Mol Cell Biol.* 2005; 6:56–68. [PubMed: 15688067]
34. Nolan K, Lacoste J, Parsons JT. Regulated expression of focal adhesion kinase-related nonkinase, the autonomously expressed C-terminal domain of focal adhesion kinase. *Mol Cell Biol.* 1999; 19:6120–6129. [PubMed: 10454559]
35. Torsoni AS, Constancio SS, Nadruz W Jr, Hanks SK, Franchini KG. Focal adhesion kinase is activated and mediates the early hypertrophic response to stretch in cardiac myocytes. *Circ Res.* 2003; 93:140–147. [PubMed: 12805241]
36. Babbitt CJ, Shai SY, Harpf AE, Pham CG, Ross RS. Modulation of integrins and integrin signaling molecules in the pressure-loaded murine ventricle. *Histochem Cell Biol.* 2002; 118:431–439. [PubMed: 12483308]
37. Holden EA Jr. Generation of different numbers of button presses under supplementary feedback conditions by educable retardates. *Percept Mot Skills.* 1975; 41:185–186. [PubMed: 1178403]
38. Lipinski CA, Loftus JC. Targeting Pyk2 for therapeutic intervention. *Expert Opin Ther Targets.* 2010; 14:95–108. [PubMed: 20001213]
39. Larjava H, Plow EF, Wu C. Kindlins: essential regulators of integrin signalling and cell-matrix adhesion. *EMBO Rep.* 2008; 9:1203–1208. [PubMed: 18997731]
40. Askar SF, Bingen BO, Schalij MJ, Swildens J, Atsma DE, Schutte CI, et al. Similar arrhythmicity in hypertrophic and fibrotic cardiac cultures caused by distinct substrate-specific mechanisms. *Cardiovasc Res.* 2013; 97:171–181. [PubMed: 22977008]
41. Frisch SM, Ruoslahti E. Integrins and anoikis. *Curr Opin Cell Biol.* 1997; 9:701–706. [PubMed: 9330874]
42. Taddei ML, Giannoni E, Fiaschi T, Chiarugi P. Anoikis: an emerging hallmark in health and diseases. *J Pathol.* 2012; 226:380–393. [PubMed: 21953325]
43. Michel JB. Anoikis in the cardiovascular system: known and unknown extracellular mediators. *Arterioscler Thromb Vasc Biol.* 2003; 23:2146–2154. [PubMed: 14551156]
44. Ding B, Price RL, Goldsmith EC, Borg TK, Yan X, Douglas PS, et al. Left ventricular hypertrophy in ascending aortic stenosis mice: anoikis and the progression to early failure. *Circulation.* 2000; 101:2854–2862. [PubMed: 10859293]
45. Holmes JW, Clarke SA, Richardson WJ. Modifying the Mechanics of Healing Infarcts: Is Better the Enemy of Good? *Journal of Molecular and Cellular Cardiology.* 2016 (in press, JMC9569).
46. Stuart SDF, De Jesus NM, Lindsey ML, Ripplinger CM. The crossroads of inflammation, fibrosis, and arrhythmia following myocardial infarction. *J Mol Cell Cardiol.* 2016 (in press, JMC9595).
47. Shinde AV, Frangogiannis NG. Fibroblasts in myocardial infarction: a role in inflammation and repair. *J Mol Cell Cardiol.* 2014; 70:74–82. [PubMed: 24321195]
48. Dobaczewski M, Gonzalez-Quesada C, Frangogiannis NG. The extracellular matrix as a modulator of the inflammatory and reparative response following myocardial infarction. *J Mol Cell Cardiol.* 2010; 48:504–511. [PubMed: 19631653]

49. MacLeod JN, Burton-Wurster N, Gu DN, Lust G. Fibronectin mRNA splice variant in articular cartilage lacks bases encoding the V, III-15, and I-10 protein segments. *J Biol Chem.* 1996; 271:18954–18960. [PubMed: 8702559]
50. Cleutjens JP, Blankesteijn WM, Daemen MJ, Smits JF. The infarcted myocardium: simply dead tissue, or a lively target for therapeutic interventions. *Cardiovasc Res.* 1999; 44:232–241. [PubMed: 10690298]
51. Beltrami CA, Finato N, Rocco M, Feruglio GA, Puricelli C, Cigola E, et al. Structural basis of end-stage failure in ischemic cardiomyopathy in humans. *Circulation.* 1994; 89:151–163. [PubMed: 8281642]
52. Nawata J, Ohno I, Isoyama S, Suzuki J, Miura S, Ikeda J, et al. Differential expression of alpha 1, alpha 3 and alpha 5 integrin subunits in acute and chronic stages of myocardial infarction in rats. *Cardiovasc Res.* 1999; 43:371–381. [PubMed: 10536667]
53. Aljamal-Naylor R, Wilson L, McIntyre S, Rossi F, Harrison B, Marsden M, et al. Allosteric modulation of beta1 integrin function induces lung tissue repair. *Adv Pharmacol Sci.* 2012; 2012:768720. [PubMed: 22505883]
54. Huang XZ, Wu JF, Cass D, Erle DJ, Corry D, Young SG, et al. Inactivation of the integrin beta 6 subunit gene reveals a role of epithelial integrins in regulating inflammation in the lung and skin. *J Cell Biol.* 1996; 133:921–928. [PubMed: 8666675]
55. Hertle MD, Kubler MD, Leigh IM, Watt FM. Aberrant integrin expression during epidermal wound healing and in psoriatic epidermis. *J Clin Invest.* 1992; 89:1892–1901. [PubMed: 1601996]
56. Sun M, Opavsky MA, Stewart DJ, Rabinovitch M, Dawood F, Wen WH, et al. Temporal response and localization of integrins beta1 and beta3 in the heart after myocardial infarction: regulation by cytokines. *Circulation.* 2003; 107:1046–1052. [PubMed: 12600920]
57. Krishnamurthy P, Subramanian V, Singh M, Singh K. Deficiency of beta1 integrins results in increased myocardial dysfunction after myocardial infarction. *Heart.* 2006; 92:1309–1315. [PubMed: 16547211]
58. Sofia RR, Serra AJ, Silva JA Jr, Antonio EL, Manchini MT, Oliveira FA, et al. Gender-based differences in cardiac remodeling and ILK expression after myocardial infarction. *Arq Bras Cardiol.* 2014; 103:124–130. [PubMed: 25098374]
59. Ding L, Dong L, Chen X, Zhang L, Xu X, Ferro A, et al. Increased expression of integrin-linked kinase attenuates left ventricular remodeling and improves cardiac function after myocardial infarction. *Circulation.* 2009; 120:764–773. [PubMed: 19687354]
60. Hannigan GE, Coles JG, Dedhar S. Integrin-linked kinase at the heart of cardiac contractility, repair, and disease. *Circ Res.* 2007; 100:1408–1414. [PubMed: 17525380]
61. Mao Q, Lin C, Gao J, Liang X, Gao W, Shen L, et al. Mesenchymal stem cells overexpressing integrin-linked kinase attenuate left ventricular remodeling and improve cardiac function after myocardial infarction. *Mol Cell Biochem.* 2014; 397:203–214. [PubMed: 25134935]
62. Nahrendorf M, Swirski FK. Innate immune cells in ischaemic heart disease: does myocardial infarction beget myocardial infarction? *Eur Heart J.* 2015
63. Heidt T, Courties G, Dutta P, Sager HB, Sebas M, Iwamoto Y, et al. Differential contribution of monocytes to heart macrophages in steady-state and after myocardial infarction. *Circ Res.* 2014; 115:284–295. [PubMed: 24786973]
64. Epelman S, Mann DL. Communication in the heart: the role of the innate immune system in coordinating cellular responses to ischemic injury. *J Cardiovasc Transl Res.* 2012; 5:827–836. [PubMed: 23054658]
65. Fang L, Moore XL, Dart AM, Wang LM. Systemic inflammatory response following acute myocardial infarction. *J Geriatr Cardiol.* 2015; 12:305–312. [PubMed: 26089856]
66. Latet SC, Hoymans VY, Van Herck PL, Vrints CJ. The cellular immune system in the post-myocardial infarction repair process. *Int J Cardiol.* 2015; 179:240–247. [PubMed: 25464457]
67. Christia P, Frangogiannis NG. Targeting inflammatory pathways in myocardial infarction. *Eur J Clin Invest.* 2013; 43:986–995. [PubMed: 23772948]
68. Chavakis T. Leucocyte recruitment in inflammation and novel endogenous negative regulators thereof. *Eur J Clin Invest.* 2012; 42:686–691. [PubMed: 22577952]

69. Hyun YM, Chung HL, McGrath JL, Waugh RE, Kim M. Activated integrin VLA-4 localizes to the lamellipodia and mediates T cell migration on VCAM-1. *J Immunol.* 2009; 183:359–369. [PubMed: 19542447]
70. Moser M, Legate KR, Zent R, Fassler R. The tail of integrins, talin, and kindlins. *Science.* 2009; 324:895–899. [PubMed: 19443776]
71. Chan JR, Hyduk SJ, Cybulsky MI. Chemoattractants induce a rapid and transient upregulation of monocyte alpha4 integrin affinity for vascular cell adhesion molecule 1 which mediates arrest: an early step in the process of emigration. *J Exp Med.* 2001; 193:1149–1158. [PubMed: 11369786]
72. Wu H, Gower RM, Wang H, Perrard XY, Ma R, Bullard DC, et al. Functional role of CD11c+ monocytes in atherogenesis associated with hypercholesterolemia. *Circulation.* 2009; 119:2708–2717. [PubMed: 19433759]
73. O'Doherty U, Peng M, Gezelter S, Swiggard WJ, Betjes M, Bhardwaj N, et al. Human blood contains two subsets of dendritic cells, one immunologically mature and the other immature. *Immunology.* 1994; 82:487–493. [PubMed: 7525461]
74. Yonekawa K, Harlan JM. Targeting leukocyte integrins in human diseases. *J Leukoc Biol.* 2005; 77:129–140. [PubMed: 15548573]
75. Van der Vieren M, Le Trong H, Wood CL, Moore PF, St John T, Staunton DE, et al. A novel leukointegrin, alpha d beta 2, binds preferentially to ICAM-3. *Immunity.* 1995; 3:683–690. [PubMed: 8777714]
76. Simpson PJ, Todd RF 3rd, Fantone JC, Mickelson JK, Griffin JD, Lucchesi BR. Reduction of experimental canine myocardial reperfusion injury by a monoclonal antibody (anti-Mo1, anti-CD11b) that inhibits leukocyte adhesion. *J Clin Invest.* 1988; 81:624–629. [PubMed: 3339135]
77. Palazzo AJ, Jones SP, Girod WG, Anderson DC, Granger DN, Lefer DJ. Myocardial ischemia-reperfusion injury in CD18- and ICAM-1-deficient mice. *Am J Physiol.* 1998; 275:H2300–H2307. [PubMed: 9843832]
78. Investigators AA, Armstrong PW, Granger CB, Adams PX, Hamm C, Holmes D Jr, et al. Pexelizumab for acute ST-elevation myocardial infarction in patients undergoing primary percutaneous coronary intervention: a randomized controlled trial. *JAMA.* 2007; 297:43–51. [PubMed: 17200474]
79. Faxon DP, Gibbons RJ, Chronos NA, Gurbel PA, Sheehan F, Investigators H-M. The effect of blockade of the CD11/CD18 integrin receptor on infarct size in patients with acute myocardial infarction treated with direct angioplasty: the results of the HALT-MI study. *J Am Coll Cardiol.* 2002; 40:1199–1204. [PubMed: 12383565]
80. Baran KW, Nguyen M, McKendall GR, Lambrew CT, Dykstra G, Palmeri ST, et al. Double-blind, randomized trial of an anti-CD18 antibody in conjunction with recombinant tissue plasminogen activator for acute myocardial infarction: limitation of myocardial infarction following thrombolysis in acute myocardial infarction (LIMIT AMI) study. *Circulation.* 2001; 104:2778–2783. [PubMed: 11733394]
81. Rusnak JM, Kopecky SL, Clements IP, Gibbons RJ, Holland AE, Peterman HS, et al. An anti-CD11/CD18 monoclonal antibody in patients with acute myocardial infarction having percutaneous transluminal coronary angioplasty (the FESTIVAL study). *Am J Cardiol.* 2001; 88:482–487. [PubMed: 11524054]
82. Walker JA, Beck GA, Campbell JH, Miller AD, Burdo TH, Williams KC. Anti-alpha4 Integrin Antibody Blocks Monocyte/Macrophage Traffic to the Heart and Decreases Cardiac Pathology in a SIV Infection Model of AIDS. *J Am Heart Assoc.* 2015; 4
83. Berk BC, Fujiwara K, Lehoux S. ECM remodeling in hypertensive heart disease. *J Clin Invest.* 2007; 117:568–575. [PubMed: 17332884]
84. Manso AM, Kang SM, Plotnikov SV, Thievessen I, Oh J, Beggs HE, et al. Cardiac fibroblasts require focal adhesion kinase for normal proliferation and migration. *Am J Physiol Heart Circ Physiol.* 2009; 296:H627–H638. [PubMed: 19136609]
85. Burgess ML, Terracio L, Hirozane T, Borg TK. Differential integrin expression by cardiac fibroblasts from hypertensive and exercise-trained rat hearts. *Cardiovasc Pathol.* 2002; 11:78–87. [PubMed: 11934598]

86. Dalla Costa AP, Clemente CF, Carvalho HF, Carvalheira JB, Nadruz W Jr, Franchini KG. FAK mediates the activation of cardiac fibroblasts induced by mechanical stress through regulation of the mTOR complex. *Cardiovasc Res.* 2010; 86:421–431. [PubMed: 20038548]
87. Clemente CF, Tornatore TF, Theizen TH, Deckmann AC, Pereira TC, Lopes-Cendes I, et al. Targeting focal adhesion kinase with small interfering RNA prevents and reverses load-induced cardiac hypertrophy in mice. *Circ Res.* 2007; 101:1339–1348. [PubMed: 17947798]
88. Balasubramanian S, Quinones L, Kasiganesan H, Zhang Y, Pleasant DL, Sundararaj KP, et al. beta3 integrin in cardiac fibroblast is critical for extracellular matrix accumulation during pressure overload hypertrophy in mouse. *PLoS One.* 2012; 7:e45076. [PubMed: 22984613]
89. Rich MW. Epidemiology, pathophysiology, and etiology of congestive heart failure in older adults. *Journal of the American Geriatrics Society.* 1997; 45:968–974. [PubMed: 9256850]
90. Edgley AJ, Krum H, Kelly DJ. Targeting fibrosis for the treatment of heart failure: a role for transforming growth factor-beta. *Cardiovasc Ther.* 2012; 30:e30–e40. [PubMed: 21883991]
91. Horn MA, Trafford AW. Aging and the cardiac extracellular matrix: novel mediators of fibrotic remodeling. *J Mol Cell Cardiol.* 2016 (in press, JMC9600).
92. Burgess ML, McCrea JC, Hedrick HL. Age-associated changes in cardiac matrix and integrins. *Mech Ageing Dev.* 2001; 122:1739–1756. [PubMed: 11557277]
93. Mamuya W, Chobanian A, Brecher P. Age-related changes in fibronectin expression in spontaneously hypertensive, Wistar-Kyoto, and Wistar rat hearts. *Circ Res.* 1992; 71:1341–1350. [PubMed: 1423932]
94. Wilson CG, Stone JW, Fowlkes V, Morales MO, Murphy CJ, Baxter SC, et al. Age-dependent expression of collagen receptors and deformation of type I collagen substrates by rat cardiac fibroblasts. *Microsc Microanal.* 2011; 17:555–562. [PubMed: 21740617]
95. Communal C, Huq F, Lebeche D, Mestel C, Gwathmey JK, Hajjar RJ. Decreased efficiency of adenovirus-mediated gene transfer in aging cardiomyocytes. *Circulation.* 2003; 107:1170–1175. [PubMed: 12615797]
96. Shai SY, Harpf AE, Babbitt CJ, Jordan MC, Fishbein MC, Chen J, et al. Cardiac myocytespecific excision of the beta1 integrin gene results in myocardial fibrosis and cardiac failure. *Circ Res.* 2002; 90:458–464. [PubMed: 11884376]
97. Russo I, Frangogiannis NG. Diabetes-associated cardiac fibrosis: cellular effectors, molecular mechanisms and therapeutic opportunities. *J Mol Cell Cardiol.* 2016 (in press, JMC9592).
98. Huynh K, Bernardo BC, McMullen JR, Ritchie RH. Diabetic cardiomyopathy: mechanisms and new treatment strategies targeting antioxidant signaling pathways. *Pharmacol Ther.* 2014; 142:375–415. [PubMed: 24462787]
99. Goyal BR, Mehta AA. Diabetic cardiomyopathy: pathophysiological mechanisms and cardiac dysfunction. *Hum Exp Toxicol.* 2013; 32:571–590. [PubMed: 23174745]
100. Mandavia CH, Aroor AR, Demarco VG, Sowers JR. Molecular and metabolic mechanisms of cardiac dysfunction in diabetes. *Life Sci.* 2013; 92:601–608. [PubMed: 23147391]
101. Yamazaki KG, Gonzalez E, Zambon AC. Crosstalk between the renin-angiotensin system and the advance glycation end product axis in the heart: role of the cardiac fibroblast. *J Cardiovasc Transl Res.* 2012; 5:805–813. [PubMed: 23054657]
102. Kelly DJ, Zhang Y, Connelly K, Cox AJ, Martin J, Krum H, et al. Tranilast attenuates diastolic dysfunction and structural injury in experimental diabetic cardiomyopathy. *Am J Physiol Heart Circ Physiol.* 2007; 293:H2860–H2869. [PubMed: 17720766]
103. Tang M, Zhang W, Lin H, Jiang H, Dai H, Zhang Y. High glucose promotes the production of collagen types I and III by cardiac fibroblasts through a pathway dependent on extracellular-signal-regulated kinase 1/2. *Mol Cell Biochem.* 2007; 301:109–114. [PubMed: 17206378]
104. Martin J, Kelly DJ, Mifsud SA, Zhang Y, Cox AJ, See F, et al. Tranilast attenuates cardiac matrix deposition in experimental diabetes: role of transforming growth factor-beta. *Cardiovasc Res.* 2005; 65:694–701. [PubMed: 15664396]
105. Talior-Volodarsky I, Connelly KA, Arora PD, Gullberg D, McCulloch CA. alpha11 integrin stimulates myofibroblast differentiation in diabetic cardiomyopathy. *Cardiovasc Res.* 2012; 96:265–275. [PubMed: 22869616]

106. Leask A. Potential therapeutic targets for cardiac fibrosis: TGFbeta, angiotensin, endothelin, CCN2, and PDGF, partners in fibroblast activation. *Circ Res.* 2010; 106:1675–1680. [PubMed: 20538689]
107. Sun Y, Weber KT. Angiotensin II receptor binding following myocardial infarction in the rat. *Cardiovasc Res.* 1994; 28:1623–1628. [PubMed: 7842454]
108. Campbell SE, Katwa LC. Angiotensin II stimulated expression of transforming growth factor-beta1 in cardiac fibroblasts and myofibroblasts. *J Mol Cell Cardiol.* 1997; 29:1947–1858. [PubMed: 9236148]
109. Schultz Jel J, Witt SA, Glascock BJ, Nieman ML, Reiser PJ, Nix SL, et al. TGF-beta1 mediates the hypertrophic cardiomyocyte growth induced by angiotensin II. *J Clin Invest.* 2002; 109:787–796. [PubMed: 11901187]
110. De Mello WC, Specht P. Chronic blockade of angiotensin II AT1-receptors increased cell-to-cell communication, reduced fibrosis and improved impulse propagation in the failing heart. *J Renin Angiotensin Aldosterone Syst.* 2006; 7:201–205. [PubMed: 17318788]
111. Shibasaki Y, Nishiue T, Masaki H, Tamura K, Matsumoto N, Mori Y, et al. Impact of the angiotensin II receptor antagonist, losartan, on myocardial fibrosis in patients with end-stage renal disease: assessment by ultrasonic integrated backscatter and biochemical markers. *Hypertens Res.* 2005; 28:787–795. [PubMed: 16471172]
112. Kassem L, Deygas M, Fattet L, Lopez J, Goulvent T, Lavergne E, et al. TIF1gamma interferes with TGFbeta1/SMAD4 signaling to promote poor outcome in operable breast cancer patients. *BMC Cancer.* 2015; 15:453. [PubMed: 26040677]
113. Graf K, Neuss M, Stawowy P, Hsueh WA, Fleck E, Law RE. Angiotensin II and alpha(v)beta(3) integrin expression in rat neonatal cardiac fibroblasts. *Hypertension.* 2000; 35:978–984. [PubMed: 10775572]
114. Kawano H, Cody RJ, Graf K, Goetze S, Kawano Y, Schnee J, et al. Angiotensin II enhances integrin and alpha-actinin expression in adult rat cardiac fibroblasts. *Hypertension.* 2000; 35:273–279. [PubMed: 10642310]
115. Bouzeghrane F, Mercure C, Reudelhuber TL, Thibault G. Alpha8beta1 integrin is upregulated in myofibroblasts of fibrotic and scarring myocardium. *J Mol Cell Cardiol.* 2004; 36:343–353. [PubMed: 15010273]
116. Carver W, Molano I, Reaves TA, Borg TK, Terracio L. Role of the alpha 1 beta 1 integrin complex in collagen gel contraction in vitro by fibroblasts. *J Cell Physiol.* 1995; 165:425–437. [PubMed: 7593221]
117. Hartner A, Cordasic N, Rascher W, Hilgers KF. Deletion of the alpha8 integrin gene does not protect mice from myocardial fibrosis in DOCA hypertension. *Am J Hypertens.* 2009; 22:92–99. [PubMed: 19008864]
118. Verma SK, Lal H, Golden HB, Gerilechaogetu F, Smith M, Guleria RS, et al. Rac1 and RhoA differentially regulate angiotensinogen gene expression in stretched cardiac fibroblasts. *Cardiovasc Res.* 2011; 90:88–96. [PubMed: 21131638]
119. Mehta PK, Griendling KK. Angiotensin II cell signaling: physiological and pathological effects in the cardiovascular system. *Am J Physiol Cell Physiol.* 2007; 292:C82–C97. [PubMed: 16870827]
120. Shimojo N, Hashizume R, Kanayama K, Hara M, Suzuki Y, Nishioka T, et al. Tenascin-C May Accelerate Cardiac Fibrosis by Activating Macrophages via the Integrin alphaVbeta3/Nuclear Factor-kappaB/Interleukin-6 Axis. *Hypertension.* 2015; 66:757–766. [PubMed: 26238448]
121. Midwood KS, Hussenet T, Langlois B, Orend G. Advances in tenascin-C biology. *Cell Mol Life Sci.* 2011; 68:3175–3199. [PubMed: 21818551]
122. Friedrichs K, Adam M, Remane L, Mollenhauer M, Rudolph V, Rudolph TK, et al. Induction of atrial fibrillation by neutrophils critically depends on CD11b/CD18 integrins. *PLoS One.* 2014; 9:e89307. [PubMed: 24558493]
123. Xie Z, Singh M, Singh K. Osteopontin modulates myocardial hypertrophy in response to chronic pressure overload in mice. *Hypertension.* 2004; 44:826–831. [PubMed: 15534078]
124. Wang KX, Denhardt DT. Osteopontin: role in immune regulation and stress responses. *Cytokine Growth Factor Rev.* 2008; 19:333–345. [PubMed: 18952487]

125. Kock LM, Schulz RM, van Donkelaar CC, Thummler CB, Bader A, Ito K. RGD-dependent integrins are mechanotransducers in dynamically compressed tissue-engineered cartilage constructs. *J Biomech.* 2009; 42:2177–2182. [PubMed: 19656515]
126. Yokosaki Y, Tanaka K, Higashikawa F, Yamashita K, Eboshida A. Distinct structural requirements for binding of the integrins α v β 6, α v β 3, α v β 5, α 5 β 1 and α 9 β 1 to osteopontin. *Matrix Biol.* 2005; 24:418–427. [PubMed: 16005200]
127. Matsui Y, Jia N, Okamoto H, Kon S, Onozuka H, Akino M, et al. Role of osteopontin in cardiac fibrosis and remodeling in angiotensin II-induced cardiac hypertrophy. *Hypertension.* 2004; 43:1195–1201. [PubMed: 15123578]
128. Collins AR, Schnee J, Wang W, Kim S, Fishbein MC, Bruemmer D, et al. Osteopontin modulates angiotensin II-induced fibrosis in the intact murine heart. *J Am Coll Cardiol.* 2004; 43:1698–1705. [PubMed: 15120833]
129. Singh K, Sirokman G, Communal C, Robinson KG, Conrad CH, Brooks WW, et al. Myocardial osteopontin expression coincides with the development of heart failure. *Hypertension.* 1999; 33:663–670. [PubMed: 10024324]
130. Ashizawa N, Graf K, Do YS, Nunohiro T, Giachelli CM, Meehan WP, et al. Osteopontin is produced by rat cardiac fibroblasts and mediates A(II)-induced DNA synthesis and collagen gel contraction. *J Clin Invest.* 1996; 98:2218–2227. [PubMed: 8941637]
131. Leask A, Abraham DJ. TGF- β signaling and the fibrotic response. *FASEB J.* 2004; 18:816–827. [PubMed: 15117886]
132. Sun Y, Zhang JQ, Zhang J, Ramires FJ. Angiotensin II, transforming growth factor- β 1 and repair in the infarcted heart. *J Mol Cell Cardiol.* 1998; 30:1559–1569. [PubMed: 9737942]
133. Ikeuchi M, Tsutsui H, Shiomi T, Matsusaka H, Matsushima S, Wen J, et al. Inhibition of TGF- β signaling exacerbates early cardiac dysfunction but prevents late remodeling after infarction. *Cardiovasc Res.* 2004; 64:526–535. [PubMed: 15537506]
134. Koitabashi N, Danner T, Zaiman AL, Pinto YM, Rowell J, Mankowski J, et al. Pivotal role of cardiomyocyte TGF- β signaling in the murine pathological response to sustained pressure overload. *J Clin Invest.* 2011; 121:2301–2312. [PubMed: 21537080]
135. Brooks WW, Conrad CH. Myocardial fibrosis in transforming growth factor β 1 heterozygous mice. *J Mol Cell Cardiol.* 2000; 32:187–195. [PubMed: 10722796]
136. Munger JS, Harpel JG, Giancotti FG, Rifkin DB. Interactions between growth factors and integrins: latent forms of transforming growth factor- β are ligands for the integrin α v β 1. *Mol Biol Cell.* 1998; 9:2627–2638. [PubMed: 9725916]
137. Gleizes PE, Munger JS, Nunes I, Harpel JG, Mazzieri R, Noguera I, et al. TGF- β latency: biological significance and mechanisms of activation. *Stem Cells.* 1997; 15:190–197. [PubMed: 9170210]
138. Wipff PJ, Hinz B. Integrins and the activation of latent transforming growth factor β 1 - an intimate relationship. *Eur J Cell Biol.* 2008; 87:601–615. [PubMed: 18342983]
139. Ludbrook SB, Barry ST, Delves CJ, Horgan CM. The integrin α v β 3 is a receptor for the latency-associated peptides of transforming growth factors β 1 and β 3. *Biochem J.* 2003; 369:311–318. [PubMed: 12358597]
140. Asano Y, Ihn H, Yamane K, Jinnin M, Mimura Y, Tamaki K. Increased expression of integrin α (v) β 3 contributes to the establishment of autocrine TGF- β signaling in scleroderma fibroblasts. *J Immunol.* 2005; 175:7708–7718. [PubMed: 16301681]
141. Mu D, Cambier S, Fjellbirkeland L, Baron JL, Munger JS, Kawakatsu H, et al. The integrin α (v) β 8 mediates epithelial homeostasis through MT1-MMP-dependent activation of TGF- β 1. *J Cell Biol.* 2002; 157:493–507. [PubMed: 11970960]
142. Munger JS, Huang X, Kawakatsu H, Griffiths MJ, Dalton SL, Wu J, et al. The integrin α v β 6 binds and activates latent TGF β 1: a mechanism for regulating pulmonary inflammation and fibrosis. *Cell.* 1999; 96:319–328. [PubMed: 10025398]
143. Wipff PJ, Rifkin DB, Meister JJ, Hinz B. Myofibroblast contraction activates latent TGF β 1 from the extracellular matrix. *J Cell Biol.* 2007; 179:1311–1323. [PubMed: 18086923]

144. Asano Y, Ihn H, Yamane K, Jinnin M, Tamaki K. Increased expression of integrin alphavbeta5 induces the myofibroblastic differentiation of dermal fibroblasts. *Am J Pathol.* 2006; 168:499–510. [PubMed: 16436664]
145. Breuss JM, Gillett N, Lu L, Sheppard D, Pytela R. Restricted distribution of integrin beta 6 mRNA in primate epithelial tissues. *J Histochem Cytochem.* 1993; 41:1521–1527. [PubMed: 8245410]
146. Hahm K, Lukashev ME, Luo Y, Yang WJ, Dolinski BM, Weinreb PH, et al. Alphav beta6 integrin regulates renal fibrosis and inflammation in Alport mouse. *Am J Pathol.* 2007; 170:110–125. [PubMed: 17200187]
147. Popov Y, Patsenker E, Stickel F, Zaks J, Bhaskar KR, Niedobitek G, et al. Integrin alphavbeta6 is a marker of the progression of biliary and portal liver fibrosis and a novel target for antifibrotic therapies. *J Hepatol.* 2008; 48:453–464. [PubMed: 18221819]
148. Wang L, Liang Y, Li Z, Cai X, Zhang W, Wu G, et al. Increase in beta1–6 GlcNAc branching caused by N-acetylglucosaminyltransferase V directs integrin beta1 stability in human hepatocellular carcinoma cell line SMMC-7721. *J Cell Biochem.* 2007; 100:230–241. [PubMed: 16924681]
149. Ma LJ, Yang H, Gaspert A, Carlesso G, Barty MM, Davidson JM, et al. Transforming growth factor-beta-dependent and -independent pathways of induction of tubulointerstitial fibrosis in beta6(–/–) mice. *Am J Pathol.* 2003; 163:1261–1273. [PubMed: 14507636]
150. Patsenker E, Popov Y, Stickel F, Jonczyk A, Goodman SL, Schuppan D. Inhibition of integrin alphavbeta6 on cholangiocytes blocks transforming growth factor-beta activation and retards biliary fibrosis progression. *Gastroenterology.* 2008; 135:660–670. [PubMed: 18538673]
151. Annes JP, Rifkin DB, Munger JS. The integrin alphaVbeta6 binds and activates latent TGFbeta3. *FEBS Lett.* 2002; 511:65–68. [PubMed: 11821050]
152. Jenkins RG, Su X, Su G, Scotton CJ, Camerer E, Laurent GJ, et al. Ligation of protease-activated receptor 1 enhances alpha(v)beta6 integrin-dependent TGF-beta activation and promotes acute lung injury. *J Clin Invest.* 2006; 116:1606–1614. [PubMed: 16710477]
153. Shi M, Zhu J, Wang R, Chen X, Mi L, Walz T, et al. Latent TGF-beta structure and activation. *Nature.* 2011; 474:343–349. [PubMed: 21677751]
154. Wei X, Wang X, Xia Y, Tang Y, Li F, Fang W, et al. Kindlin-2 regulates renal tubular cell plasticity by activation of Ras and its downstream signaling. *Am J Physiol Renal Physiol.* 2014; 306:F271–F278. [PubMed: 24226523]
155. Araya J, Cambier S, Markovics JA, Wolters P, Jablons D, Hill A, et al. Squamous metaplasia amplifies pathologic epithelial-mesenchymal interactions in COPD patients. *J Clin Invest.* 2007; 117:3551–3562. [PubMed: 17965775]
156. Kitamura H, Cambier S, Somanath S, Barker T, Minagawa S, Markovics J, et al. Mouse and human lung fibroblasts regulate dendritic cell trafficking, airway inflammation, and fibrosis through integrin alphavbeta8-mediated activation of TGF-beta. *J Clin Invest.* 2011; 121:2863–2875. [PubMed: 21646718]
157. Khan S, Lakhe-Reddy S, McCarty JH, Sorenson CM, Sheibani N, Reichardt LF, et al. Mesangial cell integrin alphavbeta8 provides glomerular endothelial cell cytoprotection by sequestering TGF-beta and regulating PECAM-1. *Am J Pathol.* 2011; 178:609–620. [PubMed: 21281793]
158. Zhou X, Murphy FR, Gehdu N, Zhang J, Iredale JP, Benyon RC. Engagement of alphavbeta3 integrin regulates proliferation and apoptosis of hepatic stellate cells. *J Biol Chem.* 2004; 279:23996–24006. [PubMed: 15044441]
159. Patsenker E, Popov Y, Stickel F, Schneider V, Ledermann M, Sagesser H, et al. Pharmacological inhibition of integrin alphavbeta3 aggravates experimental liver fibrosis and suppresses hepatic angiogenesis. *Hepatology.* 2009; 50:1501–1511. [PubMed: 19725105]
160. Asano Y, Ihn H, Jinnin M, Mimura Y, Tamaki K. Involvement of alphavbeta5 integrin in the establishment of autocrine TGF-beta signaling in dermal fibroblasts derived from localized scleroderma. *J Invest Dermatol.* 2006; 126:1761–1769. [PubMed: 16675963]
161. Asano Y, Ihn H, Yamane K, Jinnin M, Mimura Y, Tamaki K. Involvement of alphavbeta5 integrin-mediated activation of latent transforming growth factor beta1 in autocrine transforming growth

- factor beta signaling in systemic sclerosis fibroblasts. *Arthritis Rheum.* 2005; 52:2897–2905. [PubMed: 16142753]
162. Henderson NC, Arnold TD, Katamura Y, Giacomini MM, Rodriguez JD, McCarty JH, et al. Targeting of alphav integrin identifies a core molecular pathway that regulates fibrosis in several organs. *Nat Med.* 2013; 19:1617–1624. [PubMed: 24216753]
 163. Reed NI, Jo H, Chen C, Tsujino K, Arnold TD, DeGrado WF, et al. The alphavbeta1 integrin plays a critical in vivo role in tissue fibrosis. *Sci Transl Med.* 2015; 7:288ra79.
 164. Lagares D, Busnadiego O, Garcia-Fernandez RA, Kapoor M, Liu S, Carter DE, et al. Inhibition of focal adhesion kinase prevents experimental lung fibrosis and myofibroblast formation. *Arthritis Rheum.* 2012; 64:1653–1664. [PubMed: 22492165]
 165. Leask A. Focal Adhesion Kinase: A Key Mediator of Transforming Growth Factor Beta Signaling in Fibroblasts. *Adv Wound Care (New Rochelle).* 2013; 2:247–249. [PubMed: 24527346]
 166. Thannickal VJ, Lee DY, White ES, Cui Z, Larios JM, Chacon R, et al. Myofibroblast differentiation by transforming growth factor-beta1 is dependent on cell adhesion and integrin signaling via focal adhesion kinase. *J Biol Chem.* 2003; 278:12384–12389. [PubMed: 12531888]
 167. Sarrazy V, Koehler A, Chow ML, Zimina E, Li CX, Kato H, et al. Integrins alphavbeta5 and alphavbeta3 promote latent TGF-beta1 activation by human cardiac fibroblast contraction. *Cardiovasc Res.* 2014; 102:407–417. [PubMed: 24639195]
 168. Lamouille S, Xu J, Derynck R. Molecular mechanisms of epithelial-mesenchymal transition. *Nat Rev Mol Cell Biol.* 2014; 15:178–196. [PubMed: 24556840]
 169. Kim KK, Kugler MC, Wolters PJ, Robillard L, Galvez MG, Brumwell AN, et al. Alveolar epithelial cell mesenchymal transition develops in vivo during pulmonary fibrosis and is regulated by the extracellular matrix. *Proc Natl Acad Sci U S A.* 2006; 103:13180–13185. [PubMed: 16924102]
 170. Li Y, Tan X, Dai C, Stolz DB, Wang D, Liu Y. Inhibition of integrin-linked kinase attenuates renal interstitial fibrosis. *J Am Soc Nephrol.* 2009; 20:1907–1918. [PubMed: 19541809]
 171. Li Y, Dai C, Wu C, Liu Y. PINCH-1 promotes tubular epithelial-to-mesenchymal transition by interacting with integrin-linked kinase. *J Am Soc Nephrol.* 2007; 18:2534–2543. [PubMed: 17656471]
 172. Li Y, Yang J, Dai C, Wu C, Liu Y. Role for integrin-linked kinase in mediating tubular epithelial to mesenchymal transition and renal interstitial fibrogenesis. *J Clin Invest.* 2003; 112:503–516. [PubMed: 12925691]
 173. Zhao YX, Yin HQ, Yu QT, Qiao Y, Dai HY, Zhang MX, et al. ACE2 overexpression ameliorates left ventricular remodeling and dysfunction in a rat model of myocardial infarction. *Hum Gene Ther.* 2010; 21:1545–1554. [PubMed: 20507236]
 174. Rock JR, Barkauskas CE, Cronic MJ, Xue Y, Harris JR, Liang J, et al. Multiple stromal populations contribute to pulmonary fibrosis without evidence for epithelial to mesenchymal transition. *Proc Natl Acad Sci U S A.* 2011; 108:E1475–E1483. [PubMed: 22123957]
 175. Humphreys BD, Lin SL, Kobayashi A, Hudson TE, Nowlin BT, Bonventre JV, et al. Fate tracing reveals the pericyte and not epithelial origin of myofibroblasts in kidney fibrosis. *Am J Pathol.* 2010; 176:85–97. [PubMed: 20008127]
 176. Lovisa S, LeBleu VS, Tampe B, Sugimoto H, Vlodavets K, Carstens JL, et al. Epithelial-to-mesenchymal transition induces cell cycle arrest and parenchymal damage in renal fibrosis. *Nat Med.* 2015; 21:998–1009. [PubMed: 26236991]
 177. Grande MT, Sanchez-Laorden B, Lopez-Blau C, De Frutos CA, Boutet A, Arevalo M, et al. Snail1-induced partial epithelial-to-mesenchymal transition drives renal fibrosis in mice and can be targeted to reverse established disease. *Nat Med.* 2015; 21:989–997. [PubMed: 26236989]
 178. Yoshimatsu Y, Watabe T. Roles of TGF-beta signals in endothelial-mesenchymal transition during cardiac fibrosis. *Int J Inflam.* 2011; 2011:724080. [PubMed: 22187661]
 179. Widyanoto B, Emoto N, Nakayama K, Anggrahini DW, Adiarto S, Iwasa N, et al. Endothelial cell-derived endothelin-1 promotes cardiac fibrosis in diabetic hearts through stimulation of endothelial-to-mesenchymal transition. *Circulation.* 2010; 121:2407–2418. [PubMed: 20497976]

180. Zeisberg EM, Tarnavski O, Zeisberg M, Dorfman AL, McMullen JR, Gustafsson E, et al. Endothelial-to-mesenchymal transition contributes to cardiac fibrosis. *Nat Med.* 2007; 13:952–961. [PubMed: 17660828]
181. Ruiz-Villalba A, Simon AM, Pogontke C, Castillo MI, Abizanda G, Pelacho B, et al. Interacting resident epicardium-derived fibroblasts and recruited bone marrow cells form myocardial infarction scar. *J Am Coll Cardiol.* 2015; 65:2057–2066. [PubMed: 25975467]
182. Moore-Morris T, Guimaraes-Camboa N, Banerjee I, Zambon AC, Kisseleva T, Velayoudon A, et al. Resident fibroblast lineages mediate pressure overload-induced cardiac fibrosis. *J Clin Invest.* 2014; 124:2921–2934. [PubMed: 24937432]
183. Liu F, Mih JD, Shea BS, Kho AT, Sharif AS, Tager AM, et al. Feedback amplification of fibrosis through matrix stiffening and COX-2 suppression. *J Cell Biol.* 2010; 190:693–706. [PubMed: 20733059]
184. Olsen AL, Bloomer SA, Chan EP, Gaca MD, Georges PC, Sackey B, et al. Hepatic stellate cells require a stiff environment for myofibroblastic differentiation. *Am J Physiol Gastrointest Liver Physiol.* 2011; 301:G110–G118. [PubMed: 21527725]
185. Kloxin AM, Benton JA, Anseth KS. In situ elasticity modulation with dynamic substrates to direct cell phenotype. *Biomaterials.* 2010; 31:1–8. [PubMed: 19788947]
186. Galie PA, Khalid N, Carnahan KE, Westfall MV, Stegemann JP. Substrate stiffness affects sarcomere and costamere structure and electrophysiological function of isolated adult cardiomyocytes. *Cardiovasc Pathol.* 2013; 22:219–227. [PubMed: 23266222]
187. Zhao H, Li X, Zhao S, Zeng Y, Zhao L, Ding H, et al. Microengineered in vitro model of cardiac fibrosis through modulating myofibroblast mechanotransduction. *Biofabrication.* 2014; 6:045009. [PubMed: 25378063]
188. Galie PA, Westfall MV, Stegemann JP. Reduced serum content and increased matrix stiffness promote the cardiac myofibroblast transition in 3D collagen matrices. *Cardiovasc Pathol.* 2011; 20:325–333. [PubMed: 21306921]
189. Wells RG. Tissue mechanics and fibrosis. *Biochim Biophys Acta.* 2013; 1832:884–890. [PubMed: 23434892]
190. MacKenna DA, Dolfi F, Vuori K, Ruoslahti E. Extracellular signal-regulated kinase and c-Jun NH2-terminal kinase activation by mechanical stretch is integrin-dependent and matrix-specific in rat cardiac fibroblasts. *J Clin Invest.* 1998; 101:301–310. [PubMed: 9435301]
191. Gershlak JR, Black LD 3rd. Beta 1 integrin binding plays a role in the constant traction force generation in response to varying stiffness for cells grown on mature cardiac extracellular matrix. *Exp Cell Res.* 2015; 330:311–324. [PubMed: 25220424]
192. Arendt E, Ueberham U, Bittner R, Gebhardt R, Ueberham E. Enhanced matrix degradation after withdrawal of TGF-beta1 triggers hepatocytes from apoptosis to proliferation and regeneration. *Cell Prolif.* 2005; 38:287–299. [PubMed: 16202037]
193. Kulkarni AB, Huh CG, Becker D, Geiser A, Lyght M, Flanders KC, et al. Transforming growth factor beta 1 null mutation in mice causes excessive inflammatory response and early death. *Proc Natl Acad Sci U S A.* 1993; 90:770–774. [PubMed: 8421714]
194. Shull MM, Ormsby I, Kier AB, Pawlowski S, Diebold RJ, Yin M, et al. Targeted disruption of the mouse transforming growth factor-beta 1 gene results in multifocal inflammatory disease. *Nature.* 1992; 359:693–699. [PubMed: 1436033]
195. Datta A, Scotton CJ, Chambers RC. Novel therapeutic approaches for pulmonary fibrosis. *Br J Pharmacol.* 2011; 163:141–172. [PubMed: 21265830]
196. Lagares D, Kapoor M. Targeting focal adhesion kinase in fibrotic diseases. *BioDrugs.* 2013; 27:15–23. [PubMed: 23329396]
197. Schultze A, Fiedler W. Therapeutic potential and limitations of new FAK inhibitors in the treatment of cancer. *Expert Opin Investig Drugs.* 2010; 19:777–788.
198. Schultze A, Fiedler W. Clinical importance and potential use of small molecule inhibitors of focal adhesion kinase. *Anticancer Agents Med Chem.* 2011; 11:593–599. [PubMed: 21787277]
199. Wong VW, Rustad KC, Akaishi S, Sorkin M, Glotzbach JP, Januszyk M, et al. Focal adhesion kinase links mechanical force to skin fibrosis via inflammatory signaling. *Nat Med.* 2012; 18:148–152. [PubMed: 22157678]

200. Shimazaki M, Nakamura K, Kii I, Kashima T, Amizuka N, Li M, et al. Periostin is essential for cardiac healing after acute myocardial infarction. *J Exp Med*. 2008; 205:295–303. [PubMed: 18208976]

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Highlights

- Cardiac fibrosis is one of the major components of pathological cardiac remodeling
- Integrins regulate key effectors that drive fibrosis in many organs including heart
- Integrins are potential therapeutic targets to treat cardiac fibrosis

Figure 1. Integrins and Integrin-related proteins

Integrins shown spanning the cell membrane, connect and aggregate a range of adapter and signaling proteins such as integrin linked kinase (ILK), focal adhesion kinase (FAK), paxillin (Pax), vinculin (Vcl), talin (Tln), Kindlin, PINCH, Parvin, actinin and even actin. This allows both bridging of ECM to the intracellular cytoskeleton, and also allows propagation of signals bidirectionally across the cell membrane.

Figure 2. Model of mechanical activation of latent TGFβ1

TGFβ1 is secreted in a large latent complex (**LLC**) that consists of TGFβ1 associated with LAP (Latency Associated Peptide), the short latent complex (**SLC**), and LTBP-1 (Latent TGFβ1 Binding Protein 1). LAP contains the amino acid sequence motif RGD (Arg-Gly-Asp) which serves as a recognition site for several integrins. Actin / myosin-mediated cell contraction force can be transmitted to an RGD binding site in LAP through the αv integrins and induces a putative conformational change that liberates TGFβ1, activating it.

Figure 3. Myofibroblast precursors and integrin subtypes involved in end-effector regulation
Myofibroblasts can be derived from resident fibroblasts and pericytes, as well as from smooth muscle cells (SMC), endothelial cells (EC), epithelial cells and fibrocytes. As one example, in response to angiotensin II (Ang II), transforming growth factor beta-1 (TGFβ1), can increase the conversion of fibroblasts to myofibroblasts. Mechanical stress (perhaps functioning through Ang II) and ED-A fibronectin (ED-A FN) can also increase this transformation. Integrins are thus involved in TGFβ activation, in EMT regulation and in Ang II synthesis. Myofibroblasts are characterized by the presence of α-smooth muscle actin (α-SMA), stress fibers and collagen production. EMT (epithelial - mesenchymal transition); EndMT (endothelial - mesenchymal transition). + denotes factors can positively increase the transformation of fibroblasts to myofibroblasts.