Published in final edited form as:

J Neuroimmunol. 2016 February 15; 291: 59–69. doi:10.1016/j.jneuroim.2015.12.009.

IL-12R β 2 has a protective role in relapsing-remitting experimental autoimmune encephalomyelitis

Chong Xie^{1,2,3}, Bogoljub Ciric^{1,3}, Shuo Yu¹, Guang-Xian Zhang¹, and Abdolmohamad Rostami¹

¹Department of Neurology, Thomas Jefferson University, Philadelphia, PA 19107, USA

²Department of Neurology, Changhai Hospital, Second Military Medical University, Shanghai, 200433, China

Abstract

IL-12R β 2 participates in the receptors of IL-12 and IL-35, two cytokines that are involved in a variety of immune responses. In this study we evaluate the role of IL-12R β 2 in relapsing-remitting experimental autoimmune encephalomyelitis (RR-EAE). We found that the IL-12R β 2 deficient SJL/J EAE mice presented more severe symptoms and had more frequent, more severe relapses compared with wild type controls. IL-12R β 2 deficient EAE mice also had more infiltrating mononuclear cells in the central nervous system, as well as higher splenic proliferative capacity and decreased IFN- γ production at the periphery. These findings suggest a protective role of IL-12R β 2 in RR-EAE, an animal model of RR-MS, the most prevalent form of MS.

Keywords

Experimental autoimmune encephalomyelitis (EAE); IL-12R β 2; relapse; IL-12; IL-35

1. Introduction

Experimental autoimmune encephalomyelitis (EAE), an animal model of multiple sclerosis (MS), is a CD4⁺ T cell-mediated autoimmune disease (Noseworthy et al., 2000). In 2003, Cua et al. demonstrated that IL-23, but not IL-12, plays an essential role in EAE development (Cua et al., 2003). This finding dispelled the long-held view that IL-12 is critical to the development of EAE and called for a re-evaluation of its role in autoimmune central nervous system (CNS) inflammation.

IL-12 is a covalent heterodimer of p40 and p35 subunits, which form the complete cytokine IL-12p70 (Trinchieri, 1998). IL-23 is comprised of covalently bound p40 and p19 (Oppmann et al., 2000); p40 is therefore a common subunit between IL-12 and IL-23. IL-12 exerts its effects through its receptor (IL-12R), which is comprised of two chains, IL-12R β 1 and

Correspondence should be addressed to Abdolmohamad Rostami (a.m.rostami@jefferson.edu), Department of Neurology, Thomas Jefferson University, 901 Walnut Street, Suite 400, Philadelphia, Pennsylvania 19107, USA.

³Chong Xie and Bogoljub Ciric contributed equally to this work.

Conflict of interest

The authors declare that they have no conflict of interest.

IL-12R β 2 (Chua et al., 1995). IL-12R β 1 is also part of the IL-23 receptor heterodimer (Oppmann, Lesley, 2000), while IL-12R β 2 also participates in the IL-35 receptor, which is a non-covalent heterodimer of p35 and Epstein-Barr-virus-induced gene 3 (EBI3) (Collison et al., 2007). EBI3 together with p28 forms IL-27 (Pflanz et al., 2002); hence, in addition to a cytokine subunit, IL-35 and IL-27 also share a receptor subunit, gp130 (Pflanz et al., 2004), which is also part of the IL-6 receptor (Hibi et al., 1990). The members of the IL-12 family of cytokines and their receptors are shown in Supplementary Fig 1. IL-23 has a well-known essential role in EAE, while the role of IL-35 is less clear. IL-35 is produced by regulatory T cells (Tregs) and is regarded as a regulatory/anti-inflammatory cytokine (Collison, Workman, 2007, Niedbala et al., 2007). In one study systemic deficiency in IL-35 had no effect on EAE (Liu et al., 2012), while in another study lack of IL-35 production by B cells resulted in an exacerbated disease course (Shen et al., 2014). Given that both IL-12 subunits and both IL-12R subunits are shared with other cytokines or their receptors, it has been difficult to devise an approach that would address the role of IL-12; the same is true for IL-35. More data are available on the role of IL-12R β 2 in EAE when this receptor subunit mediates IL-12 signaling rather than IL-35 signaling. However, in each case the overall effect is the result of both IL-12 and IL-35 signaling.

The most relevant findings on the role of IL-12R β 2 in EAE come from a study with IL-12R β 2^{-/-} C57BL/6 mice. These mice were hypersusceptible to EAE compared with wild-type (WT) controls. Consistent with severe clinical signs, the IL-12R β 2^{-/-} C57BL/6 mice had extensive CNS demyelination and inflammation, as well as increased production of pro-inflammatory cytokines. These findings demonstrated that IL-12R β 2 has a suppressive role in chronic EAE (Zhang et al., 2003).

Factors that determine various disease courses in MS and EAE are not well understood, but it is apparent that the clinical disease course in chronic relapsing-remitting (RR) and progressive MS/EAE is governed by different mechanisms. As a consequence, a single therapy might not be effective in all disease types, as illustrated by the ineffectiveness of IFN- β in primary-progressive MS, while in RR-MS IFN- β has a beneficial effect (Rojas et al., 2009, Wingerchuk, 2008). In addition, the underlying pathogenic mechanisms that control various disease stages might be different, and treatment that is efficacious early on might be ineffective at a later disease stage (Caon, 2009). For example, IFN- β reduces relapses in RR-MS but loses most of its effectiveness when the disease becomes secondary progressive (Applebee and Panitch, 2009).

MOG₃₅₋₅₅-induced EAE in C57BL/6 mice has a chronic disease course, while the role of IL-12R β 2 signaling in EAE models with a different clinical course, such as RR-EAE, is not known. The majority of MS patients (~85%) develop RR disease (Compston and Coles, 2008), and EAE models that have RR clinical course are likely better suited for studying immunopathogenic mechanisms of RR-MS. The most commonly used RR-EAE model is in SJL mice immunized with PLP₁₃₉₋₁₅₁ peptide (Furlan et al., 2009). Histopathologically and clinically, it is similar to the RR form of MS and is used for studying relapse.

In this study, we for the first time demonstrate that IL-12R β 2 plays a strong suppressive role in RR-EAE. IL-12R β 2^{-/-} SJL mice developed markedly more severe EAE characterized by

high mortality and stronger relapses compared with WT counterparts. Given that the effects of IL-12R β 2 deficiency reflect the roles of both IL-12 and IL-35, we draw conclusions on the role of IL-12R β 2, without specifically attributing it to either of these two cytokines.

2. Materials and Methods

2.1. Mice and EAE induction

IL-12R β 2^{-/-} SJL/J mice were generated by the Jackson Laboratory (Bar Harbor, ME) using the Speed Congenic approach. Mutant IL-12R β 2 allele was derived from IL-12R β 2^{-/-} C57BL/6 mice (strain name: B6.129S1-*IL12R β 2^{tm1Jm}/J*). IL-12R β 2^{-/-} SJL/J mice were kept homozygous by brother-sister breeding. WT SJL/J mice were purchased from the Jackson Laboratory. In most experiments IL-12R β 2^{-/-} and WT mice were immunized for EAE induction with 200 μ g PLP₁₃₉₋₁₅₁ (GenScript) emulsified in complete Freund's adjuvant (CFA) containing 3 mg/ml *Mycobacterium tuberculosis* (H37Ra; Difco Laboratories). In addition, mice were given 200 ng pertussis toxin (List Biologicals Laboratories) i.p. on days 0 and 2 post-immunization (p.i.). In some experiments mice received less PLP₁₃₉₋₁₅₁ peptide; immunizing conditions are specified in figure legends associated with those experiments. Mice were graded for clinical manifestations of EAE by the following criteria: 1, tail paralysis; 2, one hind limb paralysis; 3, both hind limbs paralysis; 4, forelimb weakness or paralysis; 5, moribund or dead. All animal procedures were performed in accordance with the guidelines of the Institutional Animal Care and Committee of Thomas Jefferson University.

2.2. Mononuclear cell Preparation

Depending on the experiment, mice were sacrificed on various days p.i. After being anesthetized, mice were perfused with 20 ml phosphate-buffered saline (PBS). To isolate mononuclear cells (MNCs) from the CNS, spinal cords and brains were digested with 0.5 mg/mL Liberase TM (Roche) for 30 min at 37°C and then mechanically dissociated through a 70- μ m cell strainer. The single-cell suspension was then fractionated on a 70/30% Percoll gradient by centrifugation at 300 $\times$ g for 20 min. Cell layer at 70/30 interface was collected and viable cells were counted in 0.4% Trypan blue. For preparation of splenocytes, spleen was dissociated through a 70- μ m cell strainer and then red blood cells were lysed with Red Blood Cell Lysis Buffer (BioLegend). Splenocytes were then washed with cold medium and collected for use.

2.3. Splenocyte proliferation assay

Splenocytes were cultured in 96-well plates in 200 μ l IMDM medium supplemented with 10% Fetal Calf Serum (FCS), L-glutamine (2 mM), penicillin (100 U/ml), and streptomycin (100 μ g/ml). In the presence of 20 μ g/ml PLP₁₃₉₋₁₅₁ or 1 μ g/ml anti-CD3/CD28 antibodies, splenocytes were cultured at a density of 1 $\times$ 10⁵ cells/well, while without Ag/mitogen splenocytes were cultured at a density of 4 $\times$ 10⁵ cells/well. After 60 h of incubation at 37°C/5% CO₂, cells were pulsed for 12 h with 1 μ Ci of [³H]methylthymidine. Cells were then harvested and thymidine incorporation (cpm) was determined using a β -counter.

2.4. Cytokine measurement

Splenocytes of immunized mice were cultured at a density of 2.5×10^6 cells/ml in medium with or without 20 $\mu\text{g/ml}$ PLP_{139–151}. Supernatants were collected after 72 h of culturing. ELISA kits for measurement of IFN- γ and IL-17A concentrations were purchased from R&D System. Assays were performed according to the manufacturer's recommendation.

2.5. Flow cytometry

MNCs from CNS and spleen were stained with fluorochrome-conjugated antibodies against mouse CD4, CD11b, CD25, IL-17A, INF- γ and Gr1 (BD Biosciences). Flow cytometric analysis was performed on FACS Aria (BD Biosciences) and data were analyzed with FlowJo software (Tree Star).

2.6. Statistics

A two-tailed paired or unpaired Student's t-test was used to analyze differences between groups. A value of $P < 0.05$ was considered statistically significant.

3. Results

3.1. Characterization of IL-12R $\beta 2^{-/-}$ SJL/J mice

To study the role of IL-12R $\beta 2$ in RR-EAE we generated congenic IL-12R $\beta 2^{-/-}$ SJL/J mouse strain by crossing WT SJL/J mice and IL-12R $\beta 2^{-/-}$ C57BL/6 mice. SJL/J IL-12R $\beta 2^{-/-}$ mice were generated by the “speed congenic” approach carried out by Jackson Laboratories, Inc. We characterized basic immunological parameters of these mice; similar to IL-12R $\beta 2^{-/-}$ C57BL/6 mice (Wu et al., 2000), we did not find major defects in the immune system of their SJL/J counterparts.

IL-12 promotes IFN- γ production through IL-12R signaling (Magram et al., 1996, Trinchieri, 1994, Trinchieri and Scott, 1995). To functionally verify lack of IL-12R $\beta 2$ in IL-12R $\beta 2^{-/-}$ SJL/J mice (hereafter referred to as IL-12R $\beta 2^{-/-}$ mice), we tested the effect of recombinant IL-12 (rIL-12) on IFN- γ production by splenocytes of naïve mice activated with anti-CD3/CD28 antibodies. rIL-12 significantly increased (~4-fold) IFN- γ concentrations in cell culture supernatants of WT splenocytes, but had no effect in IL-12R $\beta 2^{-/-}$ cultures (Supplementary Fig. 2). These data clearly demonstrate the lack of IL-12R signaling in immune cells of IL-12R $\beta 2^{-/-}$ mice.

3.2. IL-12R $\beta 2^{-/-}$ mice were hypersusceptible to RR-EAE

To characterize EAE in IL-12R $\beta 2^{-/-}$ mice we immunized them and WT controls with 100 μg PLP_{139–151} emulsified in CFA. Incidence of disease was 100 % in both groups. Compared with WT mice, IL-12R $\beta 2^{-/-}$ mice had earlier (1–3 days) disease onset (Fig 1A, B) and more severe clinical disability (Fig 1B). RR-EAE clinical course in IL-12R $\beta 2^{-/-}$ mice paralleled that in WT mice, but with significantly more severity. Mortality in the IL-12R $\beta 2^{-/-}$ group was notably higher than in the WT group (2/10 WT, 6/10 IL-12R $\beta 2^{-/-}$). Because high mortality among IL-12R $\beta 2^{-/-}$ mice during acute attack precluded meaningful follow-up analyses, we induced EAE with suboptimal quantities of PLP_{139–151}, *M. tuberculosis*, and pertussis toxin. When mice were immunized with 75 μg PLP_{139–151}

(instead of 100 μ g), 2 mg/ml *M. tuberculosis* in CFA (instead of 3 mg/ml), and given 2 $\times$ 100 ng pertussis toxin (instead of 2 $\times$ 200 ng), clinical course (data not shown) and mortality (3/10 WT, 7/10 IL-12R β 2^{-/-}) remained similar to that in mice having received optimal immunization. We therefore further reduced the quantity of PLP_{139–151} (50 μ g) given to each mouse. This suboptimal immunization showed the full extent of hypersusceptibility of IL-12R β 2^{-/-} mice to EAE, as it induced mild EAE in WT mice with 100% incidence, whereas IL-12R β 2^{-/-} mice developed severe disease, also with 100% incidence (Fig 1C). Mortality in IL-12R β 2^{-/-} mice (2/15) and WT mice (0/15) was much reduced compared to those immunized with 100 ng of PLP_{139–151}.

As regards relapses, IL-12R β 2^{-/-} mice had a higher incidence of relapse (100%) and a greater number of relapses per mouse compared to WT controls. Relapses in IL-12R β 2^{-/-} mice developed earlier, were more severe and lasted longer than in WT controls. All 13 of the surviving IL-12R β 2^{-/-} mice developed 2 relapses by day 42 p.i. when the experiment was terminated, while 8/15 WT mice developed 1 relapse and none of the mice had 2 relapses.

Numbers of MNCs obtained from the CNS at different stages of disease (onset, remission, first relapse, relapsing-progressive stage) were consistent with clinical severity of disease, with 2–3 times more cells obtained from IL-12R β 2^{-/-} mice (Fig 1D).

These data demonstrate that IL-12R β 2 has a strong suppressive role in RR-EAE in SJL/J mice, similar to its suppressive role in chronic EAE in C57BL/6 mice (Zhang, Gran, 2003).

3.3. IL-12R β 2^{-/-} mice had a higher Th17/Th1 ratio among CD4⁺ T cells isolated from the CNS during EAE

Mice were immunized for EAE induction; groups of mice were sacrificed, and infiltrating MNCs in the CNS were isolated and analyzed at different time points. Once the first clinical signs of disease appeared, both groups of mice progressed to a high clinical score (3–4) within 1–2 days. In our experiments disease onset and peak thus largely overlapped, but day 11 p.i. can be viewed as disease peak rather than onset in most of our experiments. T helper type 17 (Th17) cells (IL-17A⁺IFN- γ ⁻) constituted half of all CD4⁺ T cells at day 11 in both groups of mice, while IL-17A⁺IFN- γ ⁺CD4⁺ T cells were less abundant (10–15%) (Fig. 2). Th1 cells (IL-17A⁻IFN- γ ⁺) were the least abundant (5–9%) population. Hence, IL-12R β 2^{-/-} and WT mice had similarly high proportions of IL-17A-producing CD4⁺ T cells in the CNS at EAE onset and/or peak of acute attack. The proportion of IFN- γ -producing CD4⁺ T cells was higher in WT mice, but these cells were also abundant in IL-12R β 2^{-/-} mice. During remission (day 21 p.i.), proportions of IL-17A-producing cells dropped dramatically in IL-12R β 2^{-/-} (3.5-fold) and especially in WT mice (11-fold) compared with disease peak, while proportions of Th1 cells increased twofold in both groups. In the 1st relapse (day 28 p.i.), proportions of IL-17A-producing cells increased relative to remission, but remained 2–3 times lower than at initial peak of disease. Compared with WT mice, IL-12R β 2^{-/-} mice had higher proportions of IL-17A-producing and lower proportions of IFN- γ -producing CD4⁺ T cells. Later in disease (day 50 p.i.) during the relapsing-progressive phase, IL-12R β 2^{-/-} mice still had a higher proportion of IL-17A-producing cells compared with WT mice, while proportions of IFN- γ -producing cells were similar.

Over all, at disease onset/peak proportions of IL-17A-producing CD4⁺ T cells were similar between WT and IL-12Rβ2^{-/-} mice, while at later time points, irrespective of disease phase (remission, relapse) WT mice had substantially smaller proportions (two- to threefold) of these cells. WT mice typically had higher proportions of IFN-γ-producing CD4⁺ T cells throughout disease compared with IL-12Rβ2^{-/-} mice. It should be taken into account that total numbers of MNCs and CD4⁺ T cells present in the CNS of IL-12Rβ2^{-/-} mice were at all disease stages 2–3 times higher than in WT mice. Hence, proportions of certain Th lineages among CD4⁺ T cells do not simply translate into total numbers of these cells present in the CNS, and, therefore, IL-12Rβ2^{-/-} mice had approximately two- to threefold more IL-17A-producing, and even IFN-γ producing, Th cells in the CNS throughout disease.

3.4. Splenocytes of IL-12Rβ2^{-/-} mice produced less IFN-γ

In addition to the CNS, we also evaluated PLP_{139–151}-specific Th response at the periphery of immunized mice. Splenocytes were harvested at different time points post immunization and cultured for 72 h with or without addition of PLP_{139–151}. We did not observe consistent differences in IL-17A production between splenocytes of IL-12Rβ2^{-/-} and WT mice (Fig 3). As for IFN-γ production, IL-12Rβ2^{-/-} splenocytes consistently produced 2–5 times less IFN-γ than WT splenocytes, suggesting that IL-12Rβ2^{-/-} mice developed significantly weaker PLP_{139–151}-specific Th1 responses.

We also analyzed splenocytes by flow cytometry, ex vivo, immediately after harvesting, and after 3 days of stimulation with PLP_{139–151} *in vitro*. In neither case did we observe consistent differences in numbers of IL-17A⁺ and IFN-γ⁺ Th cells between IL-12Rβ2^{-/-} and WT splenocytes at different time points p.i. (data not shown).

Compared with WT mice, IL-12Rβ2^{-/-} mice had more of both CD25⁺Foxp3⁺CD4⁺ and CD25⁻Foxp3⁺CD4⁺ cells in the spleen at each time point we analyzed (Fig 4). The percentages of CD25⁺Foxp3⁺CD4⁺ cells showed minor fluctuations over time, while percentages of CD25⁻Foxp3⁺CD4⁺ T cells steadily increased. These data show that IL-12Rβ2^{-/-} mice develop more classical CD25⁺Foxp3⁺CD4⁺ Tregs in the spleen than WT counterparts, as well as non-classical CD25⁻Foxp3⁺CD4⁺ T cells.

These data show that the lack of IL-12Rβ2 results in reduced production of IFN-γ in response to immunizing peptide, and perhaps overall Th1 response, while IL-17A production and Th17 response are only marginally increased.

Analysis of splenic cells after disease onset has only limited value, given that a large number of Th cells trafficked to the CNS, resulting in diminished numbers of these cells in the spleen.

3.5. IL-12Rβ2^{-/-} splenocytes had higher proliferative capacity

We then tested the proliferative capacity of splenocytes from immunized mice. Splenocytes prepared on days 7, 11, 21, 28, and 50 p.i. were cultured in the presence of PLP_{139–151} or anti-CD3/CD28 antibodies. Without stimulation, IL-12Rβ2^{-/-} and WT splenocytes had similarly low levels of proliferation. With PLP_{139–151} stimulation, IL-12Rβ2^{-/-} splenocytes proliferated significantly more strongly than WT splenocytes up to day 28 p.i., while on day

50 p.i. there was no difference (Fig 5). This indicates that IL-12R β 2^{-/-} mice develop stronger myelin-specific immune responses upon immunization. However, IL-12R β 2^{-/-} splenocytes also proliferated much more strongly upon stimulation with anti-CD3/CD28 antibodies, suggesting that lack of IL-12R β 2^{-/-} signaling causes abnormally high proliferative capacity of CD4⁺ T cells. As for splenocytes from naïve mice, the overall level of proliferation of IL-12R β 2^{-/-} and WT splenocytes was similar at 24–48 h. However, IL-12R β 2^{-/-} splenocytes retained the same level of proliferation at 48–72 h, while WT splenocytes exhibited a sharp decrease at that time (data not shown).

3.6. IL-12R β 2^{-/-} mice had a higher proportion of CD11b⁺Gr1⁺ cells

Compared with WT mice, IL-12R β 2^{-/-} mice had higher percentages of CD11b⁺Gr1⁺ cells in both the CNS and spleen from day 7 until day 50 p.i. (Fig 6). Percentages of these cells among splenocytes remained similar throughout disease, while in the CNS after the spike at onset/peak of clinical disease percentages of CD11b⁺Gr1⁺ cells dropped two- to fourfold at later time points. Compared with CD11b⁺Gr1⁺ cells, proportions of CD11b⁺Gr1⁻ cells among spleen and CNS cells were much smaller, and not significantly different between IL-12R β 2^{-/-} and WT cells, except on day 28 p.i., which coincides with the first relapse. At that time point, IL-12R β 2^{-/-} mice had a large proportion of CD11b⁺Gr1⁻ cells among cells obtained from the CNS. The majority of CD11b⁺Gr1⁻ cells had a lower expression of CD11b than CD11b⁺Gr1⁺ cells, which is consistent with a microglia phenotype. In our experience, the vast majority of CD11b⁺Gr1⁺ cells, both in the CNS and spleen of mice with EAE, are Ly6G⁺Ly6C⁻ neutrophils, while Ly6G⁻Ly6C⁺ monocytes represent less than 10% of CD11b⁺Gr1⁺ cells. Hence, our data suggest that in comparison with WT mice, IL-12R β 2^{-/-} mice have a higher proportion of neutrophils in the CNS and spleen. The assumption that CD11b⁺Gr1⁺ cells are predominantly neutrophils is further supported by the drop in their numbers among MNCs from the CNS (Fig. 6), as neutrophils constitute a large portion of CNS infiltrating cells at disease onset but their number decreases during disease course (Maatta et al., 1998, Wu et al., 2010).

4. Discussion

In this study we addressed the role of IL-12R β 2 in RR-EAE; IL-12R β 2^{-/-} SJL/J mice were hypersusceptible to EAE, which includes greater mortality, an earlier disease onset, more severe clinical signs, as well as more frequent and more severe relapses compared with WT controls. Consistent with a more severe clinical disease, IL-12R β 2^{-/-} mice had more infiltrating MNCs in the CNS and stronger PLP_{139–151}-specific Th responses at the periphery. These findings demonstrate that IL-12R β 2 plays a protective role in RR-EAE.

The protective role of IL-12R β 2 in EAE has been shown previously in C57BL/6 mice, which develop chronic disease course upon immunization with MOG_{35–55} peptide (Zhang, Gran, 2003). Paralleling severe EAE in IL-12R β 2^{-/-} SJL mice, IL-12R β 2^{-/-} C57BL/6 mice had earlier onset of disease, higher clinical scores, and a greater rate of mortality compared with WT controls. Additional similarities include: 2–3 times more infiltrating MNCs in the CNS, greater proliferative response of splenocytes upon stimulation with immunizing peptides, reduced production of IFN- γ and a tendency to produce more IL-17A in response to

immunizing peptides *in vitro*. Moreover, C57BL/6 mice deficient in IL-12p35, which binds to IL-12R β 2, were also susceptible to EAE and produced less IFN- γ (Gran et al., 2002).

In our study, we further noticed that IL-12R β 2^{-/-} SJL/J mice had more CD11b⁺Gr1⁺ cells in CNS and spleen. Although in this study we did not specifically analyze neutrophils, which are CD11b⁺Ly6G^{high}Ly6C^{low} (Hestdal et al., 1991), we know from our studies that the vast majority (>90%) of CD11b⁺Gr1⁺ cells in spleens and the CNS of mice with EAE are CD11b⁺Ly6G^{high}Ly6C^{low} neutrophils, while the proportion of CD11b⁺Ly6G⁻Ly6C^{high/medium} monocytes typically does not exceed 5–10%. Hence, in the absence of IL-12R β 2, proportions of neutrophils among immune cells in spleen and CNS significantly increase compared with mice with intact IL-12R β 2 signaling. Neutrophils are among the first cells recruited into the CNS (Christy et al., 2013). In the preclinical phase of EAE, neutrophils are activated and alter the integrity of the blood-brain barrier, promoting subsequent infiltration of other immune cell types (Aube et al., 2014). The proportion of neutrophils among infiltrating cells in the CNS is highest during onset of clinical disease; it remains high until disease peak then gradually declines (Wu, Cao, 2010). Possible reason for the increase in neutrophils in IL-12R β 2^{-/-} mice can be stronger Th17 responses together with marked reduction in IFN- γ production, as it has been established that Th17 cell-dominated encephalitogenic responses lead to a greater proportion of granulocytes, and especially neutrophils, in CNS infiltrates (Aube, Levesque, 2014, Maatta, Sjöholm, 1998, Steinbach et al., 2013, Wu, Cao, 2010).

Zhang et al. found that splenocytes of IL-12R β 2^{-/-} C57BL/6 EAE mice produce more granulocyte-macrophage colony-stimulating factor (GM-CSF) and express 5 times more IL-23p19 than WT splenocytes (Zhang, Gran, 2003). IL-23 and GM-CSF form a positive feedback loop, in which IL-23 produced by antigen presenting cells (APCs) promotes development of GM-CSF-producing Th17 cells, and GM-CSF secreted by these cells enhances IL-23 production by APCs (El-Behi et al., 2011). It appears that the lack of IL-12R β 2 results in greater production of IL-23 and GM-CSF, increasing the numbers of Th17 cells, which then cause more severe EAE. An additional likely reason for the increase in Th17 cells is markedly reduced IFN- γ production in IL-12R β 2^{-/-} mice, as IFN- γ antagonizes development of Th17 cells (Bettelli et al., 2006, Mangan et al., 2006, Veldhoen et al., 2006), and also suppresses GM-CSF production (Kroenke et al., 2010). Enhanced Th17 response in IL-12R β 2^{-/-} mice may also result from the absence of IL-35 signaling, given that IL-35 inhibits Th17 cell development *in vitro* and *in vivo* (Niedbala, Wei, 2007, Shen, Roch, 2014, Yang et al., 2008). IL-35 is composed of IL-12p35 and EBI3 (Collison, Workman, 2007); Yang et al. demonstrated that EBI3^{-/-} mice developed more Th17 cells *in vivo*, and splenocytes from EBI3^{-/-} mice produced more IL-17A *in vitro* (Yang, Yang, 2008). Furthermore, mice with a B-cell-restricted deficiency in IL-12p35 or EBI3, hence lacking IL-35, developed exacerbated EAE and had increased MOG_{35–55}-induced IL-17A production (Shen, Roch, 2014). Both the IL-12R β 2^{-/-} C57BL/6 and SJL/J mice had stronger Th17 responses, as evidenced by greater numbers of Th17 cells infiltrated in the CNS and more IL-17A being produced by peripheral immune cells. Th17 cells plays a critical role in EAE and it is believed that they are more pathogenic than Th1 cells (El-Behi, Ciric, 2011, Langrish et al., 2005); enhanced Th17 response may therefore be one of the reasons for more severe EAE in IL-12R β 2^{-/-} mice.

Apart from the enhanced Th17 response, higher splenic proliferative capacity may also contribute to more severe EAE in IL-12R β 2^{-/-} mice. Splenocytes of IL-12R β 2^{-/-} mice had greater proliferation upon stimulation with both antigen and mitogen than WT splenocytes, indicating an immunoregulatory role of IL-12R β 2 signaling. IL-12 can induce the proliferation of both activated CD4⁺ and CD8⁺ T cells (Gately et al., 1991), while in contrast, IL-35 inhibits T cell proliferation (Collison et al., 2010, Collison, Workman, 2007). Recombinant IL-35 potently inhibits the proliferation of effector T cells *in vitro* (Collison, Workman, 2007). The suppressive effect on T cell proliferation may also be indirect, due to the promoting effect of IL-35 on Tregs (Collison, Chaturvedi, 2010, Collison, Workman, 2007). The greater proliferative capacity of IL-12R β 2^{-/-} splenocytes may be caused by a combination of factors, such as lack of direct and indirect, via Tregs, suppressive effects of IL-35 signaling, and reduced production of IFN- γ , which suppresses CD4⁺ T cell proliferation (De Rose et al., 2004).

IFN- γ production in IL-12R β 2^{-/-} mice was lower than in WT mice, which may also have contributed to their hypersusceptibility to EAE. It has been established that IFN- γ plays a suppressive role in EAE, as mice deficient in IFN- γ or its receptor are hypersusceptible to EAE (Ferber et al., 1996, Willenborg et al., 1996). Induction of IFN- γ by administration of IL-12 during the priming phase of EAE suppressed disease development (Gran et al., 2004). Direct intrathecal delivery of IFN- γ also inhibited the course of chronic-progressive EAE and promoted apoptosis of infiltrating lymphocytes in the CNS (Furlan et al., 2001). On the contrary, injection of neutralizing antibody against IFN- γ caused an increase in disability and mortality of EAE mice (Billiau et al., 1988). Similarly, the neutralizing antibody against IFN- γ also facilitated EAE relapses in SJL/J and Biozzi ABH mice (Heremans et al., 1996). The suppressive role of IFN- γ in EAE may be mediated by multiple mechanisms, such as direct inhibition of Th17 cell development (Bettelli, Carrier, 2006), enhancement of Treg development and function (Harrington et al., 2005, Wang et al., 2006), inhibition of GM-CSF production (Codarri et al., 2011), increase in Th cell apoptosis (Refaeli et al., 2002), and reduction in iNOS expression (Igietseme, 1996, Riquelme et al., 2013). Even though IL-12R β 2^{-/-} mice had reduced production of IFN- γ , they still produced substantial quantities of this cytokine. It is therefore difficult to discern to which extent EAE phenotype of IL-12R β 2^{-/-} mice is shaped by reduced levels of IFN- γ vs. lack of IFN- γ -independent effects of IL-12 and IL-35 signaling.

It has been reported that both IL-35 and IL-12R β 2 are important for the development of Tregs and their suppressive activities (Collison, Chaturvedi, 2010, Collison, Workman, 2007, Niedbala, Wei, 2007). IL-35 can expand Tregs, increase Foxp3 expression and IL-10 production (Niedbala, Wei, 2007). IL-35 can also induce a different type of regulatory T cell called iT_R35, which does not express or require Foxp3 for its development and function and mediates suppression via IL-35 (Collison, Chaturvedi, 2010). However, in our study more CD4⁺CD25⁺Foxp3⁺ Tregs were found in IL-12R β 2^{-/-} mice, which lacked IL-35 signaling. The higher frequency of Tregs in IL-12R β 2^{-/-} EAE mice may represent a negative feedback loop against the stronger auto-immunoreaction in these mice. In addition to CD4⁺CD25⁺Foxp3⁺ Tregs, we also found CD4⁺CD25⁻Foxp3⁺ T cells in spleen. CD4⁺CD25⁻Foxp3⁺ T cells have been seen in patients with systemic lupus erythematosus (Bonelli et al., 2009, Zhang et al., 2008) and they have been associated with aging (Nishioka

et al., 2006). Phenotypic analysis showed that CD4⁺CD25⁻Foxp3⁺ T cells resemble conventional Tregs and exert lower, but still considerable, suppressive capacity on T cell proliferation (Bonelli, Savitskaya, 2009).

In summary, we evaluated the role of IL-12Rβ2 signaling in RR-EAE using IL-12Rβ2^{-/-} SJL/J mice. Our findings clearly demonstrate that IL-12Rβ2 has a protective role in this EAE model, which parallels its protective role in chronic EAE. Given that IL-12Rβ2 participates in signaling of two cytokines, IL-12 and IL-35, the potential mechanisms underlying its protective role are likely to be complex. Both these cytokines have numerous known effects, and possibly some that are yet to be discovered, making identification of those that significantly contribute to the EAE phenotype of IL-12Rβ2^{-/-} mice highly challenging.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank Katherine Regan for editorial assistance.

References

- Applebee A, Panitch H. Early stage and long term treatment of multiple sclerosis with interferon-beta. *Biologics*. 2009; 3:257–271. [PubMed: 19707414]
- Aube B, Levesque SA, Pare A, Chamma E, Kebir H, Gorina R, et al. Neutrophils mediate blood-spinal cord barrier disruption in demyelinating neuroinflammatory diseases. *Journal of immunology*. 2014; 193:2438–2454.
- Bettelli E, Carrier Y, Gao W, Korn T, Strom TB, Oukka M, et al. Reciprocal developmental pathways for the generation of pathogenic effector TH17 and regulatory T cells. *Nature*. 2006; 441:235–238. [PubMed: 16648838]
- Billiau A, Heremans H, Vandekerckhove F, Dijkmans R, Sobis H, Meulepas E, et al. Enhancement of experimental allergic encephalomyelitis in mice by antibodies against IFN-gamma. *Journal of immunology*. 1988; 140:1506–1510.
- Bonelli M, Savitskaya A, Steiner CW, Rath E, Smolen JS, Scheinecker C. Phenotypic and functional analysis of CD4⁺ CD25⁻ Foxp3⁺ T cells in patients with systemic lupus erythematosus. *Journal of immunology*. 2009; 182:1689–1695.
- Caon C. Maximising therapeutic outcomes in patients failing on current therapy. *J Neurol Sci*. 2009; 277(Suppl 1):S33–S36. [PubMed: 19200864]
- Christy AL, Walker ME, Hessner MJ, Brown MA. Mast cell activation and neutrophil recruitment promotes early and robust inflammation in the meninges in EAE. *Journal of autoimmunity*. 2013; 42:50–61. [PubMed: 23267561]
- Chua AO, Wilkinson VL, Presky DH, Gubler U. Cloning and characterization of a mouse IL-12 receptor-beta component. *Journal of immunology*. 1995; 155:4286–4294.
- Codarri L, Gyulveszi G, Tosevski V, Hesske L, Fontana A, Magnenat L, et al. RORγt drives production of the cytokine GM-CSF in helper T cells, which is essential for the effector phase of autoimmune neuroinflammation. *Nature immunology*. 2011; 12:560–567. [PubMed: 21516112]
- Collison LW, Chaturvedi V, Henderson AL, Giacomini PR, Guy C, Bankoti J, et al. IL-35-mediated induction of a potent regulatory T cell population. *Nature immunology*. 2010; 11:1093–1101. [PubMed: 20953201]

- Collison LW, Workman CJ, Kuo TT, Boyd K, Wang Y, Vignali KM, et al. The inhibitory cytokine IL-35 contributes to regulatory T-cell function. *Nature*. 2007; 450:566–569. [PubMed: 18033300]
- Compston A, Coles A. Multiple sclerosis. *Lancet*. 2008; 372:1502–1517. [PubMed: 18970977]
- Cua DJ, Sherlock J, Chen Y, Murphy CA, Joyce B, Seymour B, et al. Interleukin-23 rather than interleukin-12 is the critical cytokine for autoimmune inflammation of the brain. *Nature*. 2003; 421:744–748. [PubMed: 12610626]
- De Rose V, Cappello P, Sorbello V, Ceccarini B, Gani F, Bosticardo M, et al. IFN-gamma inhibits the proliferation of allergen-activated T lymphocytes from atopic, asthmatic patients by inducing Fas/FasL-mediated apoptosis. *Journal of leukocyte biology*. 2004; 76:423–432. [PubMed: 15123769]
- El-Behi M, Ciric B, Dai H, Yan Y, Cullimore M, Safavi F, et al. The encephalitogenicity of T(H)17 cells is dependent on IL-1- and IL-23-induced production of the cytokine GM-CSF. *Nature immunology*. 2011; 12:568–575. [PubMed: 21516111]
- Ferber IA, Brocke S, Taylor-Edwards C, Ridgway W, Dinisco C, Steinman L, et al. Mice with a disrupted IFN-gamma gene are susceptible to the induction of experimental autoimmune encephalomyelitis (EAE). *Journal of immunology*. 1996; 156:5–7.
- Furlan R, Brambilla E, Ruffini F, Poliani PL, Bergami A, Marconi PC, et al. Intrathecal delivery of IFN-gamma protects C57BL/6 mice from chronic-progressive experimental autoimmune encephalomyelitis by increasing apoptosis of central nervous system-infiltrating lymphocytes. *Journal of immunology*. 2001; 167:1821–1829.
- Furlan R, Cuomo C, Martino G. Animal models of multiple sclerosis. *Methods Mol Biol*. 2009; 549:157–173. [PubMed: 19378202]
- Gately MK, Desai BB, Wolitzky AG, Quinn PM, Dwyer CM, Podlaski FJ, et al. Regulation of human lymphocyte proliferation by a heterodimeric cytokine, IL-12 (cytotoxic lymphocyte maturation factor). *Journal of immunology*. 1991; 147:874–882.
- Gran B, Chu N, Zhang GX, Yu S, Li Y, Chen XH, et al. Early administration of IL-12 suppresses EAE through induction of interferon-gamma. *Journal of neuroimmunology*. 2004; 156:123–131. [PubMed: 15465603]
- Gran B, Zhang GX, Yu S, Li J, Chen XH, Ventura ES, et al. IL-12p35-deficient mice are susceptible to experimental autoimmune encephalomyelitis: evidence for redundancy in the IL-12 system in the induction of central nervous system autoimmune demyelination. *Journal of immunology*. 2002; 169:7104–7110.
- Harrington LE, Hatton RD, Mangan PR, Turner H, Murphy TL, Murphy KM, et al. Interleukin 17-producing CD4+ effector T cells develop via a lineage distinct from the T helper type 1 and 2 lineages. *Nature immunology*. 2005; 6:1123–1132. [PubMed: 16200070]
- Heremans H, Dillen C, Groenen M, Martens E, Billiau A. Chronic relapsing experimental autoimmune encephalomyelitis (CREAE) in mice: enhancement by monoclonal antibodies against interferon-gamma. *European journal of immunology*. 1996; 26:2393–2398. [PubMed: 8898951]
- Hestdal K, Ruscetti FW, Ihle JN, Jacobsen SE, Dubois CM, Kopp WC, et al. Characterization and regulation of RB6-8C5 antigen expression on murine bone marrow cells. *Journal of immunology*. 1991; 147:22–28.
- Hibi M, Murakami M, Saito M, Hirano T, Taga T, Kishimoto T. Molecular cloning and expression of an IL-6 signal transducer, gp130. *Cell*. 1990; 63:1149–1157. [PubMed: 2261637]
- Igietseme JU. Molecular mechanism of T-cell control of Chlamydia in mice: role of nitric oxide in vivo. *Immunology*. 1996; 88:1–5. [PubMed: 8707333]
- Kroenke MA, Chensue SW, Segal BM. EAE mediated by a non-IFN-gamma/non-IL-17 pathway. *European journal of immunology*. 2010; 40:2340–2348. [PubMed: 20540117]
- Langrish CL, Chen Y, Blumenschein WM, Mattson J, Basham B, Sedgwick JD, et al. IL-23 drives a pathogenic T cell population that induces autoimmune inflammation. *The Journal of experimental medicine*. 2005; 201:233–240. [PubMed: 15657292]
- Liu JQ, Liu Z, Zhang X, Shi Y, Talebian F, Carl JW Jr, et al. Increased Th17 and regulatory T cell responses in EBV-induced gene 3-deficient mice lead to marginally enhanced development of autoimmune encephalomyelitis. *Journal of immunology*. 2012; 188:3099–3106.
- Maatta JA, Sjöholm UR, Nygardas PT, Salmi AA, Hinkkanen AE. Neutrophils secreting tumor necrosis factor alpha infiltrate the central nervous system of BALB/c mice with experimental

- autoimmune encephalomyelitis. *Journal of neuroimmunology*. 1998; 90:162–175. [PubMed: 9817444]
- Magram J, Connaughton SE, Warriar RR, Carvajal DM, Wu CY, Ferrante J, et al. IL-12-deficient mice are defective in IFN gamma production and type 1 cytokine responses. *Immunity*. 1996; 4:471–481. [PubMed: 8630732]
- Mangan PR, Harrington LE, O'Quinn DB, Helms WS, Bullard DC, Elson CO, et al. Transforming growth factor-beta induces development of the T(H)17 lineage. *Nature*. 2006; 441:231–234. [PubMed: 16648837]
- Niedbala W, Wei XQ, Cai B, Hueber AJ, Leung BP, McInnes IB, et al. IL-35 is a novel cytokine with therapeutic effects against collagen-induced arthritis through the expansion of regulatory T cells and suppression of Th17 cells. *European journal of immunology*. 2007; 37:3021–3029. [PubMed: 17874423]
- Nishioka T, Shimizu J, Iida R, Yamazaki S, Sakaguchi S. CD4+CD25+Foxp3+ T cells and CD4+CD25-Foxp3+ T cells in aged mice. *Journal of immunology*. 2006; 176:6586–6593.
- Noseworthy JH, Lucchinetti C, Rodriguez M, Weinshenker BG. Multiple sclerosis. *The New England journal of medicine*. 2000; 343:938–952. [PubMed: 11006371]
- Oppmann B, Lesley R, Blom B, Timans JC, Xu Y, Hunte B, et al. Novel p19 protein engages IL-12p40 to form a cytokine, IL-23, with biological activities similar as well as distinct from IL-12. *Immunity*. 2000; 13:715–725. [PubMed: 11114383]
- Pflanz S, Hibbert L, Mattson J, Rosales R, Vaisberg E, Bazan JF, et al. WSX-1 and glycoprotein 130 constitute a signal-transducing receptor for IL-27. *Journal of immunology*. 2004; 172:2225–2231.
- Pflanz S, Timans JC, Cheung J, Rosales R, Kanzler H, Gilbert J, et al. IL-27, a heterodimeric cytokine composed of EBI3 and p28 protein, induces proliferation of naive CD4+ T cells. *Immunity*. 2002; 16:779–790. [PubMed: 12121660]
- Refaeli Y, Van Parijs L, Alexander SI, Abbas AK. Interferon gamma is required for activation-induced death of T lymphocytes. *The Journal of experimental medicine*. 2002; 196:999–1005. [PubMed: 12370261]
- Riquelme P, Tomiuk S, Kammler A, Fandrich F, Schlitt HJ, Geissler EK, et al. IFN-gamma-induced iNOS expression in mouse regulatory macrophages prolongs allograft survival in fully immunocompetent recipients. *Molecular therapy : the journal of the American Society of Gene Therapy*. 2013; 21:409–422. [PubMed: 22929659]
- Rojas JI, Romano M, Ciapponi A, Patrucco L, Cristiano E. Interferon beta for primary progressive multiple sclerosis. *Cochrane Database Syst Rev*. 2009;CD006643. [PubMed: 19160292]
- Shen P, Roch T, Lampropoulou V, O'Connor RA, Stervbo U, Hilgenberg E, et al. IL-35-producing B cells are critical regulators of immunity during autoimmune and infectious diseases. *Nature*. 2014; 507:366–370. [PubMed: 24572363]
- Steinbach K, Piedavent M, Bauer S, Neumann JT, Friese MA. Neutrophils amplify autoimmune central nervous system infiltrates by maturing local APCs. *Journal of immunology*. 2013; 191:4531–4539.
- Trinchieri G. Interleukin-12: a cytokine produced by antigen-presenting cells with immunoregulatory functions in the generation of T-helper cells type 1 and cytotoxic lymphocytes. *Blood*. 1994; 84:4008–4027. [PubMed: 7994020]
- Trinchieri G. Interleukin-12: a cytokine at the interface of inflammation and immunity. *Advances in immunology*. 1998; 70:83–243. [PubMed: 9755338]
- Trinchieri G, Scott P. Interleukin-12: a proinflammatory cytokine with immunoregulatory functions. *Research in immunology*. 1995; 146:423–431. [PubMed: 8839141]
- Veldhoen M, Hocking RJ, Atkins CJ, Locksley RM, Stockinger B. TGFbeta in the context of an inflammatory cytokine milieu supports de novo differentiation of IL-17-producing T cells. *Immunity*. 2006; 24:179–189. [PubMed: 16473830]
- Wang Z, Hong J, Sun W, Xu G, Li N, Chen X, et al. Role of IFN-gamma in induction of Foxp3 and conversion of CD4+ CD25- T cells to CD4+ Tregs. *The Journal of clinical investigation*. 2006; 116:2434–2441. [PubMed: 16906223]
- Willenborg DO, Fordham S, Bernard CC, Cowden WB, Ramshaw IA. IFN-gamma plays a critical down-regulatory role in the induction and effector phase of myelin oligodendrocyte glycoprotein-induced autoimmune encephalomyelitis. *Journal of immunology*. 1996; 157:3223–3227.

- Wingerchuk DM. Current evidence and therapeutic strategies for multiple sclerosis. *Semin Neurol.* 2008; 28:56–68. [PubMed: 18256987]
- Wu C, Wang X, Gadina M, O'Shea JJ, Presky DH, Magram J. IL-12 receptor beta 2 (IL-12R beta 2)-deficient mice are defective in IL-12-mediated signaling despite the presence of high affinity IL-12 binding sites. *Journal of immunology.* 2000; 165:6221–6228.
- Wu F, Cao W, Yang Y, Liu A. Extensive infiltration of neutrophils in the acute phase of experimental autoimmune encephalomyelitis in C57BL/6 mice. *Histochemistry and cell biology.* 2010; 133:313–322. [PubMed: 20063008]
- Yang J, Yang M, Htut TM, Ouyang X, Hanidu A, Li X, et al. Epstein-Barr virus-induced gene 3 negatively regulates IL-17, IL-22 and RORgamma t. *European journal of immunology.* 2008; 38:1204–1214. [PubMed: 18412165]
- Zhang B, Zhang X, Tang FL, Zhu LP, Liu Y, Lipsky PE. Clinical significance of increased CD4+CD25-Foxp3+ T cells in patients with new-onset systemic lupus erythematosus. *Annals of the rheumatic diseases.* 2008; 67:1037–1040. [PubMed: 18199598]
- Zhang GX, Gran B, Yu S, Li J, Siglienti I, Chen X, et al. Induction of experimental autoimmune encephalomyelitis in IL-12 receptor-beta 2-deficient mice: IL-12 responsiveness is not required in the pathogenesis of inflammatory demyelination in the central nervous system. *Journal of immunology.* 2003; 170:2153–2160.

Figure 1. IL-12R β 2^{-/-} SJL/J mice were more susceptible to EAE

A) IL-12R β 2^{-/-} (n=10) and WT mice (n=10) were immunized with 100 μ g of PLP₁₃₉₋₁₅₁ in CFA containing 3 mg/ml *M. tuberculosis* H37Ra. Pertussis toxin (200 ng) was given i.p. on days 0 and 2 p.i. Data represent the mean clinical scores $\pm$ SEM. One of two experiments is shown.

B) IL-12R β 2^{-/-} (n=7) and WT mice (n=7) were immunized with 100 μ g of PLP₁₃₉₋₁₅₁ in CFA containing 3 mg/ml *M. tuberculosis* H37Ra. Pertussis toxin (200 ng) was given i.p. on days 0 and 2 p.i. Data represent the mean clinical scores $\pm$ SEM.

C) IL-12R β 2^{-/-} (n=8) and WT mice (n=8) were immunized with 50 μ g of PLP₁₃₉₋₁₅₁ in CFA containing 2 mg/ml *M. tuberculosis* H37Ra. Pertussis toxin (100 ng) was given i.p. on days 0 and 2 p.i. Data represent the mean clinical scores $\pm$ SEM. One of two experiments is shown.

D) IL-12R β 2^{-/-} and WT mice were immunized with 100 μ g of PLP₁₃₉₋₁₅₁ in CFA containing 3 mg/ml *M. tuberculosis* H37Ra. Pertussis toxin (200 ng) was given i.p. on days 0 and 2 p.i. Groups of mice (n=5–10, depending on the time point) were sacrificed at the time p.i. indicated in the figure. Spinal cords and brains from mice in the same group were pooled, MNCs isolated and their numbers determined by counting under the microscope.

Figure 2. IL-12Rβ2^{-/-} mice had a higher proportion of Th17 cells and lower proportion of Th1 cells among CD4⁺ T cells isolated from the CNS during EAE
 IL-12Rβ2^{-/-} and WT mice were immunized with 100 μg of PLP₁₃₉₋₁₅₁ in CFA containing 3 mg/ml *M. tuberculosis* H37Ra. Pertussis toxin (200 ng) was given i.p. on days 0 and 2 p.i. Groups of mice (n=5–10, depending on the time point) were sacrificed at the time p.i. indicated in the figure. Spinal cords and brains from mice in the same group were pooled, MNCs isolated, incubated with PMA, ionomycin and GolgiPlug, stained, and gated CD4⁺ cells analyzed by flow cytometry. Data shown are representative of 4 experiments.

Figure 3. Splenocytes of immunized IL-12R β 2^{-/-} EAE mice produced less IFN- γ than WT splenocytes

Splenocytes harvested from PLP₁₃₉₋₁₅₁-immunized mice on days 7, 11, 21, 28, 50 p.i. were cultured for 72 h with or without 20 μ g/ml PLP₁₃₉₋₁₅₁. Concentrations of IL-17A and IFN- γ in splenic culture supernatants were measured by ELISA. Data shown are from 3 independent experiments.

Figure 4. IL-12R β 2^{-/-} EAE mice developed more Treg cells in spleen
 Splenocytes from PLP₁₃₉₋₁₅₁-immunized mice on day 7, 11, 21, 28 p.i. were isolated and stained. Expression of CD25 and Fopx3 in gated CD4⁺ T cells were analyzed by flow cytometry. Data shown originate from 3 independent experiments.

Figure 5. IL-12Rβ2^{-/-} splenocytes had higher proliferative capacity

Splenocytes harvested from PLP₁₃₉₋₁₅₁-immunized mice on days 7, 11, 21, 28, 50 p.i. were cultured (4×10^5 cells/well) in triplicates for 72 h with 20 μg/ml PLP₁₃₉₋₁₅₁ or 1 μg/ml anti-CD3/CD28 antibodies. After 60 h, cells were pulsed for 12 h with 1 μCi of [³H] methylthymidine. Data shown are from 3 independent experiments.

Figure 6. IL-12Rβ2^{-/-} mice had a higher proportion of CD11b⁺Gr1⁺ cells in the CNS and spleen
A) MNCs from the CNS of EAE mice on days 11, 21, 28 and 50 p.i. were isolated and stained for Gr1 and CD11b.

B) Splenocytes from EAE mice on days 7, 11, 21, 28 and 50 p.i. were isolated, stained for Gr1 and CD11b and analyzed by flow cytometry. One representative of three experiments is shown.