Published in final edited form as:

Int Rev Neurobiol. 2016 ; 126: 179–261. doi:10.1016/bs.irm.2016.02.017.

A Genetic Animal Model of Alcoholism for Screening Medications to Treat Addiction

Richard L. Bell, Ph.D.^{a,*}, Sheketha Hauser, Ph.D.^a, Zachary A. Rodd, Ph.D.^a, Tiebing Liang, Ph.D.^b, Youssef Sari, Ph.D.^c, Jeanette McClintick, Ph.D.^d, Shafiqur Rahman, Ph.D.^e, and Eric A. Engleman, Ph.D.^a

^aDepartment of Psychiatry, Institute of Psychiatric Research, Indiana University School of Medicine, Indianapolis, Indiana, 46202, USA

^bDepartment of Medicine, Indiana University School of Medicine, Indianapolis, Indiana, 46202, USA

^cDepartment of Pharmacology, University of Toledo, Toledo, Ohio, 43614, USA

^dDepartment of Biochemistry & Molecular Biology, Center for Medical Genomics, Indiana University School of Medicine, Indianapolis, Indiana, 46202, USA

^eDepartment of Pharmaceutical Sciences, South Dakota State University, Brookings, SD 57007, USA

Abstract

The purpose of this review is to present up-to-date pharmacological, genetic and behavioral findings from the alcohol-preferring P rat and summarize similar past work. Behaviorally, the focus will be on how the P rat meets criteria put forth for a valid animal model of alcoholism with a highlight on its use as an animal model of polysubstance abuse, including alcohol, nicotine and psychostimulants. Pharmacologically and genetically, the focus will be on the neurotransmitter and neuropeptide systems that have received the most attention: cholinergic, dopaminergic, GABAergic, glutamatergic, serotonergic, noradrenergic, corticotrophin releasing hormone, opioid, and neuropeptide Y. Herein we sought to place the P rat's behavioral and neurochemical phenotypes, and to some extent its genotype, in the context of the clinical literature. After reviewing the findings thus far, this paper discusses future directions for expanding the use of this genetic animal model of alcoholism to identify molecular targets for treating drug addiction in general.

Keywords

Animal models; alcohol; drug addiction; dependence; pharmacogenetics; pharmacogenomics; CNS neurotransmitters; proteomics; selective breeding; genomics; family history; predisposition

*Address correspondence to Richard L. Bell, Ph.D., Indiana University School of Medicine, Department of Psychiatry, Institute of Psychiatric Research, Neuroscience Research Building (NB300), 320 W. 15th Street, Indianapolis, IN 46202-2266; TEL: 317-278-8407; FAX: 317-231-0202; ribell@iupui.edu (RLBell).

The authors do not have any conflicts of interest with this material.

1. Alcohol Abuse and Dependence

Over half of adult Americans have a family member with an alcohol abuse or dependence disorder [alcohol use disorder (AUD)], although only a subset of these individuals have this across multiple generations [Research Society on Alcoholism (RSA), 2009, 2015]. Moreover, approximately 1 in 4 Americans have had an AUD during their lifetime (RSA, 2009, 2015). The repercussions of AUDs cost the US economy nearly a quarter of a trillion dollars per year (RSA, 2015). Much of this is in health care costs, with AUDs being the third leading cause of preventable death according to the Centers for Disease Control and Prevention (CDC, Mokdad, Marks, Stroup, Gerberding, 2004; also see Johnson, 2010) and a causal relationship has been established between AUDs and at least 50 different medical conditions (Reed, Page, Viken, Christian, 1996; Rehm, 2011; Rehm et al., 2003).

Alcohol dependence is a chronic, progressive, relapsing disorder that advances in stages from experimentation to dependence (Heilig & Egli, 2006; Jupp & Lawrence, 2010; Koob, 2009; Koob & LeMoal, 2008; Koob & Volkow, 2010; Spanagel, 2009; Volkow & Li, 2005). During the experimentation stage, the individual experiences the rewarding, euphoric and positive-reinforcing effects of alcohol consumption. These positive reinforcing effects are often associated with acute increases in motor and autonomic (e.g., heart rate) activity as well as pro-social behavior and are generally perceived, by the individual, as euphoric (i.e., pleasant). This learning process results in positive reinforcement; which increases the probability, frequency and magnitude of subsequent drinking behavior. However, with continued usage the individual experiences an increase in the magnitude, duration and/or frequency of dysphoria (as opposed to euphoria), such as anxiety, during periods without access to alcohol. These dysphoric effects can be physiological in nature (e.g., hangover, hyperthermia, tachycardia, etc.) or associated with behavioral sequelae, such as getting arrested for driving while intoxicated. Moreover, given this increase in dysphoria, the individual often seeks to relieve this state by returning to drinking alcohol, often to excess. Therefore, during the early stages of the disease positive reinforcement generally predominates, whereas during the later stages of the disease negative reinforcement predominates (Koob, Arends, & Le Moal, 2014; Koob, Buck et al., 2014).

Fundamentally, reinforcement results in an increase in behavior and its associated cognitive processes. Thus, increases in approach behavior are associated with positive reinforcement during initial stages of the disease, whereas increases in retreat or avoidance behavior (e.g., consuming alcohol upon waking to counter hangover effects) are associated with negative reinforcement during later stages of the disease. The roles of positive-reinforcement vs negative-reinforcement can also be characterized in terms of impulsive vs compulsive alcohol drinking (Koob, Arends, & Le Moal, 2014; Koob, Buck, et al., 2014; Koob & Le Moal, 2006; Koob & Le Moal, 2008). Within this construct, impulsive drinking leads to, and is associated with, binge drinking and intoxication (e.g., Gray & MacKillop, 2014; Hamilton, Felton, Risco, Lejuez, MacPherson, 2014; but see Irimia et al., 2013). However after chronic usage, impulsive drinking, during which an individual will putatively have some volitional control, will be replaced by compulsive drinking to mitigate physical and behavioral withdrawal from alcohol. This, in turn, leads to a preoccupation with, and an anticipation of, future alcohol consumption during alcohol withdrawal (Koob & Le Moal,

2008). Therefore, impulsive drinking and positive reinforcement predominate in the early stages of alcohol dependence, whereas compulsive drinking and negative reinforcement predominate in later stages of addiction (Koob, Arends, & Le Moal, 2014; Koob, Buck, et al., 2014; Koob & Le Moal, 2006; Koob & Le Moal, 2008). However, despite this general trend of cycles of active drinking and relapse, with a concomitant increase in alcohol, or drugs of abuse, intake and the development of tolerance to the effects of alcohol, or drugs of abuse; the progression of the disease is not necessarily linear, such that the frequency and/or duration a person experiences these cycles differs across individuals (e.g., Barker & Taylor, 2014; Mackenzie, El-Gabalaway, Chou, & Sareen, 2014; Sartor, Kranzler, & Gelernter, 2014; van Rizen & Dishion, 2014). Moreover, not all individuals who abuse alcohol, or drugs of abuse, need formal treatment to reduce or stop their excessive intake. For instance, many individuals who abused alcohol during their adolescence and emerging adulthood do not develop alcohol dependence.

Nevertheless, many individuals who abused alcohol during adolescence and emerging adulthood do develop alcohol dependence either during this developmental stage or later in life. Because of this, there is a strong emphasis to study the acute and long-term effects of alcohol and/or drug abuse during the peri-adolescent developmental window (i.e., juvenile to emerging adulthood) (e.g., Bell, Franklin, Hauser, & Engleman, 2013; Gulley and Juraska, 2013; Spear, 2010, 2014; Witt, 1994, 2006, 2010). According to the National Institute on Alcohol Abuse and Alcoholism (NIAAA, 2012), 11% of all alcohol consumed in the U.S. is done so by 12 to 20 year olds and 90% of this drinking is in the form of binges. Additionally, about a third of high school seniors report binge drinking during high school and ~75% of college students report binge drinking (Johnston, O'Malley, & Bachman, 1999). For ~30% of male college students (Wechsler, Lee, Kuo, & Lee, 2000; White, Kraus, & Swartzwelder, 2006), this behavior continues into college and the magnitude of these binges often exceeds, by to 2- to 3-fold, threshold consumption levels [5 drinks in one sitting resulting in blood alcohol concentrations (BACs) of 80 mg% or higher] put forth in NIAAA's definition of binge drinking (NIAAA National Advisory Council, 2004). Thus, binge drinking during peri-adolescence has become a serious public health concern, with research indicating it is a strong predictor of future alcohol-related problems in North America (Dawson, Grant, Stinson, & Chou, 2004; Johnston, O'Malley, Bachman, & Schulenberg, 2008; Kuntsche, Rehm, & Gmel, 2004; Presley, Meilman, & Lyster, 1994; Wechsler et al., 2000; White et al., 2006).

Other predictors of AUDs as well as their epidemiological antecedents and trajectory are the pattern of drinking (e.g., social vs binge vs continuous) and amount consumed (Flory, Lynam, Milich, Leukefield, & Clayton, 2004; Heather, Tebbutt, Mattick, & Zamir, 1993; Lancaster, 1994; Shield, Rehm, Gmel, Rehm, & Allamani, 2013; Zucker, 1995). Characterization of these predictors and their antecedents has led to the development of different typologies and/or drinking profiles as well as subcategories of severity associated with a diagnosis of an AUD in the Diagnostic and Statistical Manual of Mental Disorders, 4th and 5th Editions (DSM-4, DSM-TR-4 and DSM-5, American Psychiatric Association, 1994, 2000, 2013; Babor et al., 1992; Cloninger, 1987; Conrod, Pihl, Stewart, & Dongier, 2000; Epstein, Kahler, McCrady, Lewis, & Lewis, 1995; Lesch & Walter 1996; Moss, Chen, & Yi, 2007; Prelipceanu & Mihailescu, 2005; Preuss, Watzke, & Wurst, 2014; Windle &

Scheidt, 2004; Zucker, 1987). It has also been shown that an individual's ranking within a particular typology predicts the efficacy of certain pharmacotherapies (Cherpitel, Moskalewicz, & Swiatkiewicz, 2004; Dundon, Lynch, Pettinati, & Lipkin, 2004; Epstein et al., 1995; Forray & Sofuoglu, 2014; Hulse, 2012; Johnson, 2005, 2010; Johnson, Ait-Daoud, Ma, & Wang, 2003; Keating, 2013). Therefore, age-of-onset and pattern of drinking, recognizing that these are often correlated, have significant predictive validity for a life-time diagnosis of alcohol abuse or dependence and, in some cases, the effectiveness of pharmacotherapies to treat alcohol dependence. In this chapter, alcohol and ethanol are used interchangeably; such that in the clinical setting the term alcohol is preferentially used over ethanol, whereas the term ethanol (scientific name for the two-carbon chain alcohol) is preferentially used over alcohol in the preclinical setting.

2. Neurobehavioral Correlates with Alcohol Abuse and Dependence

Clinical and basic research indicate that 1) lower responsivity to ethanol's effects is directly associated with alcohol abuse and dependence (e.g., Crabbe, Bell & Ehlers, 2010; Draski & Deitrich, 1996; Heit et al., 2013; Morean & Corbin, 2010; Morozova, Mackay & Anholt, 2014; Schuckit & Gold, 1988; Silveri, 2012, 2015; Spear, 2010, 2014); 2) the ability to display greater levels and quicker development of tolerance (a reduction in ethanol's effects after prior treatment with ethanol) to ethanol's effects is also associated with alcohol abuse and dependence (e.g., Lê & Mayer, 1996); 3) additionally, the expression of anxiety-like behavior under basal and/or withdrawal conditions is associated with a propensity to abuse alcohol (e.g., Heilig, Egli, Crabbe, & Becker, 2010; Heilig, Thorsell, et al., 2010; Kirby, Zeeb, & Winstanley, 2011; Pautassi, Camarini, Quadros, Miczek, & Israel, 2010; Thorsell, 2010); and 4) moreover, the expression of low- to moderate-dose ethanol-induced stimulation [which is modeled in rodents by increased motor activity/approach behavior (Chappell & Weiner, 2008; Faria et al., 2008; Wise & Bozarth, 1987), aggression (Chiavegatto, Quadros, Ambar, & Miczek, 2010), and social facilitation (Varlinskaya & Spear, 2009, 2010)] is associated with excessive alcohol consumption. This behavioral phenotype may have pharmacological validity as well, such that the histaminergic (c.f., Panula & Nuutinen, 2011 and references therein) and ghrelin (c.f., Jerlhag, Landgren, Eggecioglu, Dickson, & Engel, 2011b and references therein) systems have been implicated in ethanol-induced motor activation, ethanol-induced conditioned place preference, alcohol-preference and high alcohol consumption behavior. However, there remain concerns with the translatability of ethanol-induced stimulation in rodents vs humans (e.g., Crabbe et al., 2010). For instance, other than low- to moderate-dose effects on (a) self-report (Morzorati, Ramchandani, Flury, Li, & O'Connor, 2002; Viken, Rose, Morzorati, Christian, & 2003), (b) heart rate (Finn & Justus, 1997; Peterson et al., 1996), and (c) brain activity (Lukas, Mendelson, Benedikt, & Jones, 1986; Sorbel, Morzorati, O'Connor, Li, & Christian, 1996; Trim et al., 2010), the stimulating effects of ethanol are less apparent in humans compared with rodents.

3. Neurochemical Correlates with Alcohol Abuse and Dependence

Clinical and basic research indicate that alcohol abuse and dependence are mediated in part by a number of neurobiological systems (c.f., Koob, Arends, & Le Moal, 2014; Koob, Buck,

et al., 2014; Noronha, Cui, Harris, & Crabbe, 2014; Pierce & Kenny, 2013; Robbins, Everitt, & Nutt, 2010; Self & Staley, 2010; Sommer & Spanagel, 2013; Spanagel, 2009): acetylcholine (ACh: Chatterjee & Bartlett, 2010; Davis & de Fiebre, 2006; Rahman, Engleman, & Bell, 2015, 2016; Soderpalm, Ericson, Olausson, Blomqvist, & Engel, 2000), adenosine (Filip, Zaniewska, Frankowska, Wydra, & Fuxe, 2012; Nam, Bruner, & Choi, 2013), dopamine (DA: Bhaskar & Kumar, 2014; Engel & Jerlhag, 2014; Heinz, 2002; Nutt, Lingfor-Hughes, Erritzoe, & Stokes, 2015; Soderpalm & Ericson, 2013), endocannabinoid (Moreira, Jupp, Belin, & Dalley, 2015), gamma-aminobutyric-acid (GABA: Agabio & Colombo, 2014; Kumar et al., 2009; Liang & Olsen, 2014; Maccioni & Colombo, 2009), glutamate (Barron et al., 2012; Bell et al., 2016; Davis & Wu, 2001; Gass & Olive, 2008; Rao, Bell, Engleman, & Sari, 2015), purinergic (Franklin et al., 2014), serotonin (5-HT: Engleman, Rodd, Bell, & Murphy, 2008; Hauser et al., 2014; Lovinger, 1999), melanocortin (Olney, Navarro, & Thiele, 2014), opiate (Charbogne, Kieffer, Befort, 2014; Drews & Zimmer, 1997), orexin (Baimel et al., 2015), oxytocin (Buisman-Pijlman et al., 2014), neuropeptide-Y (NPY: Heilig & Thorsell, 2002), corticotropin releasing factor (CRF: Burke & Miczek, 2014; Koob, 2010), substance P (George et al., 2008), nociceptin/orphanin FQ (NOP, N/OFQ: Economidou et al., 2008; Witkin et al., 2014); ghrelin (Jerlhag, Egecioglu, Dickson, & Engel, 2011; Jerlhag et al., 2009; Jerlhag, Landgren, et al., 2011); neurotrophic factors such as BDNF (Logrip, Janak, & Ron, 2009), and hypothalamic-pituitary-adrenal (HPA) activity including corticosteroids, etc. (Gianoulakis, Guillaume, De Waele, & Angelogianni, 1995; Keith, Roberts, Wisen, & Crabbe, 1995; Kiefer, Jahn, Otte, Nakovics, & Wiedemann, 2006; Rasmussen, Boldt, Wilkinson, & Mitton, 2002; Richardson, Lee, O'Dell, Koob, & Rivier, 2008) systems within the brain. More recently there has been a significant increase in research on the neuroimmune system, which can modulate these neurochemical/neuropeptide systems and is an important contributor to the development of addiction (Crews, Qin, Sheedy, Ventreno, & Zou, 2013; Cui, Shurtleff, & Harris, 2014; Kane et al., 2014; Robinson et al., 2014; Ward, Lallemand, & de Witte, 2014). This is one area where basic and clinical research have informed each other, such that innate differences or ethanol-induced changes in neurotransmitter, neuropeptide and neuroimmune systems of subjects genetically predisposed for excessive ethanol consumption strongly suggests these systems play a significant role in the development of alcoholism, at least in predisposed individuals (c.f., Bell et al., 2012; Robinson et al., 2014). By extension, demonstration of these innate neurotransmitter and/or neuropeptide differences in an animal model of alcoholism underscore its utility in screening the efficacy of compounds to treat alcohol dependence.

5. The Genetics of Alcoholism

Family History Positive (FHP) for alcoholism individuals are persons who have relatives that meet diagnostic criteria for AUDs. The strongest expression of correlative phenotypes (discussed later) is observed in FHP individuals with this characteristic across multiple generations, starting with the immediate family. These findings provide strong support for genetics as a mediator in the development of alcoholism (c.f., Schuckit, 2014). Epidemiologically, twin studies have yielded data indicating a strong genetic component to the development and expression of alcohol dependence. Heath (1995) provides an excellent

overview of the early Australian (e.g., Eysenck & Eysenck, 1975; Martin et al., 1985), Finnish (e.g., Kaprio, Rose, Romanov, & Koskenvuo, 1991; Partanen, Bruun, & Markkanen, 1966), London Twin Family Survey (e.g., Clifford, Hopper, Fulker, & Murray, 1984), Swedish (e.g., Cederlof, Friberg, & Lundman, 1977), US National Academy of Science/ National Research Council twin registry (e.g., Hrubec & Neel, 1978; Jablon, Neel, Gershowitz, & Atkinson, 1967), the US Vietnam Era Twin Survey (e.g., Goldberg, Eisen, True, & Henderson, 1990) and his own work with the US Virginia 30,000 Survey. What is most striking about his review of these twin registries and AUDs is that, while the US data tends to suggest that the development of AUDs in men is genetically influenced to a greater extent than that observed in women, multiple European studies suggest that the development of AUDs by women is genetically influenced to the same degree or more so than men (Heath, 1995). For further reading on the genetics of AUDs, see Hesselbrock (1995b) for an overview of early adoption studies, see Hesselbrock (1995a) for an overview of early work looking at Alcoholic Subtypes, see Cadoret (1995) for an overview of early studies on genetic correlates with other psychiatric disorders and see Schuckit (2014) for a recent review on the history of research investigating the genetics of alcohol and drug dependence.

6. Criteria for an Animal Model of Alcoholism

Different animal models have had different levels of success in research to develop treatments for both medical and psychiatric disorders (Gobrogge, 2014; Golbidi, Frisbee, & Laher, 2015; Griffin, 2002; McCairn & Isoda, 2013; McGonigle & Ruggeri, 2014; McKinney, 2001; McLarnon, 2014; Reser, 2014; Nestler & Hyman, 2010; Whiteside, Pomonis, & Kennedy, 2013). An animal model allows an experimenter to control the subject's genetic background, environmental factors and prior drug experience. In addition, it allows for the examination of neurobehavioral, neurochemical and neurophysiological correlates associated with the behavioral, physiological and neurological states being modeled, in the present case alcohol abuse and dependence. These correlates can, in turn, facilitate the development of pharmacological and/or behavioral treatments for these disorders. There have been reservations as to whether a valid animal model of alcoholism could be developed (Cicero, 1979). These concerns stemmed from the fact that, in general, heterogeneous stock rats consume only modest levels of ethanol, such that blood alcohol concentrations (BACs) achieved are modest (c.f., Bell, Rodd, Engleman, Toalston, & McBride, 2014 for a comparison of 22 different rat lines and strains). Nevertheless, certain criteria for an animal model of alcoholism have been put forth (Cicero, 1979; Lester & Freed, 1973). Briefly, these criteria are 1) the animal should readily consume ethanol under free-choice access conditions; 2) the amount of ethanol consumed should result in pharmacologically relevant BACs; 3) ethanol should be consumed for its pharmacological effects; 4) ethanol should be reinforcing, usually demonstrated through operant procedures; 5) chronic ethanol consumption should lead to the expression of metabolic and functional/ neuronal tolerance to alcohol's effects; and 6) chronic consumption should lead to dependence, as indicated by withdrawal signs after access to ethanol is terminated. Other proposed criteria for a valid animal model of alcoholism include displaying characteristics associated with relapse-like drinking, generally demonstrated by an alcohol deprivation effect (ADE: McBride & Li, 1998); as well as excessive ethanol drinking during

adolescence, such that it exceeds adult intake levels and binge-like access results in BACs > 80 mg% as well as motor impairment (Bell et al., 2011, 2013, 2014; McBride, Rodd, Bell, Lumeng, & Li, 2014).

7. A Rat Genetic Animal Model of Alcoholism

The well-documented familial incidence of alcoholism, including multiple international twin studies, indicates a strong (40–70%) genetic component (discussed above) mediates a predisposition for and the development of alcohol use disorders (AUDs) (e.g., Cloninger, 1987; Cotton, 1979; Hesselbrock, 1995a, 1995b; Schuckit, 1986). Given heterogeneous stock rats display a wide-range of ethanol-consumption levels (Richter & Campbell, 1940) and very early work on selective breeding for alcohol consumption, Williams and associates (Williams, Berry, & Beerstecher, 1949) as well as Mardones and colleagues (Mardones, Segovia, & Hederra, 1953; Mardones & Segovia-Riquelme, 1983) proposed a genetic link to ethanol intake in rodents. From their work (e.g., Mardones et al., 1953; Mardones & Segovia-Riquelme, 1983) and that of four other international sites, bidirectional selective breeding has resulted in at least six high alcohol-consuming vs. their respective low alcohol-consuming rat lines (c.f., Bell et al., 2012).

Bi-directional selective breeding is a powerful genetic tool for studying alcohol-associated phenotypes (e.g., Crabbe, 2008). Compared to pure association studies such as genome-wide association studies (GWAS) and recombinant inbred lines (RILs), selective breeding from a heterogeneous outbred stock can make low frequency/rare alleles (minor allele frequency <0.05) more common by segregating these genetic polymorphisms into the opposite extremes of the overall population. This bidirectional selection is accomplished through systematic mating of animals with similar traits (alcohol-preferring on the one hand vs. alcohol-avoiding on the other) over successive generations. Thus, the high and low lines will exhibit extreme phenotypes exceeding the range found in the original parent population. Additionally, selective breeding for any phenotypes (such as alcohol preference) is hypothesis driven and genetically correlated traits of the primary selected phenotype (presumably due to pleiotropic actions of genes: Crabbe, Phillips, Kosobud, & Belknap, 1990) can be identified for further study. A major advantage of this selection process is that the ethanol-drinking phenotype is observed without the stress of environmental manipulations.

8. The P Rat as a Genetic Animal Model of Alcoholism

The alcohol-preferring, P, and alcohol-nonpreferring, NP, rat lines were developed by bi-directional mass selection from a closed-colony of Wistar rats at the Walter Reed Army Hospital and subsequently transferred to the Indiana University School of Medicine, Indianapolis, Indiana, USA (Lumeng, Hawkins, & Li, 1977). Two metrics were used for the selection of an alcohol preference. First, the animals had to prefer an unadulterated 10% ethanol solution over water by a ratio of at least 2:1. Second, the animals had to consume at least 5 g of ethanol/kg body weight/day. The first criterion was used to prevent selecting for an artifact of body weight when using the second criterion (selection for lower body weight will appear to be equivalent to a selection for high ethanol consumption only). To place this

5 g/kg/day selection criterion in a clinical perspective, using 0.793 as the specific gravity of ethanol, 90-proof whiskey as the preferred alcoholic beverage of choice for a 70 kg male, and the fact that rats metabolize ethanol close to 1.4 times the rate of humans, 5 g/kg/day of ethanol intake by a rat would be approximately a fifth of 90-proof whiskey being consumed each day.

Regarding criteria for an animal model of alcoholism, under free-choice conditions (water and food available) P rats readily consume greater than 5 g/kg/day of ethanol, whereas NP rats consume less than 1 g/kg/day (Li, Lumeng, McBride, & Murphy, 1987). P rats readily attain pharmacologically relevant blood alcohol concentrations (BACs, 80 to 250 mg%: this would approximate 0.08 to 0.25 in forensic terms, the latter under operant binge-like conditions) (Bell et al., 2013, 2014, 2011; Bell, Rodd, Lumeng, Murphy, & McBride, 2006; Bell, Rodd, Sable, et al., 2006; Bell, Rodd, Schultz, et al., 2008; McBride, Kimpel, McClintick, Ding, Hauser, et al., 2013; McBride, Kimpel, et al., 2014; McBride, Rodd, et al., 2014; Murphy et al., 2002). Ethanol-drinking in the home-cage also results in intoxication including motor impairment, as measured with an oscillating bar apparatus (Bell, McKinzie, Murphy, & McBride, 2000; Bell et al., 2001). Moreover, P rats will display tolerance to this effect after chronic binge drinking (Bell et al., 2011). P rats operantly self-administer ethanol intragastrically for its post-ingestive effects (Murphy et al., 1988; Waller, McBride, Gatto, Lumeng, & Li, 1984). P rats will operantly self-administer ethanol using a dipper model, indicating these rats will work for access to ethanol (Files, Samson, Denning, & Marvin, 1998; Murphy, Gatto, McBride, Lumeng, & Li, 1989; Rodd et al., 2003; Rodd-Henricks et al., 2002a, 2002b; Samson, Files, Denning, & Marvin, 1998; Toalston et al., 2008) or sipper tube model (Beckwith & Czachowski, 2014; Bertholomey, Verplaetse, & Czachowski, 2013; Czachowski & Samson, 2002; Samson & Czachowski, 2003; Verplaetse, Rasmussen, Froehlich, & Czachowski, 2012; Verplaetse & Czachowski, 2015) indicating these rats will work for access to ethanol.

Whereas ethanol-naïve P and NP rats display similar levels of ethanol clearance (Li & Lumeng, 1977; Lumeng, Waller, McBride, & Li, 1982), after chronic ethanol-drinking (6–8 weeks) P rats display metabolic as well as functional tolerance to the motor impairing and aversive effects of ethanol (Gatto et al., 1987; Lumeng & Li, 1986; Stewart, McBride, Lumeng, Li, & Murphy, 1991). Moreover, similarly treated P rats also display dependence-associated signs when ethanol access is terminated (Kampov-Polevoy, Matthews, Gause, Morrow, & Overstreet, 2000; Waller, McBride, Lumeng, & Li, 1982). In addition, P rats display relapse-like drinking by exhibiting a robust alcohol deprivation effect (ADE: Rodd et al., 2003; Rodd-Henricks et al., 2001; Rodd-Henricks, McKinzie, Shaikh, et al., 2000). The ADE is a transient increase in ethanol intake after a period of ethanol withdrawal (c.f., Rodd, Bell, McKinzie, et al., 2004; Rodd, Bell, Sable, Murphy, & McBride, 2004; Vengeliene, Bilbao, & Spanagel, 2014). Regarding initial sensitivity, compared with NP rats, P rats are less sensitive to the ataxic (Bell et al., 2001) and hypothermic (Stewart, Kurtz, Zweifel, Li, & Froehlich, 1992) effects of ethanol; and P rats develop tolerance quicker to the ataxic (Bell et al., 2001) and hypnotic (Kurtz, Stewart, Zweifel, Li, & Froehlich, 1996) effects of ethanol as well. During ethanol withdrawal, P, but not NP, rats display greater acoustic startle reactivity compared with basal conditions (Chester, Blose, & Froehlich, 2004). P rats display greater low dose ethanol-induced locomotor activity, compared with NP rats (Rodd,

Bell, McKinzie, et al., 2004; Waller, Murphy, McBride, Lumeng, & Li, 1986), and display locomotor activation during ethanol drinking or self-administration (Bell, Rodd, Toalston, et al., 2008; Bell et al., 2002; Melendez et al., 2002). The latter provides additional support to the view that P rats find ethanol rewarding. Thus, the P line of rats satisfies criteria proposed by multiple authors for a valid animal model of alcoholism. The key behavioral features of this genetic rat animal model of alcoholism are outlined below:

Key features of the P genetic rat animal model of alcoholism

1. >7 g ethanol/kg body weight/day is orally consumed under home-cage, free-choice, 24 hr conditions (Bell et al., 2011, 2013, 2014; Bell, Rodd, Schultz, et al., 2008; Li et al., 1987)
2. >1 g/kg ethanol is orally consumed during the first 15-min of home-cage, limited access conditions (Bell et al., 2014; Bell, Rodd, Lumeng, et al., 2006; Murphy et al., 1986; Russell, McBride, Lumeng, Li, & Murphy, 1996)
3. Pharmacologically relevant blood alcohol concentrations (BACs: 80 to 250 mg%), which parallel those observed in alcoholics, are achieved during ethanol drinking and self-administration (Bell et al., 2011, 2014; Bell, Rodd, Lumeng, et al., 2006; Bell, Rodd, Sable, et al., 2006; Bell, Rodd, Schultz, et al., 2008; Murphy et al., 1986, 2002; Rodd et al., 2003)
4. Ethanol is self-administered intragastrically (Murphy et al., 1988; Waller et al., 1984) and intracranially (Engleman et al., 2009; Toalston et al., 2014) indicating taste and calories are not the primary motivators for this behavior
5. Ethanol is consumed and self-administered despite ethanol-induced motor impairment (Bell et al., 2011; McBride, Kimpel, McClintick, Ding, Hauser, et al., 2013)
6. Ethanol drinking and self-administration induce autonomic (heart rate) and/or behavioral (motor) activation (Bell, Rodd, Toalston, et al., 2008; Bell et al., 2002; Melendez et al., 2002)
7. Ethanol is operantly self-administered, without using fading/adaptation techniques (i.e., P rats work for access to ethanol) (Files et al., 1998; Murphy et al., 1989; Rodd et al., 2003; Rodd-Henricks et al., 2002a, 2002b; Rodd-Henricks, McKinzie, Shaikh, et al., 2000; Samson et al., 1998; Toalston et al., 2008)
8. Chronic home-cage, free-choice ethanol consumption leads to metabolic (Lumeng & Li, 1986) and functional (Gatto et al., 1987; Stewart et al., 1991; 1996) tolerance
9. Chronic home-cage, ethanol consumption leads to dependence (e.g., decreased seizure thresholds) (Kampov-Polevoy et al., 2000; Waller et al., 1982)
10. Relapse behavior is displayed under home-cage and operant conditions [e.g., expression of an alcohol deprivation effect (ADE) is observed] (Rodd et al., 2003; Rodd-Henricks et al., 2001; Rodd-Henricks, McKinzie, Shaikh, et al., 2000)
11. Serve as an animal model of adolescent binge alcohol abuse by exceeding ethanol intakes seen during adulthood, with motor impairment and BACs exceeding 80 mg

%, under both continuous and limited access conditions (Bell et al., 2011, 2014; McBride et al., 2005; McBride, Rodd, et al., 2014)

12. Self-administer or consume drugs of abuse by (a) orally consuming nicotine-adulterated solutions (Hauser et al., 2012); (b) operantly self-administering nicotine both orally (Hauser et al., 2014) and intravenously (Le et al., 2006); and (c) intracranially self-administering cocaine (Katner et al., 2011), nicotine (Deehan et al., 2015; Hauser et al., 2014), and nicotine + ethanol (Truitt et al., 2015).

9. Some Family History Positive (FHP) Correlates

The above survey of the literature concerning neurobehavioral and neurobiological phenotypes found in these selectively bred high vs. low alcohol-consuming rat lines indicates that many phenotypes present in alcohol abusing or dependent individuals are also present in these lines. For example, similar to the animal literature, clinical studies of FHP subjects (i.e., with a family history of alcoholism) report an inverse relationship between low level responsivity to ethanol and risk for the development of AUDs (c.f., Crabbe et al., 2010; Schuckit, 1994, 2009, 2014). Thus, after an ethanol challenge young adult FHP females (Eng et al., 2005; Lex et al., 1988) and males (Schuckit, 1985; Schuckit & Gold, 1988) display less body sway than family history negative (FHN) controls. An individual's level of response (low vs. high) to ethanol also influences brain regional activation following an acute ethanol challenge (Trim et al., 2010). Moreover, an individual's level of response to ethanol is associated with the long- vs. short-allele for the 5-HT transporter (*5htt*) gene and this association has significant predictive validity for alcohol intake by adolescents (Hinckers et al., 2006). As discussed earlier, the FHP, P, rat displays lower levels of ethanol-induced behavioral and physiological changes compared with FHN, NP, rats (Bell et al., 2001; Stewart et al., 1992).

Alcohol dependence is a chronic relapsing disorder, with craving and relapse often precipitated by physiological and behavioral responses (i.e., cue-reactivity) to environmental and interoceptive cues associated with alcohol and/or drugs of abuse consumption (Childress et al., 1993; Drummond et al., 1990; Greeley et al., 1993; Kaplan et al., 1983, 1985; O'Brien et al., 1992; Rajan et al., 1998; Stormark et al., 1998). In addition, ethanol's effects on heart rate-reactivity are associated with level of genetic density in FHP individuals as well dissociating FHP from FHN subjects, such that persons that are FHP display greater sensitivity to ethanol reward and display sensation-seeking behavior (Assaad et al., 2003; Conrod, Pihl, & Vassileva, 1998; Finn, Earleywine, & Pihl, 1992; Peterson, Pihl, Seguin, Finn, & Stewart, 1993). This has also been shown in cross-sensitivity by stimulant users displaying a characteristic alcohol-reward heart rate response (Brunelle, Barrett, & Pihl, 2006). It is noteworthy that both male (Bell et al., 2002) and female (Bell, Rodd, Toalston, et al., 2008) P rats display increased heart rate during ethanol drinking. In addition, this autonomic reactivity can be conditioned to the environment associated with ethanol consumption (Bell, Rodd, Toalston, et al., 2008; Bell et al., 2002). Thus, monitoring autonomic reactivity in P rats may be an important model system for testing compounds targeting craving, especially in the context of cue-reactivity.

10. Some Neurochemical, Neuropharmacological as well as Neurogenetic Correlates

The Cholinergic System

Acetylcholine is released from neurons projecting to a broad range of cortical and subcortical structures and influences cellular physiology and neuronal function throughout the brain (Newman, Gupta, Climer, Monaghan, & Hasselmo, 2012). There are two classifications of cholinergic receptors: nicotinic and muscarinic. The neuronal nicotinic acetylcholine receptors (nAChRs) belong to the family of ligand-gated ion channel receptors (Albuquerque, Pereira, Alkondon, & Rogers, 2009; Gotti et al., 2009). Nicotinic acetylcholine receptors (nAChRs) consist of 11 neuronal subunits, which are divided into 8 alpha subunits ($\alpha 2$ – $\alpha 10$) and 3 beta subunits ($\beta 2$ – $\beta 4$). nAChR subtypes with diverse subunit combinations are distributed across multiple brain regions, including the mesocorticolimbic and extended amygdala reward circuitry, where they regulate dopaminergic function. A notable difference among the receptor subtypes is that the homomeric $\alpha 7$ nAChR does not desensitize to nicotine stimulation as the heteromeric nAChRs (e.g., $\alpha 4\beta 2$ nAChR) do. The $\alpha 7$ nAChRs are located presynaptically on glutamatergic projections from the mPFC to the NAcSh. Therefore, activation $\alpha 7$ nAChRs may enhance glutamatergic excitatory drive and promote DA release in the NAc after $\alpha 4\beta 2$ receptors are desensitized. Muscarinic acetylcholine receptors (mAChRs) are G-protein coupled receptors that are widely distributed in the brain. There are 5 mAChRs neuronal subunits (M1–M5) (Bymaster, McKinzie, Felder, & Wess, 2003; Wess, 2003; Wess et al., 2003; Yamada et al., 2003). Studies examining the striatum indicated that M1 receptors are expressed on spiny projection neurons (Wang et al 2006), whereas M2/M4 receptors are primarily presynaptic autoreceptors (Yan & Surmeier, 1996; Zhang et al., 2002). As autoreceptors, M2/M4 receptors inhibit ACh release and subsequent nAChR-dependent DA release in the striatum (Shin et al., 2015). In addition, M2 receptors are located on glutamatergic terminals which inhibit its release (Hersch, Gutekunst, Rees, Heilman, & Levey, 1994). The M5 receptor is the only mAChR subtype found on midbrain DA neurons (Vilaró et al., 1990; Palacios, & Mengod, 1990; Weiner, Levey, & Brann, 1990) and these receptors modulate DA and DA/ glutamate projections from the midbrain (Shin et al., 2015).

Substantial research indicates that nAChR activity mediates, in part, the rewarding effects of drugs of abuse (Chatterjee & Bartlett, 2010; Corrigan & Coen, 1994; Ericson, Blomqvist, Engel, & Soderpalm, 1998; Hendrickson, Guildford, & Tapper, 2013; McGehee & Role, 1995; Nisell, Nomikos, & Svensson 1994; Rahman, 2013; Rahman et al., 2015, 2016; Rahman & Prendergast, 2012; Sajja, Dwivedi, & Rahman, 2010). For example, both alcohol (Brodie, Pesold, & Appel, 1990; Brodie, Shefner, & Dunwiddie, 1999) and nicotine (Calabresi, Lacey, & North, 1989; Nisell et al., 1994) activate VTA DA projection neurons and stimulation of nAChRs within the VTA modulate, at least in part, the reinforcing effects of nicotine (Corrigan & Coen, 1994; Nisell et al., 1994) and alcohol (Blomqvist, Ericson, Johnson, Engel, & Soderpalm, 1996; Ericson, Molander, Lof, Engel, & Soderpalm, 2003; Soderpalm et al., 2000). Additionally, a number of reports indicate that pre-exposure to nicotine significantly increases operant or free-choice ethanol self-administration and reinstates ethanol-seeking behavior in animal models (Bito-Onon, Simms, Chatterjee,

Holgate, & Bartlett, 2011; Hauser et al., 2012; Le et al., 2003). Moreover, co-administration of ethanol and nicotine produces an additive effect on their reinforcing effects and associated dopamine release in the nucleus accumbens (NAcb, Ericson, Lof, Stomberg, & Soderpalm, 2009; Tizabi, Bai, Copeland, & Taylor, 2007; Sajja et al., 2010).

Emerging preclinical evidence indicates that a number of ligands targeting nAChRs modulate ethanol drinking behavior. For example, mecamylamine, a non-selective nAChR antagonist or varenicline, a partial agonist at $\alpha 4\beta 2^*$ nAChRs, reduces ethanol drinking behavior by targeting nAChRs in the mesocorticolimbic dopamine system (Ericson et al., 1998; Le et al., 2000; Steensland, Simms, Holgate, Richards, & Bartlett, 2007). In addition, nAChR ligands such as cytisine, a partial agonist at $\alpha 4\beta 2^*$, and lobeline, a non-selective antagonist, were found to reduce ethanol self-administration or nicotine-induced ethanol drinking in rodents (Bell, Eiler, Cook, & Rahman, 2009; Chatterjee, Steensland, Rollema, & Bartlett, 2011; Hendrickson, Zhao-Shea, & Tapper, 2009; Sajja & Rahman, 2011, 2012, 2013). Overall, brain nAChRs have emerged as important therapeutic targets for the rewarding effects of ethanol in numerous animal models. Regarding P rats, sazetidine-A, a novel ligand that desensitizes $\alpha 4\beta 2$ nAChRs with partial agonistic activity reduces ethanol drinking by P rats (Rezvani et al., 2010). In addition, pretreatment with nicotine increases operant ethanol-self administration and relapse behavior in P rats (Hauser et al., 2012). Furthermore, P rats have higher sensitivity to the reinforcing effects of nicotine in the pVTA compared with outbred Wistar rats (Hauser et al., 2014). These reinforcing effects of nicotine can be blocked by mecamylamine, an nAChR antagonist, when injected simultaneously into the pVTA of P rats (Hauser et al., 2014), indicating that mesolimbic cholinergic activity modulates the reinforcing effects of ethanol and nicotine. More recent findings indicate that P rats will also co-administer ethanol + nicotine into the pVTA (Truitt et al., 2015) and self-administer nicotine into the NAcbSh (Deehan et al., 2015). In addition, oral binge ethanol + nicotine self-administration by P rats induces greater sensitivity to the reinforcing effects of nicotine in the NAcbSh by shifting the self-administration dose-response curve to the left (Deehan et al., 2015).

Receptor binding studies have shown that P rats have lower striatal $\alpha 7$ nAChR expression than NP rats (Tizabi et al., 2001). Additionally, there is higher expression of *Chat* (choline acetyltransferase), *Chrm3* (mACh3R), *Slc5a7* (transporter uptake for acetylcholine synthesis), *Slc18a3* (vesicular amine transport into secretory vesicles) in the NAcbSh of adult P rats compared with NP rats, whereas adult P rats have lower expression of *Chrm4* (mACh4R) in the NAcbSh than NP rats (McBride, Kimpel, McClintick, Ding, Hauser, et al., 2013; McBride, Kimpel, McClintick, Ding, Hyytia, et al., 2013). These findings suggest that the accumbal cholinergic system may be more active in P rats than NP rats. Clinical research indicates individuals with AUDs have lower ChAT activity and mAChR density in the hippocampus (Antuono, Sorbi, Bracco, Fusco, & Amaducci, 1980; Nordberg, Larsson, Perdahl, & Winblad, 1983) as well as ChAT protein expression in the basal forebrain (Ventreno, Broadwater, Liu, Spear, & Crews, 2014). However, these clinical findings do not necessarily indicate that basal levels are different from control subjects. Our microarray findings indicate that ethanol drinking by adult P rats increased *Chrm7a* expression in the NAcb (Bell, Kimpel, et al., 2009), but reduced its expression in the CeA (McBride et al., 2010) and the DRN of ethanol-drinking adolescent P rats (McClintick et al., 2015). Overall,

a modest number of differences in ACh--associated gene expression levels have been detected between P and NP rats (data for the pVTA came from McBride et al., 2012; data for the CeA and NAcSh came from McBride, Kimpel, McClintick, Ding, Hyytia, et al., 2013). Similarly, few changes in ACh-associated gene expression levels have been detected following binge-drinking in adult or adolescent P rats (adult data for the whole NAc came from Rodd et al., 2008; adult data for the NAcSh came from Bell, Kimpel, et al., 2009, McBride et al., 2010; adult data for the CeA came from McBride et al., 2010; adult data for the pVTA came from McBride, Kimpel, McClintick, Ding, Hyytia, et al., 2013; adolescent data for the CeA came from McBride, Kimpel, et al., 2014; adolescent data for the DRN came from McClintick et al., 2015) thus far (Figure 1). Taken together, these data indicate that brain cholinergic activity regulates behaviors associated with alcohol abuse and alcoholism and the P rat displays some treatment characteristics seen in the clinical population. For instance, varenicline has modest effects on reducing ethanol intake by P rats. However, the tests done in P rats thus far have not incorporated ethanol and nicotine co-abuse, which appears to be necessary to observe efficacy with varenicline in the clinical setting.

The Dopaminergic System

DAergic projections (Figure 2) emanating from the tegmentum [especially the ventral tegmental area (VTA)] and terminating in limbic, forebrain and cortical regions (e.g., the mesocorticolimbic and extended amygdala reward circuitry), are involved in the appetitive and consummatory behaviors associated with, as well as the positive and negative reinforcing properties of, addictive drugs. DA activates metabotropic receptors (D_{1-5}) that are generally classified as D_1 -like (D_1 and D_5) and D_2 -like (D_{2-4}) (Cooper, Bloom, & Roth, 2002). D_1 -like receptors are typically coupled to the activation of adenylyl cyclase whereas D_2 -like receptors are coupled to the inhibition of adenylyl cyclase (Cooper et al., 2002). DA neurotransmission is terminated via clearance by uptake through the high affinity DA transporter (DAT) (Cooper et al., 2002). The DAergic system plays a central role in the processing of natural-, alcohol- and drug of abuse-associated reward and reinforcement (e.g., Di Chiara & Imperato, 1988; McBride & Li, 1998; Melendez et al., 2002; Nogueira, Kalivas, & Lavin, 2006; Palmer, Low, Grandy, & Phillips, 2003; Volkow & Morales, 2015), with imaging studies indicating $D_2/3$ receptor dysfunction in subjects addicted to several different drugs of abuse (Cosgrove, 2010). When an individual or animal ingests alcohol and/or other drugs of abuse, DA efflux is increased in several key mesocorticolimbic brain reward centers (e.g., Brodie et al., 1990; Franklin et al., 2009; Gessa, Muntoni, Collu, Vargiu, & Mereu, 1985; Imperato & Di Chiara, 1986; Smith & Weiss, 1999; Yoshimoto & McBride, 1992). This change in DAergic activity and other associated neuroplastic changes promote further alcohol and drug taking behavior. Following prolonged alcohol and/or drug abuse, individuals may display tolerance to these DA-elevating properties. Thus, the requirement for more alcohol and/or drug taking to get the same level of initial response moves the individual further through the addiction cycle, which starts out as impulsive use for intoxication and progresses to compulsive use to avoid or reduce the negative consequences of alcohol/drug withdrawal.

Regarding P rats, these animals exhibit reduced basal NAc tissue DA levels and/or enhanced ethanol-induced extracellular DA efflux, compared to outbred rats or their alcohol non-preferring counterparts (Engleman, Ingraham, McBride, Lumeng, & Murphy, 2006; McBride, Chernet, Dyr, Lumeng, & Li, 1993; Murphy, McBride, Lumeng, & Li, 1982, 1987; Smith & Weiss, 1999; Strother, Lumeng, Li, & McBride, 2005; also see Bell et al., 2012b; Murphy et al., 2002 for reviews). DAergic neuronal activity from the VTA appears to play a major role in these alterations of mesocorticolimbic DAergic tone either basally or under ethanol-induced conditions (e.g., Engleman et al., 2011; Morzorati, 1998; Morzorati & Marunde, 2006; Morzorati, Marunde, & Downey, 2010), which is modulated in part by glutamatergic activity in the pVTA (Fitzgerald, Liu, & Morzorati, 2012). In addition, chronic or binge-like alcohol drinking by P rats reduces D₂ autoreceptor function (Engleman, McBride, Li, Lumeng, & Murphy, 2003; Engleman et al., 2000), elevates extracellular DA levels (Thielen et al., 2004) and increases DA reuptake (Sahr et al., 2004) in the NAc. Sari and colleagues (2006) also reported that long-term ethanol consumption by P rats increased D₁ and D₂ expression levels in the NAc core (NAcCo) with D₂ expression also increased in NAc shell (NAcSh). These authors indicated that intermittent periods of ethanol deprivation increased D₁ receptor expression in the amygdala and D₂ receptor expression in the caudate putamen as well.

Given the evidence for differences in, and ethanol-induced changes of, DAergic activity, it is not surprising that a number of DA-associated compounds have been tested for their effects on ethanol drinking or self-administration by this line. Systemic administration of the DA agonist ibogaine (Rezvani et al., 1995) and DAT inverse modulator amphetamine (McBride, Murphy, Lumeng, & McBride, 1990) reduced ethanol drinking. Ethanol drinking by P rats also was disrupted by systemic administration of the D₂ agonist bromocriptine (Mason et al., 1994; McBride et al., 1990; Weiss et al., 1990) and the D₃ agonist 7-OH-DPAT (Mason et al., 1997) as well as intra-VTA infusion of the D₂ agonists quinpirole (Hauser et al., 2011; Nowak, McBride, Lumeng, Li, & Murphy, 2000) and quinlorane (Nowak et al., 2000). Both systemic administration (Mason et al., 1997) and intra-bed nucleus of the stria terminalis (BNST) infusion (Eiler, Seyoum, Foster, Mailey, & June, 2003) of the D₁ antagonist SCH23390 reduced ethanol drinking and self-administration by P rats. Similarly, both systemic administration (Mason et al., 1997) and intra-BNST as well as intra-VTA infusion (Eiler et al., 2003) of the D₂ antagonist eticlopride reduced ethanol drinking and self-administration by P rats. Additionally, ethanol drinking by P rats was reduced by microinfusion of the D₂ antagonist sulpiride into the VTA (Nowak et al., 2000), NAc (Levy et al., 1991) and ventral pallidum (Melendez et al., 2005). Finally, systemic administration of the D₃ antagonist SB-277011-A (Thanos et al., 2005) as well as the DAT inhibitors GBR 12909 (McBride et al., 1990) and DOV 102,677 (Yang et al., 2012) reduced ethanol intake by this line of rats.

It is noteworthy that P rats have lower levels of D₂ receptors than NP rats in the VTA, NAc and caudate putamen (McBride, Chernet, Dyr, et al., 1993; Strother, Lumeng, Li, & McBride, 2003) and smaller subpopulations of DA projections from the VTA compared with NP rats (Zhou, Zhang, Lumeng, & Li, 1995), but no differences in D₁ or D₃ receptor levels have been detected thus far (McBride et al., 1997). Finally, differences in both mRNA and protein expression levels of alpha-synuclein (SNCA, often associated with DAergic

function) have been found between P and NP rats, such that hippocampal levels were twice as high in inbred P rats compared with inbred NP rats (Liang et al., 2003). Interestingly, clinical research indicates that *Snca* expression is associated with alcohol craving, hazardous alcohol-drinking and Post Traumatic Stress Disorder (PTSD) symptomology in hazardous drinkers (Foroud et al., 2007; Guillot, Fanning, Liang, Leventhal, & Berman, 2015; Guillot, Pang, Leventhal, Liang, & Berman, 2015). Regarding gene expression differences, similar to the cholinergic system, only a modest number of DA-associated differences have been detected between P and NP rats so far (data for the pVTA came from McBride et al., 2012; data for the CeA and NAcSh came from McBride, Kimple, McClintick, Ding, Hyytia, et al., 2013). This paucity of findings was also true for DA-associated changes following ethanol drinking, usually binge-like, by P rats (adult data for the whole NAc came from Rodd et al., 2008; adult data for the NAcSh came from Bell, Kimpel, et al., 2009; McBride et al., 2010; adult data for the CeA came from McBride et al., 2010; adult data for the pVTA came from McBride, Kimpel, McClintick, Ding, Hauser, et al., 2013; adolescent data for the CeA came from McBride, Kimpel, et al., 2014; McBride, Rodd, et al., 2014; adolescent data for the DRN came from McClintick et al., 2015) (Figure 2). In summary, whenever ethanol-induced changes were detected, the direction of change was generally an up-regulation of mRNA expression. Overall, the DAergic system is the center of the mesocorticolimbic reward circuit and it appears that effective medications for alcoholism modulate the circuit, rather than impacting the DA-system directly.

The GABAergic System

GABA is the primary inhibitory neurotransmitter in the CNS. Thus, it and its receptors are found throughout the brain (e.g., Benham, Engin, & Rudolph, 2014). In addition, there are four GABA transporters: GAT1 and GAT3 located pre-synaptically; GAT2 and GAT3 located post-synaptically; GAT1, GAT2, GAT3 and GAT4 on glia (c.f., Clausen et al., 2006; Gonzalez-Burgos, 2010; Madsen, White, & Schousboe, 2010). The GABA receptors are classified as either GABA_A or GABA_B. There are multiple subunit isoforms for GABA_A receptors [(α1–6), (β1–3), γ1–3], delta, (ρ1–3), epsilon, pi and theta], but the most common in the CNS is a pentamer comprising (α1)2(β2)2(γ2). The receptor itself is a ligand-gated chloride channel that has binding sites for GABA, benzodiazepine, picrotoxin, steroids and anesthetics. Activation of the GABA_A receptor opens the chloride channel for influx, which induces a hyperpolarized state that decreases the probability of an action potential resulting in an inhibitory state (c.f., Fritschy, Panzanelli, & Tyagarajan, 2012; McCarson & Enna, 2014). GABA_A receptors are located extra- and post-synaptically, whereas GABA_B receptors are located both pre- and post-synaptically (Hancher, Dodson, Olsen, Otis, & Wallner, 2005; Lovinger & Roberto, 2013). GABA_B receptors are G-protein coupled, heteromer, receptors with two known subunits R1 and R2, where it appears the R1 subunit binds GABA and the R2 subunit interacts with the G-protein (Gaiarsa, Kucwesi, & Porcher, 2011; Terunuma et al., 2014).

Several research groups have reported significant associations between GABA gene variants, expression levels, and activation in brain regions such as the mesocorticolimbic system and the extended amygdala (which includes substructures of the bed nucleus of the stria terminalis, amygdala, nucleus accumbens and prefrontal cortex), with high alcohol-

consuming phenotypes and risk for developing alcohol dependence in alcoholics as well as alcohol-preferring rats (Dick & Bierut, 2006; Enoch et al., 2012; Herman et al., 2012; Korpi & Sinkkonen, 2006; McBride et al., 2010; Tabakoff et al., 2009). Thus, differential GABA signaling could reflect one mechanism that predisposes individuals to consume alcohol. In addition, GABAergic activity regulates, in part, other neuromodulator systems in the mesocorticolimbic reward circuit (Eiler & June, 2007; Melis, Camarini, Ungless, & Bonci, 2002; Rahman & McBride, 2002), supporting the role of GABA in DA-associated responses to reward. Given there are genetic differences, between P and NP rats, in these other neuromodulator systems of the mesocorticolimbic reward circuit as well (e.g., Bell et al., 2012, 2016; Franklin et al., 2014; Rahman et al., 2014, 2016 and discussed herein), this gene-by-gene interactional effect may serve as another contributing factor for the development of alcohol abuse and dependence (e.g., Saba et al., 2015; Tabakoff et al., 2009). Acute alcohol experience potentiates GABA signaling and facilitates its hyperpolarizing actions (Koob, 2004). And, GABA_A and GABA_B receptors mediate some of the rewarding, reinforcing, and motivational effects of alcohol consumption and alcohol binge drinking (Eiler & June, 2007; Nowak, McBride, Lumeng, Li, & Murphy, 1998; Tanchuck, Yoneyama, Ford, Fretwell, & Finn, 2011; also see Agabio & Colombo, 2014). Systemically, the GABA_A agonist topiramate (Breslin, Johnson, & Lynch, 2010; Lynch, Bond, Breslin, & Johnson, 2011) and GABA_B agonist baclofen (Liang et al., 2006; Maccioni et al., 2012) and GABA_B positive modulators GHB (June et al., 1995), CGP7930 (Liang et al., 2006) and GS39783 (Maccioni et al., 2012) all reduced ethanol drinking and/or self-administration by P rats. Similarly, negative modulators of the benzodiazepine-site Ro 15-4513 (McBride, Murphy, Lumeng, & Li, 1988), Ro 19-4603 (June et al., 1996; June, Murphy, Mellor-Burke, Lumeng, & Li, 1994; June, Torres, et al., 1998), Ru 34000 (June, Eggers, et al., 1998), Ro 15-1788 (June et al., 1994; June, Torres, et al., 1998), CGS 8216 and ZK 93426 (June, Devaraju, et al., 1998; June, Zuccarelli, et al., 1998) all reduced ethanol intake and/or self-administration by P rats. The partial agonist/antagonist β CCt systemically also reduced ethanol self-administration by P rats (June et al., 2003). Centrally, intra-VTA infusion of the GABA_A receptor antagonists bicuculline, picrotoxin (Nowak et al., 1998) and SR95531 (Eiler & June, 2007) reduced ethanol intake or self-administration by P rats. Similarly, intra-VTA infusion of the negative modulator of the benzodiazepine-site Ru 34000 (June, Eggers, et al., 1998) as well as intra-CeA (Foster et al., 2004) and intra-VP (June et al., 2003) infusion of the partial agonist/antagonist β CCt reduced operant self-administration of ethanol by P rats. Also, shRNA-induced reductions of GABA_A- α 2-subunit and its associated toll-like receptor 4 (*Tlr4*), in the CeA of P rats, significantly reduced alcohol self-administration (Liu et al., 2011).

Previous work indicated that P rats have more GABA_A receptors in the NAcB than NP rats (Hwang, Lumeng, Wu, & Li, 1990), and P rats display a greater response to benzodiazepines in the PFC, NAcBSh, CPU, cingulate gyrus and dorsal lateral septum than NP rats as well (Thielen, McBride, Chernet, Lumeng, & Li, 1997). Additionally, as seen in Figure 3, gene expression for a number of GABA_A and GABA_B receptor subunits differ between P and NP rats (data for the pVTA came from McBride et al., 2012; data for the CeA and NAcBSh came from McBride, Kimpel, McClintick, Ding, Hyytia, et al., 2013). Also seen in Figure 3 are our findings on the role of ethanol drinking, usually in a binge-like manner, on GABA_A

and GABA_B receptor subunit gene expression (adult data for the whole NAc came from Rodd et al., 2008; adult data for the NAcSh came from Bell, Kimpel, et al., 2009, McBride et al., 2010; adult data for the CeA came from McBride et al., 2010, adult data for the pVTA came from McBride, Kimpel, McClintick, Ding, Hauser, et al., 2013; adolescent data for the CeA came from McBride, Kimpel, et al., 2014; McBride, Rodd, et al., 2014; adolescent data for the DRN came from McClintick et al., 2015). In general, where differences in mRNA expression were detected in the NAcSh and pVTA, P rats had lower GABA_B-associated levels than NP rats. Following ethanol consumption by adult P rats, the majority of detected changes in the NAcSh represented downregulated gene expression, whereas detected changes in the CeA were approximately equally down- vs up-regulated gene expression. Following ethanol drinking by adolescent P rats, practically all detected changes represented down-regulation of GABA-associated gene expression. These findings suggest lowered GABAergic function in the NAcSh and DRN following excessive ethanol intake by P rats, which would support increased excitatory/glutamatergic activity as a contributor to excessive ethanol intake (see glutamatergic section directly below).

The Glutamatergic System

The amino acid glutamate is the primary excitatory neurotransmitter in the central nervous system (CNS). Given this, it is not surprising that glutamatergic projections, transporters and receptors are found throughout the brain. As the primary excitatory neurotransmitter, glutamate plays a crucial role in neuroplasticity, learning and memory (c.f., Henley & Wilkinson, 2013; Morris, 2013; Warburton, Barker, & Brown, 2013). Glutamate interacts with both metabotropic (Grm1-Grm8) and ionotropic receptors, which include those that can bind to N-methyl-D-aspartate (NMDA) subunits [Grin1, Grin2a-Grin2d, and Grin3a-Grin3b], α -amino-3-hydroxy-5-methyl-4-isoxazole propionic acid (AMPA) subunits [Gria1-Gria4] or kainite subunits [Grik1-Grik4], for excellent reviews see (Danbolt, 2001; Nicu, Kelmendi, & Sanacora, 2010; Traynelis et al., 2010). Because of glutamate's role in excitotoxicity, extracellular glutamate must be tightly controlled (Danbolt, 2001; Sari, 2014; Wang & Qin, 2010). Multiple glutamate transporters have been implicated in this process (Anderson & Swanson, 2000; Danbolt, 2001; Gegelashvili & Schousboe, 1997, Seal & Amara, 1999). Nevertheless, the human excitatory amino acid transporter 2 (EAAT2) and its rodent analog glutamate transporter 1 (GLT1) appear to be the primary transporters performing this function (Danbolt, 2001; Mitani & Tanaka, 2003; Rothstein, Van Kammen, Levey, Martin, & Kunc, 1995; Sari, 2014). However, the glial protein cystine/glutamate exchanger (xCT) appears to exchange extracellular cystine for intracellular (glial) glutamate (Bannai et al., 1984; Bannai & Ishii, 1982), which seems counterintuitive except for the fact that cystine is converted to cysteine, among other things, and reduces oxidative stress (c.f., Ishii & Mann, 2014).

Glutamatergic activity has been shown to mediate natural as well as drug and non-drug associated reward through direct and indirect interactions with other neurotransmitter/neuromodulatory systems within the mesocorticolimbic and extended amygdala reward neurocircuitry (e.g., Carlezon & Wise, 1996; Grace, Floresco, Goto, & Lodge, 2007; Kupila et al., 2012). For instance, research with the P rat has shown that free-choice ethanol drinking results in elevated levels of extracellular glutamate in the NAc (Das, Yamamoto,

Hristov, & Sari, 2015), as well as its subregion the NAcSh, and the pVTA (Ding et al., 2013). The latter authors reported that this increase in glutamate was inversely related to decreases in glutamate clearance. Considerable research has examined the hypothesis that sensitized mesocorticolimbic and extended amygdala glutamate neurotransmission mediate in part alcohol and drug dependence. For example, it appears continued alcohol or drug intake results in a hyperglutamatergic state within mesocorticolimbic and extended amygdala reward circuits (Gass & Olive, 2008; Kryger & Wilce, 2010; Vengeliene, Bilbao, Molander, & Spanagel, 2008). Pre-clinical evidence supports clinical findings that alcohol acutely inhibits, and chronically sensitizes and upregulates glutamate neurotransmission, in brain reward regions of the mesocorticolimbic and extended amygdala circuits (e.g., Carlezon & Wise, 1996; Chandler, Newsom, Sumners, & Crews, 1993; Cui et al., 2013; Ding, Engleman, Rodd, & McBride, 2012; Floyd, Jung, & McCool, 2003; Gass & Olive, 2008; Kapasova & Szumlinski, 2008; Nevo & Hamon, 1995; Nie, Madamba, & Siggins, 1994; Nie, Yuan, Madamba, & Siggins, 1993; Tabakoff & Hoffman, 2013; Weitlauf & Woodward, 2008), which may be due, in part, to changes in glutamate clearance (Ding et al., 2013; Kapasova & Szumlinski, 2008; Othman, Sinclair, Haughey, Geiger, & Parkinson, 2002; Parks et al., 2002; Rao, Bell, et al., 2015; Sari et al., 2011; Smith, 1997; Smith & Zsigo, 1996; Thoma et al., 2011). Moreover, recent data showed that 10 weeks of operant binge-like self administration of solutions containing both ethanol and nicotine resulted in elevation of extracellular glutamate levels in the PFC (Deehan, et al., 2015).

The role of glutamate in alcohol consumption is prominent in binge-drinking as well (c.f., Bell et al., 2016; Rao, Bell, et al., 2015). Genetic animal models of alcoholism, the P rat in this case, engaging in binge-like drinking, which results in BACs of 80 mg% and higher (c.f., Bell et al., 2011, 2014), display numerous changes in glutamate receptor and/or subunits, transporters, scaffolding proteins as well as other associated gene expression levels in discrete brain regions of the mesocorticolimbic and extended amygdala circuits (Bell et al., 2016; Bell, Kimpel, et al., 2009; Coleman et al., 2011; McBride, Kimpel, et al., 2014; McBride et al., 2009, 2010; McBride, Rodd, et al., 2014; McClintick et al., 2015; Rodd et al., 2008). These changes in glutamatergic neurotransmission include enhanced receptor activation and intracellular downstream signaling cascades (Cuzzoli et al., 2009; Szumlinski et al., 2007; Tabakoff et al., 2009). In support of this contention, glutamate receptor antagonists such as acamprosate and MPEP reduce binge-like drinking dose-dependently (Grace et al., 2007; Gupta et al., 2008). Increases in excitatory neurotransmission may be greater during periods of acute ethanol withdrawal, which is commonly associated with binge drinking, compared to more protracted withdrawal periods (Ward et al., 2009). This may support the hypothesis that binge alcohol abuse increases susceptibility to alcohol-induced excitotoxic brain damage to a greater extent than continuous excessive drinking (e.g., Hunt, 1993; see also discussion and references in Bell et al., 2013). Overall, it is likely that glutamatergic neuroadaptations following repeated binge-like drinking behavior lead to a glutamate-GABA functional imbalance (Enna, 1997; Fadda & Rossetti, 1998; Szumlinski et al., 2007), and are responsible, in part, for withdrawal symptomology when ethanol access is terminated. This withdrawal symptomology in turn increases the negative reinforcement-associated properties of continued binge drinking (Everitt & Robbins, 2005; Koob & LeMoal, 2008; Robinson & Berridge, 2008). These effects are consistent with a proposed

transition from binge/impulsive alcohol drinking to habitual/compulsive drinking to dependence (c.f., Koob, 2013; Koob & Volkow, 2010).

Systemically, the GRM1 antagonist CPCCOEt did not affect operant self-administration (Schroeder, Overstreet, & Hodge, 2005a, 2005b), whereas the GRM1 antagonist JNJ 16259685 (Besheer, Faccidomo, Grondin, & Hodge, 2008a, 2008b) did significantly reduce operant self-administration by P rats. Slight procedural and/or motor effect differences may explain the difference between the former and latter findings from the same laboratory. The GRM2/3 antagonist LY404039 significantly reduced ethanol-seeking behavior as well as relapse, but did not affect the maintenance, of ethanol self-administration by P rats (Rodd et al., 2006). The GRM5 antagonists MPEP (Schroeder et al., 2005a, 2005b) and MTEP (Cowen, Djouma, & Lawrence, 2005) both reduced operant self-administration of ethanol by P rats. Intra-NAcB infusion of the GRM2/3 agonist LY379268 and GRM5 antagonist MPEP both significantly reduced operant ethanol self-administration by P rats (Besheer et al., 2010). Much more research has been done exploring the effects of manipulating GLT1 transporter levels, within subregions of the mesocorticolimbic reward circuit, on ethanol intake by P rats, to which we turn next.

The P rat has been very useful in assessing the efficacy of beta-lactam antibiotic, and similar, molecules in reducing alcohol intake, which appears to be due to a reversal of ethanol-induced down-regulation of GLT1 levels and the concomitant increase in extracellular glutamate. For example, chronic free-choice drinking by P rats reduced GLT1 expression in the NAcB but not in the PFC, although xCT (the glutamate-cystine antiporter) was reduced in both regions (Alhaddad, Das, & Sari, 2014; Sari & Sreemantula, 2012; Sari, Sreemantula, Lee, & Choi, 2013). Ceftriaxone administered peripherally reversed ethanol-induced downregulation of GLT1 (both isoforms GLT1a and GLT1b) in the PFC and NAcB as well as xCT in the NAcB, PFC and Amyg (Alhaddad, Das, et al., 2014; Rao & Sari, 2014a, 2014b). Similarly, ceftriaxone reversed ethanol-induced increases in extracellular glutamate of the NAcB (Das et al., 2015). Research with other beta-lactam antibiotics in P rats has shown that amoxicillin, augmentin and ampicillin, which have the potential to be orally administered, reduced alcohol intake and increased GLT-1 expression in PFC and NAcB of P rats (Goodwani, Rao, Bell, & Sari, 2015; Rao, Goodwani, et al., 2015). Non-antibiotic compounds with a putative ability to upregulate GLT1 expression and/or activity have been tested in P rats as well. For example, GPI-1046, an analog of FK506, significantly reduces free-choice alcohol intake by male P rats with a concomitant increase in GLT1 expression levels in the NAcB and PFC (Sari & Sreemantula, 2012). Similar results were found for MS-153 with significant decreases in ethanol intake paralleling reversals of ethanol-induced GLT1 downregulation in the NAcB, Amyg and hippocampus (Aal-Aaboda, Alhadad, Osowik, Nauli, & Sari, 2015; Alhadad, Kim, et al., 2014). Finally, it has been demonstrated that upregulation of GLT1 (including both isoforms) and xCT by ceftriaxone and MS-153 involves NF- κ B and Akt signaling pathways (Alhaddad, Kim, et al., 2014; Rao, Saternos, Goodwani, & Sari, 2015).

An early study found that, globally, CNS *Grm3* and *Grm7* and the glycine- α 1 subunit mRNA levels are lower in inbred P rats compared with inbred NP rats (Kimpel et al., 2007). More recently, Zhou and colleagues (2013) reported that P rats are homozygous for a *Grm2*

stop codon (*Grm2**407) that essentially renders them a functional KO of *Grm2*. At the same time, Meinhardt and colleagues (2013) published their own work indicating GRM2 deficits are inversely related to drug-seeking behavior. As highlighted in Bell et al. (2016), a number of site-specific differences in glutamate-associated mRNA expression levels (see Figure 4) have been observed between P and NP rats (data for the pVTA came from McBride et al., 2012; data for the CeA and NAcSh came from McBride, Kimpel, McClintick, Ding, Hyytia, et al., 2013). As with differences between P and NP rats, a number of glutamate-associated gene and/or protein expression changes are observed after ethanol drinking by P rats (adult data for the whole NAc came from Rodd et al., 2008; adult data for the NAcSh came from Bell, Kimpel, et al., 2009; McBride et al., 2010; adult data for the CeA came from McBride et al., 2010; adult data for the pVTA came from McBride, Kimpel, McClintick, Ding, Hauser, et al., 2013; adolescent data for the CeA came from McBride, Kimpel, et al., 2014; McBride, Rodd, et al., 2014; adolescent data for the DRN came from McClintick et al., 2015) (Figure 4). The protein changes after ethanol drinking in the NAc and amygdala are from Obara et al. (2009). In general, differences, between P and NP rats, in accumbal ionotropic-associated glutamate receptor subunits are equally greater or lesser than that observed in the other line. However, metabotropic glutamate receptors are, for the most part, expressed lower in the P rat than in the NP within this brain region. Following ethanol consumption by adult P rats, expression levels for metabotropic glutamate receptors remained unchanged, whereas gene expression for ionotropic-associated glutamate receptor subunits are equally up- and down-regulated in the NAc. In the CeA, the only observed difference was lower expression of *Grm2* and *Grm3* in P vs NP rats. However following ethanol drinking, all detected ionotropic subunit-associated and metabotropic glutamate receptor gene expression changes reflected up-regulation. This would suggest a strong neuroplastic response associated with ethanol-induced elevations in glutamate, such that substantial, enhanced glutamatergic neurotransmission is occurring in this brain region of the P rat. Similar observations have been seen in glutamate-associated protein levels of P rats (Obara et al., 2009) and electrophysiological activity in Marchigian sP rats (Herman et al., 2016). As discussed in our recent glutamate review (Bell et al., 2016), the findings from P rats support the hyperglutamatergic hypothesis of alcohol and drug dependence. Given this, the P rat serves as a genetic animal model of alcoholism with characteristics of glutamatergic function paralleling clinical observations including the efficacy of topiramate in significantly reducing ethanol intake.

The Serotonergic System

The neurotransmitter serotonin (5-HT) is associated with addictive behaviors, appetite regulation, behavioral inhibition, mood, and cognitive functions. Thus, dysregulation of the 5-HT system is implicated in the development of alcohol dependence. The serotonin transporter (SERT) clears 5-HT from the synapse through reuptake into the presynapse. There are seven families of 5-HT receptors (5-HT₁₋₇) and at least 14 distinct 5-HT receptor subtypes (Barnes & Sharp, 1999), which makes the task of understanding which 5-HT receptor subtypes mediate addictive behaviors a complex one. The raphe nucleus, where 5-HT neurons originate, sends 5-HT projections to numerous regions including the VTA, NAc, and PFC and studies have shown that the 5-HT system regulates DA neuronal activity in these subregions of the mesocorticolimbic system (Azmitia & Segal 1978; Halliday & Tork,

1989; Herve, Pickel, Joh, & Beaudet, 1987; Parent, Descarries, & Beaudet, 1981; Van Bockstaele, Cestari, & Pickel, 1994). For example, 5-HT activates VTA-DA neurons (Pessia, Jiang, North, & Johnson, 1994), induces DA release in VTA slices (Beart & McDonald 1982), enhances DA release in NAc when locally applied to the VTA (Guan & McBride 1989) or the dorsal raphe is activated (Yoshimoto & McBride, 1992), potentiates the excitatory actions of alcohol on VTA-DA neurons (Brodie, Trifunovic, & Shefner, 1995), and increases extracellular DA release in the PFC (Iyer & Bradberry, 1996).

Acute alcohol exposure appears to increase 5-HT activity (McBride, Chernet, Rabold, Lumeng, & Li, 1993; Smith & Weiss, 1999), whereas chronic exposure to alcohol may result in the development of tolerance to this effect (Smith & Weiss 1999). Clinical and/or pre-clinical studies have reported deficiencies of 5-HT and/or its major metabolite 5-HIAA in the brains of human alcoholics (Pivac, Muck-Seler, Mustapic, Nenadic-Sviglin, & Kozaric-Kovacic, 2004; Schmidt, Dufeu, Heinz, Kuhn, & Rommelspacher, 1997) and genetically selected alcohol-preferring rats (McBride, Chernet, Rabold, et al., 1993; Murphy et al., 1987; Strother, Chernet, Lumeng, Li, & McBride, 2001; Zhou, Bledsoe, Lumeng, & Li, 1991a, 1991b). Moreover, treatments that reduce 5-HT neurotransmission can elevate self-administration of alcohol (Ciccocioppo, Angeletti, Colombo, Gessa, & Massi, 1999; Lyness & Smith, 1992). Drug treatments with antidepressants that affect 5-HT CNS activity have been shown to reduce craving and/or symptomatic behavior associated with alcohol dependence (c.f. Goodman, 2008) and alcoholic individuals with a polymorphism of the 5-HT transporter can respond favorably to certain medication combinations (Johnson, 2010). Therefore, it has been proposed that modulation of the 5-HT system is a viable therapy for alcoholism in a sub-set of patients (Johnson 2005, 2010; Wrase, Reimold, Puls, Kienast, & Heinz, 2006). Research on the involvement of 5-HT in binge alcohol drinking has been limited, with some evidence that binge drinking induces a blunted 5-HT response in the Scheduled High Alcohol Consumption (SHAC) mouse binge drinking model (Szumlinski et al., 2007). Additionally, acute withdrawal from alcohol after binge-like exposure leads to a wide-spread reduction in 5-HT and other neurotransmitters in several brain regions including those associated with the mesocorticolimbic system (Smith, Co, McIntosh, & Cunningham, 2008). In general, these findings indicate that serotonergic treatments may disrupt binge alcohol drinking and may interfere with the progression to alcohol dependence, in certain individuals. However, actual efficacy of 5HT manipulation to treat alcohol dependence in the clinic has been rather modest and may be relevant only in certain subpopulations of alcoholics.

Systemically, the 5HT precursor, D1L-5-HTP (McBride et al., 1990), reverse SERT modulator, fenfluramine (McBride et al., 1990); as well as agonists for the 5HT₁, TFMPP (McBride et al., 1990), 5HT_{1A}, 8-OH-DPAT (McBride et al., 1990), 5HT₂, DOI (McBride et al., 1990), all reduced ethanol intake by P rats. However, most work has examined receptor antagonists, including WAY 100,635 (Zhou, McKinzie, Patel, Lumeng, & Li, 1998) which targets the 5HT_{1A} receptor; amperozide/FG 5606 (Lankford, Bjork, & Myers, 1996; Overstreet, McArthur, Rezvani, & Post, 1997), and FG 5974 (Lankford et al., 1996; Overstreet et al., 1997; Piercy, Bjork, & Myers, 1996) which target 5HT₂ receptors; as well as MDL 72222 (Rodd-Henricks, McKinzie, Edmundson, et al., 2000) and ICS 205-930 (Rodd et al., 2010; Rodd-Henricks, McKinzie, Edmundson, et al., 2000) which target 5HT₃

receptors, all of which reduce ethanol intake or the acquisition of operant ethanol self-administration by P rats. Additionally, the SERT inhibitors fluoxetine (Murphy et al., 1985, 1988; Rezvani et al., 2000; Zhou et al., 1998), fluvoxamine (Murphy et al., 1985) and DOV 102,677 (Yang et al., 2012) all reduced ethanol intake by P rats. The last compound also inhibits the norepinephrine and dopamine transporters.

A number of differences in serotonin-associated protein (Ciccocioppo, Ge, Barnes, & Cooper, 1998; McBride, Chernet, Raboid, et al., 1993; McBride et al., 1997; McBride, Guan, Chernet, Lumeng, & Li, 1994; Murphy et al., 1987; Pandey, Lumeng, & Li, 1996) and mRNA expression levels (see Figure 5) have been reported between P and NP rats (data for the pVTA came from McBride et al., 2012; data for the CeA and NAcSh came from McBride, Kimpel, McClintick, Ding, Hyytia, et al., 2013). Observed ethanol drinking-induced changes in gene expression have come mainly from our work with adolescent bingeing P rats and in the DRN, from which all 5HT projections emanate (adult data for the whole NAc came from Rodd et al., 2008; adult data for the NAcSh came from Bell, Kimpel, et al., 2009, McBride et al., 2010; adult data for the CeA came from McBride et al., 2010; adult data for the pVTA came from McBride, Kimpel, McClintick, Ding, Hauser, et al., 2013; adolescent data for the CeA came from McBride, Kimpel, et al., 2014; McBride, Rodd, et al., 2014; adolescent data for the DRN came from McClintick et al., 2015) (Figure 5). For the most part, P rats have lower levels of central 5HT, 5HIAA, and 5HT receptors than NP rats. Following ethanol drinking by both adult and adolescent P rats, usually in binge-like form, nearly all detected changes in 5HT-associated gene expression represented down-regulation. These findings provide strong evidence that serotonergic deficits predispose an individual to abuse ethanol and, without intervention, foster continued excessive drinking.

The Noradrenergic and Corticotrophin Releasing Hormone (CRH) Systems

It is widely established that the central noradrenergic system serves a global function in neuromodulation, controlling vigilance, attention, and the sleep–wake cycle as well as contributing to learning and memory processes. Neuroanatomical evidence indicates that noradrenergic system in the brain arise from the cell bodies in the locus coeruleus and project to different cerebral regions and to the spinal cord (Cooper et al., 2002). Moreover, there is a complex interaction between norepinephrine (NE) and corticotrophin releasing factor (CRF), a neuropeptide strongly associated with central autonomic and stress activity, receptors within the locus coeruleus (Reyes, Bangasser, Valentino, & Van Bockstaele, 2014). In addition to major projections to the frontal cortex, (NE) neurons project to the limbic system (Flavin & Winder, 2013), including amygdala, bed nucleus of the stria terminalis, hippocampus, and hypothalamus where it is implicated in addiction (Becker, 2012; Koob, 2013; Sofuoglu, Rosenheck, Petrakis, 2014; See also Al' Absi, 2007), anxiety (Geiger, Neufang, Stein, & Domschke, 2014), attention (Geiger et al., 2014; Hegerl & Hensch, 2014), cognition (Chandler, Waterhouse, & Gao, 2014), memory, mood (Gold, 2015), pain (Elman, Borsook, & Volkow, 2013; Strobel, Hunt, Sullivan, Sun, & Sah, 2014), post-traumatic stress disorder (PTSD, Sofuoglu et al., 2014; Wimalawansa, 2014), sleep (Zeitzer, 2013), suicide (Elman et al., 2013), and associated physiological processes (Klimek, Rajkowska, Luker, Dilley, et al., 1999; Moret & Briley, 2011).

NE in synaptic vesicles is derived from two sources. The synthesis of NE begins with the synthesis of dopamine from tyrosine and is transported into the vesicle by the vesicular monoamine transporter (VMAT). Once dopamine is synthesized and stored in synaptic vesicles, an enzyme called dopamine- β -hydroxylase further hydroxylates dopamine into NE. The synthesis of NE is different than the other neurotransmitters as they are usually made in the cytoplasm of the terminal buttons, whereas NE's final stage of synthesis occurs in synaptic vesicles. Neurotransmission is initiated by an action potential which triggers the release of NE into the synaptic cleft. Released NE interacts with multiple adrenergic receptors, including presynaptic α_2 and postsynaptic α_1 , β_1 and β_2 receptors. NE is removed from the synaptic cleft by both selective NE transporter (NET) as well nonselective transporters. NE's stimulation of α_2 -adrenergic receptors provide's feedback inhibition of further release. Cytoplasmic NE that is not sequestered in synaptic vesicles by VMAT is degraded into its metabolites by the enzyme monoamine oxidase, type a (MAO-A) (Cooper et al., 2002; Golan, Tashjian, Armstrong, & Armstrong, 2012; Krishnan & Nestler, 2008).

Evidence indicates that central NE activity modulates alcohol drinking behavior (Ehrenreich, Schuck, Stender, et al., 1997; Getachew, Hauser, Taylor, & Tizabi, 2010; See also Al' Absi, 2007). For example, selective NE uptake inhibition may normalize the behavioral and negative affective effects of alcohol (Getachew et al., 2010). Similarly, α_1 -adrenergic antagonists, α_2 -adrenergic agonists and β -adrenergic antagonists modulate alcohol drinking or associated withdrawal behavior (e.g., Gilpin & Koob, 2010; Riihioja, Jaatinen, Oksanen, et al., 1997; Walker, Rasmussen, Raskind, & Koob, 2008). There have been several studies investigating the adrenergic system's role in excessive ethanol drinking by P rats using peripheral routes of administration. Prazosin, an α_1 -adrenergic antagonist (Menkes, Baraban, & Aghajanian, 1981), reduces home cage limited access drinking (Rasmussen, Alexander, Raskind, & Froehlich, 2009), relapse drinking (Froehlich, Hausauer, Fischer, Wise, & Rasmussen, 2015), operant self-administration (Verplaetse & Czachowski, 2015, Verplaetse et al., 2012), and operant ethanol-seeking (Verplaetse & Czachowski, 2015, Verplaetse et al., 2012). It also disrupts the acquisition of excessive ethanol drinking by P rats (Froehlich, Hausauer, Federoff, Fischer, & Rasmussen, 2013). Combining prazosin and naltrexone was more effective in reducing ethanol drinking than either compound alone, at least in P rats (Froehlich, Hausauer, & Rasmussen, 2013). Additionally, tolerance did not appear to develop following repeated daily treatments (Rasmussen, Kincaid, & Froehlich, 2015). Another study found that clonidine, an α_2 -adrenergic receptor agonist, can also reduce ethanol drinking by P rats (Rasmussen, Alexander, Malone, Federoff, & Froehlich, 2014; Rasmussen, Beckwith, Kincaid, & Froehlich, 2014). In addition, the triple monoamine uptake inhibitor (i.e., DAT, NET and SERT) DOV 102,677 reduced ethanol intake by P rats (Yang et al., 2012). Paralleling the preclinical findings, prazosin also has shown promise in the treatment of AUDs in humans (e.g., Simpson, Saxon, Meredith, et al., 2009). Overall, limited data suggest the involvement of the noradrenergic system in AUDs but substantial territory still needs to be explored regarding the role of NE and stress-associated systems in alcohol dependence.

The CRF system is localized, often colocalizing with NE, in multiple brain regions associated with addiction, anxiety, consummatory behavior, sleep, stress, learning and memory (Reul & Holsboer, 2002; Sajdyk, Shekhar, & Gehlert, 2004). These brain regions

include the raphe nucleus (Lukkes et al., 2011), multiple nuclei of the hypothalamus and the amygdala (Blume et al., 2009; Campbell, Grove, & Smith, 2003; Sajdyk et al., 2004; Smialowska, Wieronska, & Wedzony, 2002), pituitary (Stanley et al., 2004), cortex and lateral septum (Miyata, Shiota, Chaki, Okuyama, & Inagami, 2001), bed nucleus of the stria terminalis and hippocampus (Van Pett et al., 2000). Moreover, CRF-R1 receptors colocalize with cholinergic, noradrenergic and DAergic neurons in many of these brain regions as well (Sauvage & Steckler, 2001). There is increasing research into the role of CRH, NPY, glucocorticoids and HPA activity (the endogenous opioid system is discussed next) in alcohol dependence. One important reason for this is the fact that most earlier research on pharmacological treatments was conducted in animal models mimicking early stages of the dependence cycle, rather than the later stages observed in dependence. In the P rat, CRF protein expression levels are lower in the PFC, pyriform cortex, hypothalamus and amygdala compared with NP rats (Ehlers et al., 1992).

A subsequent study revealed that both CRF protein and mRNA are lower in the CeA of P rats compared with NP rats (Hwang, Stewart, Zhang, Lumeng, & Li, 2004). More recent work has shown that P rats have a polymorphism in the promoter region of the *CRFR2* gene, which is not present in NP rats (Yong et al., 2014). This polymorphism is associated with lower *Crfr2* expression, especially in the amygdala; and, similar to innately reduced NPY levels in the amygdala, reduced *Crfr2* expression appears to be directly associated with higher ethanol intake and anxiety in P rats (Yong et al., 2014). Regarding CRF, ligands targeting the CRF1 receptor, such as antalarmin and MPZP, reduce ethanol drinking by P rats but dependence is a requirement to see these effects (Gilpin, Richardson, & Koob, 2008; Gilpin, Stewart, & Badia-Elder, 2008; Heilig & Egli, 2006).

Similar to observations for the cholinergic and DAergic systems, very few differences in adrenergic and CRF-associated mRNA expression differences (see Figure 6) between P and NP rats (data for the pVTA came from McBride et al., 2012; data for the CeA and NAcSh came from McBride, Kimpel, McClintick, Ding, Hyytia, et al., 2013). The same is true for ethanol drinking-induced changes in P rats (adult data for the whole NAc came from Rodd et al., 2008; adult data for the NAcSh came from Bell, Kimpel, et al., 2009; McBride et al., 2010; adult data for the CeA came from McBride et al., 2010; adult data for the pVTA came from McBride, Kimpel, McClintick, Ding, Hauser, et al., 2013; adolescent data for the CeA came from McBride, Kimpel, et al., 2014; McBride, Rodd, et al., 2014; adolescent data for the DRN came from McClintick et al., 2015) (Figure 6). However, most drinking protocols used for our recent microarray work have been binge-like and limited access in nature. Thus, the animals were probably not physically dependent on alcohol, despite being quite motivated to consume it and achieving BACs greater than 80–100 mg%.

The Opioid System(s)

There are several classes of endogenous opioids including enkephalins, endorphins, dynorphins, and endomorphins. These classes of ligands bind with some specificity to the delta, kappa and mu-receptors, respectively. One role of these peptides in the brain is to process information about rewarding stimuli, including alcohol (c.f., Oswald & Wand, 2004). Therefore, it is not surprising that these peptides have been shown to influence the

development of alcohol abuse and dependence. Opioid receptors are found pre-synaptically on DAergic neurons of the mesocorticolimbic system (e.g., within the NAc) where they control the release of DA. Thus, opioid activity, similar to the glutamatergic and GABAergic systems, modulates DA activity in this “reward” neurocircuit.

Variations in opioid-related gene expression and function may contribute to high levels of alcohol consumption as well (e.g., Marini et al., 2013). For example, high alcohol drinking rats exhibit a greater level of mu-opioid receptor (MOR)-associated and enkephalin mRNA, compared to low alcohol drinking rats (Morganstern et al., 2012). For a review of neurobiological differences in the opioid system between selectively bred high and low alcohol-consuming rats see Bell et al. (2012). A great deal of existing evidence for the role of opioids in alcohol abuse and dependence comes from pharmacological experiments using the FDA-approved treatment for alcoholism, naltrexone (ReVia) and other non-specific opioid antagonists. Naltrexone blocks alcohol-induced changes in gene transcription in several receptor systems, including the mu-opioid system. Evidence from knock-out mice lacking MORs or dynorphin suggest that MORs and kappa-opioid receptors (KORs) are involved in the rewarding or reinforcing effects of alcohol (Blednov et al., 2006; Charbogne et al., 2014; Roberts et al., 2000).

There is a substantial literature on peripheral administration of opioid-associated ligands to reduce ethanol-drinking by P rats. Essentially, peripheral naltrexone, a pan-opioid antagonist, has been tested repeatedly (Coonfield, Kiefer, Ferraro, & Sinclair, 2004; Dhaher et al., 2012; June, Grey, et al., 1998; Sable, Bell, Rodd, & McBride, 2006) along with a study testing the effects of CeA microinjections (Foster et al., 2004). These publications all reported significant reductions in ethanol intake by male and female, adolescent and adult P rats using both home-cage and operant procedures. Peripheral testing with the pan-opioid antagonists, naloxone (Badia-Elder et al., 1999; June et al., 1991) and nalmefene (June, Grey, et al., 1998), also revealed significant reductions in ethanol intake by male and female, adolescent and adult P rats using both home-cage and operant procedures. Another study reported that nalmefene microinjections into the NAcb, Hipp and VTA significantly reduced operant alcohol self-administration by adult female P rats (June et al., 2004). A study examining the acquisition, maintenance and relapse operant self-administration of ethanol by adult female P rats found that the mu opioid receptor (MOR) antagonist LY255582 significantly reduced all of these behaviors (Dhaher et al., 2012). Other studies examined the role of the delta opioid receptor (DOR) in mediating ethanol intake by adult P rats. Systemic treatment with the DOR antagonists naltriben, naltrindole and ICI 174,864 reduced home-cage ethanol drinking (Krishnan-Sarin, Jing, et al., 1995; Krishnan-Sarin, Portoghese, Li, & Froehlich, 1995) and operant self-administration (June et al., 1999) of the same.

Despite a clear role for the opioid system (including ligand-associated changes in drinking) in alcohol drinking and consummatory behavior across reinforcers, very few changes induced by ethanol and/or differences between P and NP rats have been reported to date (Figure 7). Data of line-dependent differences for the pVTA came from McBride et al., 2012 and data for the CeA and NAcbSh came from McBride, Kimpel, McClintick, Ding, Hyytia, et al., 2013. Data for ethanol drinking-induced changes in P rats (adult data for the whole NAcb came from Rodd et al., 2008; adult data for the NAcbSh came from Bell, Kimpel, et

al., 2009, McBride et al., 2010; adult data for the CeA came from McBride et al., 2010; adult data for the pVTA came from McBride, Kimpel, McClintick, Ding, Hauser, et al., 2013; adolescent data for the CeA came from McBride, Kimpel, et al., 2014; McBride, Rodd, et al., 2014; adolescent data for the DRN came from McClintick et al., 2015) (Figure 7). The findings thus far indicate that multiple opioid systems are involved in excessive ethanol drinking and modulation of this system consistently reduces ethanol intake significantly. Thus, the P rat also displays the relatively robust reduction in alcohol abuse induced by the pan-opioid receptor antagonist naltrexone, let alone similar observations for more selective mu- and delta-opioid antagonists.

The Neuropeptide Y (NPY) system

Neuropeptide Y (NPY) is a 36 amino acid peptide abundantly expressed throughout the central and peripheral nervous systems (Allen, Adrian, Allen, et al., 1983) and acts centrally on target cells through the G-protein, coupled NPY receptors Y1, Y2, and Y5 (Dumont, Satoh, Cadieux, et al., 1993; Fetissov, Kopp, & Hokfelt, 2004; Wolak et al., 2003). NPY_{Y1} cell bodies have been found in the hypothalamus, hippocampus, amygdala, brain stem nuclei and ganglions of the sympathetic and parasympathetic nervous systems. There is evidence that NPY is locally synthesized (van den Pol, 2012), therefore its expression is not dependent upon projection neurons. For the most part, NPY neurons are interneurons and found in most brain regions except for the thalamus and the cerebellum, although Y₂-receptors have been detected in the thalamus (e.g., Kaji, 2013). NPY is implicated in food intake and energy balance, anxiety, stress, autonomic function, learning and memory (c.f., Parker, 2013). Given the role of stress and anxiety (at least their comorbidity for the latter) in alcohol and drug dependence, it is not surprising that NPY, and for that matter the interaction of NPY and CRF (e.g., Thorsell, 2010), is implicated in the development and maintenance of alcohol and/or drug dependence (c.f., Al'Absi, 2007). Regarding the NPY-CRF interaction, There appears to be a confluence of activity from multiple systems in the CeA that mediates anxiety and alcohol dependence with a complex interaction of the CRF, GABAergic, glutamatergic and NPY systems (Gilpin, Herman, & Roberto, 2015; Roberto, Gilpin, & Siggins, 2012). Additionally, cAMP response element-binding protein (CREB), brain derived neurotrophic factor (BDNF), activity-regulated cytoskeleton (Arc) associated protein are involved in the modulation of CeA-NPY activity and, by extension, are involved in anxiety and alcohol dependence as well (c.f., Pandey, 2003; Wand, 2005).

Centrally, site-specific injections of NPY point to the amygdala (Gilpin, Richardson, et al., 2008; Gilpin, Stewart, et al., 2008; Pandey, Chartoff, Carlezon, et al., 2005) and hypothalamus (Gilpin, Stewart, Murphy, & Badia-Elder, 2004; Kelley, Nannini, Bratt, & Hodge, 2001) as possible primary neuroanatomical loci for NPY-induced alterations in ethanol drinking. Previous studies suggest that NPY is a major factor which distinguishes the ethanol-drinking behavior of P vs NP rats (Cowen, Chen, & Lawrence, 2004; Pandey, Carr, Heilig, Ilveskoski, & Thiele, 2003; Thiele & Badia-Elder, 2003). P rats display lower levels of NPY immunoreactivity in various regions of the brain, including the CeA, hippocampus and the FC with higher levels in the paraventricular hypothalamic nucleus and arcuate nucleus of the hypothalamus (Ehlers, Li, Lumeng, et al., 1998; Hwang, Zhang, Ehlers, Lumeng, & Li, 1999). Furthermore, decreased levels of NPY are associated with

increased anxiety in P rats (Spence et al., 2005; Stewart et al., 1993), and ICV infusion of NPY has been shown to reduce ethanol intake in the P rat (Gilpin, Stewart, Murphy, Li, & Badia-Elder, 2003) (Fig. 8). Genomically, NPY is localized in an interval that is highly associated with alcohol preference and consumption, mapping to a quantitative trait locus (QTL) with a lod score of 9.2 on rat chromosome 4, using an F2 population bred from iP and iNP rats (Bice, Foroud, Bo, et al., 1998; Carr, Foroud, Bice, et al., 1998).

11. Pharmacogenomics and Alcoholism/Addiction

The interest in using pharmacogenomic research to treat alcohol dependence spans more than a decade (e.g., Anton et al., 2008; Goldman, Oroszi, O'Malley, & Anton, 2005). Numerous studies have identified a number of single nucleotide polymorphisms that are associated with alcohol dependence and/or drug codependence including (a) CHRM2 (Luo et al., 2005; Wang et al., 2004), CHRNA4 (Kim et al., 2004), CHRNA5 (Saccone et al., 2007; Wang et al., 2009) as well as the CHRNA5-CHRNA3-CHRNA4 cluster and alcohol abuse/dependence (Schlaepfer et al., 2008); (b) DAT (Heinz, Goldman, Gallinat, Schumann, & Puls, 2004; see also Bhaskar & Kumar, 2014 for this and other DA-associated polymorphisms), DA beta hydroxylase (DBH) and alcohol dependence in women (Preuss et al., 2013), DRD3 and alcohol craving (Agrawal et al., 2013) as well as DA dysfunction and Cloninger Type I alcoholism (Leggio & Addolorato, 2008); (c) GABRA1, GABRA2, GABRB3, GABRG3 and alcohol dependence or sensitivity to its intoxicating effects during the ascending slope of the BAC curve (e.g., Bierut et al., 2010; Dick et al., 2004; Dick et al., 2006; Edenberg et al., 2004; Enoch, Schwartz, Albaugh, Virkkunen, & Goldman, 2006; Haughey et al., 2008; Noble et al., 1998); (d) GRIK3 (Grzywacz, Malecka, Suchancka, Bienkowski, Samochowiec, 2013) and GRIN2A (Domart et al., 2012) with alcohol dependence as well as GRM8 and event-related potential (ERP) theta power and alcohol dependence (Chen et al., 2009); (e) 5HT dysfunction and Cloninger Type II alcoholism (Leggio & Addolorato, 2008), HTR1A and alcohol as well as nicotine co-dependence (Zuo et al., 2013), HTR1B and alcohol as well as multiple drug abuse (Cao, LaRocque, & Li, 2013; Contini et al., 2012), HTR2A and alcohol as well as heroin abuse (Cao et al., 2014), HTR7 and alcohol dependence as well as electrophysiological measures (Zlojutro et al., 2010; Zuo et al., 2014), alcohol dependence and SERT (e.g., Heinz et al., 2004; c.f., Johnson, 2010; McHugh, Hofman, Asnaani, Sawyer, & Otto, 2010; Plemenitas et al., 2015); (f) OPRM1 and level of response to ethanol in Native Americans (Ehlers, Lind, & Wilhelmsen, 2008), OPRM1 polymorphisms and naltrexone's efficacy for treating alcohol dependence (e.g., Jonas et al., 2014), as well as PDYN and OPRK1 with alcohol dependence (Gerra et al., 2007; Williams et al., 2007; Xuei et al., 2006); (g) CRFR1 polymorphism with P3 ERP and alcohol dependence (Chen et al., 2010); and (h) NPY and its receptor's association with alcohol as well as multiple drug abuse and dependencies (Bhaskar et al., 2013; Frances et al., 2011; Okahisa et al., 2009; Sato et al., 2010; Wetherill et al., 2008). While there is overlap in the ethanol affected neurotransmitter and neuropeptide systems between the clinical alcohol dependent population and the P rat, much more research needs to be done.

12. Conclusions

The present chapter sought to present the existing neuropharmacological findings on P rats in a more holistic manner than done in the past. While previous reviews listed many differences between the P rat and its control line the NP rat, this was mostly done in tabular form or buried in the text. By using figures of the primary neurotransmitter and neuropeptide systems examined thus far in the P rat, our objective was to map the published findings in the context of the projections and/or localization of each respective neuromodulatory system. This chapter also outlines how the P rat has neurochemical, physiological and behavioral characteristics often seen in individuals with alcohol, and in some cases drug, dependence. Finally, it has been proposed that an animal model of alcoholism should display similar pharmacological efficacy as that seen in the alcohol dependent treatment population. Because excessive alcohol is under genetic control, individual differences are expected in animal models just as individual differences prevail in the clinical treatment population. Thus, an animal model of alcoholism also should display some pharmacological treatment validity (Dyr & Kostowski, 2008; Litten et al., 2012; Overstreet, Rezvani, Djouma, Parsian, & Lawrence, 2007).

For the cholinergic system, the P rat displays a modest effect of varenicline on ethanol intake. This parallels the clinical literature in that there are mixed findings for the efficacy of varenicline, especially in the context of smokers vs nonsmokers and/or co-morbid psychaitric conditions. For the DAergic system, findings with the P rat do not match clinical observations; such that whereas manipulations of the DAergic system consistently affect ethanol intake in P rats, this is not true in the treatment setting. For the GABAergic system, the results are mixed. GABAergic modulators are often used during ethanol withdrawal, but cross-tolerance with the effects of ethanol and inherent abuse liability limit their usefulness beyond acute care. One exception should be noted and that is topiramate, which has shown similar efficacy in P rats, other animal models and a tested treatment population. Topiramate, as an anticonvulsant, modulates both GABAergic and glutamatergic activity. Acamprosate, another modulator of the glutamatergic system, has demonstrated modest effects in the clinical treatment population, but only has marginal effects in P rats. Similar to the DAergic system, modulators of the serotonergic system have had limited success in the clinical setting even though robust effects are seen in P rats. Ondansetron is one exception, with mixed findings in both P rats and human laboratory subjects. In addition, it appears that variances in the SERT gene (*5htt*) can determine some pharmacotherapeutic efficacy supporting a role for pharmacogenomics. For the noradrenergic system, prazosin consistently decreases ethanol intake in P rats and promising results are being seen in human clinical studies as well. For the opioid system, naltrexone consistently reduces ethanol intake in P rats and while some clinical studies report robust effects, other clinical studies have reported modest to marginal results. Similar to observations of *5htt* polymorphisms, variances in the OPRM1, and possibly OPRK1, gene appear to have some predictive validity for naltrexone's efficacy in the treatment of alcohol dependence. A final example, which does not fall into any of the systems discussed in this chapter, is ibudilast and more selective inhibitors of phosphodiesterase 4 (PDE4, e.g., Bell et al., 2015; Franklin, Hauser, Lasek, Bell, & McBride, 2015). Robust findings in P rats and multiple other animal models parallel

early results from clinical laboratory studies. Regarding these early findings, it should be noted that in 2015 the Food and Drug Administration (FDA) gave ibudilast a fast-track designation for the treatment of methamphetamine dependence. While considerable progress has been made in the treatment of alcohol and drug dependence, considerable more work needs to be done. One future direction for research with the P rat is to determine if this genetic animal model will self-administer other drugs of abuse. Present work, and some previous work, from our laboratory indicates P rats will readily self-administer nicotine and cocaine into discrete regions of the mesocorticolimbic reward circuit. Intravenous work has not been published yet, but the results look promising for this route of administration as well. These findings suggest that the P rat may be a genetic animal model of polysubstance abuse/dependence.

To date, the vast majority of the genomic information about the P rat has come from microarray and RT-PCR techniques. Future research should use next-generation DNA sequencing to identify genomic signatures of selection between P and NP rats and next-generation RNA sequencing methodologies to analyze allele-specific expression of genes in F1 crosses of these lines (e.g., Farris & Mayfield, 2014; Wang, Kapoor, & Goate, 2012). This will help move the field from QTL analyses to quantitative trait nucleotide [QTN; i.e., (SNPs)] and quantitative trait gene (QTG) analyses (e.g., Ehlers, Walter, Dick, Buck, & Crabbe, 2010; Milner & Buck, 2010; Spence et al., 2009). By doing so, the level of genomic resolution and the power of these analyses will be exponentially increased over the existing techniques. In addition, by localizing genetic variation to genes and SNPs, research on the role of epigenomics/epigenetics (e.g., Moonat et al., 2010; Renthal & Nestler, 2009) in alcohol preference can also be advanced. These advances will allow investigators to combine traditional hypothesis-driven research based on deductive reasoning with unprejudiced genome association studies. These approaches will delineate putative neuromolecular pathways (e.g., intracellular cascades) mediating alcohol dependence and identify possible new drugable targets to prevent and/or treat alcohol abuse and dependence.

Another direction for future research is the use of emerging and evolving neuroscience methodologies to examine the role of second messenger systems, synaptic plasticity, protein-protein interactions, gene-gene interactions, and the role of noncoding RNAs (e.g., Clerget, Bouguignon-Igel, & Rederstorff, 2015; Gedik et al., 2015; Gorini, Bell, & Mayfield, 2011; Gorini, Harris, & Mayfield, 2014; Manzardo, McGuire, & Butler, 2015; Nunez et al., 2013; Ponomarev, Wang, Zhang, Harris, & Mayfield, 2012). Emerging bioinformatic strategies would synthesize the large amounts of data obtained with high throughput gene and/or protein expression techniques (e.g., Gorini et al., 2011; Gorini et al., 2014). Presently, this synthesis has started to reveal the complex neurobiology of alcoholism and the multiple roles of genetics in its development through functional and genetical genomics (e.g., Spanagel et al., 2013; Zuo et al., 2014). We believe that continued research with the P rat using these more advanced genomic, proteomic and bioinformatic techniques will yield new information on molecular substrates to target for repurposing existing FDA-approved medications, or those that are in advanced clinical trials, to treat alcohol and drug abuse/dependence.

Acknowledgments

This work was supported in part by AA013522 an INIA-Neuroimmune project (RLB) and AA020396 to EAE/RLB. The views expressed herein are solely those of the authors and do not necessarily represent the views of the National Institutes of Health (NIH) funding agencies.

References

- Aal-Aaboda M, Alhaddad H, Osowik F, Nauli SM, Sari Y. Effects of (R)-(-)-5-methyl-1-nicotinoyl-2-pyrazoline on glutamate transporter 1 and cysteine/glutamate exchanger as well as ethanol drinking behavior in male, alcohol-preferring rats. *Journal of Neuroscience Research*. 2015; 93:930–937. [PubMed: 25601490]
- Agabio R, Colombo G. GABAB receptor ligands for the treatment of alcohol use disorder: Preclinical and clinical evidence. *Frontiers in Neuroscience/Neuropharmacology*. 2014; 8:140. <http://dx.doi.org/10.3389/fnins.2014.00140>.
- Agrawal A, Wetherill L, Bucholz KK, Kramer J, Kuperman S, Lynskey MT, et al. Genetic influences on craving for alcohol. *Addictive Behaviors*. 2013; 38:1501–1508. [PubMed: 22481050]
- Al' Absi, M. *Stress and Addiction: Biological and Psychological Mechanisms*. New York, NY: Elsevier Academic Press; 2007.
- Albuquerque EX, Pereira EFR, Alkondon M, Rogers SW. Mammalian nicotinic acetylcholine receptors: from structure to function. *Physiology Reviews*. 2009; 89:73–120.
- Alhaddad H, Das SC, Sari Y. Effects of ceftriaxone on ethanol intake: a possible role for xCT and GLT-1 isoforms modulation of glutamate levels in P rats. *Psychopharmacology (Berl)*. 2014; 231:4049–4057. [PubMed: 24687412]
- Alhaddad H, Kim NT, Aal-Aaboda M, Althobaiti YS, Leighton J, Boddu SH, et al. Effects of MS-153 on chronic ethanol consumption and GLT1 modulation of glutamate levels in male alcohol-preferring rats. *Frontiers in Behavioral Neuroscience/Neuropharmacology*. 2014; 8:366.
- Allen YS, Adrian TE, Allen JM, et al. Neuropeptide Y distribution in the rat brain. *Science*. 1983; 221:877–879. [PubMed: 6136091]
- American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders*. 4. Author; Washington, DC: 1994.
- American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders-Text Revision*. 4. Author; Washington, DC: 2000.
- American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders*. 5. Author; Washington, DC: 2013.
- Anderson CM, Swanson RA. Astrocyte glutamate transport: review of properties, regulation, and physiological functions. *Glia*. 2000; 32:1–14. [PubMed: 10975906]
- Anton RF, Oroszi G, O'Malley S, Couper D, Swift R, Pettinati H, et al. An evaluation of mu-opioid receptor (OPRM1) as a predictor of naltrexone response in the treatment of alcohol dependence. *Archives of General Psychiatry*. 2008; 65:135–144. [PubMed: 18250251]
- Antuono, P.; Sorbi, S.; Bracco, L.; Fusco, T.; Amaducci, L. A discrete sampling technique in senile dementia of the Alzheimer type and alcoholic dementia: study of the cholinergic system. In: Amaducci, L.; Davison, AN.; Antuono, P., editors. *Aging of Brain and Dementia*. Vol. 1980. Raven Press; New York: 1980. p. 151-158.
- Assaad J-M, Pihl RO, Seguin JR, Nagin D, Vitaro F, Carbonneau R, et al. Aggressiveness, family history of alcoholism, and the heart rate response to alcohol intoxication. *Experimental and Clinical Psychopharmacology*. 2003; 11:158–166. [PubMed: 12755460]
- Azmithia EC, Segal M. An autoradiographic analysis of the differential ascending projections of the dorsal and median raphe nuclei in the rat. *Journal of Comparative Neurology*. 1978; 179:641–67. [PubMed: 565370]
- Babor TF, De Hofmann MI, Boca FK, Hesselbrock V, Meyer RE, Rounsaville B. Types of alcoholics, I. Evidence for an empirically derived typology based on indicator of vulnerability and severity. *Archives of General Psychiatry*. 1992; 49:599–608. [PubMed: 1637250]

- Badia-Elder NE, Mosemiller AK, Elder RL, Froehlich JC. Naloxone retards the expression of a genetic predisposition toward alcohol drinking. *Psychopharmacology*. 1999; 144:205–212. [PubMed: 10435386]
- Baimel C, Bartlett SE, Chiou LC, Lawrence AJ, Muschamp JW, Patkar O, et al. Orexin/hypocretin role in reward: implication for opioid and other addictions. *British Journal of Pharmacology*. 2015; 172:334–348. [PubMed: 24641197]
- Bannai S, Christensen HN, Vadgama JV, Ellory JC, Englesberg E, Guidotti GG, et al. Amino acid transport systems. *Nature*. 1984; 311:308. [PubMed: 6482960]
- Bannai S, Ishii T. Transport of cystine and cysteine and cell growth in cultured human diploid fibroblasts: effect of glutamate and homocysteate. *Journal of Cellular Physiology*. 1982; 112:265–272. [PubMed: 6126484]
- Barker JM, Taylor JR. Habitual alcohol seeking: modeling the transition from casual drinking to addiction. *Neuroscience and Biobehavioral Reviews*. 2014; 47:281–294. <http://dx.doi.org/10.1016/j.neubiorev.2014.08.012>. [PubMed: 25193245]
- Barnes NM, Sharp T. A review of central 5-HT receptors and their function. *Neuropharmacology*. 1999; 38:1083–1152. [PubMed: 10462127]
- Barron S, Lewis B, Wellman K, Carter M, Farook J, Ring J, et al. Polyamine modulation of NMDARs as a mechanism to reduce effects of alcohol dependence. *Recent Patents and CNS Drug Discovery*. 2012; 7:129–144. <http://dx.doi.org/10.2174/157488912800673128>.
- Beart PM, McDonald D. 5-Hydroxytryptamine and 5-hydroxytryptaminergic dopaminergic interactions in the ventral tegmental area of rat brain. *Journal of Pharmacy and Pharmacology*. 1982; 34:591–593. [PubMed: 6127387]
- Becker HC. Effects of alcohol dependence and withdrawal on stress responsiveness and alcohol consumption. *Alcohol Research*. 2012; 34:448–458. [PubMed: 23584111]
- Beckwith SW, Czachowski CL. Increased delay discounting tracks with a high ethanol-seeking phenotype and subsequent ethanol seeking but not consumption. *Alcoholism: Clinical and Experimental Research*. 2014; 38:2607–2614.
- Bell RL, Eiler WJA II, Cook JB, Rahman S. Nicotinic receptor ligands reduce ethanol intake by high alcohol-drinking HAD-2 rats. *Alcohol*. 2009; 43:581–592. [PubMed: 20004336]
- Bell, RL.; Franklin, KM.; Hauser, SR.; Engleman, EA. Next stop dependence. Binge drinking on the road to alcoholism: preclinical findings on its neurobiology from rat animal models. In: Harris, SB., editor. *Binge eating and binge drinking: Psychological, social and medical implications*. New York: Nova Science Publishers; New York, NY: 2013. p. 1-60.
- Bell RL, Hauser SR, McClintick J, Rahman S, Edenberg HJ, Szumlinski KK, et al. Ethanol-associated changes in glutamate reward neurocircuitry: a mini-review of clinical and preclinical genetic findings. *Progress in Molecular Biology and Translational Science*. 2016; 137:41–85. <http://dx.doi.org/10.1016/bs.pmbts.2015.10.018>. [PubMed: 26809998]
- Bell RL, Kimpel MW, McClintick JN, Strother WN, Carr LG, Liang T, et al. Gene expression changes in the nucleus accumbens of alcohol-preferring rats following chronic ethanol consumption. *Pharmacology Biochemistry and Behavior*. 2009; 94:131–147. <http://dx.doi.org/10.1016/j.pbb.2009.07.019>.
- Bell RL, Lopez M, Changhai C, Egli M, Johnson K, Franklin K, et al. Ibudilast reduces alcohol drinking in multiple animal models of alcohol-dependence. *Addiction Biology*. 2015; 20:38–42. <http://dx.doi.org/10.1111/adb.12106>. [PubMed: 24215262]
- Bell RL, McKinzie DL, Murphy JM, McBride WJ. Sensitivity and tolerance to the motor impairing effects of moderate doses of ethanol. *Pharmacology Biochemistry and Behavior*. 2000; 67:583–586. [http://dx.doi.org/10.1016/S0091-3057\(00\)00387-7](http://dx.doi.org/10.1016/S0091-3057(00)00387-7).
- Bell RL, Rodd ZA, Engleman EA, Toalston JE, McBride WJ. Scheduled access alcohol drinking by alcohol-preferring (P) and high alcohol-drinking (HAD) rats: Modeling adolescent and adult binge-like drinking. *Alcohol*. 2014; 48:225–234. <http://dx.doi.org/10.1016/j.alcohol.2013.10.004>. [PubMed: 24290311]
- Bell RL, Rodd ZA, Lumeng L, Murphy JM, McBride WJ. The alcohol-preferring P rat and animal models of excessive alcohol drinking. *Addiction Biology*. 2006; 11:270–288. <http://dx.doi.org/10.1111/j.1369-1600.2005.00029.x>. [PubMed: 16961759]

- Bell RL, Rodd ZA, Sable HJ, Schultz JA, Hsu CC, Lumeng L, et al. Daily patterns of ethanol drinking in peri-adolescent and adult alcohol-preferring (P) rats. *Pharmacology Biochemistry and Behavior*. 2006; 83:35–46.
- Bell RL, Rodd ZA, Schultz JA, Peper CL, Lumeng L, Murphy JM, et al. Effects of short deprivation and re-exposure intervals on the ethanol drinking behavior of selectively bred high alcohol-consuming rats. *Alcohol*. 2008; 42:407–416. <http://dx.doi.org/10.1016/j.alcohol.2008.03.130>. [PubMed: 18486429]
- Bell RL, Rodd ZA, Toalston JE, McKinzie DL, Lumeng L, Li T-K, et al. Autonomic activation associated with ethanol self-administration in adult female P rats. *Pharmacology Biochemistry and Behavior*. 2008; 91:223–232. <http://dx.doi.org/10.1016/j.pbb.2008.07.016>.
- Bell RL, Rodd ZA, Smith RJ, Toalston JE, Franklin KM, McBride WJ. Modeling binge-like ethanol drinking by peri-adolescent and adult P rats. *Pharmacology Biochemistry and Behavior*. 2011; 100:90–97. <http://dx.doi.org/10.1016/j.pbb.2011.07.017>.
- Bell RL, Rodd-Henricks ZA, Webster AA, Lumeng L, Li T-K, McBride WJ, et al. Heart rate and motor activating effects of orally self-administered ethanol in alcohol-preferring (P) rats. *Alcoholism: Clinical and Experimental Research*. 2002; 26:1162–1170. <http://dx.doi.org/10.1111/j.1530-0277.2002.tb02652.x>.
- Bell RL, Sable HJK, Colombo G, Hyytia P, Rodd ZA, Lumeng L. Animal models for medications development targeting alcohol abuse using selectively bred rat lines: neurobiological and pharmacological validity. *Pharmacology Biochemistry and Behavior*. 2012; 103:119–155. <http://dx.doi.org/10.1016/j.pbb.2012.07.007>.
- Bell RL, Stewart RB, Woods JE II, Lumeng L, Li T-K, Murphy JM, McBride WJ. Responsivity and development of tolerance to the motor impairing effects of moderate doses of ethanol in Alcohol-Preferring (P) and –Nonpreferring (NP) rat lines. *Alcoholism: Clinical and Experimental Research*. 2001; 25:644–650. <http://dx.doi.org/10.1111/j.1530-0277.2001.tb02262.x>.
- Benham RS, Engin E, Rudolph U. Diversity of neuronal inhibition: a path to novel treatments for neuropsychiatric disorders. *JAMA Psychiatry*. 2014; 71:91–93. [PubMed: 24196284]
- Bertholomey ML, Verplaetse TL, Czachowski CL. Alterations in ethanol seeking and self-administration following yohimbine in selectively bred alcohol-preferring (P) and high alcohol drinking (HAD-2) rats. *Behavioural Brain Research*. 2013; 238:252–258. [PubMed: 23103404]
- Besheer J, Faccidomo S, Grondin JJM, Hodge CW. Effects of mGlu1-receptor blockade on ethanol self-administration in inbred alcohol-preferring rats. *Alcohol*. 2008a; 42:13–20. [PubMed: 18164577]
- Besheer J, Faccidomo S, Grondin JJM, Hodge CW. Regulation of motivation to self-administer ethanol by mGluR5 in alcohol-preferring (P) rats. *Alcoholism: Clinical and Experimental Research*. 2008b; 32:209–221.
- Besheer J, Grondin JJM, Cannady R, Sharko AC, Faccidomo S, Hodge CW. Metabotropic glutamate receptor 5 activity in the nucleus accumbens is required for the maintenance of ethanol self-administration in a rat genetic model of high alcohol intake. *Biological Psychiatry*. 2010; 67:812–822. [PubMed: 19897175]
- Bhaskar LV, Kumar SA. Polymorphisms in genes encoding dopamine signaling pathway and risk of alcohol dependence: a systematic review. *Acta Neuropsychiatrica*. 2014; 26:69–80. [PubMed: 24983092]
- Bhaskar LV, Thangaraj K, Kumar KP, Pardhasaradhi G, Singh L, Rao VR. Association between neuropeptide Y gene polymorphisms and alcohol dependence: a case-control study in two independent populations. *European Addiction Research*. 2013; 19:307–313. [PubMed: 23652361]
- Bice P, Foroud T, Bo R, et al. Genomic screen for QTLs underlying alcohol consumption in the P and NP rat lines. *Mammalian Genome*. 1998; 9:949–955. [PubMed: 9880658]
- Bierut LJ, Agrawal A, Bucholz KK, Doheny KF, Laurie C, Pugh E, et al. Gene, Environment Association Studies Consortium, A genome wide association study of alcohol dependence. *Proceedings of the National Academy of Sciences, USA*. 2010; 107:5082–5087.
- Bito-Onon JJ, Simms JA, Chatterjee S, Holgate J, Bartlett SE. Varenicline, a partial agonist at neuronal nicotinic acetylcholine receptors, reduces nicotine-induced increases in 20% ethanol operant self-

administration in Sprague-Dawley rats. *Addiction Biology*. 2011; 16:440–449. [PubMed: 21392178]

- Blednov YA, Walker D, Martinez M, Harris RA. Reduced alcohol consumption in mice lacking preprodynorphin. *Alcohol*. 2006; 40:73–86. [PubMed: 17307643]
- Blomqvist O, Ericson M, Johnson DH, Engel JA, Söderpalm B. Voluntary ethanol intake in the rat: effects of nicotinic acetylcholine receptor blockade or subchronic nicotine treatment. *European Journal of Pharmacology*. 1996; 314:257–267. [PubMed: 8957244]
- Blume A, Torner L, Liu Y, Subburaju S, Aguilera G, Neumann ID. Prolactin activates mitogen-activated protein kinase signaling and corticotropin releasing hormone transcription in rat hypothalamic neurons. *Endocrinology*. 2009; 150:1841–1849. [PubMed: 19022892]
- Breslin FJ, Johnson BA, Lynch WJ. Effect of topiramate treatment on ethanol consumption in rats. *Psychopharmacology (Berl)*. 2010; 207:529–534. [PubMed: 19823810]
- Brodie MS, Pesold C, Appel SB. Ethanol directly excites dopaminergic ventral tegmental area reward neurons. *Alcoholism: Clinical and Experimental Research*. 1999; 23:1848–1852.
- Brodie MS, Shefner SA, Dunwiddie TV. Ethanol increases the firing rate of dopamine neurons of the rat ventral tegmental area in vitro. *Brain Research*. 1990; 508:65–69. [PubMed: 2337793]
- Brodie MS, Trifunovi RD, Shefner SA. Serotonin potentiates ethanol-induced excitation of ventral tegmental area neurons in brain slices from three different rat strains. *Journal of Pharmacology and Experimental Therapeutics*. 1995; 273:1139–1146. [PubMed: 7791084]
- Brunelle C, Barrett SP, Pihl RO. Psychostimulant users are sensitive to the stimulus properties of alcohol as indexed by alcohol-induced cardiac reactivity. *Psychology of Addictive Behavior*. 2006; 20:478–483.
- Buisman-Pijlman FT, Sumracki NM, Gordon JJ, Hull PR, Carter CS, Tops M. Individual differences underlying susceptibility to addiction: role for the endogenous oxytocin system. *Pharmacology, Biochemistry and Behavior*. 2014; 119:22–38.
- Burke AR, Miczek KA. Stress in adolescence and drugs of abuse in rodent models: role of dopamine, CRF, and HPA axis. *Psychopharmacology*. 2014; 231:1557–1580. [PubMed: 24370534]
- Bymaster FP, McKinzie DL, Felder CC, Wess J. Use of M1–M5 muscarinic receptor knockout mice as novel tools to delineate the physiological roles of the muscarinic cholinergic system. *Neurochemistry Research*. 2003; 28:437–42.
- Cadoret RJ. Familial transmission of psychiatric disorders associated with alcoholism. In: Begleiter, H.; Kissin, B., editors. *The Genetics of Alcoholism*. Oxford University Press; New York, NY: 1995. p. 70–81.
- Calabresi P, Lacey MG, North RA. Nicotinic excitation of rat ventral tegmental neurones in vitro studied by intracellular recording. *British Journal of Pharmacology*. 1989; 98:135–140. [PubMed: 2804543]
- Campbell RE, Grove KL, Smith MS. Distribution of corticotropin releasing hormone receptor immunoreactivity in the rat hypothalamus: coexpression in neuropeptide Y and dopamine neurons in the arcuate. *Brain Research*. 2003; 973:223–232. [PubMed: 12738066]
- Cao J, LaRocque E, Li D. Associations of the 5-hydroxytryptamine (serotonin) receptor 1B gene (HTR1B) with alcohol, cocaine, and heroin abuse. *American Journal of Medical Genetics B Neuropsychiatric Genetics*. 2013; 162B:169–176. [PubMed: 23335468]
- Cao J, Liu X, Han S, Zhang CK, Liu Z, Li D. Association of the HTR2A gene with alcohol and heroin abuse. *Human Genetics*. 2014; 133:357–365. [PubMed: 24178752]
- Carlezon WA, Wise RA. Microinjections of phencyclidine (PCP) and related drugs into nucleus accumbens shell potentiate medial forebrain bundle brain stimulation reward. *Psychopharmacology*. 1996; 128:413–420. [PubMed: 8986012]
- Carr LG, Foroud T, Bice P, et al. A quantitative trait locus for alcohol consumption in selectively bred rat lines. *Alcoholism: Clinical and Experimental Research*. 1998; 22:884–887.
- Cederlof R, Friberg L, Lundman T. The interaction of smoking, environment and heredity and their implications for disease aetiology: a report of epidemiological studies on the Swedish twin registries. *Acta Medica Scandinavia*. 1977; 612(Suppl 1):1–128.

- Chandler LJ, Newsom H, Sumners C, Crews F. Chronic ethanol exposure potentiates NMDA excitotoxicity in cerebral cortical neurons. *Journal of Neurochemistry*. 1993; 60:1578–1581. [PubMed: 8455043]
- Chandler DJ, Waterhouse BD, Gao WJ. New perspectives on catecholaminergic regulation of executive circuits: evidence for independent modulation of prefrontal functions by midbrain dopaminergic and noradrenergic neurons. *Frontiers in Neural Circuits*. 2014; 8:53. [PubMed: 24904299]
- Chappell AM, Weiner JL. Relationship between ethanol's acute locomotor effects and ethanol self-administration in male Long-Evans rats. *Alcoholism: Clinical and Experimental Research*. 2008; 32:2088–2099.
- Charbogne P, Kieffer BL, Befort K. 15 years of genetic approaches in vivo for addiction research: opioid receptor and peptide gene knockout in mouse models of drug abuse. *Neuropharmacology*. 2014; 76:204–217. [PubMed: 24035914]
- Chatterjee S, Bartlett SE. Neuronal nicotinic acetylcholine receptors as pharmacotherapeutic targets for the treatment of alcohol use disorders. *CNS Neurological Discovery of Drug Targets*. 2010; 9:60–76.
- Chen AC, Manz N, Tang Y, Rangaswamy M, Almasy L, Kuperman S, Porjesz B, et al. Single-nucleotide polymorphisms in corticotropin releasing hormone receptor 1 gene (CRHR1) are associated with quantitative trait of event-related potential and alcohol dependence. *Alcoholism: Clinical and Experimental Research*. 2010; 34:988–996.
- Chen AC, Tang Y, Rangaswamy M, Wang JC, Almasy L, Foroud T, Porjesz B, et al. Association of single nucleotide polymorphisms in a glutamate receptor gene (GRM8) with theta power of event-related oscillations and alcohol dependence. *American Journal of Medical Genetics Part B*. 2009; 150:359–368.
- Cherpitel CJ, Moskalewicz J, Swiatkiewicz G. Drinking patterns and problems in emergency services in Poland. *Alcohol and Alcoholism*. 2004; 39:256–261. [PubMed: 15082465]
- Chester JA, Blose AM, Froehlich JC. Acoustic startle reactivity during acute alcohol withdrawal in rats that differ in genetic predisposition toward alcohol drinking: effect of stimulus characteristics. *Alcoholism: Clinical and Experimental Research*. 2004; 28:677–687.
- Chiavegatto S, Quadros IMH, Ambar G, Miczek KA. Individual vulnerability to escalated aggressive behavior by a low dose of alcohol: Decreased serotonin receptor mRNA in the prefrontal cortex of male mice. *Genes Brain and Behavior*. 2010; 9:110–119.
- Childress AR, Hole AV, Ehrman RN, Robbins SJ, McLellan AT, O'Brien CP. Cue reactivity and cue reactivity interventions in drug dependence. *NIDA Research Monograph*. 1993; 137:73–95. [PubMed: 8289929]
- Ciccocioppo R, Angeletti S, Colombo G, Gessa GL, Massi M. Autoradiographic analysis of 5-HT_{2A} binding sites in the brain of Sardinian alcohol-preferring and nonpreferring rats. *European Journal of Pharmacology*. 1999; 373:13–19. [PubMed: 10408247]
- Ciccocioppo R, Ge J, Barnes NM, Cooper SJ. Central 5-HT₃ receptors in P and in AA alcohol-preferring rats: An autoradiographic study. *Brain Research Bulletin*. 1998; 46:311–315. [PubMed: 9671260]
- Cicero, TJ. A critique of animal analogues of alcoholism. In: Majchrowicz, E.; Noble, EP., editors. *Biochemistry and Pharmacology of Ethanol*. Vol. 2. Plenum Press; New York, NY: 1979. p. 533–560.
- Clausen RP, Madsen K, Larsson OM, Frolund B, Krogsgaard-Larsen P, Schousboe A. Structure-activity relationship and pharmacology of gamma-aminobutyric acid (GABA) transport inhibitors. *Advances in Pharmacology*. 2006; 54:265–284. [PubMed: 17175818]
- Clerget G, Bouguignon-Igel V, Rederstorff M. Alcoholic precipitation of small non-coding RNAs. *Methods in Molecular Biology*. 2015; 1296:11–16. [PubMed: 25791586]
- Clifford CA, Hopper JL, Fulker DW, Murray RM. A genetic and environmental analysis of a twin family study of alcohol use, anxiety, and depression. *Genetic Epidemiology*. 1984; 1:63–79. [PubMed: 6544233]
- Cloninger CR. Neurogenetic adaptive mechanisms in alcoholism. *Science*. 1987; 236:410–416. [PubMed: 2882604]

- Coleman J, Williams A, Phan TH, Mummalaneni S, Melone P, Ren Z, Lyall V, et al. Strain differences in the neural, behavioral, and molecular correlates of sweet and salty taste in naive, ethanol- and sucrose-exposed P and NP rats. *Journal of Neurophysiology*. 2011; 106:2606–2621. [PubMed: 21849614]
- Conrod PJ, Pihl RO, Stewart SH, Dongier M. Validation of a system of classifying female substance abusers on the basis of personality and motivational risk factors for substance abuse. *Psychology of Addictive Behavior*. 2000; 14:243–256.
- Conrod PJ, Pihl RO, Vassileva J. Differential sensitivity to alcohol reinforcement in groups of men at risk for distinct alcoholism subtypes. *Alcoholism: Clinical and Experimental Research*. 1998; 22:585–597.
- Contini V, Bertuzzi GP, Polina ER, Hunemeier T, Hendler EM, Hutz MH, Bau CH. A haplotype analysis is consistent with the role of functional HTR1B variants in alcohol dependence. *Drug and Alcohol Dependence*. 2012; 122:100–104. [PubMed: 22005095]
- Coonfield DL, Kiefer SW, Ferraro FM III, Sinclair JD. Ethanol palatability and consumption by high ethanol-drinking rats: manipulation of the opioid system with naltrexone. *Behavioral Neuroscience*. 2004; 118:1189–1196.
- Cooper, JR.; Bloom, FE.; Roth, RH. *Biochemical Basis of Neuropharmacology*. 8. Oxford University Press; Oxford, UK: 2002.
- Corrigall WA, Coen KM. Nicotine self-administration and locomotor activity are not modified by the 5-HT₃ antagonists ICS 205-930 and MDL 72222. *Pharmacology, Biochemistry and Behavior*. 1994; 49:67–71.
- Cosgrove KP. Imaging receptor changes in human drug abusers. *Current Topics in Behavioral Neuroscience*. 2010; 3:199–217.
- Cotton NS. The familial incidence of alcoholism. *Journal of Studies on Alcohol*. 1979; 40:89–116. [PubMed: 376949]
- Cowen MS, Chen F, Lawrence AJ. Neuropeptides: implications for alcoholism. *Journal of Neurochemistry*. 2004; 89:273–285. [PubMed: 15056272]
- Cowen MS, Djouma E, Lawrence AJ. The metabotropic glutamate 5 receptor antagonist 3-[(2-Methyl-1,3-thiazol-4-yl) ethynyl]-pyridine reduces ethanol self-administration in multiple strains of alcohol-preferring rats and regulates olfactory glutamatergic systems. *Journal of Pharmacology and Experimental Therapeutics*. 2005; 315:590–600. [PubMed: 16014750]
- Cozzoli DK, Goulding SP, Zhang P-W, Xiao B, Hu J-H, Ary AW, Szumlinski KK, et al. Binge drinking up-regulated accumbens mGluR5-Homer2-PI3K signaling: functional implications for alcoholism. *Journal of Neuroscience*. 2009; 29:8655–8668. [PubMed: 19587272]
- Crabbe JC. Neurogenetic studies of alcohol addiction. *Philosophical Transactions of the Royal Society of London B Biological Science*. 2008; 363:3201–3211.
- Crabbe JC, Bell RL, Ehlers CL. Human and laboratory rodent low response to alcohol: Is better consilience possible? *Addiction Biology*. 2010; 15:125–144. [PubMed: 20148776]
- Crabbe JC, Phillips TJ, Kosobud A, Belknap JK. Estimation of genetic correlation: interpretation of experiments using selectively bred and inbred animals. *Alcoholism: Clinical and Experimental Research*. 1990; 14:141–151.
- Crews FT, Qin L, Sheedy D, Ventreno RP, Zou J. High mobility group box 1/Toll-like receptor danger signaling increases brain neuroimmune activation in alcohol dependence. *Biological Psychiatry*. 2013; 73:602–612. [PubMed: 23206318]
- Cui C, Noronha A, Morikawa H, Alvarez VA, Stuber GD, Szumlinski KK, et al. New insights on neurobiological mechanisms underlying alcohol addiction. *Neuropharmacology*. 2013; 67:223–232. [PubMed: 23159531]
- Cui C, Shurtleff D, Harris RA. Neuroimmune mechanisms of alcohol and drug addiction. *International Review of Neurobiology*. 2014; 118:1–12. [PubMed: 25175859]
- Czachowski CL, Samson HH. Ethanol- and sucrose-reinforced appetitive and consummatory responding in HAD1, HAD2, and P rats. *Alcohol Clin Exp Res*. 26:1653–1661. [PubMed: 12436053]
- Danbolt NC. (2001) Glutamate uptake. *Progress in Neurobiology*. 2002; 65:1–105. [PubMed: 11369436]

- Das SC, Yamamoto BK, Hristov AM, Sari Y. Ceftriaxone attenuates ethanol drinking and restores extracellular glutamate concentration through normalization of GLT-1 in nucleus accumbens of male alcohol-preferring rats. *Neuropharmacology*. 2015; 97:67–74. [PubMed: 26002627]
- Davis TJ, de Fiebre CM. Alcohol's actions on neuronal nicotinic acetylcholine receptors. *Alcohol Research Health*. 2006; 29:179–185. [PubMed: 17373406]
- Davis KM, Wu JY. Role of glutamatergic and GABAergic systems in alcoholism. *Journal of Biomedical Science*. 2001; 8:7–19. [PubMed: 11173971]
- Dawson DA, Grant BF, Stinson FS, Chou PS. Another look at heavy episodic drinking and alcohol use disorders among college and noncollege youth. *Journal of Studies on Alcohol*. 2004; 65:477–488. [PubMed: 15378804]
- Deehan GA Jr, Hauser SR, Waeiss RA, Knight CP, Toalston JE, Truitt WA, et al. Co-administration of ethanol and nicotine: the enduring alterations in the rewarding properties of nicotine and glutamate activity within the mesocorticolimbic system of female alcohol-preferring (P) rats. *Psychopharmacology (Berl)*. 2015; 232:4293–4302. [PubMed: 26306917]
- Dhaheer R, Toalston JE, Hauser SR, Bell RL, McKinzie DL, McBride WJ, et al. Effects of naltrexone and LY255582 on ethanol maintenance, seeking, and relapse responding by alcohol-preferring (P) rats. *Alcohol*. 2012; 46:17–27. [PubMed: 21962974]
- Di Chiara G, Imperato A. Drugs abused by humans preferentially increase synaptic dopamine concentrations in the mesolimbic system of freely moving rats. *Proceedings of the National Academy of Sciences, USA*. 1988; 85:5274–5278.
- Dick DM, Bierut LJ. The genetics of alcohol dependence. *Current Psychiatry Reports*. 2006; 8:151–157. [PubMed: 16539893]
- Dick DM, Edenberg HJ, Xuei X, Goate A, Kuperman S, Schuckit M, et al. Association of GABRG3 with alcohol dependence. *Alcoholism: Clinical and Experimental Research*. 2004; 28:4–9.
- Dick DM, Plunkett J, Wetherill LF, Xuei X, Goate A, Hesselbrock V, et al. Association between GABRA1 and drinking behaviors in the collaborative study on the genetics of alcoholism sample. *Alcoholism: Clinical and Experimental Research*. 2006; 30:1101–1110.
- Ding ZM, Engleman EA, Rodd ZA, McBride WJ. Ethanol increases glutamate neurotransmission in the posterior ventral tegmental area of female Wistar rats. *Alcoholism: Clinical and Experimental Research*. 2012; 36:633–40.
- Ding ZM, Rodd ZA, Engleman EA, Bailey JA, Lahiri DK, McBride WJ. Alcohol drinking and deprivation alter basal extracellular glutamate concentrations and clearance in the mesolimbic system of alcohol-preferring (P) rats. *Addiction Biology*. 2013; 18:297–306. [PubMed: 23240885]
- Domart MC, Benyamina A, Lemoine A, Bourgain C, Blecha L, Debuire B, et al. Association between a polymorphism in the promoter of a glutamate receptor subunit gene (GRIN2A) and alcoholism. *Addiction Biology*. 2012; 17:783–785. [PubMed: 21507155]
- Draski, LJ.; Deitrich, RA. Initial effects of ethanol on the central nervous system. In: Deitrich, RA.; Erwin, VG., editors. *Pharmacological Effects of Ethanol on the Nervous System*. CRC Press; Boca Raton, FL: 1996. p. 227-250.
- Drews E, Zimmer A. Modulation of alcohol and nicotine responses through the endogenous opioid system. *Progress in Neurobiology*. 1997; 90:1–15. [PubMed: 19800387]
- Drummond DC, Cooper T, Glautier S. Conditioned learning in alcohol dependence: Implications for cue exposure treatment. *British Journal of Addiction*. 1990; 85:725–743. [PubMed: 2198966]
- Dumont Y, Satoh H, Cadieux A, et al. Evaluation of truncated neuropeptide Y analogues with modifications of the tyrosine residue in position 1 on Y1, Y2 and Y3 receptor sub-types. *European Journal of Pharmacology*. 1993; 238:37–45. [PubMed: 8405081]
- Dundon W, Lynch KG, Pettinati HM, Lipkin C. Treatment outcomes in type A and B alcohol dependence 6 months after serotonergic pharmacotherapy. *Alcoholism: Clinical and Experimental Research*. 2004; 28:1065–1073.
- Dyr W, Kostowski W. Warsaw high-preferring (WHP) and Warsaw low-preferring (WLP) lines of rats selectively bred for high and low voluntary ethanol intake: preliminary phenotypic characterization. *Alcohol*. 2008; 42:161–170. [PubMed: 18420111]

- Economidou D, Hansson AC, Weiss F, Terasmaa A, Sommer WH, Cippitelli A, et al. Dysregulation of nociceptin/orphanin FQ activity in the amygdala is linked to excessive alcohol drinking in the rat. *Biological Psychiatry*. 2008; 64:211–218. [PubMed: 18367152]
- Edenberg HJ, Dick DM, Xuei X, Tian H, Almasy L, Bauer LO, Begleiter H, et al. Variations in GABRA2, encoding the alpha 2 subunit of the GABA(A) receptor are associated with alcohol dependence and with brain oscillations. *American Journal of Human Genetics*. 2004; 74:705–714. [PubMed: 15024690]
- Ehlers CL, Chaplin RI, Wall TL, Lumeng L, Li T-K, Owens MJ, et al. Corticotropin releasing factor (CRF): Studies in alcohol-preferring and -nonpreferring rats. *Psychopharmacology*. 1992; 106:359–364. [PubMed: 1570383]
- Ehlers CL, Li TK, Lumeng L, et al. Neuropeptide Y levels in ethanol-naive alcohol-preferring and nonpreferring rats and in Wistar rats after ethanol exposure. *Alcoholism: Clinical and Experimental Research*. 1998; 22:1778–1782.
- Ehlers CL, Lind PA, Wilhelmsen KC. Association between single nucleotide polymorphisms in the mu opioid receptor gene (OPRM1) and self-reported responses to alcohol in American Indians. *BMC Medical Genetics*. 2008; 9:35.10.1186/1471-2350-9-35 [PubMed: 18433502]
- Ehlers CL, Walter NAR, Dick DM, Buck KJ, Crabbe JC. A comparison of selected quantitative trait loci associated with alcohol use phenotypes in humans and mouse models. *Addiction Biology*. 2010; 15:185–199. [PubMed: 20148779]
- Ehrenreich H, Schuck J, Stender N, et al. Endocrine and hemodynamic effects of stress versus systemic CRF in alcoholics during early and medium term abstinence. *Alcoholism: Clinical and Experimental Research*. 1997; 21:1285–1293.
- Eiler WA II, June HL. Blockade of GABAA receptors within the extended amygdala attenuates D2 regulation of alcohol-motivated behaviors in the ventral tegmental area of alcohol-preferring (P) rats. *Neuropharmacology*. 2007; 52:1570–1579. [PubMed: 17451754]
- Eiler WJA II, Seyoum R, Foster KL, Mailey C, June HL. D1 dopamine receptor regulates alcohol-motivated behaviors in the bed nucleus of the stria terminalis in alcohol-preferring (P) rats. *Synapse*. 2003; 48:45–56. [PubMed: 12557272]
- Elman I, Borsook D, Volkow ND. Pain and suicidality: insights from reward and addiction neuroscience. *Progress in Neurobiology*. 2013; 109:1–27. [PubMed: 23827972]
- Eng MY, Schuckit MA, Smith TL. The level of response to alcohol in daughters of alcoholics and controls. *Drug and Alcohol Dependence*. 2005; 79:83–93. [PubMed: 15943947]
- Engel JA, Jerlhag E. Alcohol: mechanisms along the mesolimbic dopamine system. *Progress in Brain Research*. 2014; 211:201–233. [PubMed: 24968782]
- Engleman EA, Ding Z-M, Oster SM, Toalston JE, Bell RL, Murphy JM, McBride WJ, Rodd ZA. Ethanol is self-administered into the nucleus accumbens shell, but not core: Evidence of genetic sensitivity. *Alcoholism: Clinical and Experimental Research*. 2009; 33:2162–2171. <http://dx.doi.org/10.1111/j.1530-0277.2009.01055.x>.
- Engleman EA, Ingraham CM, McBride WJ, Lumeng L, Murphy JM. Extracellular dopamine levels are lower in the medial prefrontal cortex of alcohol-preferring rats compared to Wistar rats. *Alcohol*. 2006; 38:5–12. [PubMed: 16762687]
- Engleman EA, Keen EJ, Tilford SS, Thielen RJ, Morzorati SL. Ethanol drinking reduces extracellular dopamine levels in the posterior ventral tegmental area of nondependent alcohol-preferring rats. *Alcohol*. 2011; 45:549–557. [PubMed: 21827929]
- Engleman EA, McBride WJ, Wilber AA, Shaikh SR, Eha RD, Lumeng L, et al. Reverse microdialysis of a dopamine uptake inhibitor in the nucleus accumbens of alcohol-preferring rats: effects on dialysate dopamine levels and ethanol intake. *Alcoholism: Clinical and Experimental Research*. 2000; 24:795–801.
- Engleman EA, McBride WJ, Li TK, Lumeng L, Murphy JM. Ethanol drinking experience attenuates (–)sulpiride-induced increases in extracellular dopamine levels in the nucleus accumbens of alcohol-preferring (P) rats. *Alcoholism: Clinical and Experimental Research*. 2003; 27:424–31.
- Engleman EA, Rodd ZA, Bell RL, Murphy JM. The role of 5-HT3 receptors in drug abuse and as a target for pharmacotherapy. *CNS Neurological Discovery Drug Targets*. 2008; 7:454–467.

- Enna SJ. GABAB receptor agonists and antagonists: pharmacological properties and therapeutic possibilities. *Expert Opinion in Investigational Drugs*. 1997; 6:1319–25.
- Enoch MA, Schwartz L, Albaugh B, Virkkunen M, Goldman D. Dimensional anxiety mediates linkage of GABRA2 haplotypes with alcoholism. *American Journal of Medical Genetics B Neuropsychiatric Genetics*. 2006; 141B:599–607. [PubMed: 16874763]
- Enoch MA, Zhou Z, Kimura M, Mash DC, Yuan Q, Goldman D. GABAergic gene expression in postmortem hippocampus from alcoholics and cocaine addicts; corresponding findings in alcohol-naïve P and NP rats. *PLoS One*. 2012; 7:e29369. [PubMed: 22253714]
- Epstein EE, Kahler CW, McCrady BS, Lewis KD, Lewis S. An empirical classification of drinking patterns among alcoholics: binge, episodic, sporadic, and steady. *Addictive Behaviors*. 1995; 20:23–41. [PubMed: 7785480]
- Ericson M, Blomqvist O, Engel JA, Söderpalm B. Voluntary ethanol intake in the rat and the associated accumbal dopamine overflow are blocked by ventral tegmental mecamylamine. *European Journal of Pharmacology*. 1998; 358:189–196. [PubMed: 9822883]
- Ericson M, Löf E, Stomberg R, Söderpalm B. The smoking cessation medication varenicline attenuates alcohol and nicotine interactions in the rat mesolimbic dopamine system. *Journal of Pharmacology and Experimental Therapeutics*. 2009; 329:225–230. [PubMed: 19126781]
- Ericson M, Molander A, Lof E, Engel JA, Soderpalm B. Ethanol elevates accumbal dopamine levels via indirect activation of ventral tegmental nicotinic acetylcholine receptors. *European Journal of Pharmacology*. 2003; 467:85–93. [PubMed: 12706460]
- Everitt BJ, Robbins TW. Neural systems of reinforcement for drug addiction: from actions to habits to compulsion. *Nature Neuroscience*. 2005; 8:1481–1489. [PubMed: 16251991]
- Eysenck, HJ.; Eysenck, SBG. *Manual of the Eysenck Personality Questionnaire*. Hodder & Stoughton; London: 1975.
- Fadda F, Rossetti ZL. Chronic ethanol consumption: from neuroadaptation to neurodegeneration. *Progress in Neurobiology*. 1998; 56:385–431. [PubMed: 9775400]
- Faria RR, Rueda AVL, Sayuri C, Soares SL, Malta MB, Carrara-Nascimento PF, et al. Environmental modulation of ethanol-induced locomotor activity: Correlation with neuronal activity in distinct brain regions of adolescent and adult Swiss mice. *Brain Research*. 2008; 1239:127–140. [PubMed: 18789904]
- Farris SP, Mayfield RD. RNA-seq reveals novel transcriptional reorganization in human alcoholic brain. *International Reviews in Neurobiology*. 2014; 116:275–300.
- Fetissov SO, Kopp J, Hokfelt T. Distribution of NPY receptors in the hypothalamus. *Neuropeptides*. 2004; 38:175–188. [PubMed: 15337370]
- Files FJ, Samson HH, Denning CE, Marvin S. Comparison of alcohol-preferring and -nonpreferring selectively bred rat lines. II. Operant self-administration in a continuous-access situation. *Alcoholism: Clinical and Experimental Research*. 1998; 22:2147–2158.
- Filip M, Zaniewska M, Frankowska M, Wydra K, Fuxe K. The importance of adenosine A(2A) receptor-dopamine D(2) receptor interaction in drug addiction. *Current Medicinal Chemistry*. 2012; 19:317–355. [PubMed: 22335511]
- Finn PR, Justus A. Physiological responses in sons of alcoholics. *Alcohol Health Research World*. 1997; 21:227. [PubMed: 15706773]
- Finn PR, Earleywine M, Pihl RO. Sensation seeking, stress reactivity, and alcohol dampening discriminate the density of a family history of alcoholism. *Alcoholism: Clinical and Experimental Research*. 1992; 16:585–590.
- Fitzgerald GJ, Liu H, Morzorati SL. Decreased sensitivity of NMDA receptors on dopaminergic neurons from the posterior ventral tegmental area following chronic nondependent alcohol consumption. *Alcoholism: Clinical and Experimental Research*. 2012; 36:1710–1719.
- Flavin SA, Winder DG. Noradrenergic control of the bed nucleus of the stria terminalis in stress and reward. *Neuropharmacology*. 2013; 70:324–330. [PubMed: 23466330]
- Flory K, Lynam D, Milich R, Leukefeld C, Clayton R. Early adolescent through young adult alcohol and marijuana use trajectories: Early predictors, young adult outcomes, and predictive utility. *Developmental Psychopathology*. 2004; 16:193–213.

- Floyd DW, Jung KY, McCool BA. Chronic ethanol ingestion facilitates N-methyl-D aspartate receptor function and expression in rat lateral/basolateral amygdala neurons. *Journal of Pharmacology and Experimental Therapeutics*. 2003; 307:1020–1029. [PubMed: 14534353]
- Foroud T, Wetherill LF, Liang T, Dick DM, Hesselbrock V, Kramer J, Edenberg HJ, et al. Association of alcohol craving with alpha-synuclein (SNCA). *Alcoholism: Clinical and Experimental Research*. 2007; 31:537–545.
- Forray A, Sofuoglu M. Future pharmacological treatments for substance use disorders. *British Journal of Clinical Pharmacology*. 2014; 77:382–400. [PubMed: 23039267]
- Foster KL, McKay PF, Seyoum R, Milbourne D, Yin W, Sarma PVVS, et al. GABA_A and opioid receptors of the central nucleus of the amygdala selectively regulates ethanol-maintained behaviors. *Neuropsychopharmacology*. 2004; 29:269–284. [PubMed: 14666116]
- Frances F, Guillen M, Verdu F, Portoles O, Castello A, Sorli JV, Corella D. The 1258 G>A polymorphism in the neuropeptide Y gene is associated with greater alcohol consumption in a Mediterranean population. *Alcohol*. 2011; 45:131–136. [PubMed: 21303710]
- Franklin KM, Asatryan L, Jakowec MW, Trudell JR, Bell RL, Davies DL. P2X4 receptors (P2X4Rs) represent a novel target for the development of drugs to prevent and/or treat alcohol use disorders. *Frontiers in Neuroscience/Neuropharmacology*. 2014; 8:176. <http://dx.doi.org/10.3389/fnins.2014.00176>.
- Franklin KM, Engleman EA, Ingraham CM, McClaren JA, Keith CM, McBride WJ, Murphy JM. A single, moderate ethanol exposure alters extracellular dopamine levels and dopamine receptor function in the nucleus accumbens of Wistar rats. *Alcoholism: Clinical and Experimental Research*. 2009; 33:1721–1730.
- Franklin KM, Hauser SR, Lasek AW, Bell RL, McBride WJ. Involvement of purinergic P2X4 receptors in alcohol intake of high alcohol-drinking (HAD) rats. *Alcoholism: Clinical and Experimental Research*. 2015; 39:2022–2031. <http://dx.doi.org/10.1111/acer.12836>.
- Fritschy J-M, Panzanelli P, Tyagarajan SK. Molecular and functional heterogeneity of GABAergic synapses. *Cellular and Molecular Life Sciences*. 2012; 69:2485–2499. [PubMed: 22314501]
- Froehlich JC, Hausauer BJ, Federoff DL, Fischer SM, Rasmussen DD. Prazosin reduces alcohol drinking throughout prolonged treatment and blocks the initiation of drinking in rats selectively bred for high alcohol intake. *Alcoholism: Clinical and Experimental Research*. 2013; 37:1552–1560.
- Froehlich JC, Hausauer BJ, Rasmussen DD. Combining naltrexone and prazosin in a single oral medication decreases alcohol drinking more effectively than does either drug alone. *Alcoholism: Clinical and Experimental Research*. 2013; 37:1763–1770.
- Froehlich JC, Hausauer B, Fischer S, Wise B, Rasmussen DD. Prazosin Reduces Alcohol Intake in an Animal Model of Alcohol Relapse. *Alcoholism: Clinical and Experimental Research*. 2015; 39:1538–1546.
- Gaiarsa J-L, Kuczewski N, Porcher C. Contribution of metabotropic GABAB receptors to neuronal network construction. *Pharmacology Therapeutics*. 2011; 132:170–179. [PubMed: 21718720]
- Gass JT, Olive MF. Glutamatergic substrates of drug addiction and alcoholism. *Biochemistry and Pharmacology*. 2008; 75:218–265.
- Gatto GJ, Murphy JM, Waller MB, McBride WJ, Lumeng L, Li T-K. Chronic ethanol tolerance through free-choice drinking in the P line of alcohol-preferring rats. *Pharmacology, Biochemistry and Behavior*. 1987; 28:111–115.
- Gedik H, Erdal ME, Yilmaz SG, Sengul C, Sengul CB, Herken H. Association of microRNA biogenesis pathway gene variants and alcohol dependence risk. *DNA and Cellular Biology*. 2015; 34:220–226.
- Gegelashvili G, Schousboe A. High affinity glutamate transporters: regulation of expression and activity. *Molecular Pharmacology*. 1997; 52:6–15. [PubMed: 9224806]
- Geiger MJ, Neufang S, Stein DJ, Domschke K. Arousal and the attentional network in panic disorder. *Human Psychopharmacology*. 2014; 29:599–603. [PubMed: 25311787]
- George DT, Gilman J, Hersch J, Thorsell A, Herion D, Geyer C, et al. Neurokinin 1 receptor antagonism as a possible therapy for alcoholism. *Science*. 2008; 319:1536–1539. [PubMed: 18276852]

- Gerra G, Leonardi C, Cortese E, D'Amore A, Lucchini A, Strepparola G, Donnini C, et al. Human kappa opioid receptor gene (OPRK1) polymorphism is associated with opiate addiction. *American Journal of Medical Genetics Part B Neuropsychiatric Genetics*. 2007; 144B:771–775.
- Gessa GL, Muntoni F, Collu M, Vargiu L, Mereu G. Low doses of ethanol activate dopaminergic neurons in the ventral tegmental area. *Brain Research*. 1985; 348:201–203. [PubMed: 2998561]
- Getachew B, Hauser SR, Taylor RE, Tizabi Y. Alcohol-induced depressive-like behavior is associated with cortical norepinephrine reduction. *Pharmacology, Biochemistry and Behavior*. 2010; 96:395–401.
- Gianoulakis, C.; Guillaume, P.; De Waele, J-P.; Angelogianni, P. Effects of stress and alcohol on the proopiomelanocortin/ β -endorphin system. In: Hunt, WA.; Zakhari, S., editors. *Stress, Gender, and Alcohol-Seeking Behavior*. National Institute on Alcohol Abuse and Alcoholism; Bethesda, MD: 1995. p. 145-165. Research Monograph No. 29
- Gilpin NW, Herman MA, Roberto M. The central amygdala as an integrative hub for anxiety and alcohol use disorders. *Biological Psychiatry*. 2015; 77:859–869. [PubMed: 25433901]
- Gilpin NW, Koob GF. Effects of β -adrenoceptor antagonists on alcohol drinking by alcohol-dependent rats. *Psychopharmacology (Berl)*. 2010; 212:431–439. [PubMed: 20676608]
- Gilpin NW, Richardson HN, Koob GF. Effects of CRF1-receptor and opioid-receptor antagonists on dependence-induced increases in alcohol drinking by alcohol-preferring (P) rats. *Alcoholism: Clinical and Experimental Research*. 2008; 32:1535–1542.
- Gilpin NW, Stewart RB, Murphy JM, Badia-Elder NE. Neuropeptide Y in the paraventricular nucleus of the hypothalamus increases ethanol intake in high- and low-alcohol-drinking rats. *Alcoholism: Clinical and Experimental Research*. 2004; 28:1492–1498.
- Gilpin NW, Stewart RB, Badia-Elder NE. Neuropeptide Y administration into the amygdala suppresses ethanol drinking in alcohol-preferring (P) rats following multiple deprivations. *Pharmacology, Biochemistry and Behavior*. 2008; 90:470–474.
- Gilpin NW, Stewart RB, Murphy JM, Li TK, Badia-Elder NE. Neuropeptide Y reduces oral ethanol intake in alcohol-preferring (P) rats following a period of imposed ethanol abstinence. *Alcoholism: Clinical and Experimental Research*. 2003; 27:787–794.
- Gobrogge KL. Sex, drugs, and violence: neuromodulation of attachment and conflict in voles. *Current Topics in Behavioral Neuroscience*. 2014; 17:229–264. http://dx.doi.org/10.1007/7854_2013_264.
- Golan, DE.; Tashjian, AH.; Armstrong, EJ.; Armstrong, AW. *Principles of Pharmacology: The Pathophysiologic Basis of Drug Therapy*. 3. Lippincott Williams & Wilkins; Philadelphia, PA: 2012.
- Golbidi S, Frisbee JC, Laher I. Chronic stress impacts the cardiovascular system: animal models and clinical outcomes. *American Journal of Physiology Heart Circulation and Physiology*. 2015; 308:H1476–H1498. <http://dx.doi.org/10.1152/ajpheart.00859.2014>.
- Gold PW. The organization of the stress system and its dysregulation in depressive illness. *Molecular Psychiatry*. 2015; 20:32–47. [PubMed: 25486982]
- Goldberg J, Eisen SA, True WR, Henderson WG. A twin study of the effects of the Vietnam conflict on alcohol drinking patterns. *American Journal of Public Health*. 1990; 80:570–574. [PubMed: 2327534]
- Goldman D, Oroszi G, O'Malley S, Anton R. COMBINE genetics study: the pharmacogenetics of alcoholism treatment response: genes and mechanisms. *Journal of Studies on Alcohol*. 2005; 15(Suppl):56–64.
- Gonzalez-Burgos G. GABA transporter GAT1: a crucial determinant of GABAB receptor activation in cortical circuits? *Advances in Pharmacology*. 2010; 58:175–204. [PubMed: 20655483]
- Goodman A. Neurobiology of addiction. An integrative review. *Biochemistry and Pharmacology*. 2008; 75:266–322.
- Goodwani S, Rao PS, Bell RL, Sari Y. Amoxicillin and amoxicillin/clavulanate reduce ethanol intake and increase GLT-1 expression as well as AKT phosphorylation in mesocorticolimbic regions. *Brain Research*. 2015; 1622:397–408. [PubMed: 26168897]
- Gorini, G.; Bell, RL.; Mayfield, RD. Molecular targets of alcohol action: Translational research for pharmacotherapy development and screening. In: Rahman, S., editor. *Progress in Molecular*

Biology and Translational Science: The Brain as a Drug Target. Vol. 98. Elsevier Science; New York, NY: p. 293-347.

- Gorini G, Harris RA, Mayfield RD. Proteomic approaches and identification of novel therapeutic targets for alcoholism. *Neuropsychopharmacology*. 2014; 39:104–130. [PubMed: 23900301]
- Gotti C, Clementi F, Fornari A, Gaimarri A, Guiducci S, Manfredi I, et al. Structural and functional diversity of native brain nicotinic receptors. *Biochemistry and Pharmacology*. 2009; 78:703–711.
- Grace AA, Floresco SB, Goto Y, Lodge DJ. Regulation of firing of dopaminergic neurons and control of goal-directed behaviors. *Trends in Neuroscience*. 2007; 30:220–227.
- Gray JC, MacKillop J. Interrelationships among individual differences in alcohol demand, impulsivity, and alcohol misuse. *Psychology of Addictive Behaviors*. 2014; 28:282–287. <http://dx.doi.org/10.1037/a0032766>. [PubMed: 23915370]
- Greeley JD, Swift W, Prescott J, Heather N. Reactivity to alcohol-related cues in heavy and light drinkers. *Journal of Studies on Alcohol*. 1993; 54:359–368. [PubMed: 8487545]
- Griffin JF. A strategic approach to vaccine development: Animal models, monitoring vaccine efficacy, formulation and delivery. *Advances in Drug Delivery Reviews*. 2002; 54:851–861.
- Grzywacz A, Małecka I, Suchanecka A, Bie kowski P, Samochowiec J. Family-based and case-control study of glutamate receptor GRIK3 Ser310Ala polymorphism in alcohol dependence. *European Addiction Research*. 2013; 19:55–59. [PubMed: 23006490]
- Guan XM, McBride WJ. Serotonin microinfusion into the ventral tegmental area increases accumbens dopamine release. *Brain Research Bulletin*. 1989; 23:541–547. [PubMed: 2575444]
- Guillot CR, Fanning JR, Liang T, Leventhal AM, Berman ME. An α -synuclein gene (SNCA) polymorphism moderates the association of PTSD symptomatology with hazardous alcohol use, but not with aggression-related measures. *Journal of Anxiety Disorders*. 2015; 30:41–47. [PubMed: 25594371]
- Guillot CR, Pang RD, Leventhal AM, Liang T, Berman ME. The association of SNCA with hazardous alcohol use is mediated by impulsivity. *Psychiatry Research*. 2015; 30:226, 523–524.
- Gulley JM, Juraska JM. The effects of abused drugs on adolescent development of corticolimbic circuitry and behavior. *Neuroscience*. 2013; 249:3–20. [PubMed: 23711583]
- Gupta T, Syed YM, Revis AA, Miller SA, Martinez M, Cohn KA, et al. Acute effects of acamprosate and MPEP on ethanol drinking-in-the-dark in male C57BL/6J mice. *Alcoholism: Clinical and Experimental Research*. 2008; 32:1992–1998.
- Halliday GM, Törk I. Serotonin-like immunoreactive cells and fibres in the rat ventromedial mesencephalic tegmentum. *Brain Research Bulletin*. 1989; 22:725–735. [PubMed: 2736398]
- Hamilton KR, Felton JW, Risco CM, Lejuez CW, MacPherson L. Brief report: the interaction of impulsivity with risk-taking is associated with early alcohol use initiation. *Journal of Adolescence*. 2014; 37:1253–1256. <http://dx.doi.org/10.1016/j.adolescence.2014.08.013>. [PubMed: 25278465]
- Hanchar HJ, Dodson PD, Olsen RW, Otis TS, Wallner M. Alcohol-induced motor impairment caused by extrasynaptic GABA(A) receptor activity. *Nature Neuroscience*. 2005; 8:339–345. [PubMed: 15696164]
- Haughey HM, Ray LA, Finan P, Villanueva R, Niculescu M, Hutchison KE. Human gamma-aminobutyric acid A receptor alpha2 gene moderates the acute effects of alcohol and brain mRNA expression. *Genes Brain and Behavior*. 2008; 7:447–454.
- Hauser SR, Ding ZM, Getachew B, Toalston JE, Oster SM, McBride WJ, Rodd ZA. The posterior ventral tegmental area mediates alcohol-seeking behavior in alcohol-preferring rats. *Journal of Pharmacology and Experimental Therapeutics*. 2011; 336:857–865. [PubMed: 21148248]
- Hauser SR, Getachew B, Oster SM, Dhaher R, Ding Z-M, Bell RL, et al. Nicotine modulates alcohol-seeking and relapse by alcohol-preferring (P) rats in a time dependent manner. *Alcoholism: Clinical and Experimental Research*. 2012; 36:43–54.
- Hauser SR, Hedlund PB, Roberts AJ, Sari Y, Bell RL, Engleman EA. The 5-HT-7 receptor as a potential target for treating drug and alcohol abuse. *Frontiers in Neuroscience/Neuropharmacology*. 2014; 8:448. <http://dx.doi.org/10.3389/fnins.2014.00448>.
- Heath AC. Genetic influences on drinking behavior in humans. In: Begleiter, H.; Kissin, B., editors. *The Genetics of Alcoholism*. Oxford University Press; New York, NY: 1995. p. 82-121.

- Heather N, Tebbutt JS, Mattick RP, Zamir R. Development of a scale for measuring impaired control over alcoholism: a preliminary report. *Journal of Studies on Alcohol*. 1993; 54:700–709. [PubMed: 8271806]
- Hegerl U, Hensch T. The vigilance regulation model of affective disorders and ADHD. *Neuroscience and Biobehavioral Reviews*. 2014; 44:45–57. [PubMed: 23092655]
- Heilig M, Egli M. Pharmacological treatment of alcohol dependence: Target symptoms and target mechanisms. *Pharmacology and Therapeutics*. 2006; 111:855–876. [PubMed: 16545872]
- Heilig M, Thorsell A. Brain neuropeptide Y (NPY) in stress and alcohol dependence. *Reviews in Neuroscience*. 2002; 13:85–94.
- Heilig M, Egli M. Pharmacological treatment of alcohol dependence: Target symptoms and target mechanisms. *Pharmacology and Therapeutics*. 2006; 111:855–876. [PubMed: 16545872]
- Heilig M, Egli M, Crabbe JC, Becker HC. Acute withdrawal, protracted abstinence and negative affect in alcoholism: are they linked? *Addiction Biology*. 2010; 15:169–184. [PubMed: 20148778]
- Heilig M, Thorsell A, Sommer WH, Hansson AC, Ramchandani VA, George DT, et al. Translating the neuroscience of alcoholism into clinical treatments: from blocking the buzz to curing the blues. *Neuroscience and Biobehavioral Reviews*. 2010; 35:334–344. [PubMed: 19941895]
- Heinz A. Dopaminergic dysfunction in alcoholism and schizophrenia—psychopathological and behavioral correlates. *European Psychology Journal of the Association of European Psychiatry*. 2002; 17:9–16.
- Heinz A, Goldman D, Gallinat J, Schumann G, Puls I. Pharmacogenetic insights to monoaminergic dysfunction in alcohol dependence. *Psychopharmacology*. 2004; 174:561–570. [PubMed: 15148564]
- Heit C, Dong H, Chen Y, Thompson DC, Deitrich RA, Vasiliou VK. The role of CYP2E1 in alcohol metabolism and sensitivity in the central nervous system. *Sub-Cellular Biochemistry*. 2013; 67:235–247. [PubMed: 23400924]
- Hendrickson LM, Guildford MJ, Tapper AR. Neuronal nicotinic acetylcholine receptors: common molecular substrates of nicotine and alcohol dependence. *Frontiers in Psychiatry*. 2013; 4:29. [PubMed: 23641218]
- Hendrickson LM, Zhao-Shea R, Tapper AR. Modulation of ethanol drinking-in-the-dark by mecamylamine and nicotinic acetylcholine receptor agonists in C57BL/6J mice. *Psychopharmacology*. 2009; 204:563–572. [PubMed: 19247637]
- Henley JM, Wilkinson KA. AMPA receptor trafficking and the mechanisms underlying synaptic plasticity and cognitive aging. *Dialogs in Clinical Neuroscience*. 2013; 15:11–27.
- Herman MA, Kallupi M, Luu G, Oleata CS, Heilig M, Koob GF, et al. Enhanced GABAergic transmission in the central nucleus of the amygdala of genetically selected Marchigian Sardinian rats: Alcohol and CRF effects. *Neuropharmacology*. 2012; 67C:337–348. [PubMed: 23220399]
- Herman MA, Varodayan FP, Oleata CS, Luu G, Kirson D, Heilig M, et al. Glutamatergic transmission in the central nucleus of the amygdala is selectively altered in Marchigian Sardinian alcohol-preferring rats: alcohol and CRF effects. *Neuropharmacology*. 2016; 102:21–31. [PubMed: 26519902]
- Hersch SM, Gutekunst CA, Rees HD, Heilman CJ, Levey AI. Distribution of m1-m4 muscarinic receptor proteins in the rat striatum: Light and electron microscopic immunocytochemistry using subtype-specific antibodies. *Journal of Neuroscience*. 1994; 14:3351–3363. [PubMed: 8182478]
- Herve D, Pickel VM, Joh TH, Beaudet A. Serotonin axon terminals in the ventral tegmental area of the rat: fine structure and synaptic input to dopaminergic neurons. *Brain Research*. 1987; 435:71–83. [PubMed: 2892580]
- Hesselbrock, MN. Genetic determinants of alcoholic subtypes. In: Begleiter, H.; Kissin, B., editors. *The Genetics of Alcoholism*. Oxford University Press; New York, NY: 1995. p. 40–69.
- Hesselbrock, VM. The genetic epidemiology of alcoholism. In: Begleiter, H.; Kissin, B., editors. *The Genetics of Alcoholism*. Oxford University Press; New York, NY: 1995. p. 17–39.
- Hinckes AS, Laucht M, Schmidt MH, Mann KF, Schumann G, Schuckit MA, et al. Low level of response to alcohol as associated with serotonin transporter genotype and high alcohol intake in adolescents. *Biological Psychiatry*. 2006; 60:282–287. [PubMed: 16497275]

- Hrubec Z, Neel JV. The National Academy of Sciences--National Research Council Twin Registry: ten years of operation. *Progress in Clinical Biological Research*. 1978; 24(Pt B):153–172. [PubMed: 569304]
- Hulse GK. Improving clinical outcomes for naltrexone as a management of problem alcohol use. *British Journal of Clinical Pharmacology*. 2012; 76(5):632–641. <http://dx.doi.org/10.1111/j.1365-2125.2012.04452.x>. [PubMed: 22946873]
- Hunt WA. Neuroscience research: how has it contributed to our understanding of alcohol abuse and alcoholism? A review. *Alcoholism: Clinical and Experimental Research*. 1993; 17:1055–1065.
- Hwang BH, Lumeng L, Wu J-Y, Li T-K. Increased number of GABAergic terminals in the nucleus accumbens is associated with alcohol preference in rats. *Alcoholism: Clinical and Experimental Research*. 1990; 14:503–507.
- Hwang BH, Stewart R, Zhang J-K, Lumeng L, Li T-K. Corticotropin-releasing factor gene expression is down-regulated in the central nucleus of the amygdala of alcohol-preferring rats which exhibit high anxiety: a comparison between rat lines selectively bred for high and low alcohol preference. *Brain Research*. 2004; 1026:143–150. [PubMed: 15476706]
- Hwang BH, Zhang JK, Ehlers CL, Lumeng L, Li TK. Innate differences of neuropeptide Y (NPY) in hypothalamic nuclei and central nucleus of the amygdala between selectively bred rats with high and low alcohol preference. *Alcoholism: Clinical and Experimental Research*. 1999; 23:1023–1030.
- Imperato A, Di Chiara G. Preferential stimulation of dopamine release in the nucleus accumbens of freely moving rats by ethanol. *Journal of Pharmacology and Experimental Therapeutics*. 1986; 239:219–228. [PubMed: 3761194]
- Irimia C, Wiskerke J, Natividad LA, Polis IY, de Vries TJ, Pattij T, Parsons LH. Increased impulsivity in rats as a result of repeated cycles of alcohol intoxication and abstinence. *Addiction Biology*. 2013; 20:263–274. <http://dx.doi.org/10.1111/adb.12119>. [PubMed: 24341858]
- Ishii T, Mann GE. Redox status in mammalian cells and stem cells during culture in vitro: critical roles of Nrf2 and cystine transporter activity in the maintenance of redox balance. *Redox Biology*. 2014; 2:786–794. [PubMed: 25009780]
- Iyer RN, Bradberry CW. Serotonin-mediated increase in prefrontal cortex dopamine release: pharmacological characterization. *Journal of Pharmacology and Experimental Therapeutics*. 1996; 277:40–47. [PubMed: 8613947]
- Jablon S, Neel JV, Gershowitz H, Atkinson GF. The NAS-NRC twin panel: methods of construction of the panel, zygosity diagnosis, and proposed use. *American Journal of Human Genetics*. 1967; 19:133–161. [PubMed: 6067093]
- Jerlhag E, Egicioglu E, Dickson S, Engel JA. Glutamatergic regulation of ghrelin-induced activation of the mesolimbic dopamine system. *Addiction Biology*. 2011; 16:82–91. [PubMed: 20579004]
- Jerlhag E, Egicioglu E, Landgren S, Salome N, Heilig M, Moechars D, et al. Requirement of central ghrelin signaling for alcohol reward. *Proceedings of the National Academy of Sciences, USA*. 2009; 106:11318–11323.
- Jerlhag E, Landgren S, Egicioglu E, Dickson SL, Engel JA. The alcohol-induced locomotor stimulation and accumbal dopamine release is suppressed in ghrelin knockout mice. *Alcohol*. 2011; 45:341–347. [PubMed: 21145690]
- Johnson BA. Recent advances in the development of treatments for alcohol and cocaine dependence. *CNS Drugs*. 2005; 19:873–896. [PubMed: 16185095]
- Johnson BA. Medication treatment of different types of alcoholism. *American Journal of Psychiatry*. 2010; 167:630–639. [PubMed: 20516163]
- Johnson BA, Ait-Daoud N, Ma JZ, Wang Y. Ondansetron reduces mood disturbance among biologically predisposed, alcohol-dependent individuals. *Alcoholism: Clinical and Experimental Research*. 2003; 27:1773–1779.
- Johnston, LD.; O'Malley, PM.; Bachman, JG. National Survey Results on Drug Use from the Monitoring the Future Study, 1975–1997. National Institute on Drug Abuse; Rockville, MD: 1999.

- Johnston, LD.; O'Malley, PM.; Bachman, JG.; Schulenberg, JE. National Survey Results on Drug Use from the Monitoring the Future Study, 1975–2007. Volume I: Secondary school students. National Institute on Drug Abuse; Bethesda, MD: 2008. NIH Publication No. 08-6418A
- Jonas DE, Amick HR, Feltner C, Wines R, Shanahan E, Rowe CJ, Garbutt JC. Genetic polymorphisms and response to medications for alcohol use disorders: a systematic review and meta-analysis. *Pharmacogenomics*. 2014; 15:1687–1700. [PubMed: 25410894]
- June HL, Cummings R, Eiler WJA II, Foster KL, McKay PF, Seyoum R, et al. Central opioid receptors differentially regulate the nalmefene-induced suppression of ethanol- and saccharin-reinforced behaviors in alcohol-preferring (P) rats. *Neuropsychopharmacology*. 2004; 29:285–299. [PubMed: 14603266]
- June HL, Devaraju SL, Eggers MW, Williams JA, Cason CR, Greene TL, et al. Benzodiazepine receptor antagonists modulate the actions of ethanol in alcohol-preferring and –nonpreferring rats. *European Journal of Pharmacology*. 1998; 342:139–151. [PubMed: 9548379]
- June HL, Eggers MW, Warren-Reese C, DeLong J, Ricks-Cord A, Durr LF, et al. The effects of the novel benzodiazepine inverse agonist RU 34000 on ethanol-maintained behaviors. *European Journal of Pharmacology*. 1998; 350:151–158. [PubMed: 9696402]
- June HL, Foster KL, McKay PF, Seyoum R, Woods JE II, Harvey SC, et al. The reinforcing properties of alcohol are mediated by GABA_{A1} receptors in the ventral pallidum. *Neuropsychopharmacology*. 2003; 28:2124–2137. [PubMed: 12968126]
- June HL, Greene TL, Murphy JM, Hite ML, Williams JA, Cason CR, et al. Effects of the benzodiazepine inverse agonist RO19-4603 alone and in combination with the benzodiazepine receptor antagonists flumazenil, ZK 93426 and CGS 8216, on ethanol in take in alcohol-preferring (P) rats. *Brain Research*. 1996; 734:19–34. [PubMed: 8896804]
- June HL, Grey C, Warren-Reese C, Durr LF, Ricks-Cord A, Johnson A, et al. The opioid receptor antagonist nalmefene reduces responding maintained by ethanol presentation: Preclinical studies in ethanol-preferring and outbred Wistar rats. *Alcoholism: Clinical and Experimental Research*. 1998; 22:2174–2185.
- June HL, Lummis GH, Colker RE, Moore TO, Lewis MJ. Ro 15-4513 attenuates the consumption of ethanol in deprived rats. *Alcoholism: Clinical and Experimental Research*. 1991; 15:406–411.
- June HL, McCane SR, Zink RW, Portoghese PS, Li T-K, Froehlich JC. The delta-2 opioid receptor antagonist naltriben reduces motivated responding for ethanol. *Psychopharmacology*. 1999; 147:81–89. [PubMed: 10591872]
- June HL, Murphy JM, Mellor-Burke JJ, Lumeng L, Li T-K. The benzodiazepine inverse agonist RO19-4603 exerts prolonged and selective suppression of ethanol intake in alcohol-preferring (P) rats. *Psychopharmacology*. 1994; 115:325–331. [PubMed: 7871072]
- June HL, Torres L, Cason CR, Hwang BH, Braun MR, Murphy JM. The novel benzodiazepine inverse agonist RO19-4603 antagonizes ethanol motivated behaviors: Neuropharmacological studies. *Brain Research*. 1998; 784:256–275. [PubMed: 9518641]
- June HL, Williams JA, Cason CR, Devaraju S, Lin M, Murphy JM, et al. Low doses of gamma-hydroxybutyric acid (GHB) attenuate ethanol intake in alcohol-preferring (P) rats. *Alcoholism: Clinical and Experimental Research*. 1995; 19:14A.
- June HL, Zuccarelli D, Torres L, Craig KS, DeLong J, Allen A, et al. High-affinity benzodiazepine antagonists reduce responding maintained by ethanol presentation in ethanol-preferring rats. *Journal of Pharmacology and Experimental Therapeutics*. 1998; 284:1006–1014. [PubMed: 9495861]
- Jupp B, Lawrence AJ. New horizons for therapeutics in drug and alcohol abuse. *Pharmacology and Therapeutics*. 2010; 125:138–168. [PubMed: 19917308]
- Kaji, H. Neuropeptide Y and its receptors: molecular structure and pathophysiological role in food intake and energy homeostasis. In: Parker, SL., editor. *Neuropeptide Y: Molecular Structure, Role in Food Intake and Direct/Indirect Effects*. Nova Biomedical; New York, NY: 2013. p. 39-82.
- Kampov-Polevoy AB, Matthews DB, Gause L, Morrow AL, Overstreet DH. P rats develop physical dependence on alcohol via voluntary drinking: Changes in seizure thresholds, anxiety, and patterns of alcohol drinking. *Alcoholism: Clinical and Experimental Research*. 2000; 24:278–284.

- Kane CJ, Phelan KD, Douglas JC, Wagoner G, Johnson JW, Xu J, et al. Effects of ethanol on immune response in the brain: region-specific changes in adolescent versus adult mice. *Alcoholism: Clinical and Experimental Research*. 2014; 38:384–391.
- Kapasova Z, Szumlinski KK. Strain differences in alcohol induced neurochemical plasticity: a role for accumbens glutamate in alcohol intake. *Alcoholism: Clinical and Experimental Research*. 2008; 32:617–631.
- Kaplan RF, Cooney NL, Baker LH, Gillespie RA, Meyer RE, Pomerleau OF. Reactivity to alcohol-related cues: Physiological and subjective responses in alcoholics and nonproblem drinkers. *Journal of Studies on Alcohol*. 1985; 46:267–272. [PubMed: 4033125]
- Kaplan RF, Meyer RE, Stroebe CF. Alcohol dependence and responsivity to an ethanol stimulus as predictors of alcohol consumption. *British Journal of Addiction*. 1983; 78:259–267. [PubMed: 6578828]
- Kaprio J, Rose RJ, Romanov K, Koskenvuo M. Genetic and environmental determinants of use and abuse of alcohol: the Finnish twin cohort studies. *Alcohol and Alcoholism*. 1991; 1(Suppl 1): 131–136. [PubMed: 1845528]
- Katner SN, Oster SM, Ding ZM, Deehan GA Jr, Toalston JE, Hauser SR, et al. Alcohol-preferring (P) rats are more sensitive than Wistar rats to the reinforcing effects of cocaine self-administered directly into the nucleus accumbens shell. *Pharmacology, Biochemistry and Behavior*. 2011; 99:688–695.
- Keating GM. Nalmefene: a review of its use in the treatment of alcohol dependence. *CNS Drugs*. 2013; 27:761–772. [PubMed: 23949908]
- Keith, LD.; Roberts, A.; Wisen, KM.; Crabbe, JC. Stress, Gender, and Alcohol-Seeking Behavior. Hunt, WA.; Zakhari, S., editors. National Institute on Alcohol Abuse and Alcoholism; Bethesda, MD: 1995. p. 181-196. Research Monograph No. 29
- Kelley SP, Nannini MA, Bratt AM, Hodge CW. Neuropeptide-Y in the paraventricular nucleus increases ethanol self-administration. *Peptides*. 2001; 22:515–522. [PubMed: 11287109]
- Kimpel MW, Strother WN, McClintick JN, Carr LG, Liang T, Edenberg HJ, McBride WJ. Functional gene expression differences between inbred alcohol-preferring and -non-preferring rats in five brain regions. *Alcohol*. 2007; 41:95–132. [PubMed: 17517326]
- Kiefer F, Jahn H, Otte C, Nakovics H, Wiedemann K. Effects of treatment with acamprosate on β -endorphin plasma concentration in humans with high alcohol preference. *Neuroscience Letters*. 2006; 404:103–106. [PubMed: 16782270]
- Kim SA, Kim JW, Song JY, Park S, Lee HJ, Chung JH. Association of polymorphisms in nicotinic acetylcholine receptor alpha 4 subunit gene (CHRNA4), mu-opioid receptor gene (OPRM1), and ethanol-metabolizing enzyme genes with alcoholism in Korean patients. *Alcohol*. 2004; 34:115–120. [PubMed: 15902904]
- Kirby LG, Zeeb FD, Winstanley CA. Contributions of serotonin in addiction vulnerability. *Neuropharmacology*. 2011; 61:421–432. [PubMed: 21466815]
- Klimek V, Rajkowska G, Luker SN, Dilley G, Ordway GA, et al. Brain Noradrenergic Receptors in Major Depression. *Neuropsychopharmacology*. 1999; 21:69–81. [PubMed: 10379521]
- Koob GF. A role for GABA mechanisms in the motivational effects of alcohol. *Biochemistry and Pharmacology*. 2004; 68:1515–1525.
- Koob GF. Dynamics of neuronal circuits in addiction: Reward, antireward, and emotional memory. *Pharmacopsychiatry*. 2009; 42(Suppl 1):S32–S41. [PubMed: 19434554]
- Koob GF. The role of CRF and CRF-related peptides in the dark side of addiction. *Brain Research*. 2010; 1314:3–14. [PubMed: 19912996]
- Koob GF. Theoretical frameworks and mechanistic aspects of alcohol addiction: alcohol addiction as a reward deficit disorder. *Current Topics in Behavioral Neuroscience*. 2013; 13:3–30.
- Koob GF.; Arends, MA.; Le Moal, M. *Drugs, Addiction, and the Brain*. Elsevier Academic Press; New York, NY: 2014.
- Koob GF, Buck CL, Cohen A, Edwards S, Park PE, Schlosburg JE, et al. Addiction as a stress surfeit disorder. *Neuropharmacology*. 2014; 76:370–382. [PubMed: 23747571]
- Koob, GF.; Le Moal, M. *Neurobiology of Addiction*. Elsevier Academic Press; New York: 2006.

- Koob GF, LeMoal M. Neurobiological mechanisms for opponent motivational processes in addiction. *Philosophical Transactions of the Royal Society of London Part B Biological Sciences*. 2008; 363:3113–3123.
- Koob GF, Volkow ND. Neurocircuitry of addiction. *Neuropsychopharmacology*. 2010; 35:217–238. [PubMed: 19710631]
- Korpi ER, Sinkkonen ST. GABA(A) receptor subtypes as targets for neuropsychiatric drug development. *Pharmacology and Therapeutics*. 2006; 109:12–32. [PubMed: 15996746]
- Krishnan V, Nestler EJ. The molecular neurobiology of depression. *Nature*. 2008; 455:894–902. [PubMed: 18923511]
- Krishnan-Sarin S, Jing S-L, Kurtz DL, Zweifel M, Portoghesi PS, Li T-K, et al. The delta opioid receptor antagonist naltrindole attenuates both alcohol and saccharin intake in rats selectively bred for alcohol preference. *Psychopharmacology*. 1995; 120:177–185. [PubMed: 7480550]
- Krishnan-Sarin S, Portoghesi PS, Li T-K, Froehlich JC. The delta2-opioid receptor antagonist naltriben selectively attenuates alcohol intake in rats bred for alcohol preference. *Pharmacology, Biochemistry and Behavior*. 1995; 52:153–159.
- Kumar S, Porcu P, Werner DF, Matthews DB, Diaz-Granados JL, Helfand RS, et al. The role of GABA-A receptors in the acute and chronic effects of ethanol: A decade of progress. *Psychopharmacology*. 2009; 205:529–564. [PubMed: 19455309]
- Kuntsche E, Rehm J, Gmel G. Characteristics of binge drinkers in Europe. *Social Science of Medicine*. 2004; 59:113–127.
- Kupila J, Karkkainen O, Laukkanen V, Tupala E, Tiihonen J, Storvik M. mGluR1/5 receptor densities in the brains of alcoholic subjects: a whole hemisphere autoradiography study. *Psychiatry Research*. 2013; 212:245–250. [PubMed: 23149043]
- Kurtz DL, Stewart RB, Zweifel M, Li T-K, Froehlich JC. Genetic differences in tolerance and sensitization to the sedative/hypnotic effects of alcohol. *Pharmacology, Biochemistry and Behavior*. 1996; 53:585–591.
- Kryger R, Wilce PA. The effects of alcoholism on the human basolateral amygdala. *Neuroscience*. 2010; 167:361–371. [PubMed: 20153402]
- Lancaster FE. Gender differences in the brain: implications for the study of human alcoholism. *Alcoholism: Clinical and Experimental Research*. 1994; 18:740–746.
- Lankford MF, Bjork AK, Myers RD. Differential efficacy of serotonergic drugs FG5974, FG5893, and amperozide in reducing alcohol drinking in P rats. *Alcohol*. 1996; 13:399–404. [PubMed: 8836330]
- Lê AD, Corrigall WA, Harding JW, Juzysch W, Li T-K. Involvement of nicotinic receptors in alcohol self-administration. *Alcoholism: Clinical and Experimental Research*. 2000; 24:155–63.
- Lê AD, Li Z, Funk D, Shram M, Li T-K, Shaham Y. Nicotine increases alcohol self-administration and reinstates alcohol seeking in rats. *Psychopharmacology*. 2003; 168:216–221. [PubMed: 12536264]
- Lê AD, Li Z, Funk D, Shram M, Li TK, Shaham Y. Increased vulnerability to nicotine self-administration and relapse in alcohol-naïve offspring of rats selectively bred for high alcohol intake. *Journal of Neuroscience*. 2006; 26:1872–1879. [PubMed: 16467536]
- Lê, AD.; Mayer, JM. Aspects of alcohol tolerance in humans and experimental animals. In: Deitrich, RA.; Erwin, VG., editors. *Pharmacological Effects of Ethanol on the Nervous System*. CRC Press; Boca Raton, FL: 1996. p. 251-268.
- Leggio L, Addolorato G. Serotonin transporter (SERT) brain density and neurobiological Cloninger subtypes model: a lesson by human autoradiography studies. *Alcohol and Alcoholism*. 2008; 43:148–150. [PubMed: 18211952]
- Lesch OM, Walter H. Subtypes of alcoholism and their role in therapy. *Alcohol and Alcoholism*. 1996; 31(Suppl 1):63–67.
- Lester D, Freed EX. Criteria for an animal model of alcoholism. *Pharmacology, Biochemistry and Behavior*. 1973; 1:103–107.
- Levy AD, Murphy JM, McBride WJ, Lumeng L, Li T-K. Microinjection of sulpiride into the nucleus accumbens increases ethanol intake of alcohol-preferring (P) rats. *Alcohol and Alcoholism*. 1991; 1(Suppl 1):417–420. [PubMed: 1845571]

- Lex BW, Lukas SE, Greenwald NE, Mendelson HH. Alcohol-induced changes in body sway in women at risk for alcoholism: A pilot study. *Journal of Studies on Alcohol*. 1988; 49:346–356. [PubMed: 3172783]
- Li, T-K.; Lumeng, L. Alcohol metabolism of inbred strains of rats with alcohol preference and non-preference. In: Thurman, RG.; Williamson, JR.; Drott, H.; Chance, B., editors. *Alcohol and Aldehyde Metabolizing Systems*. Vol. 3. Academic Press; New York, NY: 1977. p. 625-633.
- Li T-K, Lumeng L, McBride WJ, Murphy JM. Rodent lines selected for factors affecting alcohol consumption. *Alcohol and Alcoholism*. 1987; 1(Suppl):91–96. [PubMed: 3426760]
- Liang JH, Chen F, Krstew E, Cowen MS, Carroll FY, Crawford D, et al. The GABA_B receptor allosteric modulator CGP7930, like baclofen, reduces operant self-administration of ethanol in alcohol-preferring rats. *Neuropharmacology*. 2006; 50:632–9. [PubMed: 16406445]
- Liang J, Olsen RW. Alcohol use disorders and current pharmacological therapies: the role of GABA(A) receptors. *Acta Pharmacologica Sinica*. 2014; 35:981–993. [PubMed: 25066321]
- Liang T, Spence J, Liu L, Strother WN, Chang HW, Ellison JA, et al. α -synuclein maps to a quantitative trait locus for alcohol preference and is differentially expressed in alcohol-preferring and –nonpreferring rats. *Proceedings of the National Academy of Sciences, USA*. 2003; 100:4690–4695.
- Litten RZ, Egli M, Heilig M, Cui C, Fertig JB, Ryna ML, et al. Medications development to treat alcohol dependence: a vision for the next decade. *Addiction Biology*. 2012; 17:513–527. [PubMed: 22458728]
- Liu J, Yang AR, Kelly T, Puche A, Esoga C, June HL Jr, et al. Binge alcohol drinking is associated with GABAA α 2-regulated toll-like receptor 4 (TLR4) expression in the central amygdala. *Proceedings of the National Academy of Sciences, USA*. 2011; 108:4465–4470.
- Logrip ML, Janak PH, Ron D. Escalating ethanol intake is associated with altered corticostriatal BDNF expression. *Journal of Neurochemistry*. 2009; 109:1459–1468. [PubMed: 19453942]
- Lovinger DM. 5-HT₃ receptors and the neural actions of alcohols: an increasingly exciting topic. *Neurochemistry International*. 1999; 35:125–130. [PubMed: 10405996]
- Lovinger, DM.; Roberto, M. *Behavioral Neurobiology of Alcohol Addiction*. Springer; Berlin, Heidelberg: 2013. Synaptic effects induced by alcohol.
- Lukas SE, Mendelson JH, Benedikt RA, Jones B. EEG alpha activity increases during transient episodes of ethanol-induced euphoria. *Pharmacology, Biochemistry and Behavior*. 1986; 25:889–895.
- Lukkes JL, Staub DR, Dietrich A, Truitt W, Neufeld-Cohen A, Chen A, et al. Topographical distribution of corticotropin-releasing factor type 2 receptor-like immunoreactivity in the rat dorsal raphe nucleus: co-localization with tryptophan hydroxylase. *Neuroscience*. 2011; 183:47–63. [PubMed: 21453754]
- Lumeng, L.; Hawkins, TD.; Li, T-K. New strains of rats with alcohol preference and non-preference. In: Thurman, RG.; Williamson, JR.; Drott, H.; Chance, B., editors. *Alcohol and Aldehyde Metabolizing Systems*. Vol. 3. Academic Press; New York, NY: 1977. p. 537-544.
- Lumeng L, Li T-K. The development of metabolic tolerance in the alcohol-preferring P rats: Comparison of forced and free-choice drinking of ethanol. *Pharmacology, Biochemistry and Behavior*. 1986; 25:1013–1020.
- Lumeng L, Waller MB, McBride WJ, Li T-K. Different sensitivities to ethanol in alcohol-preferring and –nonpreferring rats. *Pharmacology, Biochemistry and Behavior*. 1982; 16:125–130.
- Luo X, Kranzler HR, Zuo L, Wang S, Blumberg HP, Gelernter J. CHRM2 gene predisposes to alcohol dependence, drug dependence and affective disorders: results from an extended case-control structured association study. *Human Molecular Genetics*. 2005; 14:2421–2434. [PubMed: 16000316]
- Lynch WJ, Bond C, Breslin FJ, Johnson BA. Severity of drinking as a predictor of efficacy of the combination of ondansetron and topiramate in rat models of ethanol consumption and relapse. *Psychopharmacology (Berl)*. 2011; 217:3–12. [PubMed: 21424693]
- Lyness WH, Smith F. Influence of dopaminergic and serotonergic neurons on intravenous ethanol self-administration in the rat. *Pharmacology, Biochemistry and Behavior*. 1992; 42:187–192.

- Maccioni P, Colombo G. Role of the GABA-B receptor in alcohol-seeking and drinking behavior. *Alcohol*. 2009; 43:555–558. [PubMed: 19913200]
- Maccioni P, Zaru A, Loi B, Lobina C, Carai MAM, Gessa GL, et al. Comparison of the effect of the GABA_B receptor agonist, baclofen, and the positive allosteric modulator of the GABA_B receptor, GS39783, on alcohol self-administration in three different lines of alcohol-preferring rats. *Alcoholism: Clinical and Experimental Research*. 2012; 36:1748–1766.
- Mackenzie CS, El-Gabalaway R, Chou K-L, Sareen J. Prevalence and predictors of persistent versus remitting mood, anxiety, and substance disorders in a national sample of older adults. *American Journal of Geriatric Psychiatry*. 2014; 22(9):854–865. <http://dx.doi.org/10.1016/j.jagp.2013.02.007>. [PubMed: 23800537]
- Madsen KK, White HS, Schousboe A. Neuronal and non-neuronal GABA transporters as targets for antiepileptic drugs. *Pharmacology and Therapeutics*. 2010; 125:394–401. [PubMed: 20026354]
- Manzardo AM, McGuire A, Butler MG. Clinically relevant genetic biomarkers from the brain in alcoholism with representation on high resolution chromosome ideograms. *Gene*. 2015; 560:184–194. [PubMed: 25655461]
- Mardones J, Segovia N, Hederra A. Heredity of experimental alcohol preference in rats. II. Coefficient of heredity. *Quarterly Journal of Studies on Alcohol*. 1953; 14:1–2. [PubMed: 13037991]
- Mardones J, Segovia-Riquelme N. Thirty-two years of selection of rats by ethanol preference: UChA and UChB strains. *Neurobehavioral Toxicology and Teratology*. 1983; 5:171–178. [PubMed: 6683362]
- Marini V, Fucile C, Zuccoli ML, Testino G, Sumberaz A, Robbiano L, et al. Involvement of the mu-opioid receptor gene polymorphism A118G in the efficacy of detoxification of alcohol dependent patients. *Addictive Behaviors*. 2013; 38:1669–1671. [PubMed: 23254216]
- Martin NG, Oakeshott JG, Gibson JB, Starmer GA, Perl J, Wilks AV. A twin study of psychomotor and physiological responses to an acute dose of alcohol. *Behavior Genetics*. 1985; 15:305–347. [PubMed: 4041178]
- Mason GA, Rezvani AH, Grady DR, Garbutt JC. The subchronic effects of the TRH analog TA-0910 and bromocriptine on alcohol preference in alcohol-preferring rats: development of tolerance and cross-tolerance. *Alcoholism: Clinical and Experimental Research*. 1994; 18:1196–1201.
- Mason GA, Rezvani AH, Overstreet DH, Hamed M, Walker CH, Yang Y, Garbutt JC. Involvement of dopamine D2 receptors in the suppressive effect of the thyrotropin-releasing hormone analog TA-0910 on alcohol intake in alcohol-preferring rats. *Alcoholism: Clinical and Experimental Research*. 1997; 21:1623–1629.
- McBride WJ, Bell RL, Rodd ZA, Strother, WN, Murphy, JM. Adolescent alcohol drinking and its long-range consequences: Studies with animal models. In: Galanter, M., editor. *Alcohol Problems in Adolescents and Young Adults: Epidemiology, Neurobiology, Prevention, and Treatment*. Springer; New York, NY: 2005. p. 123–142.
- McBride WJ, Chernet E, Dyr W, Lumeng L, Li T-K. Densities of dopamine D2 receptors are reduced in CNS regions of alcohol-preferring P rats. *Alcohol*. 1993; 10:387–390. [PubMed: 8216885]
- McBride WJ, Chernet E, Rabold JA, Lumeng L, Li T-K. Serotonin-2 receptors in the CNS of alcohol-preferring and -nonpreferring rats. *Pharmacology, Biochemistry and Behavior*. 1993; 46:631–636.
- McBride WJ, Chernet E, Russell RN, Wong DT, Guan X-M, Lumeng L, et al. Regional CNS densities of monoamine receptors in alcohol-naïve alcohol-preferring P and -nonpreferring NP rats. *Alcohol*. 1997; 14:141–148. [PubMed: 9085714]
- McBride WJ, Guan X-M, Chernet E, Lumeng L, Li T-K. Regional serotonin_{1A} receptors in the CNS of alcohol-preferring and -nonpreferring rats. *Pharmacology, Biochemistry and Behavior*. 1994; 49:7–12.
- McBride WJ, Kimpel MW, McClintick JN, Ding Z-M, Hauser SR, Edenberg HJ, et al. Changes in gene expression within the ventral tegmental area following repeated excessive binge-like alcohol drinking by alcohol-preferring (P) rats. *Alcohol*. 2013; 47:367–380. <http://dx.doi.org/10.1016/j.alcohol.2013.04.002>. [PubMed: 23714385]
- McBride WJ, Kimpel MW, McClintick JN, Ding Z-M, Hyytia P, Colombo G, et al. Gene expression in the ventral tegmental area of 5 pairs of rat lines selectively bred for high or low ethanol

consumption. *Pharmacology, Biochemistry and Behavior*. 2012; 102:275–285. <http://dx.doi.org/10.1016/j.pbb.2012.04.016>.

- McBride WJ, Kimpel MW, McClintick JN, Ding Z-M, Hyytia P, Colombo G, et al. Gene expression within the extended amygdala of 5 pairs of rat lines selectively bred for high or low ethanol consumption. *Alcohol*. 2013; 47:517–529. <http://dx.doi.org/10.1016/j.alcohol.2013.08.004>. [PubMed: 24157127]
- McBride WJ, Kimpel MW, McClintick JN, Ding Z-M, Edenberg HJ, Liang T, et al. Changes in gene expression within the extended amygdala following binge-like alcohol drinking by adolescent alcohol-preferring (P) rats. *Pharmacology, Biochemistry and Behavior*. 2014; 117:52–60.
- McBride WJ, Kimpel MW, Schultz JA, McClintick JN, Edenberg HJ, Bell RL. Changes in gene expression in regions of the extended amygdala of alcohol-preferring rats after binge-like alcohol drinking. *Alcohol*. 2010; 44:171–183. [PubMed: 20116196]
- McBride WJ, Li TK. Animal models of alcoholism: neurobiology of high alcohol-drinking behavior in rodents. *Critical Reviews in Neurobiology*. 1998; 12:339–369. [PubMed: 10348615]
- McBride WJ, Murphy JM, Lumeng L, Li T-K. Effects of Ro 15-4513, fluoxetine and desipramine on the intake of ethanol, water and food by the alcohol-preferring (P) and –nonpreferring (NP) lines of rats. *Pharmacology, Biochemistry and Behavior*. 1988; 30:1045–1050.
- McBride WJ, Murphy JM, Lumeng L, Li T-K. Serotonin, dopamine and GABA involvement in alcohol drinking of selectively bred rats. *Alcohol*. 1990; 7:199–205. [PubMed: 2184832]
- McBride WJ, Rodd ZA, Bell RL, Lumeng L, Li T-K. The alcohol-preferring (P) and high-alcohol-drinking (HAD) rats—Animal models of alcoholism. *Alcohol*. 2014; 48:209–215. <http://dx.doi.org/10.1016/j.alcohol.2013.09.044>. [PubMed: 24268381]
- McBride WJ, Schultz JA, Kimpel MW, McClintick JN, Wang M, You J, Rodd ZA. Differential effects of ethanol in the nucleus accumbens shell of alcohol-preferring (P), alcohol non-preferring (NP) and Wistar rats: a proteomic study. *Pharmacology, Biochemistry and Behavior*. 2009; 92:304–313.
- McCairn KW, Isoda M. Pharmacological animal models of tic disorders. *International Reviews in Neurobiology*. 2013; 112:179–203. <http://dx.doi.org/10.1016/B978-0-12-411546-0.00007-X>.
- McCarson KE, Enna SJ. GABA pharmacology: the search for analgesics. *Neurochemistry Research*. 2014; 39:1948–1963.
- McClintick JN, McBride WJ, Bell RL, Ding Z-M, Liu Y, Xuei X, Edenberg HJ. Gene expression changes in serotonin, GABA-A receptors, neuropeptides and ion channels in the dorsal raphe nucleus of adolescent alcohol-preferring (P) rats following binge-like alcohol drinking. *Pharmacology, Biochemistry and Behavior*. 2015; 129:87–96.
- McGehee DS, Role LW. Physiological diversity of nicotinic acetylcholine receptors expressed by vertebrate neurons. *Annual Reviews in Physiology*. 1995; 57:521–546.
- McGonigle P, Ruggeri B. Animal models of human disease: challenges in enabling translation. *Biochemistry and Pharmacology*. 2014; 87(1):162–171. <http://dx.doi.org/10.1016/j.bcp.2013.08.006>.
- McHugh RK, Hofmann SG, Asnaani A, Sawyer AT, Otto MW. The serotonin transporter gene and risk for alcohol dependence: a meta-analytic review. *Drug and Alcohol Dependence*. 2010; 108:1–6. [PubMed: 20060655]
- McKinney WT. Overview of the past contributions of animal models and their changing place in psychiatry. *Seminars in Clinical Psychiatry*. 2001; 6:68–78.
- McLarnon, JG. Correlated inflammatory responses and neurodegeneration in peptide-injected animal models of Alzheimer's disease; *BioMedical Research International*. 2014. p. 923670 <http://dx.doi.org/10.1155/2014/923670>
- Meinhardt MW, Hansson AC, Perreau-Lenz S, Bauder-Wenz C, Stahlin O, Heilig M, et al. Rescue of infralimbic mGluR2 deficit restores control over drug-seeking behavior in alcohol Dependence. *Journal of Neuroscience*. 2013; 33:2794–2806. [PubMed: 23407939]
- Melendez RI, Rodd ZA, McBride WJ, Murphy JM. Dopamine receptor regulation of ethanol intake and extracellular dopamine levels in the ventral pallidum of alcohol preferring (P) rats. *Drug and Alcohol Dependence*. 2005; 77:293–301. [PubMed: 15734229]

- Melendez RI, Rodd-Henricks ZA, Engleman EA, Li T-K, McBride WJ, Murphy JM. Microdialysis of dopamine in the nucleus accumbens of alcohol-preferring (P) rats during anticipation and operant self-administration of ethanol. *Alcoholism: Clinical and Experimental Research*. 2002; 26:318–325.
- Melis M, Camarini R, Ungless MA, Bonci A. Long-lasting potentiation of GABAergic synapses in dopamine neurons after a single in vivo ethanol exposure. *Journal of Neuroscience*. 2002; 22:2074–2082. [PubMed: 11896147]
- Menkes DB, Baraban JM, Aghajanian GK. Prazosin selectively antagonizes neuronal responses mediated by alpha1-adrenoceptors in brain. *Naunyn-Schmiedeberg's Archives in Pharmacology*. 1981; 317:273–275.
- Milner LC, Buck KJ. Identifying quantitative trait loci (QTL) and genes (QTGs) for alcohol-related phenotypes in mice. *International Reviews in Neurobiology*. 2010; 91:173–204.
- Mitani A, Tanaka K. Functional changes of glial glutamate transporter GLT-1 during ischemia: an in vivo study in the hippocampal CA1 of normal mice and mutant mice lacking GLT-1. *Journal of Neuroscience*. 2003; 23:7176–7182. [PubMed: 12904478]
- Miyata I, Shiota C, Chaki S, Okuyama S, Inagami T. Localization and characterization of a short isoform of the corticotropin-releasing factor receptor type 2 alpha (CRF(2)alpha-tr) in the rat brain. *Biochemistry and Biophysics Research Communications*. 2001; 280:553–557.
- Mokdad A, Marks J, Stroup D, Gerberding J. Actual causes of death in the United States, 2000. *JAMA*. 2004; 291:1238–1245. [PubMed: 15010446]
- Moonat S, Starkman BG, Sakharkar A, Pandey SC. Neuroscience of alcoholism: molecular and cellular mechanisms. *Cellular and Molecular Life Sciences*. 2010; 67:73–88. [PubMed: 19756388]
- Morean ME, Corbin WR. Subjective response to alcohol: a critical review of the literature. *Alcoholism: Clinical and Experimental Research*. 2010; 34:385–395. <http://dx.doi.org/10.1111/j.1530-0277.2009.01103.x>.
- Moreira FA, Jupp B, Belin D, Dalley JW. Endocannabinoids and striatal function: implications for addiction-related behaviors. *Behavioral Pharmacology*. 2015; 26:59–72.
- Moret C, Briley M. The importance of norepinephrine in depression. *Neuropsychiatry Disorders and Treatment*. 2011; 7(Suppl 1):9–13.
- Morganstern I, Liang S, Ye Z, Karatayev O, Leibowitz SF. Disturbances in behavior and cortical enkephalin gene expression during the anticipation of ethanol in rats characterized as high drinkers. *Alcohol*. 2012; 46:559–568. [PubMed: 22703995]
- Morozova TV, Mackay TFC, Anholt RRH. Genetics and genomics of alcohol sensitivity. *Molecular Genetics and Genomics*. 2014; 289:253–269. <http://dx.doi.org/10.1007/s00438-013-0808-y>. [PubMed: 24395673]
- Morris RGM. NMDA receptors and memory encoding. *Neuropharmacology*. 2013; 74:32–40. [PubMed: 23628345]
- Morzorati SL. VTA dopamine neuron activity distinguishes alcohol-preferring (P) rats from Wistar rats. *Alcoholism: Clinical and Experimental Research*. 1998; 22:854–857.
- Morzorati SL, Marunde RL. Comparison of VTA dopamine neuron activity in lines of rats selectively bred to prefer or avoid alcohol. *Alcoholism: Clinical and Experimental Research*. 2006; 30:991–997.
- Morzorati SL, Marunde RL, Downey D. Limited access to ethanol increases the number of spontaneously active dopamine neurons in the posterior ventral tegmental area of nondependent P rats. *Alcohol*. 2010; 44:257–264. [PubMed: 20682193]
- Morzorati SL, Ramchandani VA, Flury L, Li T-K, O'Connor S. Self-reported subjective perception of intoxication reflects family history of alcoholism when breath alcohol levels are constant. *Alcoholism: Clinical and Experimental Research*. 2002; 26:1299–1306.
- Moss HB, Chen CM, Yi H-Y. Subtypes of alcohol dependence in a nationally representative sample. *Drug and Alcohol Dependence*. 2007; 91:149–158. [PubMed: 17597309]
- Murphy JM, Gatto GJ, McBride WJ, Lumeng L, Li T-K. Operant responding for oral ethanol in the alcohol-preferring P and alcohol-nonpreferring NP lines of rats. *Alcohol*. 1989; 6:127–131. [PubMed: 2713085]

- Murphy JM, Gatto GJ, Waller MB, McBride WJ, Lumeng L, Li T-K. Effects of scheduled access on ethanol intake by the alcohol-preferring (P) line of rats. *Alcohol*. 1986; 3:331–336. [PubMed: 3778650]
- Murphy JM, McBride WJ, Lumeng L, Li T-K. Contents of monoamines in forebrain regions of alcohol-preferring (P) and –nonpreferring (NP) lines of rats. *Pharmacology, Biochemistry and Behavior*. 1987; 26:389–392.
- Murphy JM, McBride WJ, Lumeng L, Li T-K. Regional brain levels of monoamines in alcohol-preferring and –nonpreferring rats. *Pharmacology, Biochemistry and Behavior*. 1982; 16:145–149.
- Murphy JM, Stewart RB, Bell RL, Badia-Elder NE, Carr LG, McBride WJ, et al. Phenotypic and genotypic characterization of the Indiana University rat lines selectively bred for high and low alcohol preference. *Behavior Genetics*. 2002; 32:363–388. [PubMed: 12405517]
- Murphy JM, Waller MB, Gatto GJ, McBride WJ, Lumeng L, Li T-K. Monoamine uptake inhibitors attenuate ethanol intake in alcohol-preferring (P) rats. *Alcohol*. 1985; 2:349–352. [PubMed: 3925968]
- Murphy JM, Waller MB, Gatto GJ, McBride WJ, Lumeng L, Li T-K. Effects of fluoxetine on the intragastric self-administration of ethanol in the alcohol preferring P line of rats. *Alcohol*. 1988; 5:283–286. [PubMed: 3265874]
- Nam HW, Bruner RC, Choi DS. Adenosine signaling in striatal circuits and alcohol use disorders. *Molecules and Cells*. 2013; 36:195–202. [PubMed: 23912595]
- Nestler EJ, Hyman SE. Animal models of neuropsychiatric disorders. *Nature Neuroscience*. 2010; 13:1161–1169.
- Nevo I, Hamon M. Neurotransmitter and neuromodulatory mechanisms involved in alcohol abuse and alcoholism. *Neurochemistry International*. 1995; 26:305–342. [PubMed: 763325]
- Newman EL, Gupta K, Climer JR, Monaghan CK, Hasselmo ME. Cholinergic modulation of cognitive processing: insights drawn from computational models. *Frontiers in Behavioral Neuroscience*. 2012; 6:24. [PubMed: 22707936]
- NIAAA. Underage drinking. NIAAA Fact Sheet. Available at http://pubs.niaaa.nih.gov/publications/UnderageDrinking/Underage_Fact.pdf. Published March 2012
- NIAAA National Advisory Council. NIAAA Newsletter. 2004. NIAAA Council approves definition of binge drinking. *Author*, 3:5
- Niciu MJ, Kelmendi B, Sanacora G. Overview of glutamatergic neurotransmission in the nervous system. *Pharmacology, Biochemistry and Behavior*. 2012; 100:656–664.
- Nie Z, Madamba SG, Siggins GR. Ethanol inhibits glutamatergic neurotransmission in nucleus accumbens neurons by multiple mechanisms. *Journal of Pharmacology and Experimental Therapeutics*. 1994; 271:1566–1573. [PubMed: 7527857]
- Nie Z, Yuan X, Madamba SG, Siggins GR. Ethanol decreases glutamatergic synaptic transmission in rat nucleus accumbens in vitro: naloxone reversal. *Journal of Pharmacology and Experimental Therapeutics*. 1993; 266:1705–1712. [PubMed: 8396641]
- Nisell M, Nomikos GG, Svensson TH. Systemic nicotine-induced dopamine release in the rat nucleus accumbens is regulated by nicotinic receptors in the ventral tegmental area. *Synapse*. 1994; 16:36–44. [PubMed: 8134899]
- Noble EP, Zhnag X, Ritchie T, Lawford BR, Grosser SC, Young RM, Sparkes RS. D2 dopamine receptor and GABA(A) receptor beta 3 subunit genes and alcoholism. *Psychiatry Research*. 1998; 81:133–147. [PubMed: 9858031]
- Nogueira L, Kalivas PW, Lavin A. Long-term neuroadaptations produced by withdrawal from repeated cocaine treatment: role of dopaminergic receptors in modulating cortical excitability. *Journal of Neuroscience*. 2006; 26:12308–12313. [PubMed: 17122056]
- Nordberg A, Larsson C, Perdahl E, Winblad B. Changes in cholinergic activity in human hippocampus following chronic alcohol abuse. *Pharmacology, Biochemistry and Behavior*. 1983; 18:379–400.
- Noronha, ABC.; Cui, C.; Harris, RA.; Crabbe, JC. *Neurobiology of Alcohol Dependence*. Elsevier Academic Press; New York, NY: 2014.

- Nowak KL, McBride WJ, Lumeng L, Li TK, Murphy JM. Involvement of dopamine D2 autoreceptors in the ventral tegmental area on alcohol and saccharin intake of the alcohol preferring P rat. *Alcoholism: Clinical and Experimental Research*. 2000; 24:476–483.
- Nowak KL, McBride WJ, Lumeng L, Li TK, Murphy JM. Blocking GABA(A) receptors in the anterior ventral tegmental area attenuates ethanol intake of the alcohol-preferring P rat. *Psychopharmacology*. 1998; 139:108–116. [PubMed: 9768548]
- Nunez YO, Truitt JM, Gorini G, Ponomareva ON, Bldenov YA, Harris RA, Mayfield RD. Positively correlated miRNA-mRNA regulatory networks in mouse frontal cortex during early stages of alcohol dependence. *BMC Genomics*. 2013; 14:725. [PubMed: 24148570]
- Nutt DJ, Lingfor-Hughes A, Erritzoe D, Stokes PRA. The dopamine theory of addiction: 40 years of highs and lows. *Nature Reviews in Neuroscience*. 2015; 16:305–312. [PubMed: 25873042]
- Obara I, Bell RL, Goulding SP, Reyes CM, Larson LA, Ary AW, Truitt WA, Szumlinski KK. Differential effects of chronic ethanol consumption and withdrawal on Homer/glutamate receptor expression in subregions of the accumbens and amygdala of P rats. *Alcoholism: Clinical and Experimental Research*. 2009; 33:1924–1934. <http://dx.doi.org/10.1111/j.1530-0277.2009.01030.x>.
- O'Brien, CP.; Childress, AR.; McLellan, AT.; Ehrman, R. A learning model of addiction. In: O'Brien, CP.; Jaffe, JH., editors. *Addictive States*. Raven Press; New York, NY: 1992. p. 157-177.
- Okahisa Y, Ujike H, Kotaka T, Morita Y, Kodama M, Iwata N, Iyo M, et al. Association between neuropeptide Y gene and its receptor Y1 gene and methamphetamine dependence. *Psychiatry and Clinical Neuroscience*. 2009; 63:417–422.
- Olney JJ, Navarro M, Thiele TE. Targeting central melanocortin receptors: A promising novel approach for treating alcohol abuse disorders. *Frontiers in Neuroscience/Neuropharmacology*. 2014; 8:128. <http://dx.doi.org/10.3389/fnins.2014.00128>.
- Oswald LM, Wand GS. Opioids and alcoholism. *Physiology and Behavior*. 2004; 81:339–358. [PubMed: 15159175]
- Othman T, Sinclair CJ, Haughey N, Geiger JD, Parkinson FE. Ethanol alters glutamate but not adenosine uptake in rat astrocytes: evidence for protein kinase C involvement. *Neurochemistry Research*. 2002; 27:289–296.
- Overstreet DH, McArthur RA, Rezvani AH, Post C. Selective inhibition of alcohol intake in diverse alcohol-preferring rats strains by 5-HT2A antagonists amperozide and FG 5974. *Alcoholism: Clinical and Experimental Research*. 1997; 21:1448–1454.
- Overstreet DH, Rezvani AH, Djouma E, Parsian A, Lawrence AJ. Depressive-like behavior and high alcohol drinking co-occur in the FH/WJD rat but appear to be under independent genetic control. *Neuroscience and Biobehavioral Reviews*. 2007; 31:103–114. [PubMed: 16982094]
- Palmer AA, Low MJ, Grandy DK, Phillips TJ. Effects of a Drd2 deletion mutation on ethanol-induced locomotor stimulation and sensitization suggest a role for epistasis. *Behavior Genetics*. 2003; 33:311–324. [PubMed: 12837020]
- Pandey SC. Anxiety and alcohol abuse disorders: a common role for CREB and its target, the neuropeptide Y gene. *Trends in Pharmacological Science*. 2003; 24:455–460.
- Pandey SC, Carr LG, Heilig M, Ilveskoski E, Thiele E. Neuropeptide Y and alcoholism: genetic, molecular, and pharmacological evidence. *Alcoholism: Clinical and Experimental Research*. 2003; 27:149–154.
- Pandey SC, Chartoff EH, Carlezon WA Jr, et al. CREB gene transcription factors: role in molecular mechanisms of alcohol and drug addiction. *Alcoholism: Clinical and Experimental Research*. 2005; 29:176–184.
- Pandey SC, Lumeng L, Li T-K. Serotonin_{2C} receptors and serotonin_{2C} receptor-mediated phosphoinositide hydrolysis in the brain of alcohol-preferring and alcohol-nonpreferring rats. *Alcoholism: Clinical and Experimental Research*. 1996; 20:1038–1042.
- Panula P, Nuutinen S. Histamine and H₃ receptor in alcohol-related behaviors. *Journal of Pharmacology and Experimental Therapeutics*. 2011; 336:9–16. [PubMed: 20864504]
- Parent A, Descarries L, Beaudet A. Organization of ascending serotonin systems in the adult rat brain. A radioautographic study after intraventricular administration of [³H] 5 hydroxytryptamine. *Neuroscience*. 1981; 6:115–138. [PubMed: 6164006]

- Parker, SL. Neuropeptide Y: Molecular Structure, Role in Food Intake and Direct/Indirect Effects. Nova Biomedical; New York, NY: 2013.
- Parks MH, Dawant BM, Riddle WR, Hartmann SL, Dietrich MS, Nickel MK, et al. Longitudinal brain metabolic characterization of chronic alcoholics with proton magnetic resonance spectroscopy. *Alcoholism: Clinical and Experimental Research*. 2002; 26:1368–1380.
- Partanen, J.; Bruun, K.; Markkanen, T. Alcohol Research in the Northern Countries. The Finnish Foundation for Alcohol Studies. Vol. 14. Amquist & Wiksell; Stockholm: 1966. Inheritance of drinking behavior: a study on intelligence, personality, and use of alcohol of adult twins.
- Pautassi RM, Camarini R, Quadros IM, Miczek KA, Israel Y. Genetic and environmental influences on ethanol consumption: perspectives from preclinical research. *Alcoholism: Clinical and Experimental Research*. 2010; 34:976–987.
- Pessia M, Jiang ZG, North RA, Johnson SW. Actions of 5-hydroxytryptamine on ventral tegmental area neurons of the rat in vitro. *Brain Research*. 1994; 654:324–330. [PubMed: 7987681]
- Peterson JB, Pihl RO, Gianoulakis C, Conrod P, Finn PR, Stewart SH, et al. Ethanol-induced change in cardiac and endogenous opiate function and risk for alcoholism. *Alcoholism: Clinical and Experimental Research*. 1996; 20:1542–1552.
- Peterson JB, Pihl RO, Seguin JR, Finn PR, Stewart SH. Heart-rate reactivity and alcohol consumption among sons of male alcoholics and sons of non-alcoholics. *Journal of Psychiatry Neuroscience*. 1993; 18:190–198. [PubMed: 8373755]
- Pierce, RC.; Kenny, PJ. *Addiction*. Cold Spring Harbor Laboratory Press; New York, NY: 2013.
- Piercy KT, Bjork AK, Myers RD. The mixed 5HT_{1A}/2A receptor drug FG5938 suppresses alcohol drinking while enhancing feeding in P rats. *Alcohol*. 1996; 13:521–527. [PubMed: 8888950]
- Pivac N, Mück-Seler D, Mustapi M, Nenadi S, Vligin K, Kozari D, Kovaci D. Platelet serotonin concentration in alcoholic subjects. *Life Sciences*. 2004; 76:521–531. [PubMed: 15556165]
- Plemenitas A, Kastelic M, Porcelli S, Serretti A, Dolžan V, Kores Plesnicar B. Alcohol dependence and genetic variability in the serotonin pathway among currently and formerly alcohol-dependent males. *Neuropsychobiology*. 2015; 72:57–64. [PubMed: 26352193]
- Ponomarev I, Wang S, Zhang L, Harris RA, Mayfield RD. Gene coexpression networks in human brain identify epigenetic modifications in alcohol dependence. *Journal of Neuroscience*. 2012; 32:1884–1897. [PubMed: 22302827]
- Prelicpeanu D, Mihailescu R. Non A non B variants in the binary typology of alcoholics. *Addictive Disorders and Treatment*. 2005; 4:149–156.
- Presley CA, Meilman PW, Lyster R. Development of the Core Alcohol and Drug Survey: Initial findings and future directions. *Journal of American College Health*. 1994; 42:248–255. [PubMed: 8046164]
- Preuss UW, Watzke S, Wurst FM. Dimensionality and stages of severity of DSM-5 criteria in an international sample of alcohol-consuming individuals. *Psychiatry and Medicine*. 2014; 44:3303–3314. <http://dx.doi.org/10.1017/S0033291714000889>.
- Preuss UW, Wurst FM, Ridinger M, Rujescu D, Fehr C, Koller G, et al. Association of functional DBH genetic variants with alcohol dependence risk and related depression and suicide attempt phenotypes: results from a large multicenter association study. *Drug and Alcohol Dependence*. 2013; 133:459–467. [PubMed: 23906995]
- Rahman S. Nicotinic Receptors as Therapeutic Targets for Drug Addictive Disorders. *CNS Neurology and Discovery of Drug Targets*. 2013; 12:633–640.
- Rahman S, Engleman EA, Bell RL. Nicotinic receptor modulation to treat alcohol and drug dependence. *Frontiers in Neuroscience/Neuropharmacology*. 2014; 8:426. <http://dx.doi.org/10.3389/fnins.2014.00426>.
- Rahman S, Engleman EA, Bell RL. Nicotinic receptor modulation to treat alcohol and drug dependence. *Frontiers in Neuroscience/Neuropharmacology*. 2015; 8:426. <http://dx.doi.org/10.3389/fnins.2014.00426>.
- Rahman S, Engleman EA, Bell RL. Recent advances in nicotinic receptor signaling in alcohol abuse and alcoholism. *Progress in Molecular Biology and Translational Science*. 2016; 137:183–201. <http://dx.doi.org/10.1016/bs.pmbts.2015.10.004>. [PubMed: 26810002]

- Rahman S, McBride WJ. Involvement of GABA and cholinergic receptors in the nucleus accumbens on feedback control of somatodendritic dopamine release in the ventral tegmental area. *Journal of Neurochemistry*. 2002; 80:646–654. [PubMed: 11841572]
- Rahman S, Prendergast MA. Brain cholinergic receptor system as a target for treating alcohol dependence. *Recent Patents in CNS Drug Discovery*. 2012; 7:145–150. [PubMed: 22574675]
- Rajan I, Naga Venkatesha Musthy PJ, Ramakrishnan AG, Gangadhar BN. Heart rate variability as an index of cue reactivity in alcoholics. *Biological Psychiatry*. 1998; 43:544–546. [PubMed: 9547935]
- Rao P, Sari Y. Effectiveness of Ceftriaxone Treatment in Preventing Relapse-like Drinking Behavior Following Long-term Ethanol Dependence in P Rats. *Journal of Addiction Research and Therapeutics*. 2014; 5:183. <http://dx.doi.org/10.4172/2155-6105.1000183>.
- Rao PS, Bell RL, Engleman EA, Sari Y. Targeting glutamate uptake to treat alcohol use disorders. *Frontiers in Neuroscience/Neuropharmacology*. 2015; 9:144.
- Rao PS, Goodwani S, Bell RL, Wei Y, Boddu SH, Sari Y. Effects of ampicillin, cefazolin and cefoperazone treatments on GLT-1 expressions in the mesocorticolimbic system and ethanol intake in alcohol-preferring rats. *Neuroscience*. 2015; 295:164–174. [PubMed: 25813713]
- Rao PS, Sari Y. Effects of ceftriaxone on chronic ethanol consumption: a potential role for xCT and GLT1 modulation of glutamate levels in male P rats. *Journal of Molecular Neuroscience*. 2014; 54:71–77. [PubMed: 24535561]
- Rao PS, Saternos H, Goodwani S, Sari Y. Effects of ceftriaxone on GLT1 isoforms, xCT and associated signaling pathways in P rats exposed to ethanol. *Psychopharmacology (Berl)*. 2015; 232:2333–2342. [PubMed: 25619881]
- Rasmussen DD, Alexander L, Malone J, Federoff D, Froehlich JC. The α_2 adrenergic receptor agonist, clonidine, reduces alcohol drinking in alcohol-preferring (P) rats. *Alcohol*. 2014; 48:543–549. [PubMed: 25085719]
- Rasmussen DD, Alexander LL, Raskind MA, Froehlich JC. The α_1 -adrenergic receptor antagonist, prazosin, reduces alcohol drinking in alcohol-preferring (P) rats. *Alcoholism: Clinical and Experimental Research*. 2009; 33:264–272.
- Rasmussen DD, Beckwith LE, Kincaid CL, Froehlich JC. Combining the α_1 -adrenergic receptor antagonist, prazosin, with the β -adrenergic receptor antagonist, propranolol, reduces alcohol drinking more effectively than either drug alone. *Alcoholism: Clinical and Experimental Research*. 2014; 38:1532–1539.
- Rasmussen DD, Boldt BM, Wilkinson CW, Mitton DR. Chronic daily ethanol and withdrawal: 3. Forebrain pro-opiomelanocortin gene expression and implications for dependence, relapse, and deprivation effect. *Alcoholism: Clinical and Experimental Research*. 2002; 26:534–546.
- Rasmussen DD, Kincaid CL, Froehlich JC. Prazosin + naltrexone decreases alcohol drinking more effectively than does either drug alone in P rats with a protracted history of extensive voluntary alcohol drinking, dependence, and multiple withdrawals. *Alcoholism: Clinical and Experimental Research*. 2015; 39:1832–1841.
- Reed T, Page WF, Viken RJ, Christian JC. Genetic predisposition to organ-specific endpoints of alcoholism. *Alcoholism: Clinical and Experimental Research*. 1996; 20:1528–1533.
- Rehm J. The risks associated with alcohol use and alcoholism. *Alcohol Health Research World*. 2011; 34:135–143.
- Rehm J, Room R, Graham K, Monteiro M, Gmel G, Sempos CT. The relationship of average volume of alcohol consumption and patterns of drinking to burden of disease—an overview. *Addiction*. 2003; 98:1209–1228. [PubMed: 12930209]
- Renthal W, Nestler EJ. Histone acetylation in drug addiction. *Seminars in Cellular and Developmental Biology*. 2009; 20:387–394.
- Research Society on Alcoholism. *Impact of Alcoholism and Alcohol Induced Disease on America*. Author; Austin, TX: 2009.
- Research Society on Alcoholism. *Impact of Alcoholism and Alcohol Induced Disease on America*. Author; Austin, TX: 2015.

- Reser JE. Solitary mammals provide an animal model for autism spectrum disorders. *Journal of Comparative Psychology*. 2014; 128(1):99–113. <http://dx.doi.org/10.1037/a0034519>. [PubMed: 24188618]
- Reul JM, Holsboer F. Corticotropin-releasing factor receptors 1 and 2 in anxiety and depression. *Current Opinions in Pharmacology*. 2002; 2:23–33.
- Reyes BA, Bangasser DA, Valentino RJ, Van Bockstaele EJ. Using high resolution imaging to determine trafficking of corticotropin-releasing factor receptors in noradrenergic neurons of the rat locus coeruleus. *Life Sciences*. 2014; 112:2–9. [PubMed: 25058917]
- Rezvani AH, Overstreet DH, Lee Y-W. Attenuation of alcohol intake by ibogaine in three strains of alcohol-preferring rats. *Pharmacology, Biochemistry and Behavior*. 1995; 52:615–620.
- Rezvani AH, Overstreet DH, Mason GA, Janowsky DS, Hamed M, Clark E, Yang Y. Combination pharmacotherapy: a mixture of small doses of naltrexone, fluoxetine, and a thyrotropin-releasing hormone analogue reduces alcohol intake in three strains of alcohol-preferring rats. *Alcohol and Alcoholism*. 2000; 35:76–83. [PubMed: 10684782]
- Rezvani AH, Slade S, Wells C, Petro A, Lumeng L, Li T-K, et al. Effects of sazetidine-A, a selective alpha-4-beta-2 nicotinic acetylcholine receptor desensitizing agent, on alcohol and nicotine self-administration in selectively bred alcohol-preferring (P) rats. *Psychopharmacology*. 2010; 211:161–174. [PubMed: 20535453]
- Richardson HN, Lee SY, O'Dell LE, Koob GF, Rivier CL. Alcohol self-administration acutely stimulates the hypothalamic-pituitary-adrenal axis, but alcohol dependence leads to a dampened neuroendocrine state. *European Journal of Neuroscience*. 2008; 28:1641–1653. [PubMed: 18979677]
- Richter CP, Campbell KH. Alcohol taste thresholds and concentrations of solutions preferred by rats. *Science*. 1940; 9:507–508. [PubMed: 17847448]
- Riihioja P, Jaatinen P, Oksanen H, et al. Dexmedetomidine alleviates ethanol withdrawal symptoms in the rat. *Alcohol*. 1997; 14:537–544. [PubMed: 9401667]
- Robbins, TW.; Everitt, BJ.; Nutt, DJ. *The Neurobiology of Addiction: New Vistas*. Oxford University Press; New York, NY: 2010.
- Roberto M, Gilpin NW, Siggins GR. The central amygdala and alcohol: role of gamma-aminobutyric acid, glutamate, and neuropeptides. *Cold Spring Harbor Perspectives in Medicine*. 2012; 2:a012195. [PubMed: 23085848]
- Roberts AJ, Heyser CJ, Cole M, Griffin P, Koob GF. Excessive ethanol drinking following a history of dependence: Animal model of allostasis. *Neuropsychopharmacology*. 2000; 22:581–594. [PubMed: 10788758]
- Robinson G, Most D, Ferguson LB, Mayfield J, Harris RA, Blednov YA. Neuroimmune pathways in alcohol consumption: evidence from behavioral and genetic studies in rodents and humans. *International Reviews in Neurobiology*. 2014; 118:13–39.
- Robinson TE, Berridge KC. The incentive sensitization theory of addiction: some current issues. *Philosophical Transactions of the Royal Society of London Part B Biological Science*. 2008; 363:3137–3146.
- Rodd ZA, Bell RL, Kuc KA, Murphy JM, Lumeng L, Li T-K, et al. Effects of Repeated Alcohol Deprivations on Operant Ethanol Self-Administration by Alcohol-Preferring (P) Rats. *Neuropsychopharmacology*. 2003; 28:1614–1621. [PubMed: 12799615]
- Rodd ZA, Bell RL, McKinzie DL, Webster AA, Murphy JM, Lumeng L, et al. Low-dose stimulatory effects of ethanol during adolescence in rat lines selectively-bred for high alcohol intake. *Alcoholism: Clinical and Experimental Research*. 2004; 28:535–543.
- Rodd ZA, Bell RL, Oster SM, Toalston JE, Pommer TJ, McBride WJ, et al. Serotonin-3 receptors in the posterior ventral tegmental area regulate ethanol self-administration of alcohol-preferring (P) rats. *Alcohol*. 2010; 44:245–255. [PubMed: 20682192]
- Rodd ZA, Bell RL, Sable HJ, Murphy JM, McBride WJ. Recent advances in animal models of alcohol craving and relapse. *Pharmacology, Biochemistry and Behavior*. 2004; 79:439–450.
- Rodd ZA, Kimpel MW, Edenberg HJ, Bell RL, Strother WN, McClintick JN, et al. Differential gene expression in the nucleus accumbens with ethanol self-administration in inbred alcohol-preferring rats. *Pharmacology, Biochemistry and Behavior*. 2008; 89:481–498.

- Rodd ZA, McKinzie DL, Bell RL, McQueen VK, Murphy JM, Schoepp DD, et al. The metabotropic glutamate 2/3 receptor agonist LY404039 reduces alcohol-seeking but not alcohol self-administration in alcohol-preferring (P) rats. *Behavioural Brain Research*. 2006; 171:207–215. [PubMed: 16678921]
- Rodd-Henricks ZA, Bell RL, Kuc KA, Murphy JM, McBride WJ, Lumeng L, et al. Effects of ethanol exposure on subsequent acquisition and extinction of ethanol self-administration and expression of alcohol-seeking behavior in adult alcohol-preferring (P) rats: I. Periadolescent exposure. *Alcoholism: Clinical and Experimental Research*. 2002; 26:1632–1641.
- Rodd-Henricks ZA, Bell RL, Kuc KA, Murphy JM, McBride WJ, Lumeng L, et al. Effects of ethanol exposure on subsequent acquisition and extinction of ethanol self-administration and expression of alcohol-seeking behavior in adult alcohol-preferring (P) rats: II. Adult exposure. *Alcoholism: Clinical and Experimental Research*. 2002; 26:1642–1652.
- Rodd-Henricks ZA, Bell RL, Murphy JM, McBride WJ, Lumeng L, Li T-K. Effects of concurrent access to multiple ethanol concentrations and repeated deprivations on alcohol intake of alcohol-preferring (P) rats. *Alcoholism: Clinical and Experimental Research*. 2001; 24:747–753.
- Rodd-Henricks ZA, McKinzie DL, Edmundson VE, Dagon CL, Murphy JM, McBride WJ, et al. Effects of 5-HT₃ receptor antagonists on daily alcohol intake under acquisition, maintenance, and relapse conditions in alcohol-preferring (P) rats. *Alcohol*. 2000; 21:73–85. [PubMed: 10946160]
- Rodd-Henricks ZA, McKinzie DL, Shaikh SR, Murphy JM, McBride WJ, Lumeng L, et al. The alcohol deprivation effect is prolonged in the alcohol preferring (P) rat following repeated deprivations. *Alcoholism: Clinical and Experimental Research*. 2000; 24:8–16.
- Rothstein JD, Van Kammen M, Levey AI, Martin LJ, Kuncel RW. Selective loss of glial glutamate transporter GLT-1 in amyotrophic lateral sclerosis. *Annals in Neurology*. 1995; 38:73–84.
- Russell RN, McBride WJ, Lumeng L, Li T-K, Murphy JM. Apomorphine and 7-OH DPAT reduce ethanol intake of P and HAD rats. *Alcohol*. 1996; 13:515–519. [PubMed: 8888949]
- Saba LM, Flink SL, Vanderlinden LA, Israel Y, Tampier L, Colombo G, et al. The sequenced rat brain transcriptome, its use in identifying networks predisposing alcohol consumption. *The FEBS Journal*. 2015; 282:3556–3578. <http://dx.doi.org/10.1111/febs.13358>. [PubMed: 26183165]
- Sable HJK, Bell RL, Rodd ZA, McBride WJ. Effects of naltrexone on the acquisition of alcohol intake in male and female periadolescent and adult alcohol-preferring (P) rats. *International Journal of Adolescent Medicine and Health*. 2006; 18:139–149. [PubMed: 16639868]
- Saccone SF, Hinrichs AL, Saccone NL, Chase GA, Konvicka K, Madden PA, ... Bierut LJ, et al. Cholinergic nicotinic receptor genes implicated in a nicotine dependence association study targeting 348 candidate genes with 3713 SNPs. *Human Molecular Genetics*. 2007; 16:36–49. [PubMed: 17135278]
- Sahr AE, Thielen RJ, Lumeng L, Li T-K, McBride WJ. Long-lasting alterations of the mesolimbic dopamine system after periadolescent ethanol drinking by alcohol-preferring rats. *Alcoholism: Clinical and Experimental Research*. 2004; 28:702–711.
- Sajdyk TJ, Shekhar A, Gehlert DR. Interactions between NPY and CRF in the amygdala to regulate emotionality. *Neuropeptides*. 2004; 38:225–234. [PubMed: 15337374]
- Sajja RK, Dwivedi C, Rahman S. Nicotinic ligands modulate ethanol-induced dopamine functions in mice. *Pharmacology*. 2010; 86:168–173. [PubMed: 20714207]
- Sajja RK, Rahman S. Lobeline and cytosine reduce voluntary ethanol drinking behavior in male C57BL/6J mice. *Progress in Neuropsychopharmacology and Biological Psychiatry*. 2011; 35:257–264.
- Sajja RK, Rahman S. Neuronal nicotinic receptor ligands modulate chronic nicotine-induced ethanol consumption in C57BL/6J mice. *Pharmacology, Biochemistry and Behavior*. 2012; 102:36–43.
- Sajja RK, Rahman S. Nicotinic receptor partial agonists modulate alcohol deprivation effect in C57BL/6J mice. *Pharmacology, Biochemistry and Behavior*. 2013; 110:161–167.
- Sajja RK, Dwivedi C, Rahman S. Nicotinic ligands modulate ethanol-induced dopamine functions in mice. *Pharmacology*. 2010; 86:168–173. [PubMed: 20714207]
- Samson HH, Czachowski CL. Behavioral measures of alcohol self-administration and intake control: rodent models. *International Reviews in Neurobiology*. 2003; 54:107–143.

- Samson HH, Files FJ, Denning C, Marvin S. Comparison of alcohol-preferring and -nonpreferring selectively bred rat lines. I. Ethanol initiation and limited access operant self-administration. *Alcoholism: Clinical and Experimental Research*. 1998; 22:2133–2146.
- Sari Y. Role of glutamate transporter 1 in the attenuation of alcohol intake. *Frontiers in Neuroscience/Neuropharmacology*. 2014; 8:Article 200.
- Sari Y, Bell RL, Zhou FC. Effects of chronic alcohol and repeated deprivations on dopamine D1 and D2 receptor levels in the extended amygdala of inbred alcohol-preferring rats. *Alcoholism: Clinical and Experimental Research*. 2006; 30:46–56.
- Sari Y, Sakai M, Weedman JM, Rebec GV, Bell RL. Ceftriaxone, a beta-lactam antibiotic, reduces ethanol consumption in alcohol-preferring rats. *Alcohol and Alcoholism*. 2011; 46:239–246. <http://dx.doi.org/10.1093/alcalc/agr023>. [PubMed: 21422004]
- Sari Y, Sreemantula SN. Neuroimmunophilin GPI-1046 reduces ethanol consumption in part through activation of GLT1 in alcohol-preferring rats. *Neuroscience*. 2012; 227:327–335. [PubMed: 23059796]
- Sari Y, Sreemantula SN, Lee MR, Choi DS. Ceftriaxone treatment affects the levels of GLT1 and ENT1 as well as ethanol intake in alcohol-preferring rats. *Journal of Molecular Neuroscience*. 2013; 51:779–787. [PubMed: 23893122]
- Sartor CE, Kranzler HR, Gelernter J. Rate of progression from first use to dependence on cocaine or opioids: a cross-substance examination of associated demographic, psychiatric, and childhood risk factors. *Addictive Behaviors*. 2014; 39:473–479. <http://dx.doi.org/10.1016/j.addbeh.2013.10.021>. [PubMed: 24238782]
- Sato N, Kageyama S, Chen R, Suzuki M, Mori H, Tanioka F, Sugimura H, et al. Association between neuropeptide Y receptor 2 polymorphism and the smoking behavior of elderly Japanese. *Journal of Human Genetics*. 2010; 55:755–760. [PubMed: 20811389]
- Sauvage M, Steckler T. Detection of corticotropin-releasing hormone receptor 1 immunoreactivity in cholinergic, dopaminergic and noradrenergic neurons of the murine basal forebrain and brainstem nuclei--potential implication for arousal and attention. *Neuroscience*. 2001; 104:643–652. [PubMed: 11440798]
- Schlaepfer IR, Hoft NR, Collins AC, Corley RP, Hewitt JK, Hopfer CJ, et al. Ehringer MA. The CHRNA5/A3/B4 gene cluster variability as an important determinant of early alcohol and tobacco initiation in young adults. *Biological Psychiatry*. 2008; 63:1039–1046. [PubMed: 18163978]
- Schmidt LG, Dufeu P, Heinz A, Kuhn S, Rommelspacher H. Serotonergic dysfunction in addiction: effects of alcohol, cigarette smoking and heroin on platelet 5-HT content. *Psychiatry Research*. 1997; 72:177–185. [PubMed: 9406907]
- Schroeder JP, Overstreet DH, Hodge CW. The mGluR5 antagonist MPEP decreases operant ethanol self-administration during maintenance and after repeated alcohol deprivations in alcohol-preferring (P) rats. *Psychopharmacology*. 2005a; 179:262–270. [PubMed: 15717208]
- Schroeder JP, Overstreet DH, Hodge CW. The neuropeptide-Y Y5 receptor antagonist L152,804 decreases alcohol self-administration in inbred alcohol-preferring (iP) rats. *Alcohol*. 2005b; 36:179–186. [PubMed: 16377459]
- Schuckit MA. A brief history of research on the genetics of alcohol and other drug use disorders. *Journal of Studies on Alcohol and Drugs*. 2014; (Suppl 17):59–67. [PubMed: 24565312]
- Schuckit MA. Ethanol-induced changes in body sway in men of high alcoholism risk. *Archives in General Psychiatry*. 1985; 42:3375–3379.
- Schuckit MA. Genetic aspects of alcoholism. *Annals in Emergency Medicine*. 1986; 15:991–996.
- Schuckit MA. Low level of response to alcohol as a predictor of future alcoholism. *American Journal of Psychiatry*. 1994; 151:184–189. [PubMed: 8296886]
- Schuckit MA. An overview of genetic influences in alcoholism. *Journal of Substance Abuse and Treatment*. 2009; 36:S5–S14.
- Schuckit MA, Gold EO. A simultaneous evaluation of multiple markers of ethanol/placebo challenges in sons of alcoholics and controls. *Archives of General Psychiatry*. 1988; 45:211–216. [PubMed: 3422553]

- Seal RP, Amara SG. Excitatory amino acid transporters: a family in flux. *Annual Reviews in Pharmacology and Toxicology*. 1999; 39:431–456.
- Self, DW.; Staley, JK. *Behavioral Neuroscience of Drug Addiction*. Springer; New York, NY: 2010.
- Shield KD, Rehm J, Gmel G, Rehm MX, Allamani A. Alcohol consumption, alcohol dependence, and related mortality in Italy in 2004: effects of treatment-based interventions on alcohol dependence. *Substance Abuse Treatment and Prevention Policy*. 2013; 8:21. <http://dx.doi.org/10.1186/1747-597X-8-21>.
- Shin JH, Adrover MF, Wess J, Alvarez VA. Muscarinic regulation of dopamine and glutamate transmission in the nucleus accumbens. *Proceedings of the National Academy of Sciences, USA*. 2015; 112:8124–8129.
- Silveri MM. Adolescent brain development and underage drinking in the United States: identifying risks of alcohol use in college populations. *Harvard Reviews in Psychiatry*. 2012; 20:189–200. <http://dx.doi.org/10.3109/10673229.2012.714642>.
- Silveri MM. GABAergic contributions to alcohol responsivity during adolescence: insights from preclinical and clinical studies. *Pharmacology and Therapeutics*. 2014; 143:197–216. <http://dx.doi.org/10.1016/j.pharmthera.2014.03.001>. [PubMed: 24631274]
- Simpson TL, Saxon AJ, Meredith CW, et al. A pilot trial of the alpha-1 adrenergic antagonist, prazosin, for alcohol dependence. *Alcoholism: Clinical and Experimental Research*. 2009; 33:255–263.
- Smialowska M, Wieronska JM, Wedzony K. A search for colocalization of mglu1a receptors with CRF or NPY in the rat brain amygdala. *Folia Histochemica et Cytobiologica*. 2002; 40:153–154. [PubMed: 12056621]
- Smith AD, Weiss F. Ethanol exposure differentially alters central monoamine neurotransmission in alcohol-preferring versus -nonpreferring rats. *Journal of Pharmacology and Experimental Therapeutics*. 1999; 288:1223–1228. [PubMed: 10027862]
- Smith JE, Co C, McIntosh S, Cunningham CC. Chronic binge-like moderate ethanol drinking in rats results in widespread decreases in serotonin, dopamine, and norepinephrine turnover rates reversed by ethanol intake. *Journal of Neurochemistry*. 2008; 105:2134–2155. [PubMed: 18284613]
- Smith TL. Regulation of glutamate uptake in astrocytes continuously exposed to ethanol. *Life Sciences*. 1997; 61:2499–2505. [PubMed: 9416769]
- Smith TL, Zsigo A. Increased Na(+)-dependent high affinity uptake of glutamate in astrocytes chronically exposed to ethanol. *Neuroscience Letters*. 1996; 218:142–144. [PubMed: 8945748]
- Soderpalm B, Ericson M. Neurocircuitry involved in the development of alcohol addiction: the dopamine system and its access points. *Current Topics in Behavioral Neuroscience*. 2013; 13:127–161.
- Soderpalm B, Ericson M, Olausson P, Blomqvist O, Engel JA. Nicotinic mechanisms involved in the dopamine activating and reinforcing properties of ethanol. *Behavioural Brain Research*. 2000; 113:85–96. [PubMed: 10942035]
- Sofuoglu M, Rosenheck R, Petrakis I. Pharmacological treatment of comorbid PTSD and substance use disorder: recent progress. *Addictive Behaviors*. 2014; 39:428–433. [PubMed: 24035645]
- Sommer, WH.; Spanagel, R. *Behavioral Neurobiology of Alcohol Addiction*. Springer; New York, NY: 2013.
- Sorbel J, Morzorati S, O'Connor S, Li T-K, Christian JC. Alcohol effects on the heritability of EEG spectral power. *Alcoholism: Clinical and Experimental Research*. 1996; 20:1523–1527.
- Spanagel R. Alcoholism: A systems approach from molecular physiology to addictive behavior. *Physiology Reviews*. 2009; 89:649–705.
- Spanagel R, Durstewitz D, Hansson A, Heinz A, Kiefer F, Kohr G, Derks EM, et al. A systems medicine research approach for studying alcohol addiction. *Addiction Biology*. 2013; 18:883–896. [PubMed: 24283978]
- Spear, LP. *The Behavioral Neuroscience of Adolescence*. W.W. Norton; New York, NY: 2010.
- Spear LP. Adolescents and alcohol: acute sensitivities, enhanced intake, and later consequences. *Neurotoxicology and Teratology*. 2014; 41:51–59. <http://dx.doi.org/10.1016/j.ntt.2013.11.006>. [PubMed: 24291291]

- Spence JP, Liang T, Habegger K, Carr LG. Effect of polymorphism on expression of the neuropeptide Y gene in inbred alcohol-preferring and –nonpreferring rats. *Neuroscience*. 2005; 131:871–876. [PubMed: 15749341]
- Spence JP, Liang T, Liu L, Johnson PL, Foroud T, Carr LG, Shekhar A. From QTL to candidate gene: a genetic approach to alcoholism research. *Current Drugs of Abuse Reviews*. 2009; 2:127–134.
- Stanley SA, Murphy KG, Bewick GA, Kong WM, Opacka-Juffry J, Gardiner JV, et al. Regulation of rat pituitary cocaine- and amphetamine-regulated transcript (CART) by CRH and glucocorticoids. *American Journal of Physiology and Endocrinology Metabolism*. 2004; 287:E583–E590.
- Steensland P, Simms JA, Holgate J, Richards JK, Bartlett SE. Varenicline, an alpha4beta2 nicotinic acetylcholine receptor partial agonist, selectively decreases ethanol consumption and seeking. *Proceedings of the National Academy of Sciences, USA*. 2007; 104:12518–12523.
- Stewart RB, Gatto GJ, Lumeng L, Li T-K, Murphy JM. Comparison of alcohol-preferring (P) and nonpreferring (NP) rats on tests of anxiety and for the anxiolytic effects of ethanol. *Alcohol*. 1993; 10:1–10. [PubMed: 8095393]
- Stewart RB, Kurtz DL, Zweifel M, Li T-K, Froehlich JC. Differences in the hypothermic response to ethanol in rats selectively bred for oral ethanol preference and nonpreference. *Psychopharmacology*. 1992; 106:169–174. [PubMed: 1549644]
- Stewart RB, McBride WJ, Lumeng L, Li T-K, Murphy JM. Chronic alcohol consumption in alcohol-preferring P rats attenuates subsequent conditioned taste aversion produced by ethanol injections. *Psychopharmacology*. 1991; 105:530–534. [PubMed: 1771221]
- Stewart RB, Murphy JM, McBride WJ, Lumeng L, Li T-K. Place conditioning with alcohol in alcohol-preferring and –nonpreferring rats. *Pharmacology, Biochemistry and Behavior*. 1996; 53:487–491.
- Stormark KM, Laberg JC, Nordby H, Hugdahl K. Heart rate responses indicate locked-in attention in alcoholics immediately prior to drinking. *Addictive Behaviors*. 1998; 23:251–255. [PubMed: 9573428]
- Strobel C, Hunt S, Sullivan R, Sun J, Sah P. Emotion regulation of pain: the role of noradrenaline in the amygdala. *Science China. Life Sciences*. 2014; 57:384–390. [PubMed: 24643418]
- Strother WN, Chernet EJ, Lumeng L, Li T-K, McBride WJ. Regional central nervous system densities of delta-opioid receptors in alcohol-preferring P, alcohol-nonpreferring NP, and unselected Wistar rats. *Alcohol*. 2001; 25:31–38. [PubMed: 11668015]
- Strother WN, Lumeng L, Li T-K, McBride WJ. Dopamine and serotonin content in select brain regions of weanling and adult alcohol drinking rat lines. *Pharmacology, Biochemistry and Behavior*. 2005; 80:229–237.
- Strother WN, Lumeng L, Li T-K, McBride WJ. Regional CNS densities of serotonin1A and dopamine D2 receptors in peri-adolescent alcohol-preferring P and alcohol-nonpreferring NP rat pups. *Pharmacology, Biochemistry and Behavior*. 2003; 74:335–342.
- Szumliński KK, Diab ME, Friedman R, Henze LM, Lominac KD, Bowers MS. Accumbens neurochemical adaptations produced by binge-like alcohol consumption. *Psychopharmacology*. 2007; 190:415–431. [PubMed: 17225170]
- Tabakoff B, Hoffman PL. The neurobiology of alcohol consumption and alcoholism: an integrative history. *Pharmacology, Biochemistry and Behavior*. 2013; 113:20–37.
- Tabakoff B, Saba L, Printz M, Flodman P, Hodgkinson C, ... Goldman Hoffman PL, et al. Genetical genomic determinants of alcohol consumption in rats and humans. *BMC Biology*. 2009; 7:70. <http://dx.doi.org/10.1186/1741-7007-7-70>. [PubMed: 19874574]
- Tanchuck MA, Yoneyama N, Ford MM, Fretwell AM, Finn DA. Assessment of GABA-B, metabotropic glutamate, and opioid receptor involvement in an animal model of binge drinking. *Alcohol*. 2011; 45:33–44. [PubMed: 20843635]
- Terunuma M, Revilla-Sanchez R, Quadros IM, Deng Q, Deeb TZ, Lumb M, et al. Postsynaptic GABAB receptor activity regulates excitatory neuronal architecture and spatial memory. *Journal of Neuroscience*. 2014; 34:804–816. [PubMed: 24431439]
- Thanos PK, Katana JM, Ashby CR Jr, Michaelides M, Gardner EL, Heidbreder CA, et al. The selective dopamine D3 receptor antagonist SB-277011-A attenuates ethanol consumption in ethanol

preferring (P) and non-preferring (NP) rats. *Pharmacology, Biochemistry and Behavior*. 2005; 81:190–197.

- Thiele E, Badia-Elder NE. A role for neuropeptide Y in alcohol intake control: evidence from human and animal research. *Physiology and Behavior*. 2003; 79:95–101. [PubMed: 12818714]
- Thielen RJ, Engleman EA, Rodd ZA, Murphy JM, Lumeng L, Li T-K, McBride WJ. Ethanol drinking and deprivation alter dopaminergic and serotonergic function in the nucleus accumbens of alcohol-preferring rats. *Journal of Pharmacology and Experimental Therapeutics*. 2004; 309:216–225. [PubMed: 14718606]
- Thielen RJ, McBride WJ, Chernet E, Lumeng L, Li T-K. Regional densities of benzodiazepine sites in the CNS of alcohol-naïve P and NP rats. *Pharmacology, Biochemistry and Behavior*. 1997; 57:875–882.
- Thoma R, Mullins P, Ruhl D, Monnig M, Yeo RA, Caprihan A, Gasparovic C, et al. Perturbation of the glutamate-glutamine system in alcohol dependence and remission. *Neuropsychopharmacology*. 2011; 36:1359–1365. [PubMed: 21389979]
- Thorsell A. Brain neuropeptide Y and corticotropin-releasing hormone in mediating stress and anxiety. *Experimental Biology and Medicine*. 2010; 235:1163–1167. [PubMed: 20881320]
- Tizabi Y, Bai L, Copeland RL Jr, Taylor RE. Combined effects of systemic alcohol and nicotine on dopamine release in the nucleus accumbens shell. *Alcohol and Alcoholism*. 2007; 42:413–416. [PubMed: 17686828]
- Tizabi Y, Getachew B, Davila-Garcia M, Taylor RE. Alcohol preference: association with reduced striatal nicotinic receptors. *Alcohol and Alcoholism*. 2001; 36:318–322. [PubMed: 11468132]
- Thorsell A. Brain neuropeptide Y and corticotropin-releasing hormone in mediating stress and anxiety. *Experimental Biology and Medicine*. 2010; 235:1163–1167. [PubMed: 20881320]
- Toalston JE, Deehan JA Jr, Hauser SR, Engleman EA, Bell RL, Murphy JM, et al. The reinforcing properties and neurochemical response of ethanol within the posterior ventral tegmental area are enhanced in adulthood by peri-adolescent ethanol consumption. *The Journal of Pharmacology and Experimental Therapeutics*. 2014; 351:317–326. <http://dx.doi.org/10.1124/jpet.114.218172>. [PubMed: 25150280]
- Toalston JE, Oster SM, Kuc KA, Ding Z-M, Pommer TJ, Murphy JM, et al. Effects of alcohol and saccharin deprivations on concurrent ethanol and saccharin operant self-administration by alcohol-preferring (P) rats. *Alcohol*. 2008; 42:277–284. [PubMed: 18400451]
- Traynelis SF, Wollmuth LP, McBain CJ, Menniti FS, Vance KM, Ogden KK, ... Dingledine R, et al. Glutamate receptor ion channels: structure, regulation, and function. *Pharmacology Reviews*. 2010; 62:405–496.
- Trim RS, Simmons AN, Tolentino NJ, Hall SA, Matthews SC, Robinson SK, ... Schuckit MA, et al. Acute ethanol effects on brain activation in low- and high-level responders to alcohol. *Alcoholism: Clinical and Experimental Research*. 2010; 34:1162–1170.
- Truitt WA, Hauser SR, Deehan GA Jr, Toalston JE, Wilden JA, Bell RL, et al. Ethanol and nicotine interaction within the posterior ventral tegmental area in male and female alcohol-preferring rats: evidence of synergy and differential gene activation in the nucleus accumbens shell. *Psychopharmacology (Berl)*. 2015; 232:639–649. [PubMed: 25155311]
- Van Bockstaele EJ, Cestari DM, Pickel VM. Synaptic structure and connectivity of serotonin terminals in the ventral tegmental area: potential sites for modulation of mesolimbic dopamine neurons. *Brain Research*. 1994; 647:307–322. [PubMed: 7522922]
- Van Pett K, Viau V, Bittencourt JC, Chan RK, Li HY, Arias C, et al. Distribution of mRNAs encoding CRF receptors in brain and pituitary of rat and mouse. *Journal of Comparative Neurology*. 2000; 428:191–212. [PubMed: 11064361]
- Van Rizen MJ, Dishion TJ. Adolescent deviant peer clustering as an amplifying mechanism underlying the progression from early substance use to late adolescent dependence. *Journal of Child Psychology and Psychiatry*. 2014; 55(10):1153–1161. <http://dx.doi.org/10.1111/jcpp.12211>. [PubMed: 24673521]
- Varlinskaya EI, Spear LP. Ethanol-induced social facilitation in adolescent rats: Role of endogenous activity at mu opioid receptors. *Alcoholism: Clinical and Experimental Research*. 2009; 33:991–1000.

- Varlinskaya EI, Spear LP. Sensitization to social anxiolytic effects of ethanol in adolescent and adult Sprague-Dawley rats after repeated ethanol exposure. *Alcohol*. 2010; 44:99–110. [PubMed: 20113878]
- Vengeliene V, Bilbao A, Molander A, Spanagel R. Neuropharmacology of alcohol addiction. *British Journal of Pharmacology*. 2008; 154:299–315. [PubMed: 18311194]
- Vengeliene V, Bilbao A, Spanagel R. The alcohol deprivation effect model for studying relapse behavior: a comparison between rats and mice. *Alcohol*. 2014; 48:313–320. [PubMed: 24811155]
- Verplaetse TL, Czachowski CL. Low-dose prazosin alone and in combination with propranolol or naltrexone: effects on ethanol and sucrose seeking and self-administration in the P rat. *Psychopharmacology*. 2015; 232:2647–2657. [PubMed: 25743758]
- Verplaetse TL, Rasmussen DD, Froehlich JC, Czachowski CL. Effects of prazosin, an α 1-adrenergic receptor antagonist, on the seeking and intake of alcohol and sucrose in alcohol-preferring (P) rats. *Alcoholism: Clinical and Experimental Research*. 2012; 36:881–886.
- Vetreno RP, Broadwater M, Liu W, Spear LP, Crews FT. Adolescent, but not adult, binge ethanol exposure leads to persistent global reductions of choline acetyltransferase expressing neurons in brain. *PLoS One*. 2014; 9:e113421. [PubMed: 25405505]
- Viken RJ, Rose RJ, Morzorati SL, Christian JC, Li T-K. Subjective intoxication in response to alcohol challenge: heritability and covariation with personality, breath alcohol level, and drinking history. *Alcoholism: Clinical and Experimental Research*. 2003; 27:795–803.
- Vilaró MT, Palacios JM, Mengod G. Localization of m5 muscarinic receptor mRNA in rat brain examined by in situ hybridization histochemistry. *Neuroscience Letters*. 1990; 114:154–159. [PubMed: 2395528]
- Volkow ND, Li T-K. Drugs and alcohol: Treating and preventing abuse, addiction and their medical consequences. *Pharmacology and Therapeutics*. 2005; 108:3–17. [PubMed: 16098597]
- Volkow ND, Morales M. The Brain on Drugs: From Reward to Addiction. *Cell*. 2015; 162:712–725.10.1016/j.cell.2015.07.046 [PubMed: 26276628]
- Walker BM, Rasmussen DD, Raskind MA, Koob GF. Alpha1-noradrenergic receptor antagonism blocks dependence-induced increases in responding for ethanol. *Alcohol*. 2008; 42:91–97. [PubMed: 18358987]
- Waller MB, McBride WJ, Gatto GJ, Lumeng L, Li T-K. Intragastric self-administration of ethanol by ethanol-preferring and –nonpreferring lines of rats. *Science*. 1984; 225:78–80. [PubMed: 6539502]
- Waller MB, McBride WJ, Lumeng L, Li T-K. Induction of dependence on ethanol by free-choice drinking in alcohol-preferring rats. *Pharmacology, Biochemistry and Behavior*. 1982; 16:501–507.
- Waller MB, Murphy JM, McBride WJ, Lumeng L, Li T-K. Effect of low dose ethanol on spontaneous motor activity in alcohol-preferring and –nonpreferring lines of rats. *Pharmacology, Biochemistry and Behavior*. 1986; 24:617–623.
- Wand G. The anxious amygdala: CREB signaling and predisposition to anxiety and alcoholism. *Journal of Clinical Investigations*. 2005; 115:2697–2699.
- Wang JC, Gruzca R, Cruchaga C, Hinrichs AL, Bertelsen S, Budde JP, ... Goate AM, et al. Genetic variation in the CHRNA5 gene affects mRNA levels and is associated with risk for alcohol dependence. *Molecular Psychiatry*. 2009; 14:501–510. [PubMed: 18414406]
- Wang JC, Hinrichs AL, Stock H, Budde J, Allen R, Bertelsen S, ... Bierut LJ, et al. Evidence of common and specific genetic effects: association of the muscarinic acetylcholine receptor M2 (CHRM2) gene with alcohol dependence and major depressive syndrome. *Human Molecular Genetics*. 2004; 13:1903–1911. [PubMed: 15229186]
- Wang JC, Kapoor M, Goate AM. The genetics of substance dependence. *Annual Reviews in Genomics and Human Genetics*. 2012; 13:241–261.
- Wang Y, Qin ZH. Molecular and cellular mechanisms of excitotoxic neuronal death. *Apoptosis*. 2010; 15:1382–1402.10.1007/s10495-010-0481-0 [PubMed: 20213199]

- Wang Z, et al. Dopaminergic control of corticostriatal long-term synaptic depression in medium spiny neurons is mediated by cholinergic interneurons. *Neuron*. 2006; 50:443–452. [PubMed: 16675398]
- Warburton EC, Barker GRI, Brown MW. Investigations into the involvement of NMDA mechanisms in recognition memory. *Neuropharmacology*. 2013; 74:41–47. [PubMed: 23665343]
- Ward RJ, Colivicchi MA, Allen R, Schol F, Lallemand F, de Witte P, et al. Neuro-inflammation induced in the hippocampus of ‘binge drinking’ rats may be mediated by elevated extracellular glutamate content. *Journal of Neurochemistry*. 2009; 111:1119–1128. [PubMed: 19765190]
- Ward RJ, Lallemand F, de Witte P. Influence of adolescent heavy session drinking on the systemic and brain innate immune system. *Alcohol and Alcoholism*. 2014; 49:193–197. [PubMed: 24532587]
- Wechsler H, Lee JE, Kuo M, Lee H. College binge drinking in the 1990s: a continuing problem. Results of the Harvard School of Public Health 1999 College Alcohol Study. *Journal of American College Health*. 2000; 48:199–210. [PubMed: 10778020]
- Weiner DM, Levey AI, Brann MR. Expression of muscarinic acetylcholine and dopamine receptor mRNAs in rat basal ganglia. *Proceedings National Academy of Sciences, USA*. 1990; 87:7050–7054.
- Weiss F, Mitchiner M, Bloom FE, Koob GF. Free-choice ethanol responding for ethanol versus water in alcohol-preferring (P) and unselected Wistar rats is differentially modified by naloxone, bromocriptine, and methysergide. *Psychopharmacology*. 1990; 101:178–186. [PubMed: 2349359]
- Weitlauf C, Woodward JJ. Ethanol selectively attenuates NMDAR-mediated synaptic transmission in the prefrontal cortex. *Alcoholism: Clinical and Experimental Research*. 2008; 32:690–698.
- Wess J. Novel insights into muscarinic acetylcholine receptor function using gene targeting technology. *Trends in Pharmacological Sciences*. 2003; 24:414–20. [PubMed: 12915051]
- Wess J, Duttaroy A, Zhang W, Gomeza J, Cui Y, Miyakawa T, ... Yamada M, et al. M1–M5 muscarinic receptor knockout mice as novel tools to study the physiological roles of the muscarinic cholinergic system. *Receptors and Channels*. 2003; 9:279–290. [PubMed: 12893539]
- Wetherill L, Schuckit MA, Hesselbrock V, Xuei X, Liang T, Dick DM, ... Foroud T, et al. Neuropeptide Y receptor genes are associated with alcohol dependence, alcohol withdrawal phenotypes, and cocaine dependence. *Alcoholism: Clinical and Experimental Research*. 2008; 32:2031–2040.
- White AM, Kraus CL, Swartzwelder H. Many college freshmen drink at levels far beyond the binge threshold. *Alcoholism: Clinical and Experimental Research*. 2006; 30:1006–1010.
- Whiteside GT, Pomonis JD, Kennedy JD. An industry perspective on the role and utility of animal models of pain in drug discovery. *Neuroscience Letters*. 2013; 557(Pt. A):65–72. <http://dx.doi.org/10.1016/j.neulet.2013.08.033>. [PubMed: 23994390]
- Williams RJ, Berry LJ, Beerstecher E Jr. Individual metabolic patterns, alcoholism, genotrophic diseases. *Proceedings National Academy of Sciences, USA*. 1949; 35:265–271.
- Williams TJ, LaForge KS, Gordon D, Bart G, Kellog S, Ott J, Kreek MJ. Prodynorphin gene promoter repeat associated with cocaine/alcohol codependence. *Addiction Biology*. 2007; 12:496–502. [PubMed: 17559549]
- Wimalawansa SJ. Mechanisms of developing post-traumatic stress disorder: new targets for drug development and other potential interventions. *CNS Neurology Drug Targets*. 2014; 13:807–816.
- Windle M, Scheidt DM. Alcoholic subtypes: are two sufficient? *Addiction*. 2004; 99:1508–1519. [PubMed: 15585042]
- Wise RA, Bozarth MA. A psychomotor stimulant theory of addiction. *Psychology Reviews*. 1987; 94:469–492.
- Witkin JM, Statnick MA, Rorick-Kehn LM, Pintar JE, Ansonoff M, Chen Y, et al. The biology of nociceptin/orphanin FQ (N/OFQ) related to obesity, stress, anxiety, mood, and drug dependence. *Pharmacology and Therapeutics*. 2014; 141:283–299. [PubMed: 24189487]
- Witt ED. Mechanisms of alcohol abuse and alcoholism in adolescents: a case for developing animal models. *Behavioral and Neural Biology*. 1994; 62:168–177. [PubMed: 7857239]
- Witt, ED. Neurobiology. In: Galanter, M., editor. *Alcohol Problems in Adolescents and Young Adults*. Springer; New York, NY: 2006. p. 119–122.

- Witt ED. Research on alcohol and adolescent brain development: opportunities and future directions. *Alcohol*. 2010; 44:119–124. [PubMed: 20113880]
- Wolak ML, DeJoseph MR, Cator AD, Mokashi AS, Brownfield MS, Urban JH. Comparative distribution of neuropeptide Y Y1 and Y5 receptors in the rat brain by using immunohistochemistry. *Journal of Comparative Neurology*. 2003; 464:285–311. [PubMed: 12900925]
- Wrase J, Reimold M, Puls I, Kienast T, Heinz A. Serotonergic dysfunction: brain imaging and behavioral correlates. *Cognitive and Affective Behavioral Neuroscience*. 2006; 6:53–61.
- Xuei X, Dick D, Flury-Wetherill L, Tian HJ, Agrawal A, Bierut L, ... Edenberg HJ, et al. Association of the kappa-opioid system with alcohol dependence. *Molecular Psychiatry*. 2006; 11:1016–1024. [PubMed: 16924269]
- Yamada M, Basile AS, Fedorova I, Zhang W, Duttaroy A, Cui Y, ... Wess J, et al. Novel insights into M5 muscarinic acetylcholine receptor function by the use of gene targeting technology. *Life Sciences*. 2003; 74:345–353. [PubMed: 14607263]
- Yan Z, Surmeier DJ. Muscarinic (m2/m4) receptors reduce N- and P-type Ca²⁺ currents in rat neostriatal cholinergic interneurons through a fast, membrane- delimited, G-protein pathway. *Journal of Neuroscience*. 1996; 16:2592–2604. [PubMed: 8786435]
- Yang ARST, Yi HS, Warnock KT, Mamczarz J, June HL Jr, Mallick N, ... June HL Sr, et al. Effects of the triple monoamine uptake inhibitor DOV 102,677 on alcohol-motivated responding and antidepressant activity in alcohol-preferring (P) rats. *Alcoholism: Clinical and Experimental Research*. 2012; 36:863–873.
- Yong W, Spence JP, Eskay R, Fitz SD, Damadzic R, Lai D, ... Liang T, et al. Alcohol-preferring rats show decreased corticotropin-releasing hormone-2 receptor expression and differences in HPA activation compared to alcohol-nonpreferring rats. *Alcoholism: Clinical and Experimental Research*. 2014; 38:1275–1283.
- Yoshimoto K, McBride WJ. Regulation of nucleus accumbens dopamine release by the dorsal raphe nucleus in the rat. *Neurochemistry Research*. 1992; 17:401–407.
- Zeitler JM. Control of sleep and wakefulness in health and disease. *Progress in Molecular Biology and Translational Sciences*. 2013; 119:137–154.
- Zhang W, et al. Characterization of central inhibitory muscarinic autoreceptors by the use of muscarinic acetylcholine receptor knock-out mice. *Journal of Neurosciences*. 2002; 22:1709–1717.
- Zhou FC, Bledsoe S, Lumeng L, Li T-K. Immunostained serotonergic fibers are decreased in selected brain regions of alcohol-preferring rats. *Alcohol*. 1991a; 8:425–431. [PubMed: 1723600]
- Zhou FC, Bledsoe S, Lumeng L, Li T-K. Serotonergic immunostained terminal fibers are lower in selected forebrain regions of alcohol-preferring rats. *Alcohol*. 1991b; 8:1–7. [PubMed: 2006978]
- Zhou FC, McKinzie DL, Patel TD, Lumeng L, Li T-K. Additive reduction of alcohol drinking by 5-HT_{1A} antagonist WAY 100635 and serotonin uptake blocker fluoxetine in alcohol-preferring P rats. *Alcoholism: Clinical and Experimental Research*. 1998; 22:266–269.
- Zhou FC, Zhang JK, Lumeng L, Li T-K. Mesolimbic dopamine system in alcohol-preferring rats. *Alcohol*. 1995; 12:403–412. [PubMed: 8519434]
- Zhou Z, Karlsson C, Liang T, Xiong W, Kimura M, Tapocik JD, ... Goldman D, et al. Loss of metabotropic glutamate receptor 2 escalates alcohol consumption. *Proceedings of the National Academy of Sciences, USA*. 2013; 110:16963–16968.
- Zlojutro M, Manz N, Rangaswamy M, Xuei X, Flury-Wetherill L, Koller D, Almasy L, et al. Genome-wide association study of theta band event-related oscillations identifies serotonin receptor gene HTR7 influencing risk of alcohol dependence. *American Journal of Medical Genetics Part B*. 2010; 156:44–58.
- Zucker, RA. Alcohol and addictive behaviors. In: Rivers, PC., editor. *Nebraska Symposium on Motivation*. Vol. 34. University of Nebraska Press; Lincoln, NE: 1987. p. 27–84.
- Zucker, RA. Pathways to alcohol problems and alcoholism: A developmental account of the evidence for multiple alcoholisms and for contextual contributions to risk. In: Zucker, RA.; Zucker; Boyd, GM.; Howard, J., editors. *The Development of Alcohol Problems: Exploring the Biopsychosocial*

- Matrix of Risk. Rockville, MD: U.S. Dept. of Health and Human Services, National Institute on Alcohol Abuse and Alcoholism; 1995. p. 255-289. NIAAA Research Monograph 26
- Zuo L, Lu L, Tan Y, Pan X, Cai Y, Wang X, ... Luo X, et al. Genome-wide association discoveries of alcohol dependence. *American Journal of Addiction*. 2014; 23:526–539.
- Zuo L, Zhang XY, Wang F, Li CS, Lu L, Ye L, et al. Genome-wide significant association signals in IPO11-HTR1A region specific for alcohol and nicotine codependence. *Alcoholism: Clinical and Experimental Research*. 2013; 37:730–739.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Figure 1.

Innate differences in gene expression between P and NP rats, ethanol-drinking induced changes in gene expression of P rats as well as pharmacological changes in cholinergic activity and/or effects. AcbSh = nucleus accumbens shell; Amyg = amygdala; CeA = central amygdala; Ctx = cortex; CP = caudate-putamen; DB = diagonal band of Broca; DMTN = dorsal medial thalamic nucleus; DRN = dorsal raphe nucleus; FC = frontal cortex; Hipp = hippocampus; Hyp = hypothalamus; LC = locus coeruleus; LDT = lateral dorsal tegmentum; LHab = lateral habenula; MS = medial septum; NAc = nucleus accumbens; NB = nucleus basalis; OB = olfactory bulb; OT = olfactory tubercle; PAG = periaqueductal grey; PFC = prefrontal cortex; PPT = pedunculopontine tegmentum; SC = superior colliculus; SN = substantia nigra; VP = ventral pallidum; VTA = ventral tegmental area. Dark arrows indicate cholinergic projections. Up-arrow indicates an increase, whereas down-arrow indicates a decrease in expression levels, activity or consumption. Short filled arrow = interneuron.

Figure 2.

Innate differences in gene (*italics*) and/or protein expression between P and NP rats, ethanol-drinking induced changes in gene expression of P rats as well as pharmacological changes in dopaminergic activity and/or effects. Arc = arcuate nucleus of the hypothalamus; BLA = basolateral amygdala; BNST = bednucleus of the stria terminalis; ICSA = intracranial self-administration; IA = intercalated amygdala; LA = lateral amygdala; LH = lateral hypothalamus; NAcbCo = nucleus accumbens core; Pit = pituitary; pVTA = posterior ventral tegmental area. For other abbreviations see Figure 1 legend. Dark arrows indicate dopaminergic projections. Up-arrow indicates an increase, whereas a down-arrow indicates a decrease in expression levels, activity or consumption/self-administration. *, indicates multiple studies.

Figure 3.

Innate differences in gene (*italics*) and/or protein expression between P and NP rats, ethanol-drinking induced changes in gene expression of P rats as well as pharmacological changes in GABAergic activity and/or effects. BDZ = benzodiazepine; Cg = cingulate cortex; GP = globus pallidus; HAB = habenula; mTEG = medial tegmentum; SEPT = septum; THAL = thalamus; wAmyg = whole amygdala; wNAcb = whole nucleus accumbens. For other abbreviations see Figures 1 and 2 legend. Dark arrows indicate GABAergic projections. Up-arrow indicates an increase, whereas a down-arrow indicates a decrease in expression levels, activity or consumption/self-administration. Short filled arrow = interneuron.

Figure 4. Innate differences in gene (*Italics*) and/or protein (Upper Case) expression between P and NP rats, ethanol-drinking induced changes in gene expression of P rats as well as pharmacological changes in glutamatergic activity and/or effects. WD = withdrawal. For other abbreviations see Figures 1, 2 and 3 legends. Dark arrows indicate glutamatergic projections. Up-arrow indicates an increase, whereas a down-arrow indicates a decrease in expression levels, activity or consumption/self-administration.

Figure 5.

Innate differences in gene (*italics*) and/or protein expression between P and NP rats, ethanol-drinking induced changes in gene expression of P rats as well as pharmacological changes in serotonergic activity and/or effects. mAcB = medial nucleus accumbens; LA = lateral amygdala; PC = pyriform cortex cortex; RN = raphe nuclei; TC = temporal cortex; vHipp = ventral hippocampus. For other abbreviations see Figures 1, 2, 3 and 4 legends. Dark arrows indicate serotonergic projections. Up-arrow indicates an increase, whereas a down-arrow indicates a decrease in expression levels, activity or consumption/self-administration.

Figure 6. Innate differences in gene (*italics*) and/or protein expression between P and NP rats, ethanol-drinking induced changes in gene expression of P rats as well as pharmacological changes in noradrenergic and corticotrophic releasing factor-system activity and/or effects. ACTH = adrenal corticotrophic hormone; MC4R = melanocortin-4 receptor. For other abbreviations see Figures 1, 2, 3, 4 and 5 legends. Dark, solid arrows indicate noradrenergic projections. Dark, thick, dashed arrows indicate CRF projections. Dark, thin, dashed arrows point to nuclei with glucocorticoid binding sites and activity. Up-arrow indicates an increase, whereas a down-arrow indicates a decrease in expression levels, activity or consumption/self-administration.

Figure 7.

Innate differences in gene (first letter upper case others lower case and in italics) and/or protein expression between P and NP rats, ethanol-drinking induced changes in gene expression of P rats as well as pharmacological changes in opioid-system activity and/or effects. DOR = delta opioid receptor; KOR = kappa opioid receptor; MOR = mu opioid receptor; Pdyn = prodynorphin; Penk = proenkephalin; POMC = proopiomelanocortin. For other abbreviations see Figures 1, 2, 3, 4, 5 and 6 legends. When multiple opioid precursors (PDYN, PENK or POMC) and/or opioid receptors (DOR, KOR or MOR) are localized in the brain region, the order of density are presented from highest (top) to lowest (bottom). Dark, solid arrows indicate opioid projections. Short light-shaded arrows represent opioid containing interneurons. Up-arrow indicates an increase, whereas a down-arrow indicates a decrease in expression levels, activity or consumption/self-administration.

Figure 8.

Innate differences in gene (*italics*) and/or protein expression between P and NP rats, ethanol-drinking induced changes in gene expression of P rats as well as pharmacological changes in the neuropeptide Y (NPY)-system activity and/or effects. ARC = activity-regulated cytoskeleton-associated protein; BDNF = brain derived neurotrophic factor. For abbreviations see Figures 1, 2, 3, 4, 5, 6 and 7 legends. Dark, solid arrows indicate NPY projections. It is important to remember that NPY is also synthesized locally in the brain. Up-arrow indicates an increase, whereas a down-arrow indicates a decrease in expression levels, activity or consumption/self-administration.