Acknowledgements

We thank the clinical research coordinators of the study management centre (see Supplementary material online, Appendix B) for their invaluable help in checking and correcting the enormous data.

Funding

This study was supported by the Research Institute for Production Development in Kyoto, Japan.

Conflict of interest: none declared.

References

- Cappato R, Calkins H, Chen SA, Davies W, Iesaka Y, Kalman J, Kim YH, Klein G, Natale A, Packer D, Skanes A, Ambrogi F, Biganzoli E. Updated worldwide survey on the methods, efficacy, and safety of catheter ablation for human atrial fibrillation. *Circ Arrhythm Electrophysiol* 2010;**3**:32–38.
- Oral H, Knight BP, Tada H, Ozaydin M, Chugh A, Hassan S, Scharf C, Lai SW, Greenstein R, Pelosi F Jr, Strickberger SA, Morady F. Pulmonary vein isolation for paroxysmal and persistent atrial fibrillation. *Circulation* 2002;**105**:1077–1081.
- Takahashi A, Iesaka Y, Takahashi Y, Takahashi R, Kobayashi K, Takagi K, Kuboyama O, Nishimori T, Takei H, Amemiya H, Fujiwara H, Hiraoka M. Electrical connections between pulmonary veins: implication for ostial ablation of pulmonary veins in patients with paroxysmal atrial fibrillation. *Circulation* 2002;**105**:2998–3003.
- Oral H, Chugh A, Yoshida K, Sarrazin JF, Kuhne M, Crawford T, Chalfoun N, Wells D, Boonyapisit W, Veerareddy S, Billakanty S, Wong WS, Good E, Jongnarangsin K, Pelosi F Jr, Bogun F, Morady F. A randomized assessment of the incremental role of ablation of complex fractionated atrial electrograms after antral pulmonary vein isolation for long-lasting persistent atrial fibrillation. *J Am Coll Cardiol* 2009;**53**:782–789.
- Haïssaguerre M, Jais P, Shah DC, Takahashi A, Hocini M, Quiniou G, Garrigue S, Le Mouroux A, Le Métayer P, Clémenty J. Spontaneous initiation of atrial fibrillation by ectopic beats originating in the pulmonary veins. *N Engl J Med* 1998;**339**:659–666.
- Arentz T, Macle L, Kalusche D, Hocini M, Jais P, Shah D, Haïssaguerre M. 'Dormant' pulmonary vein conduction revealed by adenosine after ostial radiofrequency catheter ablation. *J Cardiovasc Electrophysiol* 2004;**15**:1041–1047.
- Tritto M, De Ponti R, Salerno-Urriarte JA, Spadacini G, Marazzi R, Moretti P, Lanzotti M. Adenosine restores atrio-venous conduction after apparently successful ostial isolation of the pulmonary veins. *Eur Heart J* 2004;**25**:2155–2163.
- Hachiya H, Hirao K, Takahashi A, Nagata Y, Suzuki K, Maeda S, Sasaki T, Kawabata M, Isebe M, Iesaka Y. Clinical implications of reconnection between the left atrium and isolated pulmonary veins provoked by adenosine triphosphate after extensive encircling pulmonary vein isolation. *J Cardiovasc Electrophysiol* 2007;**18**:392–398.
- Matsuo S, Yamane T, Date T, Inada K, Kanzaki Y, Tokuda M, Shibayama K, Miyazaki S, Miyazaki H, Sugimoto K, Mochizuki S. Reduction of AF recurrence after pulmonary vein isolation by eliminating ATP-induced transient venous reconnection. *J Cardiovasc Electrophysiol* 2007;**18**:704–708.
- Kumagai K, Naito S, Nakamura K, Hayashi T, Fukazawa R, Sato C, Takemura N, Miki Y, Fuke E, Tanaka Y, Hori Y, Goto K, Iwamoto J, Aonuma K, Oshima S, Taniguchi K. ATP-induced dormant pulmonary veins originating from the carina region after circumferential pulmonary vein isolation of atrial fibrillation. *J Cardiovasc Electrophysiol* 2010;**21**:494–500.
- Macle L, Khairy P, Weerasooriya R, Novak P, Verma A, Willems S, Arentz T, Deisenhofer I, Veenhuyzen G, Scavée C, Jais P, Puererfellner H, Levesque S, Andrade JG, Rivard L, Guerra PG, Dubuc M, Thibault B, Talajic M, Roy D, Nattel S, ADVANCE trial investigators. Adenosine-guided pulmonary vein isolation for the treatment of paroxysmal atrial fibrillation: an international, multicentre, randomised superiority trial. *Lancet* 2015;**9994**:672–679.
- Ouyang F, Bänsch D, Ernst S, Schaumann A, Hachiya H, Chen M, Chun J, Falk P, Khanedani A, Antz M, Kuck KH. Complete isolation of left atrium surrounding the pulmonary veins: new insights from the double-Lasso technique in paroxysmal atrial fibrillation. *Circulation* 2004;**110**:2090–2096.
- Yamane T, Matsuo S, Date T, Lellouche N, Hioki M, Narui R, Ito K, Tanigawa S, Yamashita S, Tokuda M, Yoshida H, Inada K, Shibayama K, Miyazaki S, Miyazaki H, Abe K, Sugimoto K, Yoshimura M. Repeated provocation of time- and ATP-induced early pulmonary vein reconnections after PVI: eliminating paroxysmal AF in a single procedure. *Circ Arrhythm Electrophysiol* 2011;**4**:601–608.
- le Polain de Waroux JB, Weerasooriya R, Anvardeen K, Barbraud C, Marchandise S, De Meester C, Goesaert C, Reis I, Scavée C. Low contact force and force-time integral predict early recovery and dormant conduction revealed by adenosine after pulmonary vein isolation. *Europace* 2015;**17**:877–883.
- Andrade JG, Monir G, Pollak SJ, Khairy P, Dubuc M, Roy D, Talajic M, Deyell M, Rivard L, Thibault B, Guerra PG, Nattel S, Macle L. Pulmonary vein isolation using 'contact force' ablation: the effect on dormant conduction and long-term freedom from recurrent atrial fibrillation—a prospective study. *Heart Rhythm* 2014;**11**:1919–1924.
- Miyazaki S, Taniguchi H, Nakamura H, Hachiya H, Ichihara N, Araki M, Kuroi A, Takagi T, Iwasawa J, Hirao K, Iesaka Y. Adenosine triphosphate test after cryothermal pulmonary vein isolation: creating contiguous lesions is essential for eliminating dormant conduction. *J Cardiovasc Electrophysiol* 2015 [Epub ahead of print].

Corrigendum

doi:10.1093/eurheartj/ehv247

Online publish-ahead-of-print 10 June 2015

Corrigendum to: Incremental prognostic utility of coronary CT angiography for asymptomatic patients based upon extent and severity of coronary artery calcium: results from the COronary CT Angiography EvaluationN For Clinical Outcomes InteRnationalMulticenter (CONFIRM) Study [Eur Heart J (2015) 36 (8): 501–508; doi:10.1093/eurheartj/ehv358].

The funding information was incorrectly supplied and should be updated as follows. The Publisher apologises for this mistake. This has also been updated online:

NRF/MSIP This research was supported by Leading Foreign Research Institute Recruitment Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Science, ICT & Future Planning (MSIP) (2012027176)

Published on behalf of the European Society of Cardiology. All rights reserved. © The Author 2015. For permissions please email: journals.permissions@oup.com.