Plasma F₂-isoprostane class VI isomers at 12–18 weeks of pregnancy are associated with later occurrence of preeclampsia

Jean-François Bilodeau^{a,f,*}, Shu Qin Wei^d, Jessica Larose^a, Karine Greffard^{b,c}, Vanessa Moisan^a, Francois Audibert^d, William D. Fraser^d, and Pierre Julien^{b,c,e}

^aAxe reproduction, santé de la mère et de l'enfant, CRCHU de Québec, Québec, Canada

^bAxe endocrinologie et néphrologie, CRCHU de Québec, Québec, Canada

^cCentre de Recherche en endocrinologie moléculaire et oncologique et en génomique humaine (CREMOGH), Université Laval, Québec, Canada

^dDepartment of Obstetrics and Gynecology, CRCHU Sainte-Justine and University of Montreal, Montreal, Quebec, Canada

^eDépartement de Médecine, Faculté de médecine, Université Laval, Québec, Canada

^fDépartement d'Obstétrique, Gynécologie et Reproduction, Faculté de médecine, Université Laval, Québec, Canada

Abstract

Preeclampsia (PE) has long been associated with early oxidative stress whereas the symptoms occur later in pregnancy. We have hypothesized that the oxidative stress in PE, as characterized by the presence of F₂-isoprostanes (F₂-isoPs) isomers in late pregnancy, should already be present in plasma at the first regular visit of the obstetrical follow-up. There are 64 possible isomers of F₂-isoPs derived from the oxidation of arachidonic acid (AA) but only one of these isomers has been investigated so far in PE, the classical 8-iso-PGF_{2α}. Here, we have investigated two regioisomers of class III (8-iso-15(R)-PGF_{2α} and 8-iso-PGF_{2α}) and a mix of two isomers of class VI (±)5-iPF_{2α}-VI in plasma samples collected prospectively at 12–18 weeks from normotensive controls (n=60) and pregnant mothers who developed PE later in pregnancy (n=33). The plasma samples were subjected to alkaline hydrolysis followed by liquid-liquid extraction to extract total F₂-isoPs for later quantification by HPLC-MS/MS. The F₂-isoPs were normalized to either plasma volume or polyunsaturated fatty acids (PUFAs) levels measured by GC-FID in plasma phospholipids. Early in pregnancy, only the class VI F₂-isoPs isomers were found in concentrations significantly higher in women developing PE later in pregnancy (+ 13%; p=0.0074). Normalisation of F₂-isoPs to their substrate, AA, or the omega-3 to omega-6 ratio improved the predictability of PE as determined by receiver operating characteristic (from area under the curve of 0.67 to 0.68 and 0.70 respectively). Interestingly, omega-3 fatty acids were 25% higher in the control group than the PE group (P = 0.0225). Total omega-6 PUFAs correlated to F₂-Isop isomers only in cases of PE (r >

*Corresponding author: Centre de recherche du CHU de Québec, Axe reproduction, santé de la mère et de l'enfant, 2705, boulevard Laurier, local T3-67, Québec (Qc), Canada G1V 4G2, Tel: 1 (418) 525-4444 x 46153, Fax: 1 (418) 654-2765, jean-francois.bilodeau@crchudequebec.ulaval.ca.

Conflict of interest

Related to patent WO 2013-170369 A1, Early predictive markers of pre-eclampsia.

0.377; $P > 0.03$, Spearman correlation). In sum, this study indicates that specific isomers of class VI are significant predictors of PE. This work also suggests that F₂-isoPs isomers are not all generated and eliminated at the same extent and are influenced by the PUFA composition of plasma phospholipids.

Keywords

oxidative stress; pregnancy; mass spectrometry; isoprostanes; polyunsaturated fatty acids; pre-eclampsia

Introduction

The classical 8-iso PGF_{2α}, an F₂-isoprostanes (F₂-isoPs), is currently the most stable and reliable indicators of oxidative stress and lipid peroxidation [1–3]. F₂-isoPs are prostaglandin (PG) F₂-like isomers, generated by the spontaneous oxidation of arachidonic acid (AA), an *omega*-6 polyunsaturated fatty acids (PUFAs), contained in various phospholipid membranes [4]. F₂-isoPs are carried by lipoproteins, mainly HDL, in the blood circulation [5]. There are 64 possible isomers of F₂-isoPs classified in 4 classes of regioisomers (III, IV, V, VI) that comprises 16 members each [6, 7]. The most studied isomer is the 8-iso PGF_{2α} (or 8-epi-PGF_{2α}, iPF_{2α}-III). The 8-iso-PGF_{2α} is a vasoconstrictor similar in action to thromboxane (TX) A₂, when released from membranes by the action of phospholipases A₂ [8, 9]. The other 63 F₂-IsoP isomers are much less (or for most, not at all) investigated in pregnancy. Indeed, we pioneered the detection of the abundant class VI isomers and reported the first basal values in uncomplicated pregnancies by HPLC-MS/MS [2, 10].

Uncomplicated pregnancy is associated with a mild physiological increase in oxidative stress to cope with placental and fetal development throughout pregnancy [1]. Indeed, markers of oxidative stress, such as F₂-isoPs, reach their highest level between 37 and 41, just before birth [11]. Pathological pregnancies, such as preeclampsia (PE), expressed an even higher oxidative stress response especially in severe cases, as demonstrated using free 8-iso-PGF_{2α} as plasma biomarker [12].

Only few studies have looked prospectively at the potential link between PE and oxidative stress markers, such as F₂-isoPs. One report showed a urinary increase of the 8-iso-PGF_{2α} as early as the 15th week of gestation in women who will later develop PE (5-fold higher risk) compared to controls [11]. Accordingly, another study reported 8-iso-PGF_{2α} levels at 1.7-fold higher in the amniotic fluid of PE than normotensive pregnancies at an average of 17.5 weeks of gestation [13]. Notably, only one isomer was investigated in these studies. We believe that F₂-isoP isomers are not all generated and eliminated to the same extent. We hypothesized that certain F₂-isoPs isomers could be more specific to PE than the classical 8-iso-PGF_{2α}. Therefore, we propose that the concomitant assessment of several F₂-isoPs by HPLC-MS/MS should provide more knowledge on the specific oxidative stress appearing in PE. We also theorized that the plasma fatty acid composition should also affect the generation of specific F₂-isoPs or compete for the generation of other oxidation products. The objective of this study was thus to quantify two isomers of class III (8-iso-15(R)-PGF_{2α}

and 8-iso-PGF_{2α}) and a mix of two isomers of class VI, (±5)-iPF_{2α}-VI (or iPF_{2α}-VI + 5-iPF_{2α}-VI), in plasma samples collected prospectively at 12–18 weeks from normotensive controls (n=60) and women that will later develop PE (n=33). Concurrently, we quantified *omega-3* and *omega-6* fatty acids in order to correlate the levels of PUFAs with those of F₂-isoPs.

Materials and methods

Materials

The F₂-isoPs isomers, 8-iso-15(R)-PGF_{2α}, 8-iso-PGF_{2α}, (±5)-iPF_{2α}-VI (or iPF_{2α}-VI + 5-iPF_{2α}-VI), and their corresponding deuterated standards; 8-iso-PGF_{2α}-d4 and (±5)-iPF_{2α}-VI-d11 were purchased from Cayman Chemical (Ann Arbor, MI, USA). Butylated hydroxytoluene (BHT) was bought from Sigma-Aldrich (Oakville, ON, Canada). The standards for fatty acid profile analysis, FAME 37 mix, 18:2 cis/trans mix 18:3 cis/trans mix, 22:5, 9*t*-14:1, 9*t*-C16:1 and several 18:1 fatty acids (6*c*-, 11*c*-, 12*c*-, 13*c*-, 6*t*-, 11*t*-18:1) were all purchased from Supelco Inc (Bellefonte, PA, USA). A mixture of 31 fatty acids (GLC-411) was obtained from NuChek Prep Inc (Elysian, MN, USA). All other reagents and solvents were HPLC grade and were purchased from VWR International (Ville Mont-Royal, QC, Canada).

Patient's characteristics and sample processing

Patients' clinical data and biological samples were obtained from a biobank, the Maternal and Infant Research on Oxidative Stress (MIROS cohort) described previously [14]. Plasma from sixty women with uncomplicated pregnancies were recruited prospectively early between the 12th to the 18th weeks of pregnancy from a subset of the 665 possible uncomplicated pregnancies recruited by the MIROS study group in 17 Canadian centers over a 26-month period. Similarly 33 PE were obtained from the same biobank. The respective institutions approved the protocol and informed consents were obtained from all pregnant women. PE was defined as gestational hypertension with proteinuria. Proteinuria was defined as a urine protein dipstick test 2+, or the urinary excretion of 0.3 g in a 24-hour urine collection [15]. Severe PE was defined according to the criteria of the Canadian Hypertension Society Consensus Conference Report [15]. Early on-set PE was defined as delivery before 34 weeks of gestational age [16]. Venous blood was collected into EDTA tubes and plasma samples were immediately separated by centrifugation at 500 × *g* for 10 minutes at 4°C. Centrifuged plasma samples were rapidly frozen and stored at –80°C [14]. Butylated hydroxytoluene (BHT) was added (0.1%) before aliquoting [10].

F₂-isoPs analysis by HPLC-MS/MS

Total F₂-isoPs (free + esterified) were extracted from plasma after alkaline hydrolysis using a liquid-liquid extraction as described in details previously [10]. Briefly, the chromatography was carried out with a Shimadzu Prominence system (Columbia, MD, USA) linked to a 3200 QTRAP® LC/MS/MS system from AB Sciex (Concord, ON, Canada) operated in the negative mode. A Shim-Pack XR-ODS column (3.0 × 100 mm, 2.2 μm) from Shimadzu was used. The column temperature was controlled at 30°C and the injection volume was 40 μL. The chromatographic separation was done using a gradient of three solvents at 0.4 mL/min

as described earlier (method II [2]). The mobile phase A was 0.01% acetic acid in water, B was 0.01% acetic acid in acetonitrile and C was 0.01% acetic acid in methanol. The F₂-isoPs and their corresponding internal standards were monitored in multiple-reaction monitoring (MRM) mode at the transitions 353.3 → 193.2 and 357.3 → 197.2 m/z respectively for class III, and at the transitions 353.0 → 115.0 and 364.6 → 115.0 m/z respectively for class VI. Quantification of F₂-isoPs was performed using the Analyst® 1.4.2 Software (AB Sciex) [10].

Determination of plasma fatty acid profile by gas chromatography

Plasma fatty acids were isolated according to a method previously described [17]. Briefly, a solution of chloroform:methanol (2:1, by volume) was used to extract lipids from plasma. Then, phospholipids were separated by thin layer chromatography using a mix of isopropyl ether:acetic acid (96:4) as eluent and phospholipid fatty acids were methylated following a transesterification reaction using a mix of methanol:benzene (4:1) and acetyl chloride at 95°C for 1.5 hour. Methylated fatty acids were finally analysed by gas chromatography coupled with a flame ionisation detector (GC-FID) as explained elsewhere [2, 17]. The peroxidation index (PI) and the unsaturation index (UI) were calculated as described elsewhere [18].

Statistical analyses

Statistical analyses were performed with GraphPad Prism 6.0f (GraphPad Software Inc., 2013, San Diego, USA). Normality was verified using the D'Agostino & Pearson omnibus test. Students t-test was used to compare the groups. Since normality was not achieved for most of the data, the Mann-Whitney test was used to compare levels of both, F₂-isoPs and fatty acids. The Spearman's rank correlation coefficient was used to study relationship between variables. The predictive nature of F₂-isoPs was determined using the receiver operating characteristic curve analysis. A *p*-value lower than 0.05 (two-tailed) was considered statistically significant.

Results

Clinical features of the cohort used (biobank)

The clinical characteristics of the pregnant's cohort used in this study was published in details previously [14]. Briefly, the PE group was constituted of 33 pregnancies in which, 21 pregnancies were obtained from a predetermined high-risk group according to one or many of the following risk factors; chronic hypertension, prepregnancy diabetes, multiple pregnancy, or a history of PE. The mean gestational age at delivery was 36± 3 weeks (mean ± SD). Of note, 22 pregnancies qualified as severe and 11 pregnancies are classified as early on-set PE (< 34 weeks at delivery) [16]. Attempt to stratify according to the two latter factors did not yield significance in our statistical analysis (results not shown). The PE group was matched with a control group of 60 pregnancies in order to avoid any statistical differences for maternal age, education, ethnic origin, marital status, prenatal vitamin supplementation, smoking or season when blood was drawn. The pre-pregnancy body mass index (BMI) and weight gain were not statistically different between the control and the PE

groups (results not shown). Only gestational age at delivery (term pregnancy) was statistically different ($P < 0.05$).

F₂-isoPs in the plasma of women that will later develop or not PE

Table 1 shows the F₂-isoPs levels of class III and VI isomers measured by HPLC-MS/MS using maternal plasma at 12–18 weeks of gestation obtained from normotensive controls and mothers who later developed PE. The data were normalized to plasma volume or AA content, the substrate generating F₂-isoPs. Only class VI isomers, iPF_{2α}-VI + 5-iPF_{2α}-VI or (±)5-iPF_{2α}-VI were 13% and 19% higher in the PE group compared to controls when normalized with plasma volume or AA content ($P = 0.0074$ and $P = 0.0040$ respectively). These class VI isomers are a mix of two isomers and are sold as one standard for the purpose of quantification: (±)5-iPF_{2α}-VI. Both class III F₂-isoP isomers, 8-iso-15(R)-PGF_{2α} and 8-iso-PGF_{2α}, were not significantly different between groups. The sums of the class III isomers are 222 ± 18 and 250 ± 27 pg/ml for normotensives and PE respectively (mean $\pm$ SEM; $P = 0.36$). The sums of all F₂-isoP isomers investigated here are 371 ± 20 and 428 ± 33 pg/ml respectively for the normotensive and PE groups ($P = 0.12$).

Plasma PUFAs in pregnancy

Table 2 is reporting the levels of PUFAs in plasma phospholipids at 12–18 weeks of gestation in normotensive subjects and mothers who will later develop PE. The levels of total *omega-6* PUFAs, comprising AA, the substrate of F₂-isoPs, are not different between groups ($P > 0.68$). However, the level of *omega-3* fatty acids was significantly lower in the group that will develop PE compared to normotensive controls (81 ± 20 vs 71 ± 19 µg/ml; $P = 0.0225$). This last difference affects indirectly the *omega-3* to *omega-6* ratio and the calculated PI and UI indexes ($P < 0.041$).

Correlation between F₂-isoPs and PUFAs

Spearman's rank correlation coefficients were used to study at 12–18 weeks of pregnancies, the associations between F₂-isoPs and plasma phospholipid PUFAs in normotensive mothers and the group that will develop PE group (Table 3). Interestingly, all F₂-isoPs isomers correlated with either, AA or total *omega-6* PUFAs in PE ($r > 0.377$, $P > 0.0031$), but this phenomenon was not observed in the normotensive controls. The class VI isomers were better correlated than class III isomers in the group that will later develop PE, though ($r > 0.531$, $P > 0.001$). In contrast, the control group was not correlated to *omega-6* PUFAs (Table 3). In normotensive pregnancies, total *omega-3* PUFAs were negatively correlated with the class VI isomers whereas the *omega-3* to *omega-6* ratio correlated with both the classical 8-iso-PGF_{2α} and the class VI isomers. Of note, F₂-isoPs were not correlated to UI and PI indexes for all study groups.

F₂-isoPs and PE prediction

A receiver operating characteristic (ROC) curve analysis was performed in order to determine how predictive were class VI F₂-isoPs for PE (Figure 1). The iPF_{2α}-VI + 5-iPF_{2α}-VI used alone yielded to an area under the curve (AUC) of 0.67 ± 0.06 (mean $\pm$ SEM; $P = 0.0079$). The normalization to AA and the *omega-3* to *omega-6* ratio improved slightly the

test toward a fair prediction, the AUC being 0.68 ± 0.06 ($P = 0.0044$) and 0.70 ± 0.06 respectively.

Discussion

We reported relatively low levels of total F₂-isoPs (free + esterified) between 105 µg/ml for class III isomers to 152 µg/ml for class VI at 12–18 weeks in normotensive pregnancies. However, these levels are in the upper range of reported levels for the classical 8-iso-PGF_{2α} from healthy non-pregnant individuals, which were between 40–170 pg/mL in plasma [19, 20]. Indeed, an uneventful pregnancy is considered a mild oxidative stress [21], which may explained these slightly higher levels. Compared to levels determined early in pregnancy (12–18 weeks) as shown in the current study, our previous data obtained from two independent cohorts suggest a 1.4-fold higher level for 8-iso-PGF_{2α} levels and a 2.4-fold higher level for iPF_{2α}-VI + 5-iPF_{2α}-VI at the end of pregnancy [2]. These findings are in accordance with study showing that plasma 8-iso-PGF_{2α} levels are relatively low early in pregnancy with a significant increase of approximately two-fold from 15–20 weeks to 37–41 weeks of pregnancy [52].

Only class VI F₂-isoP isomers were significantly higher in the group developing PE compared to normotensive controls. To our knowledge, this is the first prospective report of plasma isoPs in pregnant mothers developing PE later in pregnancy. These data reveal a statistically significant increase, even though modest (13%), between the control normotensive group and mothers developing PE later in pregnancy, suggesting a slight maternal systemic increase of oxidative stress already detectable in the early stage of undiagnosed PE. The normalization of the data using substrate concentration (AA) in phospholipids did not alter this conclusion, but shows instead a 19% increase. The source of this increased class VI F₂-isoPs early in maternal blood is still undetermined, but might originate from the abnormally developed placenta rather than the maternal endothelial dysfunction at this stage of the pathology [1, 22]. This current discovery would not have been possible without the simultaneous analysis of several isomers of F₂-isoPs by HPLC-MS/MS. The current study confirms that specific F₂-isoP isomers may respond differently in various pathologies. Indeed, more than eight F₂-isoP isomers were shown to be increased in urine of hypercholesterolemic patients, whereas only three isomers, including the classical 8-iso-PGF_{2α} and the 8-iso-15(R)-PGF_{2α}, were found to be increased in cases of congestive heart failure [23].

According to the present data, the prominence of class III F₂-isoP isomers, like 8-iso-PGF_{2α}, in the early stage of PE appears to be much less striking than anticipated based on numerous previous studies reporting class III F₂-isoP isomers as accurate indicators of oxidative stress in many pathologies [1–3]. Furthermore, class VI isomers are rarely studied and we are pioneering their measurements in pregnancy [2, 10]. Other prospective studies on PE have looked at concentrations of urinary and amniotic fluid 8-iso-PGF_{2α} using enzyme immunoassays. Interestingly, elevated urinary excretion of 8-iso-PGF_{2α} at 15 weeks was associated with a 5-fold higher risk of PE in a cohort of 307 pregnant women. However, 24-hour urinary excretion of 8-iso-PGF_{2α} cannot be directly compared to plasma values of total F₂-isoPs (free + esterified) measured at a given time point. Indeed, the difference with our

study could be related to the increased elimination of F₂-isoPs in urine rather than their total concentrations at any given time in blood plasma [1].

In another prospective study, levels of amniotic fluid 8-iso-PGF_{2α} were 1.74-fold higher in women known to later develop PE compared to controls. The PE and control groups were both constituted of 85 women. The area under curve (AUC) of the ROC curve indicated 0.81 for the prediction of PE [13]. However, there are at least three major differences in this report compared to our study: biological samples were, in the average, obtained 3 weeks later, secondly biological specimen were collected from a very different ethnic population (Caucasian vs Asian), and thirdly the collection of amniotic fluid at the very proximity of the source of oxidative stress, the placenta. These may explain the more pronounced difference between PE and the control groups using a less specific marker for PE such as the 8-iso-PGF_{2α}.

F₂-isoPs originates from the oxidation of AA, an *omega*-6 PUFAs. However, we did not observe any difference for either *omega*-6 or AA concentrations between the study groups. This could suggest that substrate availability was not an issue for F₂-isoPs formation. Interestingly, *omega*-3 level was significantly lower in the group later developing PE compared to controls. It is believed that *omega*-3 fatty acids are more susceptible to the oxidation than *omega*-6 fatty acids. Of note, oxidation of *omega*-3 PUFAs leads, among others, to the formation F₃-isoPs and F₄-isoPs (neuroprostanes) derived from eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA) respectively [1]. This suggests that higher *omega*-3 level in the control group could somehow protect against the oxidation of *omega*-6 fatty acids, explaining the lower concentrations of AA derived F₂-isoPs in the control group. Indirect evidence of the latter came from nutritional studies. In kidney transplant patients, supplementation with fish oil rich *omega*-3 fatty acids for periods of 3 and 6 months caused a significant decrease in the level of plasma 8-iso-PGF_{2α} when compared to the untreated (placebo) group [24]. In a 6-week placebo-controlled trial, 4 g daily of either EPA or DHA in overweight dyslipidemic men caused 24% and 14% decreases in plasma 8-iso-PGF_{2α} [25]. The same *omega*-3 treatment when used in treated-hypertensive type 2 diabetic patients caused reductions in the plasma 8-iso-PGF_{2α} levels of 19% and 23% using EPA or DHA in comparison to the olive oil control group. Interestingly, correction for AA content did not change the conclusion of these two last independent trials [25].

Since the study group developing PE showed lower level of *omega*-3 fatty acids, higher level of *omega*-6 fatty acids and, likely, higher oxidative stress than normotensive controls, we believe that correlation between F₂-isoPs and its substrate AA is achieved under PE conditions. Further, AA constitutes a large proportion (27%) of the *omega*-6 fatty acids that are also correlated with the AA-derived isoPs only in the group known to develop PE later in pregnancy. It is worth noting that oxidation of AA may also yield to the formation of PGD₂/E₂-like (D₂/E₂-IsoPs) and thromboxane-like (isothromboxanes) end-products in competition with the F₂-isoPs metabolites [26]. The protective effect of *omega*-3, as emphasized by their negative correlation with class VI isoPs, is only observed in uneventful pregnancy with mild oxidative stress. This is in accordance with the reported absence of

correlation between AA and F₂-isoPs in healthy individuals followed over a 4-month period [2727].

Conclusions

In summary, this study reveals that isomers of class VI, ($\pm$)5-iPF_{2 α} -VI, are better predictors of PE at 12–18 weeks of gestation than class III isomers such as the classical 8-iso-PGF_{2 α} . This also suggests that F₂-isoPs isomers are not all generated and/or eliminated at the same rate during the pathogenesis of PE. This work also underlines the possible antioxidative role of *omega*-3 PUFAs on the production of AA-derived F₂-IsoPs and the pathogenesis of PE.

Acknowledgments

The authors would like to thank Ms. Line Berthiaume and M. François Cadelis for their help with the extraction of fatty acids. This work was supported by a grant from the Canadian Institutes of Health research (CIHR, grant No. 84219 to J.-F.B and 78879 to W.D.F., P.J. and F.A. for the MIROS cohort). Jessica Larose was a recipient of a Fonds de Recherche en Santé du Québec (FRSQ) studentship award.

Source of support: CIHR, grant No. 111271 and 78879.

Abbreviations

AA	arachidonic acid
AUC	area under the curve
BHT	butylated hydroxytoluene
EPA	eicosapentaenoic acid
DHA	docosahexaenoic acid
F₂-isoP	F ₂ -isoprostane
GC-FID	gas chromatography coupled to a flame ionisation detector
HPLC-MS/MS	high performance liquid chromatography coupled to tandem mass spectrometry
IS	internal standard
PI	peroxidation index
PUFAs	polyunsaturated fatty acids
PG	prostaglandins
PLA₂	phospholipase A ₂
ROC	receiver operating characteristic
ROS	Reactive oxygen species
TX	Thromboxane

UI unsaturation index

References

1. Bilodeau JF. Review: Maternal and placental antioxidant response to preeclampsia - Impact on vasoactive eicosanoids. *Placenta*. 2014; 35S:S32–S38.
2. Larose J, Julien P, Greffard K, Fraser WD, Audibert F, Wei SQ, Bilodeau JF. F2-isoprostanes are correlated with trans fatty acids in the plasma of pregnant women. *Prostaglandins, leukotrienes, and essential fatty acids*. 2014; 91:243–249.
3. Milne GL, Dai Q, Roberts LJ 2nd. The isoprostanes-25 years later. *Biochimica et biophysica acta*. 2014
4. Song WL, Paschos G, Fries S, Reilly MP, Yu Y, Rokach J, Chang CT, Patel P, Lawson JA, Fitzgerald GA. Novel eicosapentaenoic acid-derived F3-isoprostanes as biomarkers of lipid peroxidation. *J Biol Chem*. 2009; 284:23636–23643. [PubMed: 19520854]
5. Proudfoot JM, Barden AE, Loke WM, Croft KD, Puddey IB, Mori TA. HDL is the major lipoprotein carrier of plasma F2-isoprostanes. *J Lipid Res*. 2009; 50:716–722. [PubMed: 19050315]
6. Taber DF, Morrow JD, Roberts LJ 2nd. A nomenclature system for the isoprostanes. *Prostaglandins*. 1997; 53:63–67. [PubMed: 9112285]
7. Rokach J, Khanapure SP, Hwang SW, Adiyaman M, Lawson JA, FitzGerald GA. Nomenclature of isoprostanes: a proposal. *Prostaglandins*. 1997; 54:853–873. [PubMed: 9533181]
8. Bode-Boger SM, Boger RH, Kuhn M, Radermacher J, Frolich JC. Recombinant human erythropoietin enhances vasoconstrictor tone via endothelin-1 and constrictor prostanoids. *Kidney international*. 1996; 50:1255–1261. [PubMed: 8887285]
9. Aristegui PM, Enero MA. Endothelium and Ca²⁺ in the prostaglandin F2 alpha potentiation of vasoconstrictor responses. *Eur J Pharmacol*. 1990; 184:1–7. [PubMed: 2120075]
10. Larose J, Julien P, Bilodeau JF. Analysis of F2-isoprostanes in plasma of pregnant women by HPLC-MS/MS using a column packed with core-shell particles. *J Lipid Res*. 2013; 54:1505–1511. [PubMed: 23431046]
11. Scholl TO, Leskiw M, Chen X, Sims M, Stein TP. Oxidative stress, diet, and the etiology of preeclampsia. *Am J Clin Nutr*. 2005; 81:1390–1396. [PubMed: 15941892]
12. McKinney ET, Shouri R, Hunt RS, Ahokas RA, Sibai BM. Plasma, urinary, and salivary 8-epi-prostaglandin f2alpha levels in normotensive and preeclamptic pregnancies. *Am J Obstet Gynecol*. 2000; 183:874–877. [PubMed: 11035329]
13. Wang CN, Chen JY, Sabu S, Chang YL, Chang SD, Kao CC, Peng HH, Chueh HY, Chao AS, Cheng PJ, Lee YS, Chi LM, Wang TH. Elevated amniotic fluid F(2)-isoprostane: a potential predictive marker for preeclampsia. *Free Radic Biol Med*. 2011; 50:1124–1130. [PubMed: 21277370]
14. Wei SQ, Audibert F, Hidioglou N, Sarafin K, Julien P, Wu Y, Luo ZC, Fraser WD. Longitudinal vitamin D status in pregnancy and the risk of pre-eclampsia. *BJOG : an international journal of obstetrics and gynaecology*. 2012; 119:832–839. [PubMed: 22462640]
15. Helewa ME, Burrows RF, Smith J, Williams K, Brain P, Rabkin SW. Report of the Canadian Hypertension Society Consensus Conference: 1. Definitions, evaluation and classification of hypertensive disorders in pregnancy. *Canadian Medical Association journal*. 1997; 157:715–725. [PubMed: 9307560]
16. Raymond D, Peterson E. A critical review of early-onset and late-onset preeclampsia. *Obstetrical & gynecological survey*. 2011; 66:497–506. [PubMed: 22018452]
17. Counil E, Julien P, Lamarche B, Chateau-Degat ML, Ferland A, Dewailly E. Association between trans-fatty acids in erythrocytes and pro-atherogenic lipid profiles among Canadian Inuit of Nunavik: possible influences of sex and age. *Br J Nutr*. 2009; 102:766–776. [PubMed: 19445816]
18. Viviani Anselmi C, Ferreri C, Novelli V, Roncarati R, Bronzini R, Marchese G, Somalvico F, Condorelli G, Montenero AS, Puca AA. Fatty acid percentage in erythrocyte membranes of atrial flutter/fibrillation patients and controls. *Journal of interventional cardiac electrophysiology : an international journal of arrhythmias and pacing*. 2010; 27:95–99. [PubMed: 20162444]

19. Bastani NE, Gundersen TE, Blomhoff R. Determination of 8-epi PGF(2 α) concentrations as a biomarker of oxidative stress using triple-stage liquid chromatography/tandem mass spectrometry. *Rapid Commun Mass Spectrom*. 2009; 23:2885–2890. [PubMed: 19670343]
20. Schwedhelm E, Boger RH. Application of gas chromatography-mass spectrometry for analysis of isoprostanes: their role in cardiovascular disease. *Clin Chem Lab Med*. 2003; 41:1552–1561. [PubMed: 14708879]
21. Morris JM, Gopaul NK, Endresen MJ, Knight M, Linton EA, Dhir S, Anggard EE, Redman CW. Circulating markers of oxidative stress are raised in normal pregnancy and pre-eclampsia. *Br J Obstet Gynaecol*. 1998; 105:1195–1199. [PubMed: 9853769]
22. Siddiqui IA, Jaleel A, Tamimi W, Al Kadri HM. Role of oxidative stress in the pathogenesis of preeclampsia. *Arch Gynecol Obstet*. 2010; 282:469–474. [PubMed: 20549510]
23. Li H, Lawson JA, Reilly M, Adiyaman M, Hwang SW, Rokach J, FitzGerald GA. Quantitative high performance liquid chromatography/tandem mass spectrometric analysis of the four classes of F(2)-isoprostanes in human urine. *Proc Natl Acad Sci U S A*. 1999; 96:13381–13386. [PubMed: 10557329]
24. Ramezani M, Nazemian F, Shamsara J, Koohrokhi R, Mohammadpour AH. Effect of omega-3 fatty acids on plasma level of 8-isoprostane in kidney transplant patients. *Journal of renal nutrition : the official journal of the Council on Renal Nutrition of the National Kidney Foundation*. 2011; 21:196–199.
25. Mas E, Woodman RJ, Burke V, Puddey IB, Beilin LJ, Durand T, Mori TA. The omega-3 fatty acids EPA and DHA decrease plasma F(2)-isoprostanes: Results from two placebo-controlled interventions. *Free Radic Res*. 2010; 44:983–990. [PubMed: 20540666]
26. Belik J, Gonzalez-Luis GE, Perez-Vizcaino F, Villamor E. Isoprostanes in fetal and neonatal health and disease. *Free Radic Biol Med*. 2010; 48:177–188. [PubMed: 19854268]
27. Yu X, Ho CS, Lam CW. Biological variation of plasma F2-isoprostane-III and arachidonic acid in healthy individuals. *Clin Chem*. 2004; 50:1428–1430. [PubMed: 15277353]

Figure 1.

Receiver operating characteristic (ROC) curves for ± 5 iPF_{2 α} -VI (iPF_{2 α} -VI + 5-iPF_{2 α} -VI) used for PE prediction normalized to plasma volume (—; area = 0.67) or arachidonic acid (AA) (—; area = 0.68) or Total ω -3/total ω -6 ratio (····; area = 0.70). Total omega-3 fatty acids (ω -3) = 18:3 + 18:4 + 20:3 + 20:4 + 20:5 + C22:5 + C22:6. Total omega-6 fatty acids (ω -6) = 18:1 + 18:2 + 18:3 + 20:2 + 20:3 + 20:4 + 22:2 + 22:4 + 22:5. n=60 controls, n=33 PE.

TABLE 1

F₂-isoprostanes plasma levels at the first visit (12–18 weeks) of pregnancy between normotensive controls and pregnant women that will later develop preeclampsia (PE).

	F ₂ -Isoprostanes levels ^I			
	pg/ml of plasma		pg/μg of arachidonic acid	
	Normotensives (n = 60)	toward PE (n = 33)	Normotensives (n = 60)	toward PE (n = 33)
<i>Class III</i>				
8-iso-15(R)-PGF _{2α}	105 [69, 161]	95 [67, 185]	0.83 [0.42, 1.13]	0.70 [0.48, 1.26]
8-iso-PGF _{2α}	118 [0, 199]	147 [0, 222]	0.85 [0, 1.36]	0.94 [0, 1.51]
<i>Class VI</i>				
iPF _{2α} -VI + 5-iPF _{2α} -VI	152 [121, 168]	172 [137, 209][*]	1.02 [0.81, 1.27]	1.21 [1.03, 1.49]^{**}

^I Values are medians and quartiles [Q1, Q3].

^{*} Significantly different from normotensives pregnancies (Mann-Whitney test, $P = 0.0074$).

^{**} Significantly different from normotensives pregnancies (Mann-Whitney test, $P = 0.0040$).

TABLE 2

Total omega-6, total omega-3 fatty acid content and ratio, unsaturation (UI) and peroxidation (PI) indexes at the first obstetrical visit (12–18 weeks) between normotensive controls and pregnancies that will later develop preeclampsia (PE).

	Fatty acid levels ($\mu\text{g/mL}$ of plasma)		
	Normotensives (n = 60)	toward PE (n = 33)	P
Total ω -6 ¹	486 [449, 544]	501 [444, 556]	0.8688
20:4 ω -6 (AA)	142 [128, 163]	138 [118, 172]	0.6777
Total ω -3 ²	83.2 [64.5, 92.5] (81 ± 20)	66 [59, 84] (71 ± 19)	0.0301[*] (0.0225[*])
<i>Ratio and indexes</i>			
ω -3/ ω -6	0.16 [0.13, 0.18]	0.14 [0.12, 0.17]	0.0382[*]
UI	144 [139, 149] (144 ± 6)	141 [137, 144] (141 ± 7)	0.0559 [§] (0.0320[*])
PI	115 [107, 124] (116 ± 10)	112 [103, 118] (111 ± 11)	0.0569 [§] (0.0406[*])

Values are medians and quartiles [Q1, Q3] and ($mean \pm SD$) if the normality postulate is achieved.

^{*} Significantly different from normotensives ($P < 0.05$; Mann-Whitney test or (*student t-test*)).

[§] Tendency, ($P < 0.1$; Mann-Whitney test)

¹ Total omega-6 fatty acids = 18:2 + 18:3 + 20:2 + 20:3 + arachidonic acid (AA; 20:4) + 22:2 + 22:4 + 22:5.

² Total omega-3 fatty acids = 18:3 + 18:4 + 20:3 + 20:4 + 20:5 + 22:5 + 22:6.

Unsaturation index (UI) = (%Monoenoic x 1) + (%Dienoic x 2) + (%Trienoic x 3) + (%Tetraenoic x 4) + (%Pentaenoic x 5) + (%Hexaenoic x 6).

Peroxidation index (PI) = (%Monoenoic x 0.025) + (%Dienoic x 1) + (%Trienoic x 2) + (%Tetraenoic x 4) + (%Pentaenoic x 6) + (%Hexaenoic x 8).

TABLE 3

Correlations between F₂-isoprostanes and arachidonic acid (AA), total omega-3, total omega-6 fatty acids content or ratio in the plasma at the first visit (12–18 weeks) of pregnancy between normotensive controls and pregnant women that will later develop preeclampsia (PE).

	20:4 ω -6 (AA)	Total ω -6 ¹	Total ω -3 ²	ω -3/ ω -6
<i>Normotensives</i>				
8-iso-15(R)-PGF _{2α}	ns	ns	ns	ns
8-iso-PGF _{2α}	ns	ns	ns	$r = -0.286$ ($p = 0.028$ *)
iPF _{2α} -VI + 5-iPF _{2α} -VI	ns	ns	$r = -0.275$ ($p = 0.034$ *)	$r = -0.336$ ($p = 0.009$ *)
<i>Toward PE</i>				
8-iso-15(R)-PGF _{2α}	$r = 0.412$ ($p = 0.017$ *)	$r = 0.447$ ($p = 0.009$ *)	ns	ns
8-iso-PGF _{2α}	$r = 0.377$ ($p = 0.031$ *)	$r = 0.480$ ($p = 0.005$ *)	ns	ns
iPF _{2α} -VI + 5-iPF _{2α} -VI	$r = 0.531$ ($p = 0.001$ *)	$r = 0.551$ ($p = 0.001$ *)	ns	ns

Values (r) are Spearman correlation coefficients (n=60 for normotensives and n=33 for PE).

* Significantly different ($P < 0.05$).

ns = non-significant ($P > 0.1$).

¹Total omega-6 fatty acids = 18:2 + 18:3 + 20:2 + 20:3 + 20:4 + 22:2 + 22:4 + 22:5

²Total omega-3 fatty acids = 18:3 + 18:4 + 20:3 + 20:4 + 20:5 + 22:5 + 22:6