Published in final edited form as:

Behav Genet. 2016 January ; 46(1): 4–19. doi:10.1007/s10519-015-9761-3.

Gene–Environment Interplay in Physical, Psychological, and Cognitive Domains in Mid to Late Adulthood: Is *APOE* a Variability Gene?

Chandra A. Reynolds¹, Margaret Gatz^{2,3}, Kaare Christensen^{4,5}, Lene Christiansen⁴, Anna K. Dahl Aslan^{6,3}, Jaakko Kaprio⁷, Tellervo Korhonen^{8,9}, William S. Kremen¹⁰, Robert Krueger¹¹, Matt McGue^{11,4}, Jenae M. Neiderhiser¹², Nancy L. Pedersen^{3,2}, and for the IGEMS consortium

Chandra A. Reynolds: chandra.reynolds@ucr.edu

¹Department of Psychology, University of California Riverside, 900 University Ave., Riverside, CA 92521, USA ²Department of Psychology, University of Southern California, Los Angeles, CA 90089, USA ³Department of Medical Epidemiology and Biostatistics, Karolinska Institute, 17177 Stockholm, Sweden ⁴Epidemiology, Biostatistics and Bio-demography, Institute of Public Health, University of Southern Denmark, 5000 Odense C, Denmark ⁵Department of Clinical Genetics and Department of Clinical Biochemistry and Pharmacology, Odense University Hospital, Odense, Denmark ⁶Institute of Gerontology, School of Health and Welfare, Jönköping University, Jönköping, Sweden ⁷Department of Public Health & Institute for Molecular Medicine FIMM, University of Helsinki, 00014 Helsinki, Finland ⁸Department of Public Health, University of Helsinki, 00014 Helsinki, Finland ⁹Institute of Public Health and Clinical Nutrition, University of Eastern Finland, 70211 Kuopio, Finland ¹⁰Department of Psychiatry, University of California San Diego, La Jolla, CA 92093, USA ¹¹Department of Psychology, University of Minnesota, Minneapolis, MN 55455, USA ¹²Department of Psychology, The Pennsylvania State University, University Park, PA, USA

Abstract

Despite emerging interest in gene–environment interaction (G×E) effects, there is a dearth of studies evaluating its potential relevance apart from specific hypothesized environments and biometrical variance trends. Using a monozygotic within-pair approach, we evaluated evidence of G×E for body mass index (BMI), depressive symptoms, and cognition (verbal, spatial, attention, working memory, perceptual speed) in twin studies from four countries. We also evaluated

Correspondence to: Chandra A. Reynolds, chandra.reynolds@ucr.edu.

Members of the consortium on Interplay of Genes and Environment across Multiple Studies (IGEMS) are listed in Appendix.

Electronic supplementary material The online version of this article (doi:10.1007/s10519-015-9761-3) contains supplementary material, which is available to authorized users.

Compliance with Ethical Standards

Conflict of interest Dr. Korhonen has served as a consultant on nicotine dependence for Pfizer (Finland) in 2011–2015.

Research involving Human Participants and/or Animals All procedures performed in studies involving human participants were in accordance with the ethical standards of the respective institutional and/or national research committees for each participating study providing archival data, and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.

Informed consent Informed consent was obtained from all individual participants from their respective parent study.

whether *APOE* is a ‘variability gene’ across these measures and whether it partly represents the ‘G’ in G×E effects. In all three domains, G×E effects were pervasive across country and gender, with small-to-moderate effects. Age-cohort trends were generally stable for BMI and depressive symptoms; however, they were variable—with both increasing and decreasing age-cohort trends—for different cognitive measures. Results also suggested that *APOE* may represent a ‘variability gene’ for depressive symptoms and spatial reasoning, but not for BMI or other cognitive measures. Hence, additional genes are salient beyond *APOE*.

Keywords

Gene–environment interaction; Twins; BMI; Depression; Cognitive performance; *APOE*; Variability gene

Introduction

Emerging evidence suggests that gene–environment interplay, including gene–environment interactions (G×E), may contribute to multiple life domains. Here we focus on measures sampled from three domains: physical [body mass index (BMI)]; psychological (depressive symptoms); and cognitive (verbal, spatial, attention/working memory, perceptual speed). The role of G×E among these domains has been variously studied by examining the interaction of a specific environmental exposure with a specific gene variant, how genetic variance may differ due to a specific exposure, or how environmental variance may differ as a function of a specific gene variant. For each of these domains, the current paper concerns establishing evidence of G×E, evaluating age-cohort differences in G×E effects, and testing whether *APOE* is a variability gene, i.e., in phenotypes where there is evidence of G×E, whether sensitivity to environmental influences varies with *APOE* gene variants.

As concerns obesity, a variety of twin studies have shown how genetic risk for obesity-related traits may be mitigated (or facilitated) by specific environmental factors. For example, in a Danish Twin Registry study, higher education levels corresponded with substantially reduced genetic variance, as well as shared and nonshared environmental variance, for BMI in women, with similar reductions in shared and nonshared environmental variance for BMI in men (Johnson et al. 2011). Vigorous exercise has also been associated with reduced genetic variance in BMI (McCaffery et al. 2009) in middle-aged men, and higher levels of physical activity have been associated with reduced genetic variance for BMI, waist-hip ratio, and percent body fat (Mustelin et al. 2009; Silventoinen et al. 2009) in young adult twins from Finland and adult twins from Denmark.

In the psychological domain, lower SES indexed by income level has been associated with magnified total variance for internalizing psychopathology in middle adulthood, indexed by major depression, generalized anxiety disorder, panic attacks, and neuroticism (South and Krueger 2011). However, the moderation of total variance for internalizing psychopathology was mainly due to magnification of unique environmental variance at the lowest SES levels. The finding of moderation of internalizing psychopathology via SES in middle adulthood builds on earlier work evaluating G×E for depression and indices of adversity (see Rutter 2012; Rutter and Silberg 2002). In particular, a greater risk of depression has been observed

in the presence of a combination of prior stress, particularly childhood maltreatment, and a variant in the serotonin transporter gene promoter region (5-HTTLPR) (Caspi et al. 2003; Karg et al. 2011), although not all studies replicate this finding (see Duncan and Keller 2011).

A potential signal of the presence of G×E for adult cognitive performance has come from observations that unique environmental influences may accelerate in importance with age across multiple cognitive tests (Pahlen et al. under review; Reynolds et al. 2005, 2007), although others have reported stability of twin similarity on a cognitive composite score (McGue and Christensen 2013). Moreover, twin studies examining G×E for mid to late adult cognition are limited compared to childhood and early adulthood. There is some evidence that higher levels of childhood SES are associated with greater genetic influences on general cognitive ability (Turkheimer and Horn 2014), although this effect has not been observed when assessed in adulthood (Grant et al. 2010). In adult male twins, across greater years of parental education, total variance and particularly common environmental variance for word recognition was reduced; whereas genetic variance was relatively stable (Kremen et al. 2005). A personality trait, Experience Seeking (ES), a subscale of the Sensation Seeking Scale (Dutch translation) has been evaluated as a moderator of genetic and environmental variance in cognitive ability in an adult twin sample with results suggesting reduced genetic variance but increased nonshared environmental variance at the highest levels of ES (Vinkhuyzen et al. 2012).

‘Agnostic’ tests of G×E

Typically G×E is tested with a selected environmental feature or exposure, or a specific gene target in mind, or both. However, an agnostic test has been available, without identified genes or environments, as first proposed by Fisher (1925; see Martin et al. 1983 for correction). Specifically, Fisher delineated a test of heterogeneity that relies on evaluating monozygotic (MZ) within-pair differences (Fisher 1925; Martin et al. 1983), i.e., the test compares mean squared pair differences for a trait with the mean absolute pair differences squared. The extent to which these values differ supports a mixture of distributions of the within-pair differences rather than one distribution of differences and suggests there is possible G×E interaction. This indicates a differential sensitivity of genotypes to environments such that the MZ pair differences, which reflect nonshared environment, vary according to particular genotypes. MZ within-pair approaches are rarely used (Cornes et al. 2008; Martin 2000; Martin et al. 1983; Reynolds et al. 2007; Surakka et al. 2012), particularly since the advent of genome-wide genotyping, but such an approach may usefully quantify the extent of heterogeneity and identify the likely presence of G×E. Coupling an agnostic general test with potential genetic markers using MZ pairs can be more powerful than evaluating G×E in population-based samples of unrelated individuals (Visscher and Posthuma 2010).

Variability genes, i.e., the ‘G’ in G×E

A significant Fisher test of heterogeneity could indicate the presence of G×E interaction, i.e., differential sensitivity of particular genotypes to particular environments, or could reflect a shared environment by nonshared environment interaction, C×E. To support that an observed

significant heterogeneity test is due to $G \times E$, it is useful to consider measured genes that may explain such heterogeneity (Berg et al. 1989; Martin 2000; Martin et al. 1983). The genes of interest may be regarded as ‘variability genes’ (Berg et al. 1989), i.e., genes that are associated with trait variation and not simply associated with trait mean (Martin 2000). *APOE* may be of particular interest in this regard. The *APOE* gene, coding for the major cholesterol transporter in the brain, and its $\epsilon 4$ haplotype in particular, has demonstrated associations with cognitive decline, Alzheimer’s disease (AD) and dementia (e.g., Bennet et al. 2010; Davies et al. 2014; Reynolds et al. 2006; Schellenberg and Montine 2012). In addition, *APOE* has also shown some evidence of associations with, or moderation of, risk factors that are predictive of cognitive decline and dementia, including BMI (e.g., Besser et al. 2014; Keller et al. 2011) and depression (e.g., Karlsson et al. 2015; Skoog et al. 2015).

APOE has shown evidence that it may act as a variability gene; that is, the effects of environmental risk and protective factors have been shown to differ according to *APOE* genotype. For example, MZ twin pairs who were *APOE* $\epsilon 4$ —were more variable in their semantic memory trajectories, whereas those who were $\epsilon 4+$ were less variable (Reynolds et al. 2007). Additionally, individuals with particular *APOE* haplotypes may be differentially sensitive to dietary and exercise interventions, albeit not consistently (Brown et al. 2013a; Carvalho-Wells et al. 2012; Gomez-Pinilla and Hillman 2013; Hotting and Roder 2013). For example, in those who lead sedentary lives, amyloid burden is greater for those with $\epsilon 4+$ compared to other *APOE* haplotypes, whereas for those who engage in physical activity, amyloid burden does not vary across *APOE* haplotypes (Brown et al. 2013b; Head et al. 2012). Moreover, a recent experimental study in sedentary women suggested a particular benefit of acute exercise to $\epsilon 4+$ carriers on a cognitive inhibition task (Stroop) in comparison to a spatial attention task (Posner) that engages the prefrontal region to a lesser extent, but no benefit accrued for non- $\epsilon 4$ individuals across tasks (De Marco et al. 2015). MZ twin pair differences in semantic memory change have also been associated with twin-pair differences in depressive symptoms but in this case only among non- $\epsilon 4$ individuals (Reynolds et al. 2007). Thus, taken together, emerging evidence across multiple traits and domains supports the role of *APOE* as a variability gene and suggest that the associations of *APOE* may be complex and depend in part on environmental factors. Indeed, for BMI *APOE* may show differing patterns of evidence for sensitivity, as compared to cognition or depression traits.

The aims of the current study were to evaluate general evidence of $G \times E$ for BMI, depressive symptoms, and cognitive performance in twin studies participating in the Interplay of Genes and Environment across Multiple Studies (IGEMS) consortium (Pedersen et al. 2013). We further considered whether there were age-cohort trends in $G \times E$. Once general evidence for $G \times E$ was evaluated, we considered specific genetic aspects further, by testing the extent to which *APOE* was a variability gene across these traits. That is, we evaluated whether different *APOE* haplotypes were more or less sensitive to environmental factors and thereby showed differences in the variance of pair differences in depressive symptoms, BMI and cognitive performance.

Methods

Samples

The current analysis sample includes individuals from up to nine twin studies representing four countries: the United States, Sweden, Denmark and Finland, from the IGEMS consortium (Pedersen et al. 2013). The primary analyses considered complete MZ twin pairs to evaluate heterogeneity of within-pair differences and homogeneity of within-pair variance by *APOE* haplotypes (see Table 1). Each of the respective studies described below obtained approvals by their Institutional Review Boards, or equivalent, to carry out the original data collection, obtaining informed consent from participants as required.

USA—Data were available from the Vietnam Era Twin Study of Aging (VETSA) (Kremen et al. 2013), Minnesota Twin Study of Adult Development and Aging (MTSADA) (Finkel et al. 1995), and the Midlife Development in the United States (MIDUS) twin study (Kendler et al. 2000; Radler 2014). The VETSA study included only male twin pairs (51–60 years), while from MTSADA (25–92 years) and MIDUS (34–82 years) we included same-sex male and female pairs.

Sweden—Data were available from three population-based samples of same-sex male and female twins that originated from the Swedish Twin Registry (Lichtenstein et al. 2006; Magnusson et al. 2013): the Swedish Adoption/Twin Study of Aging (SATSA) (Pedersen et al. 1991), the Origins of Variance in the Oldest-Old (OCTO-twin) (McClearn et al. 1997), and the Twin-Offspring Study in Sweden (TOSS) (Neiderhiser et al. 2007). Data for SATSA twins (39–88 years) came from the first available questionnaire or in-person testing wave, available during one of 6 respective assessment waves. Data on OCTO-twin participants (79–99 years) came from the first assessment. Twin data from the parent generation (32–60 years) of the TOSS study were used in the current study.

Denmark—The Longitudinal Study of Aging Danish Twins (LSADT) (70–100 years) and the Middle Aged Danish Twins (MADT) (45–68 years) included pairs drawn from the Danish Twin Register (McGue and Christensen 2013; Skytthe et al. 2013). Data from the first assessment wave were used in the present study.

Finland—The Finnish Adult Twin Cohort (FTC; Kaprio and Koskenvuo 2002) sample included data from the fourth assessment wave of twins born 1945–1957, done as a postal questionnaire survey in 2011 to 2012 (Kaprio 2013).

Measures

All studies had data from at least one of the following three domains.

BMI—BMI was computed in standard fashion as weight, measured in kilograms, divided by height squared, measured in meters (kg/m^2). BMI scores were adjusted for self-report versus measured assessments (Johnson et al. 2012) given that self-reports are biased towards over-reporting of height yet under-reporting of weight (Dahl et al. 2010), i.e., $\text{Adjusted BMI} = 0.35 + 1.038 * (\text{BMI}_{\text{self-rept}})$. Studies in the current analysis with measured height and weight

assessments included OCTO-Twin and VETSA, the remainder of the studies provided self-reported data. Prior to analysis, BMI scores were rank-normalized to reduce non-normality (c.f., Reynolds et al. 2007; Surakka et al. 2012).

Depression—Depressive symptoms were measured with either the Center for Epidemiologic Studies Depression (CESD) scale (Radloff 1977) or the Cambridge Mental Disorders of the Elderly Examination (CAMDEX) as modified by McGue and Christensen (McGué and Christensen 1997). To create a common metric, both scales were collected from a separate crosswalk sample, and item response theory methods were applied in order to compare items from the two measures and create a conversion table between the scales (Gatz et al. in press). We retained those items from both CESD and CAMDEX that loaded on the respective affect and somatic subscales. The co-calibrated score is expressed in CAMDEX units, such that the total score can range from 16 for someone who endorses no symptoms of depression to 46. After harmonization, scores were rank-normalized to reduce non-normality.

Cognitive performance—Five measures of cognitive ability spanning four cognitive domains were considered in the current study: verbal (Synonyms), spatial (Block Design), attention and working memory (Digit Span Forward and Backward), and perceptual speed (Symbol Digit). Each measure was available in at least two studies. Number of individuals available for each test was therefore variable, reflecting the differential availability of the tests across studies. Cognitive tests and harmonization procedures have been described previously (Pahlen et al. under review). In short, those in the analysis sample completed at least one of the cognitive tests and scored 24 or above on the Mini-Mental State Exam (MMSE; Folstein et al. 1975); a total of 7.3 % of the total sample were excluded based on the MMSE criteria. Scores were residualized for sex and transformed to T-score scaling ($M = 50.0$ and $SD = 10.0$) against the reference age group 50 to 59.99 years (Pahlen et al. under review) and subjected to winsorizing within age group for values falling outside of ± 3 SDs. Prior to within-pair analyses, scores were rank-normalized to reduce non-normality.

Genotyping

APOE haplotypes were available for a subset of studies and were categorized as $\epsilon 2+$ ($\epsilon 22$, $\epsilon 23$, $\epsilon 24$), $\epsilon 33$, and $\epsilon 4+$ ($\epsilon 34$, $\epsilon 44$). Samples with MZ pairs and genotyping included: VETSA (US) SATSA and OCTO-Twin (Swedish), MADT and LSADT (Danish). Genotyping procedures for VETSA, SATSA and OCTO-Twin have been described elsewhere (Reynolds et al. 2013; Schultz et al. 2008). For the Danish samples, *APOE* haplotypes were formed from two genotyped SNPs, rs429358 and rs7412, that for MADT were based on TaqMan[®] SNP Genotyping Assays (Applied Biosystems, Foster City, CA, USA) and for LSADT were based on custom-designed assays.

APOE haplotype frequencies are reported in supplementary Table 1 (Table S1). Hardy–Weinberg Equilibrium based on computations for a three allele system were calculated for each study and met ($p = 0.121$). MZ twins who were not directly genotyped were assigned their cotwin's value.

Statistical analysis

We evaluated the presence of G×E by applying a test of mixture distributions of MZ within-pair differences overall, and separately by country, sex, and age group. Given that the data are cross-sectional such that age group and birth cohort are unable to be dissociated, we refer to age group as age-cohort. Specifically, we applied a test first proposed by Fisher (Fisher 1925; Martin et al. 1983). The test evaluates the difference between mean squared pair differences for a trait and the mean absolute pair differences squared as follows (Fisher 1925; Martin et al. 1983):

$$\Delta = \overline{d^2} - \frac{\pi}{2} \overline{d}^2 \quad (1.0)$$

and corresponding standard error as (Fisher 1925; Martin et al. 1983):

$$se = \frac{\overline{d^2}}{\sqrt{n}} (.5321) \quad (1.1)$$

A one-tailed *t* test was used to evaluate significance (Δ / se), given that the expected values were assumed to be positive (Martin et al. 1983), with *df* equal to the number of pairs minus 1.0. To address multiple testing, we conducted false-discovery rate (FDR) tests (Benjamini and Hochberg 1995; Weinkauff 2012) and provided Holm-Bonferroni adjusted *p* values as well for each set of tests by trait (Gaetano 2013; Holm 1979).

In addition, effect size *r*s were calculated from the *t* statistics (Rosenthal 1991) to consider the potential impact of G×E across country, sex, and age-cohort, apart from power considerations:

$$ESr = \sqrt{\frac{t^2}{(t^2 + df)}} \quad (2)$$

A measure of the heterogeneity of effect size *r*s (ESrs) were calculated according to the Chi square test outlined by Snedecor & Cochran (1989; as cited in Rosenthal 1991).

In a subset of available samples, we considered measured genes to substantiate G×E and not C×E. Specifically, heterogeneity of variance by *APOE* haplotype was evaluated using SAS Proc Mixed (SAS Inc, Cary, NC) specifying between and within pair random effects. Analyses of within-pair variation were adjusted for average effects of *APOE* haplotype, country, sex and age. A series of model constraints were tested on within pair variances, considering *APOE* haplotype differences within and across country or sex. Given the potential differential regional and within-country impact of *e4* on health outcomes, such as mortality (Ewbank 2004) and Alzheimer's disease (Ward et al. 2012), as well as differential impact of *APOE* on cognitive outcomes for women versus men (Altmann et al. 2014; Damoiseaux et al. 2012; Farrer et al. 1997), we evaluated whether *APOE* effects could be

generalized. Hence, we tested whether within-pair variances for each *APOE* haplotype could be constrained: (1) across men and women within country (i.e., $\epsilon_{2+m} = \epsilon_{2+f}$, $\epsilon_{33m} = \epsilon_{33f}$, $\epsilon_{4+m} = \epsilon_{4+f}$), and (2) across country (i.e., $\epsilon_{2+US} = \epsilon_{2+SWE} = \epsilon_{2+DEN}$, etc.). Last, we tested whether within-pair variances could be constrained equal within country across the three *APOE* haplotype groups (i.e., $\epsilon_{2+} = \epsilon_{33+} = \epsilon_{4+}$) to evaluate the significance of an *APOE* effect on variability. Sensitivity analyses considered adjustments when dropping individuals with the *APOE* ϵ_{24} haplotype. We did not evaluate age trends in within pair variances by *APOE* haplotype, primarily due to the reductions in sample sizes of those with both phenotypic data and genotyping and to resultant confounding of age-cohort and country.

Follow-up tests of association at the mean level based on *APOE* haplotype were undertaken in SAS Proc Mixed (SAS Inc, Cary, NC) allowing for within and between pair variances to differ by country; analyses adjusted for average effects of age, sex and country. Specifically, we tested whether entering the *APOE* haplotype (ϵ_{2+} , ϵ_{33} , ϵ_{4+}) led to a significant improvement in fit based on a two-degree of freedom test.

Results

Fisher heterogeneity test

The full sample heterogeneity tests for BMI included 3550 complete MZ pairs and for depressive symptoms 3508 MZ pairs. For the cognitive measures, 2338 MZ pairs had at least one cognitive test where both members participated and met MMSE criterion; test availability across studies the analysis samples ranged from 390 to 1727 MZ pairs. The Fisher (1925) test suggested significant within-pair heterogeneity in the full sample for BMI, $p = 3.54E-34$, and depressive symptoms, $p = 1.99E-41$ (see Table 2), with significant within-pair heterogeneity for each age-cohort ($p = 6.87E-03$; see Table 2), as well as both sexes and all four countries ($p = 3.90E-04$; see supplement Table S2). Overall effect size r s (ESRs) were small for both BMI and depressive symptoms (median = .19, .21, respectively). Effect sizes were consistent across age-cohort groups for both BMI and depressive symptoms [$\chi^2(4) = 2.55$, $p = 6.36E-01$] (see Table 2). BMI showed consistent small ESRs across country [$\chi^2(3) = 3.68$, $p = 3.68E-01$]. Although depressive symptoms showed small and significant evidence for G×E for each country, the ESRs were significantly variable with lower effect sizes for Sweden and Finland and higher effects for US and Denmark [$\chi^2(3) = 18.77$, $p = 3.06E-04$] (see supplement Table S2).

For cognitive performance, G×E was suggested in the full sample ($p = 2.16E-04$) (see Table 2). The ESRs were small, ranging from .12 to .23, and were not significantly heterogeneous from one another [$\chi^2(4) = 7.71$, $p = 1.03E-01$] (see Table 2, supplement Table S2). As depicted in Fig. 1, three prototypical age-cohort trends in ESRs were noticeable: (a) Block Design represented a linear pattern of increasingly stronger effect sizes across age groups; (b) Digits Backward represented a nonlinear u-shaped pattern with peaks before age of 50 (ESR = .27) and after age of 80 (ESR = .39) with a similar trend for Digits Forward (not shown), and (c) Symbol Digit displayed a pattern of decreasing effect sizes with age-cohort, with the peak at ages 50–59 (ESR = .22). The pattern for Synonyms was less consistent and is not shown in Fig. 1 (but see Table 2). The FDR tests and Holm-Bonferroni adjusted p -values generally supported the age-based patterns described in terms of significance (see

Table 2); however, heterogeneity tests of ESrs among age-cohorts suggested that only Digits Backward reached significance [$\chi^2(4) = 10.14, p = 3.81E-02$], with a trend effect in Digits Forward ($p = 5.07E-02$).

G×E was indicated on all cognitive measures both for women ($p = 5.48E-03$) and for men ($p = 4.56E-05$), apart from Synonyms ($p = 6.26E-02$). For all five measures, there was evidence of significant heterogeneity of within pair differences across all countries, although for Symbol Digit, only Denmark showed a significant effect ($p = 8.62E-12$; ESr = .23), and for Synonyms, only Sweden ($p = 1.11E-03$; ESr = .13) (see supplemental Table S2).

Measured G×E: APOE

In the primary analyses of APOE as a variability gene, we focused on testing for heterogeneity in the variance of pair differences among *APOE* haplotypes evaluating whether variances could be constrained by country and sex, adjusting for average effects of age, sex and *APOE* haplotypes on the trait scores (see Table 3). We did not evaluate age trends in within pair variances by *APOE* haplotype for BMI, depression or for cognition, primarily due to the reductions in sample sizes of those with phenotypic data and genotyping and consequent confounding of age-cohort and country. Moreover, we note that the general age-cohort consistency of the evidence for G×E observed for BMI and depressive symptoms. Significant findings are described further below. Analyses of mean level associations (i.e., whether individuals score higher or lower on the trait on average) are reported (see Table 4), with no significant associations observed; description of mean trends is provided below for traits showing significant evidence for *APOE* × variance effects. Dropping *APOE* ε24 individuals from the analysis did not alter any of the conclusions.

BMI—Variances of absolute pair differences by *APOE* haplotype could not be constrained across sex within country [$\chi^2(6) = 19.64, p = 3.21E-03$] (see Table 3 for within pair variance estimates and test statistics); hence, further analyses were conducted separately for men and women. Nonsignificant country differences in within pair variances within *APOE* haplotype were observed for men and women ($p = 9.31E-02$). In addition, within pair variances could be constrained across *APOE* haplotype within country ($p = 7.22E-01$). In sum, within pair variances for the *APOE* haplotypes differed between men and women, but across country the *APOE* haplotype effects were not statistically different from each other. Hence, there was no support for an *APOE* effect on within pair variability, but there was heterogeneity of within pair variances across men and women suggesting that female pairs are more variable than male pairs in terms of the degree to which twins differ from their cotwin in BMI.

Depressive symptoms—Variances of absolute pair differences by *APOE* haplotype could be constrained across sex within country [$\chi^2(6) = 2.05, p = 9.15E-01$]; hence analyses were conducted collapsing men and women together (see Table 3). Haplotype-based within pair variances could not be constrained across country [$\chi^2(6) = 44.99, p = 4.70E-08$]. Thus, haplotype-based within pair variances were allowed to vary within country and significant differences by *APOE* haplotype were observed [$\chi^2(6) = 19.78, p = 3.04E-03$]. Figure 2a indicates that *APOE* effects could be observed in the US and in the Swedish samples, with

smaller variances of pair differences for *APOE* ε4+ compared to larger variances for *APOE* ε33 and ε2+. This pattern suggests that those with *APOE* ε4+ may be less affected by environmental factors compared to the other haplotypes. Last, we followed up these variance tests of within-pair differences to consider whether *APOE* effects were evident for average depressive symptom scores, with no significant differences observed ($p = 2.83\text{E-}01$).

Cognitive performance—Among the five cognitive measures considered, only Block Design showed evidence of significant haplotype differences in within pair variances (see Table 3). Variances by *APOE* haplotype could be constrained across sex [$\chi^2(3) = 5.76$, $p = 1.24\text{E-}01$]; hence, analyses were conducted collapsing men and women together. As Block Design and *APOE* genotyping were only available in two Swedish samples, no country comparisons could be conducted. Significant differences in within pair variances by *APOE* haplotype were observed [$\chi^2(2) = 11.91$, $p = 2.60\text{E-}03$]. Smaller within pair variances of pair differences for *APOE* ε4+ versus larger variances for *APOE* ε2+ were observed (see Fig. 2b). This pattern indicates that those with *APOE* ε4+ may be less affected by environmental factors compared to those with *APOE* ε33 and *APOE* ε2+, and is consistent with the overall pattern observed for depressive symptoms above. Last, we followed up these within-pair variance tests to consider whether *APOE* effects were evident for average Block Design performance scores, and no significant differences were observed ($p = 2.49\text{E-}01$).

Discussion

We evaluated general evidence of G×E for BMI, depressive symptoms, and cognitive performance in twin studies from four countries, i.e., US, Sweden, Denmark, and Finland. We further evaluated whether *APOE* is a variability gene across these traits and represents, in part, the G in the G×E effects. We observed that across physical, psychological, and cognitive domains, G×E was pervasive across country and sex showing small to moderate effect sizes. While modest, the presence of these effects across domains argues for the importance of more routinely considering gene–environment interaction in biometric models. Generally stable age-cohort trends were observed for BMI and depressive symptoms. However, age-cohort trends varied by cognitive trait domains with some showing decreasing G×E effects and some showing increasing G×E effects. Last, *APOE* may represent one variability gene for depressive symptoms and spatial reasoning, but not for BMI or other cognitive tests. Hence additional variability genes are salient beyond *APOE*.

BMI

BMI evidenced small G×E effects, and these effects were consistent across country, sex, and age-cohort. This is perhaps not surprising in that the candidate G×E studies evaluating education or exercise on genetic variations in BMI have reported G×E in samples from various countries represented in our study (US, Denmark, Finland; Johnson et al. 2011; Lajunen et al. 2012; McCaffery et al. 2009; Mustelin et al. 2009; Silventoinen et al. 2009; Silventoinen et al. 2004). Others have suggested that the genetic variance for BMI may be increasing in later born Swedish cohorts (Rokholm et al. 2011), perhaps suggesting a complex cohort/generational G×E given changing dietary and activity patterns amongst others. Further examinations of longitudinal data across multiple cohorts would be

informative as to the extent to which G×E for BMI is dynamic across age versus birth cohort.

Despite agnostic evidence of G×E, no *APOE* associations were observed with within-pair variability for BMI. Prior studies have noted interactions of *APOE* with BMI, obesity, or of BMI variants (e.g., *FTO*) with outcomes such as metabolic traits (Elosua et al. 2003), dementia risk (Keller et al. 2011) or dementia progression (Besser et al. 2014). However, GWAS have not observed direct genetic association of *APOE* with mean BMI (Locke et al. 2015). Nonetheless, our lack of findings of *APOE* in the current analysis suggests that other variability genes, e.g., perhaps based on a polygenic risk score of 97 BMI loci (Locke et al. 2015), are relevant to pursue given evidence of G×E we observed in the agnostic Fisher analysis.

Depressive symptoms

Depressive symptoms showed consistently small but significant G×E effect sizes for sex and age-cohort, with lower effect sizes for Sweden and Finland and higher for US and Denmark. Our findings of ubiquitous small G×E effects furthers earlier evidence there is not simply an effect of the environment (E) on depressive symptom levels but that there is genetically influenced sensitivity to environmental factors that may foster (or mitigate) depression (c.f., Kendler et al. 1995).

We observed associations of *APOE* with within-pair variability in depression symptoms but no effect on mean depression scores. Results varied across country; evidence for *APOE* as the ‘G’ in G×E was found for the U.S. and Sweden, but not the Danish sample. Indeed, *APOE* associations with average depression symptoms and risk for a diagnosis of depression have been mixed across studies, perhaps due to differential population effects or study designs (Skoog et al. 2015). *APOE* has been associated with depressive symptomatology and depression diagnosis in late adulthood in a prospective study of Swedish individuals even when excluding prevalent or incident dementia cases (Skoog et al.). Other comparably sized (or larger) cross-sectional and longitudinal studies have not found such effects (e.g., Locke et al. 2013; Schultz et al. 2008; Surtees et al. 2009); however, the average sample age tended to be between ages 55 and 61, suggesting that the association of *APOE* and depressive symptoms may tend towards older adults.

Our results suggest that the effect of *APOE* on depression would appear to lie, not in main effects, but in the role of *APOE* in magnifying or reducing the effects of environmental risk factors for depressive symptoms. Specifically, MZ pairs carrying the $\epsilon 4$ haplotypes showed the smallest within-pair differences while those carrying the $\epsilon 2$ haplotypes the largest within-pair differences in depression scores. Hence, the depressive symptoms experienced by those with *APOE* $\epsilon 4$ + may be less driven by environmental factors, and more by familial or endogenous factors, compared to depressive symptoms experienced by those with other *APOE* haplotypes. Together with the observed age-cohort trends, such an interpretation would be consistent with the role of vascular factors and white matter changes in late onset depression (Nebes et al. 2001; Taylor et al. 2013).

Cognition

Different cognitive performance domains showed different patterns of results with respect to the agnostic Fisher G×E tests, with the pattern possibly reflecting the difference between age-sensitive cognitive tests versus more age-robust tests. The most age-sensitive test, perceptual speed indexed by Symbol Digit task performance, showed peak G×E effects in the younger age-cohorts compared to later age cohorts; whereas tests of attention, working memory, and spatial performance showed higher G×E in later age-cohorts. These latter tests tend to show later declines, accelerating across the adult lifespan (Salthouse 2009; Schaie 1994). We note that the complexity of findings underscores the need to consider specific cognitive abilities beyond general measures of ability.

In the *APOE* analyses, where we adjusted for age given the restricted sample size, we observed an effect for the spatial task, Block Design, but no other tasks. For Block Design, as for depressive symptoms, those with *APOE* ε4 + may be less affected by environmental factors compared to the other *APOE* haplotypes. It is worth noting that Block Design performance may be a salient predictor of subsequent cognitive dysfunction (e.g., Andel et al. 2001; Bozoki et al. 2001; Hamilton et al. 2008; Tabert et al. 2006). Hence those at risk for dysfunction or decline may show relatively less sensitivity to environmental factors compared to those without this risk allele, whose performance does reflect environmental influences.

The lack of association of *APOE* with variability for other cognitive measures could be viewed as puzzling. *APOE* associations with cognitive performance levels in non-demented adults have been mixed overall. However, we note that age-related change may be more salient than cross-sectional differences in performance level in terms of gene associations (e.g., Davies et al. 2014; Finkel et al. 2011; Salmon et al. 2013) as well as observing G×E effects (Reynolds et al. 2007). For example, in longitudinal work in SATSA using the within MZ pair methods, we observed significant G×E effects on semantic, episodic, and working memory trajectory features (e.g., linear and nonlinear change) but negligible effects on overall performance level (Reynolds et al. 2007). Hence, longitudinal examinations may reveal unique effects not apparent in baseline performance data. Another interpretation, given the longitudinal findings, might suggest that effects may not show up strongly until later ages. If age is adjusted for, then age periods where *APOE* or another gene or genes have a particular effect may be missed.

The smaller within-pair differences for those with *APOE* ε4 may seem to be counter-intuitive given that in some instances ε4 individuals may show greater rather than lesser sensitivity to particular environments that are relevant to brain reserve, not only dietary and exercise factors as mentioned above (Brown et al. 2013a, b; Carvalho-Wells et al. 2012; De Marco et al. 2015; Head et al. 2012), but also head injury and neuropsychological functioning and dementia (e.g., Sundstrom et al. 2004; Sundstrom et al. 2007; Tang et al. 1996) and combat exposure and PTSD (Kimbrel et al. 2015; Lyons et al. 2013). While a diathesis-stress model would expect ε4 always to act in the same direction, others have proposed the concept of a plasticity gene (Belsky et al. 2009; Belsky and Pluess 2009). Such an interpretation would be consistent with smaller within-pair differences for ε4 and greater

sensitivity to some exposures or contexts but lessened sensitivity to other exposures or contexts.

Strengths, limitations, and future directions

The strengths of the current study include the relatively large samples of MZ pairs and the ability to evaluate (and replicate) G×E trends in physical, psychological, and cognitive domains across up to four countries, by sex, and age cohorts. Moreover, in a subset of studies we were able to evaluate a well-characterized gene, *APOE*, as a potential variability gene. The primary limitation was that a single-occasion was available for evaluation of G×E for BMI, depressive symptoms and cognition. Moreover, not all studies had available *APOE* genotyping, hampering age-cohort investigations. Moreover, we had a limited set of cognitive measures and, hence, future studies would benefit from inclusion of measures of executive function and episodic memory.

Overall, future research directions should consider the possible measured environmental factors, i.e., the ‘E’ in G×E, given that G×E was ubiquitously observed albeit with generally small impact. Indeed, particularly for depression and spatial reasoning, the impact of any measured environmental factors may be modified by the *APOE* gene.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

IGEMS is supported by the National Institutes of Health Grant no. R01 AG037985. SATSA was supported by Grants R01 AG04563, R01 AG10175, the MacArthur Foundation Research Network on Successful Aging, the Swedish Council For Working Life and Social Research (FAS) (97:0147:1B, 2009-0795) and Swedish Research Council (825-2007-7460, 825-2009-6141). OCTO-Twin was supported by Grant R01 AG08861. Gender was supported by the MacArthur Foundation Research Network on Successful Aging, The Axel and Margaret Ax:son Johnson's Foundation, The Swedish Council for Social Research, and the Swedish Foundation for Health Care Sciences and Allergy Research. TOSS was supported by Grant R01 MH54610 from the National Institute of Health. The Danish Twin Registry is supported by Grants from The National Program for Research Infrastructure 2007 from the Danish Agency for Science and Innovation, the Velux Foundation and the US National Institute of Health (P01 AG08761). The Minnesota Twin Study of Adult Development and Aging was supported by NIA Grant R01 AG 06886. VETSA was supported by National Institute of Health Grants NIA R01 AG018384, R01 AG018386, R01 AG022381, and R01 AG022982, and, in part, with resources of the VA San Diego Center of Excellence for Stress and Mental Health. The Cooperative Studies Program of the Office of Research & Development of the United States Department of Veterans Affairs has provided financial support for the development and maintenance of the Vietnam Era Twin (VET) Registry. This MIDUS study was supported by the John D. and Catherine T. MacArthur Foundation Research Network on Successful Midlife Development and by National Institute on Aging Grant AG20166. The Finnish Twin Cohort study has been supported by Academy of Finland Center of Excellence in Complex Disease Genetics (Grant numbers: 213506, 129680), the Academy of Finland (Grants 265240, 263278 & 264146 to JK) and ENGAGE – European Network for Genetic and Genomic Epidemiology, FP7-HEALTH-F4-2007, Grant agreement number 201413. The content of this manuscript is solely the responsibility of the authors and does not necessarily represent the official views of the NIA/NIH, or the VA.

Appendix

Members of the consortium on Interplay of Genes and Environment across Multiple Studies (IGEMS) include: Nancy L. Pedersen (Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Stockholm, Sweden, and Department of Psychology, University of Southern California, Los Angeles, CA), Kaare Christensen (Department of

Epidemiology, University of Southern Denmark, Odense, Denmark), Anna K. Dahl Aslan (Institute of Gerontology, School of Health Sciences, Jönköping University, Jönköping, Sweden), Deborah Finkel (Department of Psychology, Indiana University Southeast, New Albany, IN), Carol E. Franz (Department of Psychiatry, University of California, San Diego, La Jolla, CA), Margaret Gatz (Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Stockholm, Sweden, and Department of Psychology, University of Southern California, Los Angeles, CA), Briana N. Horwitz (Department of Psychology, California State University, Fullerton, CA), Boo Johansson (Department of Psychology, University of Gothenburg, Gothenburg, Sweden), Wendy Johnson (Department of Psychology and Centre for Cognitive Ageing and Cognitive Epidemiology, University of Edinburgh, Edinburgh, UK), Jaakko Kaprio, Department of Public Health, University of Helsinki, Helsinki, Finland), William S. Kremen (Department of Psychiatry, University of California, San Diego, and VA Center of Excellence for Stress and Mental Health, La Jolla, CA), Robert Krueger (Department of Psychology, University of Minnesota, Minneapolis, MN), Michael J. Lyons (Department of Psychological and Brain Sciences, Boston University, Boston, MA), Matt McGue (Department of Psychology, University of Minnesota, Minneapolis, MN), Jenae M. Neiderhiser (Department of Psychology, The Pennsylvania State University, University Park, PA), Matthew S. Panizzon (Department of Psychiatry, University of California, San Diego, La Jolla, CA), Inge Petersen (Department of Epidemiology, University of Southern Denmark, Odense, Denmark), and Chandra A. Reynolds (Department of Psychology, University of California-Riverside, Riverside, CA).

References

- Altmann A, Tian L, Henderson VW, Greicius MD. Alzheimer's Disease Neuroimaging Initiative Investigators. Sex modifies the APOE-related risk of developing Alzheimer disease. *Ann Neurol*. 2014; 75(4):563–573. [PubMed: 24623176]
- Andel R, Gatz M, Pedersen NL, Reynolds CA, Johansson B, Berg S. Deficits in controlled processing may predict dementia: a twin study. *J Gerontol B*. 2001; 56B(6):P347.
- Belsky J, Pluess M. Beyond diathesis stress: differential susceptibility to environmental influences. *Psychol Bull*. 2009; 135(6):885–908. [PubMed: 19883141]
- Belsky J, Jonassaint C, Pluess M, Stanton M, Brummett B, Williams R. Vulnerability genes or plasticity genes? *Mol Psychiatry*. 2009; 14(8):746–754. [PubMed: 19455150]
- Benjamini Y, Hochberg Y. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *J Royal Stat Soc B*. 1995; 57(1):289–300.
- Bennet AM, Reynolds CA, Gatz M, Blennow K, Pedersen NL, Prince JA. Pleiotropy in the presence of allelic heterogeneity: alternative genetic models for the influence of APOE on serum LDL, CSF amyloid-beta42, and dementia. *J Alzheimer's Dis*. 2010; 22(1):129–134. [PubMed: 20847432]
- Berg K, Kondo I, Drayna D, Lawn R. "Variability gene" effect of cholesteryl ester transfer protein (CETP) genes. *Clin Genet*. 1989; 35(6):437–445. [PubMed: 2567644]
- Besser LM, Gill DP, Monsell SE, Brenowitz W, Meranus DH, Kukull W, Gustafson DR. Body mass index, weight change, and clinical progression in mild cognitive impairment and Alzheimer disease. *Alzheimer Dis Assoc Disord*. 2014; 28(1):36–43. [PubMed: 24126214]
- Bozoki A, Giordani B, Heidebrink JL, Berent S, Foster NL. Mild cognitive impairments predict dementia in nondemented elderly patients with memory loss. *Arch Neurol*. 2001; 58(3):411–416. [PubMed: 11255444]
- Brown BM, Peiffer JJ, Martins RN. Multiple effects of physical activity on molecular and cognitive signs of brain aging: can exercise slow neurodegeneration and delay Alzheimer's disease? *Mol Psychiatry*. 2013a; 18(8):864–874. [PubMed: 23164816]

- Brown BM, Peiffer JJ, Taddei K, Lui JK, Laws SM, Gupta VB, Taddei T, Ward VK, Rodrigues MA, Burnham S, Rainey-Smith SR, Villemagne VL, Bush A, Ellis KA, Masters CL, Ames D, Macaulay SL, Szoek C, Rowe CC, Martins RN. Physical activity and amyloid-beta plasma and brain levels: results from the Australian imaging, biomarkers and lifestyle study of ageing. *Mol Psychiatry*. 2013b; 18(8):875–881. [PubMed: 22889922]
- Carvalho-Wells AL, Jackson KG, Lockyer S, Lovegrove JA, Minihane AM. APOE genotype influences triglyceride and C-reactive protein responses to altered dietary fat intake in UK adults. *Am J Clin Nutr*. 2012; 96(6):1447–1453. [PubMed: 23134888]
- Caspi A, Sugden K, Moffitt TE, Taylor A, Craig IW, Harrington H, McClay J, Mill J, Martin J, Braithwaite A, Poulton R. Influence of life stress on depression: moderation by a polymorphism in the 5-HTT gene. *Science*. 2003; 301(5631):386–389. [PubMed: 12869766]
- Cornes BK, Lind PA, Medland SE, Montgomery GW, Nyholt DR, Martin NG. Replication of the association of common rs9939609 variant of FTO with increased BMI in an Australian adult twin population but no evidence for gene by environment (G [times] E) interaction. *Int J Obes*. 2008; 33(1):75–79.
- Dahl AK, Hassing LB, Fransson EI, Pedersen NL. Agreement between self-reported and measured height, weight and body mass index in old age—a longitudinal study with 20 years of follow-up. *Age Ageing*. 2010; 39(4):445–451. [PubMed: 20453247]
- Damoiseaux JS, Seeley WW, Zhou J, Shirer WR, Coppola G, Karydas A, Rosen HJ, Miller BL, Kramer JH, Greicius MD. gender modulates the APOE ε4 effect in healthy older adults: convergent evidence from functional brain connectivity and spinal fluid tau levels. *J Neurosci*. 2012; 32(24):8254–8262. [PubMed: 22699906]
- Davies G, Harris SE, Reynolds CA, Payton A, Knight HM, Liewald DC, Lopez LM, Luciano M, Gow AJ, Corley J, Henderson R, Murray C, Pattie A, Fox HC, Redmond P, Lutz MW, Chiba-Falek O, Linnertz C, Saith S, Haggarty P, McNeill G, Ke X, Ollier W, Horan M, Roses AD, Ponting CP, Porteous DJ, Tenesa A, Pickles A, Starr JM, Whalley LJ, Pedersen NL, Pendleton N, Visscher PM, Deary IJ. A genome-wide association study implicates the APOE locus in nonpathological cognitive ageing. *Mol Psychiatry*. 2014; 19(1):76–87. [PubMed: 23207651]
- De Marco M, Clough PJ, Dyer CE, Vince RV, Waby JS, Midgley AW, Venneri A. Apolipoprotein E epsilon4 allele modulates the immediate impact of acute exercise on prefrontal function. *Behav Genet*. 2015; 45(1):106–116. [PubMed: 25218559]
- Duncan LE, Keller MC. A critical review of the first 10 years of candidate gene-by-environment interaction research in psychiatry. *Am J Psychiatry*. 2011; 168(10):1041–1049. [PubMed: 21890791]
- Elosua R, Demissie S, Cupples LA, Meigs JB, Wilson PW, Schaefer EJ, Corella D, Ordovas JM. Obesity modulates the association among APOE genotype, insulin, and glucose in men. *Obes Res*. 2003; 11(12):1502–1508. [PubMed: 14694215]
- Ewbank DC. The APOE gene and differences in life expectancy in Europe. *J Gerontol A*. 2004; 59(1):B16–B20.
- Farrer LA, Cupples LA, Haines JL, Hyman B, Kukull WA, Mayeux R, Myers RH, Pericak-Vance MA, Risch N, van Duijn CM. Effects of age, sex, and ethnicity on the association between apolipoprotein E genotype and Alzheimer disease. A meta-analysis. APOE and Alzheimer disease meta analysis consortium. *JAMA*. 1997; 278(16):1349–1356. [PubMed: 9343467]
- Finkel D, Whitfield K, McGue M. Genetic and environmental influences on functional age: a twin study. *J Gerontol B Psychol Sci Soc Sci*. 1995; 50(2):P104–P113. [PubMed: 7757832]
- Finkel D, Reynolds CA, Larsson M, Gatz M, Pedersen NL. Both odor identification and ApoE-epsilon4 contribute to normative cognitive aging. *Psychol Aging*. 2011; 26(4):872–883. [PubMed: 21517181]
- Fisher RA. The resemblance between twins, a statistical examination of Lauterbach's measurements. *Genetics*. 1925; 10:569–579. [PubMed: 17246289]
- Folstein MF, Folstein SE, McHugh PR. "Mini-mental state": a practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res*. 1975; 12(3):189–198. [PubMed: 1202204]

- Gaetano, J. Holm-Bonferroni Sequential Correction: An EXCEL Calculator. 1.1.xlsx HsccB (ed). 2013. https://www.researchgate.net/publication/236969037_Holm-Bonferroni_Sequential_Correction_An_EXCEL_Calculator
- Gatz M, Reynolds CA, Finkel D, Hahn CJ, Zhou Y, Zavala C. for the IGEMS Consortium. Data harmonization in aging research: Not so fast. *Experimental aging research*. in press
- Gomez-Pinilla F, Hillman C. The influence of exercise on cognitive abilities. *Compr Physiol*. 2013; 3(1):403–428. [PubMed: 23720292]
- Grant MD, Kremen WS, Jacobson KC, Franz C, Xian H, Eisen SA, Toomey R, Murray RE, Lyons MJ. Does parental education have a moderating effect on the genetic and environmental influences of general cognitive ability in early adulthood? *Behav Genet*. 2010; 40(4):438–446. [PubMed: 20300818]
- Hamilton JM, Salmon DP, Galasko D, Raman R, Emond J, Hansen LA, Masliah E, Thal LJ. Visuospatial deficits predict rate of cognitive decline in autopsy-verified dementia with Lewy bodies. *Neuropsychology*. 2008; 22(6):729–737. [PubMed: 18999346]
- Head D, Bugg JM, Goate AM, Fagan AM, Mintun MA, Benzinger T, Holtzman DM, Morris JC. Exercise engagement as a moderator of the Effects of APOE genotype on amyloid deposition. *Arch Neurol*. 2012; 69(5):636–643. [PubMed: 22232206]
- Holm S. A simple sequentially rejective multiple test procedure. *Scand J Stat*. 1979; 6(2):65–70.
- Hotting K, Roder B. Beneficial effects of physical exercise on neuroplasticity and cognition. *Neurosci Biobehav Rev*. 2013; 37(9 Pt B):2243–2257. [PubMed: 23623982]
- Johnson W, Dahl A. for the IGEMS consortium. Education moderates genetic and environmental influences on body mass index: Findings from the consortium on Interplay of Genes and Environment across Multiple Studies (IGEMS). Gerontological Society of America, San Diego. 2012
- Johnson W, Kyvik KO, Skytthe A, Deary IJ, Sorensen TI. Education modifies genetic and environmental influences on BMI. *PLoS One*. 2011; 6(1):e16290. [PubMed: 21283825]
- Kaprio J. The finnish twin cohort study: an update. *Twin Res Hum Genet*. 2013; 16(1):157–162. [PubMed: 23298696]
- Kaprio J, Koskenvuo M. Genetic and environmental factors in complex diseases: the older Finnish Twin Cohort. *Twin Res*. 2002; 5(5):358–365. [PubMed: 12537859]
- Karg K, Burmeister M, Shedden K, Sen S. The serotonin transporter promoter variant (5-HTTLPR), stress, and depression meta-analysis revisited: evidence of genetic moderation. *Arch Gen Psychiatry*. 2011; 68(5):444–454. [PubMed: 21199959]
- Karlsson IK, Bennet AM, Ploner A, Andersson TML, Reynolds CA, Gatz M, Pedersen NL. Apolipoprotein E ε4 genotype and the temporal relationship between depression and dementia. *Neurobiol Aging*. 2015; 36:1751–1756. [PubMed: 25670333]
- Keller L, Xu W, Wang HX, Winblad B, Fratiglioni L, Graff C. The obesity related gene, FTO, interacts with APOE, and is associated with Alzheimer's disease risk: a prospective cohort study. *J Alzheimer's Dis*. 2011; 23(3):461–469. [PubMed: 21098976]
- Kendler KS, Kessler RC, Walters EE, MacLean C, Neale MC, Heath AC, Eaves LJ. Stressful life events, genetic liability, and onset of an episode of major depression in women. *Am J Psychiatry*. 1995; 152(6):833–842. [PubMed: 7755111]
- Kendler KS, Thornton LM, Gilman SE, Kessler RC. Sexual orientation in a U.S. national sample of twin and nontwin sibling pairs. *Am J Psychiatry*. 2000; 157(11):1843–1846. [PubMed: 11058483]
- Kimbrel NA, Hauser MA, Garrett M, Ashley-Koch A, Liu Y, Dennis MF, Klein RC, Veterans Affairs Mid-Atlantic Mental Illness Research Clinical E, Center W, Beckham JC. Effect of the APOE epsilon4 allele and combat exposure on PTSD among Iraq/Afghanistan-Era Veterans. *Depression and anxiety*. 2015
- Kremen WS, Jacobson KC, Xian H, Eisen SA, Waterman B, Toomey R, Neale MC, Tsuang MT, Lyons MJ. Heritability of word recognition in middle-aged men varies as a function of parental education. *Behav Genet*. 2005; 35(4):417–433. [PubMed: 15971023]
- Kremen WS, Franz CE, Lyons MJ. VETSA: the Vietnam era twin study of aging. *Twin Res Hum Genet*. 2013; 16(1):399–402. [PubMed: 23110957]

- Lajunen HR, Kaprio J, Rose RJ, Pulkkinen L, Silventoinen K. Genetic and environmental influences on BMI from late childhood to adolescence are modified by parental education. *Obesity* (Silver Spring). 2012; 20(3):583–589. [PubMed: 21996670]
- Lichtenstein P, Sullivan PF, Cnattingius S, Gatz M, Johansson S, Carlstrom E, Bjork C, Svartengren M, Wolk A, Klareskog L, de Faire U, Schalling M, Palmgren J, Pedersen NL. The Swedish twin registry in the third millennium: an update. *Twin Res Hum Genet*. 2006; 9(6):875–882. [PubMed: 17254424]
- Locke DE, Dueck AC, Stonnington CM, Knopman DS, Geda YE, Caselli RJ. Depressive symptoms in healthy apolipoprotein E epsilon4 carriers and noncarriers: a longitudinal study. *J Clin Psychiatry*. 2013; 74(12):1256–1261. [PubMed: 24434095]
- Locke AE, Kahali B, Berndt SI, Justice AE, Pers TH, Day FR, Powell C, Vedantam S, Buchkovich ML, Yang J, Croteau-Chonka DC, Esko T, Fall T, Ferreira T, Gustafsson S, Kutalik Z, Luan J, Magi R, Randall JC, Winkler TW, Wood AR, Workalemahu T, Faul JD, Smith JA, Hua Zhao J, Zhao W, Chen J, Fehrmann R, Hedman AK, Karjalainen J, Schmidt EM, Absher D, Amin N, Anderson D, Beekman M, Bolton JL, Bragg-Gresham JL, Buyske S, Demirkan A, Deng G, Ehret GB, Feenstra B, Feitosa MF, Fischer K, Goel A, Gong J, Jackson AU, Kanoni S, Kleber ME, Kristiansson K, Lim U, Lotay V, Mangino M, Mateo Leach I, Medina-Gomez C, Medland SE, Nalls MA, Palmer CD, Pasko D, Pechlivanis S, Peters MJ, Prokopenko I, Shungin D, Stancakova A, Strawbridge RJ, Ju Sung Y, Tanaka T, Teumer A, Trompet S, van der Laan SW, van Setten J, Van Vliet-Ostaptchouk JV, Wang Z, Yengo L, Zhang W, Isaacs A, Albrecht E, Arnlöv J, Arscott GM, Attwood AP, Bandinelli S, Barrett A, Bas IN, Bellis C, Bennett AJ, Berne C, Blagieva R, Bluher M, Bohringer S, Bonnycastle LL, Bottcher Y, Boyd HA, Bruinenberg M, Caspersen IH, Ida Chen YD, Clarke R, Daw EW, de Craen AJ, Delgado G, Dimitriou M, Doney AS, Eklund N, Estrada K, Eury E, Folkersen L, Fraser RM, Garcia ME, Geller F, Giedraitis V, Gigante B, Go AS, Golay A, Goodall AH, Gordon SD, Gorski M, Grabe HJ, Grallert H, Grammer TB, Grassler J, Gronberg H, Groves CJ, Gusto G, Haessler J, Hall P, Haller T, Hallmans G, Hartman CA, Hassinen M, Hayward C, Heard-Costa NL, Helmer Q, Hengstenberg C, Holmen O, Hottenga JJ, James AL, Jeff JM, Johansson A, Jolley J, Juliusdottir T, Kinnunen L, Koenig W, Koskenvuo M, Kratzer W, Laitinen J, Lamina C, Leander K, Lee NR, Lichtner P, Lind L, Lindstrom J, Sin Lo K, Lobbens S, Lorbeer R, Lu Y, Mach F, Magnusson PK, Mahajan A, McArdle WL, McLachlan S, Menni C, Merger S, Mihailov E, Milani L, Moayyeri A, Monda KL, Morken MA, Mulas A, Muller G, Muller-Nurasyid M, Musk AW, Nagaraja R, Nothen MM, Nolte IM, Pilz S, Rayner NW, Renstrom F, Rettig R, Ried JS, Ripke S, Robertson NR, Rose LM, Sanna S, Scharnagl H, Scholtens S, Schumacher FR, Scott WR, Seufferlein T, Shi J, Vernon Smith A, Smolonska J, Stanton AV, Steinthorsdottir V, Stirrups K, Stringham HM, Sundstrom J, Swertz MA, Swift AJ, Syvanen AC, Tan ST, Tayo BO, Thorand B, Thorleifsson G, Tyrer JP, Uh HW, Vandenput L, Verhulst FC, Vermeulen SH, Verweij N, Vonk JM, Waite LL, Warren HR, Waterworth D, Weedon MN, Wilkens LR, Willenborg C, Wilsaard T, Wojczynski MK, Wong A, Wright AF, Zhang Q, LifeLines Cohort S, Brennan EP, Choi M, Dastani Z, Drong AW, Eriksson P, Franco-Cereceda A, Gadin JR, Gharavi AG, Goddard ME, Handsaker RE, Huang J, Karpe F, Kathiresan S, Keildson S, Kiryluk K, Kubo M, Lee JY, Liang L, Lifton RP, Ma B, McCarroll SA, McKnight AJ, Min JL, Moffatt MF, Montgomery GW, Murabito JM, Nicholson G, Nyholt DR, Okada Y, Perry JR, Dorajoo R, Reinmaa E, Salem RM, Sandholm N, Scott RA, Stolk L, Takahashi A, Tanaka T, Van't Hooft FM, Vinkhuyzen AA, Westra HJ, Zheng W, Zondervan KT, Consortium AD, Group A-BW, Consortium CAD, Consortium CK, Gluc Icbp, Investigators M, Mu TC, Consortium MI, Consortium P, ReproGen C, Consortium G, International Endogene C, Heath AC, Arveiler D, Bakker SJ, Beilby J, Bergman RN, Blangero J, Bovet P, Campbell H, Caulfield MJ, Cesana G, Chakravarti A, Chasman DI, Chines PS, Collins FS, Crawford DC, Cupples LA, Cusi D, Danesh J, de Faire U, den Ruijter HM, Dominiczak AF, Erbel R, Erdmann J, Eriksson JG, Farrall M, Felix SB, Ferrannini E, Ferrieres J, Ford I, Forouhi NG, Forrester T, Franco OH, Gansevoort RT, Gejman PV, Gieger C, Gottesman O, Gudnason V, Gyllenstein U, Hall AS, Harris TB, Hattersley AT, Hicks AA, Hindorf LA, Hingorani AD, Hofman A, Homuth G, Hovingh GK, Humphries SE, Hunt SC, Hypponen E, Illig T, Jacobs KB, Jarvelin MR, Jockel KH, Johansen B, Jousilahti P, Jukema JW, Julia AM, Kaprio J, Kastelein JJ, Keinänen-Kiukkaanniemi SM, Kiemeny LA, Knekt P, Kooner JS, Kooperberg C, Kovacs P, Kraja AT, Kumari M, Kuusisto J, Lakka TA, Langenberg C, Le Marchand L, Lehtimäki T, Lyssenko V, Mannisto S, Marette A, Matise TC, McKenzie CA, McKnight B, Moll FL, Morris AD, Morris AP, Murray JC, Nelis M, Ohlsson C, Oldehinkel AJ,

- Ong KK, Madden PA, Pasterkamp G, Peden JF, Peters A, Postma DS, Pramstaller PP, Price JF, Qi L, Raitakari OT, Rankinen T, Rao DC, Rice TK, Ridker PM, Rioux JD, Ritchie MD, Rudan I, Salomaa V, Samani NJ, Saramies J, Sarzynski MA, Schunkert H, Schwarz PE, Sever P, Shuldiner AR, Sinisalo J, Stolk RP, Strauch K, Tonjes A, Tregouet DA, Tremblay A, Tremoli E, Virtamo J, Vohl MC, Volker U, Waeber G, Willemssen G, Witteman JC, Zillikens MC, Adair LS, Amouyel P, Asselbergs FW, Assimes TL, Bochud M, Boehm BO, Boerwinkle E, Bornstein SR, Bottinger EP, Bouchard C, Cauchi S, Chambers JC, Chanock SJ, Cooper RS, de Bakker PI, Dedoussis G, Ferrucci L, Franks PW, Froguel P, Groop LC, Haiman CA, Hamsten A, Hui J, Hunter DJ, Hveem K, Kaplan RC, Kivimaki M, Kuh D, Laakso M, Liu Y, Martin NG, Marz W, Melbye M, Metspalu A, Moebus S, Munroe PB, Njolstad I, Oostra BA, Palmer CN, Pedersen NL, Perola M, Perusse L, Peters U, Power C, Quertermous T, Rauramaa R, Rivadeneira F, Saaristo TE, Saleheen D, Sattar N, Schadt EE, Schlessinger D, Slagboom PE, Snieder H, Spector TD, Thorsteinsdottir U, Stumvoll M, Tuomilehto J, Uitterlinden AG, Uusitupa M, van der Harst P, Walker M, Wallaschofski H, Wareham NJ, Watkins H, Weir DR, Wichmann HE, Wilson JF, Zanen P, Borecki IB, Deloukas P, Fox CS, Heid IM, O'Connell JR, Strachan DP, Stefansson K, van Duijn CM, Abecasis GR, Franke L, Frayling TM, McCarthy MI, Visscher PM, Scherag A, Willer CJ, Boehnke M, Mohlke KL, Lindgren CM, Beckmann JS, Barroso I, North KE, Ingelsson E, Hirschhorn JN, Loos RJ, Speliotes EK. Genetic studies of body mass index yield new insights for obesity biology. *Nature*. 2015; 518(7538):197–206. [PubMed: 25673413]
- Lyons MJ, Genderson M, Grant MD, Logue M, Zink T, McKenzie R, Franz CE, Panizzon M, Lohr JB, Jerskey B, Kremen WS. Gene–environment interaction of ApoE genotype and combat exposure on PTSD. *Am J Med Gen B Neuropsych Gen*. 2013; 162B(7):762–769.
- Magnusson PK, Almqvist C, Rahman I, Ganna A, Viktorin A, Walum H, Halldner L, Lundstrom S, Ullen F, Langstrom N, Larsson H, Nyman A, Gumpert CH, Rastam M, Anckarsater H, Cnattingius S, Johannesson M, Ingelsson E, Klareskog L, de Faire U, Pedersen NL, Lichtenstein P. The Swedish twin registry: establishment of a biobank and other recent developments. *Twin Res Hum Genet*. 2013; 16(1):317–329. [PubMed: 23137839]
- Martin, NG. Gene–environment interaction and twin studies. In: Spector, T.; Snieder, H.; Gregor, Mac-A., editors. *Advances in twin and sib-pair analysis*. London: Greenwich Medical Media Ltd; 2000. p. 143-150.
- Martin NG, Rowell DM, Whitfield JB. Do the MN and Jk systems influence environmental variability in serum lipid levels? *Clin Genet*. 1983; 24(1):1–14. [PubMed: 6684513]
- McCaffery JM, Papandonatos GD, Bond DS, Lyons MJ, Wing RR. Gene X environment interaction of vigorous exercise and body mass index among male Vietnam-era twins. *Am J Clin Nutr*. 2009; 89(4):1011–1018. [PubMed: 19225119]
- McCleary GE, Johansson B, Berg S, Pedersen NL, Ahern F, Petrill SA, Plomin R. Substantial genetic influence on cognitive abilities in twins 80 or more years old. *Science*. 1997; 276(5318):1560–1563. [PubMed: 9171059]
- McGue M, Christensen K. Genetic and environmental contributions to depression symptomatology: evidence from Danish twins 75 years of age and older. *J Abnorm Psychol*. 1997; 106(3):439–448. [PubMed: 9241945]
- McGue M, Christensen K. Growing old but not growing apart: Twin similarity in the latter half of the lifespan. *Behav Gen*. 2013; 43:1–12.
- Mustelin L, Silventoinen K, Pietilainen K, Rissanen A, Kaprio J. Physical activity reduces the influence of genetic effects on BMI and waist circumference: a study in young adult twins. *Int J Obes*. 2009; 33(1):29–36.
- Nebes RD, Vora JJ, Meltzer CC, Fukui MB, Williams RL, Kamboh MI, Saxton J, Houck PR, DeKosky ST, Reynolds CF. Relationship of deep white matter hyperintensities and apolipoprotein E genotype to depressive symptoms in older adults without clinical depression. *Am J Psychiatry*. 2001; 158(6):878–884. [PubMed: 11384894]
- Neiderhiser JA, Reiss D, Lichtenstein P, Spotts EL, Ganiban J. Father-adolescent relationships and the role of genotype-environment correlation. *J Fam Psychol*. 2007; 21(4):560–571. [PubMed: 18179328]
- Pahlen S, Hamdi NR, Dahl Aslan AK, Horwitz BN, Panizzon MS, Petersen I, Zavala C, Christensen K, Finkel D, Franz CE, Gatz M, Johnson W, Kremen WS, Krueger RF, Neiderhiser JM, Reynolds

CA, Pedersen NL, McGue M. Widespread evidence of age-moderation of genetic and environmental contributions to cognitive functioning in mid- and late-life. *Psychol Sci.* under review

Pedersen NL, McClearn GE, Plomin R, Nesselroade JR, Berg S, DeFaire U. The Swedish Adoption twin study of aging: an update. *Acta Genet Med Gemellol.* 1991; 40(1):7–20. [PubMed: 1950353]

Pedersen NL, Christensen K, Dahl AK, Finkel D, Franz CE, Gatz M, Horwitz BN, Johansson B, Johnson W, Kremen WS, Lyons MJ, Malmberg B, McGue M, Neiderhiser JM, Petersen I, Reynolds CA. IGEMS: the consortium on interplay of genes and environment across multiple studies. *Twin Res Hum Genet.* 2013; 16(1):481–489. [PubMed: 23186995]

Radler BT. The midlife in the United States (MIDUS) series: A national longitudinal study of health and well-being. *Open health data.* 2014; 2(1):e3. [PubMed: 25558376]

Radloff LS. The CES-D scale a self-report depression scale for research in the general population. *Appl Psychol Meas.* 1977; 1(3):385–401.

Reynolds CA, Finkel D, McArdle JJ, Gatz M, Berg S, Pedersen NL. Quantitative genetic analysis of latent growth curve models of cognitive abilities in adulthood. *Dev Psychol.* 2005; 41(1):3–16. [PubMed: 15656733]

Reynolds CA, Prince JA, Feuk L, Brookes AJ, Gatz M, Pedersen NL. Longitudinal memory performance during normal aging: twin association models of APOE and other Alzheimer candidate genes. *Behav Genet.* 2006; 36(2):185–194. [PubMed: 16402284]

Reynolds CA, Gatz M, Berg S, Pedersen NL. Genotype-environment interactions: cognitive aging and social factors. *Twin Res Hum Genet.* 2007; 10(2):241–254. [PubMed: 17564514]

Reynolds CA, Zavala C, Gatz M, Vie L, Johansson B, Malmberg B, Ingelsson E, Prince JA, Pedersen NL. Sortilin receptor 1 predicts longitudinal cognitive change. *Neurobiol Aging.* 2013; 34(6): 1710.e1711–1710.e1718. [PubMed: 23318115]

Rokholm B, Silventoinen K, Tynelius P, Gamborg M, Sørensen TIA, Rasmussen F. Increasing genetic variance of body mass index during the swedish obesity epidemic. *PLoS One.* 2011; 6(11):e27135. [PubMed: 22087252]

Rosenthal, R. Meta-analytic procedures for social research. Newbury Park: Sage; 1991.

Rutter M. Achievements and challenges in the biology of environmental effects. *Proc Natl Acad Sci USA.* 2012; 109(2):17149–17153. [PubMed: 23045650]

Rutter M, Silberg J. Gene-environment interplay in relation to emotional and behavioral disturbance. *Annu Rev Psychol.* 2002; 53:463–490. [PubMed: 11752493]

Salmon DP, Ferris SH, Thomas RG, Sano M, Cummings JL, Sperling RA, Petersen RC, Aisen PS. Age and APOE genotype influence rate of cognitive decline in non-demented elderly. *Neuropsychology.* 2013

Salthouse TA. When does age-related cognitive decline begin? *Neurobiol Aging.* 2009; 30(4):507–514. [PubMed: 19231028]

Schaie KW. The course of adult intellectual development. *Am Psychol.* 1994; 49(4):304–313. [PubMed: 8203802]

Schellenberg GD, Montine TJ. The genetics and neuropathology of Alzheimer's disease. *Acta Neuropathol.* 2012; 124:305–323. [PubMed: 22618995]

Schultz MR, Lyons MJ, Franz CE, Grant MD, Boake C, Jacobson KC, Xian H, Schellenberg GD, Eisen SA, Kremen WS. Apolipoprotein E genotype and memory in the sixth decade of life. *Neurology.* 2008; 70(19 Pt 2):1771–1777. [PubMed: 18235080]

Silventoinen K, Sarlio-Lahteenkorva S, Koskenvuo M, Lahelma E, Kaprio J. Effect of environmental and genetic factors on education-associated disparities in weight and weight gain: a study of Finnish adult twins. *Am J Clin Nutr.* 2004; 80(4):815–822. [PubMed: 15447885]

Silventoinen K, Hasselbalch AL, Lallukka T, Bogl L, Pietilainen KH, Heitmann BL, Schousboe K, Rissanen A, Kyvik KO, Sorensen TI, Kaprio J. Modification effects of physical activity and protein intake on heritability of body size and composition. *Am J Clin Nutr.* 2009; 90(4):1096–1103. [PubMed: 19710197]

Skoog I, Waern M, Duberstein P, Blennow K, Zetterberg H, Börjesson-Hanson A, Östling S, Guo X, Kern J, Gustafson D, Gudmundsson P, Marlow T, Kern S. A 9-year prospective population-based

study on the association between the APOE*E4 allele and late-life depression in Sweden. *Biol Psychiatry*. 2015

- Skytthe A, Christiansen L, Kyvik KO, Bodker FL, Hvidberg L, Petersen I, Nielsen MM, Bingley P, Hjelmborg J, Tan Q, Holm NV, Vaupel JW, McGue M, Christensen K. The Danish Twin Registry: linking surveys, national registers, and biological information. *Twin Res Hum Genet*. 2013; 16(1): 104–111. [PubMed: 23084092]
- South SC, Krueger RF. Genetic and environmental influences on internalizing psychopathology vary as a function of economic status. *Psychol Med*. 2011; 41(1):107–117. [PubMed: 20236567]
- Sundstrom A, Marklund P, Nilsson LG, Cruts M, Adolfsson R, Van Broeckhoven C, Nyberg L. APOE influences on neuropsychological function after mild head injury: within-person comparisons. *Neurology*. 2004; 62(11):1963–1966. [PubMed: 15184597]
- Sundstrom A, Nilsson LG, Cruts M, Adolfsson R, Van Broeckhoven C, Nyberg L. Increased risk of dementia following mild head injury for carriers but not for non-carriers of the APOE epsilon4 allele. *Int Psychoger*. 2007; 19(1):159–165.
- Surakka I, Whitfield JB, Perola M, Visscher PM, Montgomery GW, Falchi M, Willemsen G, de Geus EJ, Magnusson PK, Christensen K, Sorensen TI, Pietilainen KH, Rantanen T, Silander K, Widen E, Muilu J, Rahman I, Liljedahl U, Syvanen AC, Palotie A, Kaprio J, Kyvik KO, Pedersen NL, Boomsma DI, Spector T, Martin NG, Ripatti S, Peltonen L, Genom EP. A genome-wide association study of monozygotic twin-pairs suggests a locus related to variability of serum high-density lipoprotein cholesterol. *Twin Res Hum Genet*. 2012; 15(6):691–699. [PubMed: 23031429]
- Surtees PG, Wainwright NWJ, Bowman R, Luben RN, Wareham NJ, Khaw K-T, Bingham SA. No association between APOE and major depressive disorder in a community sample of 17,507 adults. *J Psychiatr Res*. 2009; 43(9):843–847. [PubMed: 19135213]
- Tabert MH, Manly JJ, Liu X, Pelton GH, Rosenblum S, Jacobs M, Zamora D, Goodkind M, Bell K, Stern Y, Devanand DP. Neuropsychological prediction of conversion to Alzheimer disease in patients with mild cognitive impairment. *Arch Gen Psychiatry*. 2006; 63(8):916–924. [PubMed: 16894068]
- Tang MX, Maestre G, Tsai WY, Liu XH, Feng L, Chung WY, Chun M, Schofield P, Stern Y, Tycko B, Mayeux R. Effect of age, ethnicity, and head injury on the association between APOE genotypes and Alzheimer's disease. *Ann N Y Acad Sci*. 1996; 802:6–15. [PubMed: 8993479]
- Taylor WD, Aizenstein HJ, Alexopoulos GS. The vascular depression hypothesis: mechanisms linking vascular disease with depression. *Mol Psychiatry*. 2013; 18(9):963–974. [PubMed: 23439482]
- Turkheimer, E.; Horn, EE. Interactions between socioeconomic status and components of variation in cognitive ability. In: Finkel, D.; Reynolds, CA., editors. *Behavior genetics of cognition across the lifespan*. Riverside: Springer; 2014. p. 41-68.
- Vinkhuyzen AA, van der Sluis S, Posthuma D. Interaction between experience seeking and genetic and environmental influences on general cognitive ability. *Twin Res Hum Genet*. 2012; 15(1):87–96. [PubMed: 22784458]
- Visscher PM, Posthuma D. Statistical power to detect genetic Loci affecting environmental sensitivity. *Behav Genet*. 2010; 40(5):728–733. [PubMed: 20428936]
- Ward A, Crean S, Mercaldi CJ, Collins JM, Boyd D, Cook MN, Arrighi HM. Prevalence of apolipoprotein E4 genotype and homozygotes (APOE e4/4) among patients diagnosed with Alzheimer's disease: a systematic review and meta-analysis. *Neuroepidemiology*. 2012; 38(1):1–17. [PubMed: 22179327]
- Weinkauff, M. BenjaminiHochberg.xlsx (ed). 2012. False discovery rate Excel spreadsheet (41,86 KB) —Marum. <https://www.marum.de/Binaries/Binary745/BenjaminiHochberg.xlsx>

Fig. 1.
Effect size r (ES r) for evidence for mixture distribution suggesting possible G×E:
representative cognitive tests

Fig. 2. Variance of absolute MZ within pair differences adjusted for age by *APOE*: **a** depressive symptoms, **b** block design

Table 1

MZ pairs contributing to G×E Analyses

Study	Country	Sex	BMI		Depressive Sx				Cognitive (1+ test)					
			N	N _{AOPE}	Mean age	SD	N	N _{AOPE}	Mean age	SD	N	N _{AOPE}	Mean age	SD
VETSA	USA	M	349	340	55.35	2.53	346	337	55.36	2.52	347	339	55.33	2.49
MTSADA	USA	M	69	–	56.70	12.26	117	–	58.98	9.93	66	–	56.19	12.12
MIDUS	USA	F	150	–	54.33	13.35	210	–	56.65	12.33	149	–	54.41	13.08
		M	132	–	45.56	11.46	33	–	54.70	11.70	83	–	55.19	11.12
SATSA	SWE	F	155	–	44.35	12.39	48	–	52.63	10.98	96	–	52.87	11.56
		M	112	52	55.69	13.87	102	53	57.00	13.40	59	52	62.69	7.45
Octo-Twin	SWE	F	138	70	59.80	13.59	116	72	59.97	13.28	83	73	64.15	9.03
		M	41	37	82.81	2.47	47	43	82.59	2.90	42	38	82.88	2.63
TOSS	SWE	F	66	62	83.48	2.95	67	63	83.19	3.27	55	50	82.83	2.31
		M	120	–	46.48	4.50	101	–	46.57	4.53	121	–	46.50	4.49
MADT	DEN	F	104	–	43.40	5.27	213	–	43.69	4.64	259	–	43.68	4.66
		M	335	191	56.57	6.41	333	191	56.54	6.40	330	189	56.45	6.40
LSADT	DEN	F	327	198	56.36	6.41	328	197	56.36	6.41	326	195	56.33	6.41
		M	171	52	75.05	4.57	172	51	74.92	4.54	132	37	74.31	3.73
FTC	FIN	F	274	98	76.05	4.79	266	97	75.87	4.73	190	77	74.95	4.03
		M	405	–	60.14	3.69	407	–	59.71	3.70	–	–	–	–
Total pairs	–	F	602	–	59.85	3.69	602	–	59.38	3.69	–	–	–	–
		–	3550	1100	–	–	3508	1104	–	–	2338	1050	–	–

Sx Symptoms, USA United States of America, SWE Sweden, DEN Denmark, FIN Finland

Table 2
Test of mixture distributions of MZ within pair differences in full sample and by age-cohort

	$\bar{d}$	$\overline{d^2}$	delta	se	t	p	p'	ESr
BMI (Npair)								
Full sample (3550)	0.61	0.65	0.07	0.01	12.26	3.54E-34	4.25E-33	0.20
<50 years (643)	0.51	0.45	0.04	0.01	4.65	2.05E-06	8.21E-06	0.18
50-59 (1351)	0.60	0.64	0.07	0.01	7.93	2.36E-15	2.12E-14	0.21
60-69 (929)	0.61	0.63	0.05	0.01	4.49	3.97E-06	1.19E-05	0.15
70-79 (429)	0.69	0.84	0.09	0.02	4.09	2.60E-05	5.21E-05	0.19
80+ (198)	0.76	1.01	0.10	0.04	2.49	6.87E-03	6.87E-03	0.17
Depressive Sx (Npair)								
Full sample (3508)	0.81	1.19	0.14	0.01	13.61	1.99E-41	2.39E-40	0.22
<50 years (550)	0.80	1.12	0.13	0.03	4.99	4.06E-07	2.03E-06	0.21
50-59 (1326)	0.81	1.16	0.14	0.02	7.98	1.57E-15	1.25E-14	0.21
60-69 (995)	0.77	1.08	0.14	0.02	7.88	4.37E-15	3.06E-14	0.24
70-79 (436)	0.92	1.51	0.17	0.04	4.52	3.91E-06	1.57E-05	0.21
80+ (201)	0.86	1.30	0.14	0.05	2.80	2.81E-03	2.81E-03	0.19
Synonyms (Npair)								
Full sample (912)	0.64	0.68	0.04	0.01	3.53	2.16E-04	2.16E-03	0.12
<50 years (318)	0.56	0.50	0.01	0.01	0.92	1.79E-01	3.59E-01	0.05
50-59 years (463)	0.69	0.79	0.03	0.02	1.74	4.13E-02	2.39E-01	0.08
60-69 years (48)	0.57	0.64	0.13	0.05	2.71	4.64E-03	3.25E-02	0.37
70-79 years (32)	0.64	0.62	-0.01	0.06	-0.19	5.75E-01	5.75E-01	0.03
80+ years (51)	0.72	0.94	0.13	0.07	1.79	3.98E-02	2.39E-01	0.25
Block design (Npair)								
Full sample (393)	0.56	0.57	0.07	0.02	4.61	2.75E-06	2.75E-05	0.23
<50 years (60)	0.46	0.37	0.04	0.03	1.43	7.85E-02	1.57E-01	0.18
50-59 years (85)	0.47	0.37	0.03	0.02	1.36	8.94E-02	1.57E-01	0.15
60-69 years (124)	0.59	0.60	0.06	0.03	1.98	2.47E-02	1.23E-01	0.18
70-79 years (37)	0.59	0.66	0.11	0.06	1.84	3.69E-02	1.23E-01	0.29
80+ years (87)	0.68	0.82	0.09	0.05	1.89	3.08E-02	1.23E-01	0.20

	$\bar{d}$	$\overline{d^2}$	delta	se	t	p	p'	ESr
Digits forward (Npair)								
Full Sample (1551)	0.83	1.21	0.12	0.02	7.28	2.60E-13	2.86E-12	0.18
<50 years (124)	0.91	1.39	0.07	0.07	1.13	1.31E-01	1.94E-01	0.10
50-59 years (695)	0.80	1.13	0.12	0.02	5.31	7.54E-08	6.78E-07	0.20
60-69 years (285)	0.79	1.01	0.04	0.03	1.30	9.69E-02	1.94E-01	0.08
70-79 years (313)	0.94	1.51	0.12	0.05	2.56	5.52E-03	1.66E-02	0.14
80+ years (134)	0.80	1.26	0.26	0.06	4.46	8.67E-06	5.20E-05	0.36
Digits backward (Npair)								
Full sample (1722)	0.83	1.21	0.14	0.02	8.81	1.51E-18	1.66E-17	0.21
<50 years (194)	0.82	1.25	0.18	0.05	3.84	8.23E-05	3.20E-04	0.27
50-59 years (745)	0.82	1.18	0.12	0.02	5.38	4.87E-08	3.41E-07	0.19
60-69 years (323)	0.82	1.11	0.06	0.03	1.83	3.42E-02	3.42E-02	0.10
70-79 years (327)	0.88	1.39	0.16	0.04	3.82	8.01E-05	3.20E-04	0.21
80+ years (133)	0.74	1.10	0.25	0.05	4.92	1.28E-06	7.70E-06	0.39
Symbol digit (Npair)								
Full sample (1256)	0.58	0.57	0.04	0.01	5.17	1.39E-07	1.39E-06	0.14
<50 years (190)	0.54	0.51	0.04	0.02	2.26	1.26E-02	7.56E-02	0.16
50-59 years (360)	0.59	0.61	0.07	0.02	4.20	1.66E-05	1.49E-04	0.22
60-69 years (371)	0.59	0.58	0.03	0.02	1.86	3.18E-02	1.59E-01	0.10
70-79 years (256)	0.57	0.54	0.03	0.02	1.85	3.25E-02	1.59E-01	0.12
80+ years (79)	0.63	0.63	0.01	0.04	0.33	3.70E-01	8.92E-01	0.04

D absolute pair difference, $\overline{d^2} = \frac{\pi - d^2}{2}$; $se = \frac{\overline{d^2}}{\sqrt{n}}$ (*.5321*), ESr effect size $r = \sqrt{\frac{t^2}{(t^2 + df)}}$, where $d \neq N_{pair} - 1$. p -values are based on one-tailed t -tests; bolded p values are significant according to FDR tests. $p' = \text{Holm-Bonferroni sequentially adjusted } p \text{ values, where bolded are significant}$

Table 3

Homogeneity of within pair variance by *APOE*

Measure	Country	Within pair σ^2		Likelihood ratio tests						Total Npair				
		APOE		Equate males & females										
		e2+	e33	e4+	χ^2	df	p	χ^2	df		p	χ^2	df	p
BMI	USA (m)	0.39	0.30	0.31	19.64	6	3.21E-03	10.85	6	9.31E-02	3.67	6	7.22E-01	672
	SWE (m)	0.32	0.26	0.32	-	-	-	-	-	-	-	-	-	-
	DEN (m)	0.24	0.19	0.24	-	-	-	-	-	-	-	-	-	-
	SWE (f)	0.26	0.45	0.52	-	-	-	4.50	3	2.12E-01	6.53	4	1.63E-01	428
	DEN (f)	0.30	0.31	0.40	-	-	-	-	-	-	-	-	-	-
	USA	0.91	0.44	0.39	2.05	6	9.15E-01	44.99	6	4.70E-08	19.78	6	3.04E-03	1104
Depressive Sx	SWE	0.55	0.47	0.41	-	-	-	-	-	-	-	-	-	-
	DEN	0.76	0.71	0.91	-	-	-	-	-	-	-	-	-	-
	USA	0.45	0.43	0.38	1.91	3	5.92E-01	1.62	3	6.54E-01	1.77	4	7.79E-01	506
Synonyms	SWE	0.32	0.40	0.32	-	-	-	-	-	-	-	-	-	-
	SWE	0.40	0.33	0.15	5.76	3	1.24E-01	-	-	-	11.91	2	2.60E-03	203
	USA	0.51	0.46	0.50	3.32	6	7.68E-01	9.24	6	1.60E-01	4.58	6	5.98E-01	1038
Digits forward	SWE	0.53	0.66	0.63	-	-	-	-	-	-	-	-	-	-
	DEN	0.66	0.64	0.49	-	-	-	-	-	-	-	-	-	-
	USA	0.43	0.63	0.52	6.73	6	3.46E-01	10.04	6	1.23E-01	6.71	6	3.48E-01	1031
Digits back	SWE	0.32	0.47	0.44	-	-	-	-	-	-	-	-	-	-
	DEN	0.57	0.51	0.58	-	-	-	-	-	-	-	-	-	-
	SWE	0.24	0.30	0.31	6.03	6	4.20E-01	0.08	3	9.94E-01	1.71	4	7.88E-01	618
Symbol digit	DEN	0.26	0.31	0.31	-	-	-	-	-	-	-	-	-	-

USA United States of America, SWE Sweden, DEN Denmark, m male, f female, Sx symptoms. Random effects model adjusted for average effect of *APOE* haplotype, country, sex and age

Table 4

Average effects of *APOE*: adjusted for age, sex and country

Measure	<i>APOE</i> haplotype			Test of <i>APOE</i> effect			Npair
	e2+	e33	e4+	χ^2	df	p	
BMI	-0.19	-0.12	-0.13	1.04	2	5.94E-01	1100
Depressive Sx	-0.62	-0.53	-0.50	2.53	2	2.83E-01	1104
Synonyms	-0.06	0.09	0.03	2.33	2	3.12E-01	506
Block design	0.04	0.29	0.26	2.78	2	2.49E-01	364
Digits forward	-0.04	-0.02	0.09	3.98	2	1.37E-01	1038
Digits backward	0.12	0.08	0.18	2.81	2	2.45E-01	1031
Symbol digit	0.17	0.06	0.18	3.78	2	1.51E-01	618

Sx symptoms. Random effects models adjusted for average effects of age (centered at 65 years), sex (males = -0.5, females = +0.5), and country (reference = Denmark)