Published in final edited form as:

Int J Indig Health. 2015 ; 10(2): 88–101.

Formative Evaluation to Assess Communication Technology Access and Health Communication Preferences of Alaska Native People

Renee F. Robinson, Denise A. Dillard, Vanessa Y. Hiratsuka, Julia J. Smith, Steve Tierney, Jaedon P. Avey, and Dedra S. Buchwald

Abstract

Objective—Information technology can improve the quality, safety, and efficiency of healthcare delivery by improving provider and patient access to health information. We conducted a nonrandomized, cross-sectional, self-report survey to determine whether Alaska Native and American Indian (AN/AI) people have access to the health communication technologies available through a patient-centered medical home.

Methods—In 2011, we administered a self-report survey in an urban, tribally owned and operated primary care center serving AN/AI adults. Patients in the center's waiting rooms completed the survey on paper; center staff completed it electronically.

Results—Approximately 98% ($n = 654$) of respondents reported computer access, 97% ($n = 650$) email access, and 94% ($n = 631$) mobile phone use. Among mobile phone users, 60% had Internet access through their phones. Rates of computer access ($p = .011$) and email use ($p = .005$) were higher among women than men, but we found no significant gender difference in mobile phone access to the Internet or text messaging. Respondents in the oldest age category (65–80 years of age) were significantly less likely to anticipate using the Internet to schedule appointments, refill medications, or communicate with their health care providers (all $p < .001$).

Conclusion—Information on use of health communication technologies enables administrators to deploy these technologies more efficiently to address health concerns in AN/AI communities. Our results will drive future research on health communication for chronic disease screening and health management.

Keywords

Healthcare communication; technology access; colorectal cancer; disease screening; Alaska

Introduction

Southcentral Foundation (SCF), a nonprofit tribal healthcare organization, serves more than 65,000 Alaska Native and American Indian (AN/AI) people living in SCF's region of Alaska, including Anchorage, Alaska and 60 rural villages (D. Eby, 1998; D. K. Eby, 2007).

SCF is an urban, tribally owned and operated health center that provides prepaid primary care services. Given its service population, SCF is committed to recruiting AN/AI employees, who make up 54% of a total workforce of approximately 1,400. Through patient engagement, SCF has developed a unique healthcare delivery system, the Nuka System of Care, based on the patient-centered medical home model (D. K. Eby, 2007; Thompson, 2008). This system involves patients, their families, and their providers in every stage of healthcare delivery, while taking into account cultural traditions, social circumstances, family situations, personal preferences and values, and lifestyles (Burke, Menachemi, & Brooks, 2005; Sepucha et al., 2008; Thompson, 2008).

A focus on patients drives clinical practice, system redesign, and program implementation at the SCF. To bring tangible improvements in healthcare, new technologies must be acceptable to the target population, which includes providers as well as patients, and must respond to community healthcare needs (Brown, Copas, Stephenson, Gilleran, & Ross, 2008; Sahn, MacCurtain, Hayden, Roche, & Richards, 2009). Equally important, technological innovation must address the so-called digital divide, which is defined as inequalities in access to information and communication technology stemming from socioeconomic status and geographic residence (Hsu et al., 2005; Kind, Huang, Farr, & Pomerantz, 2005; Wagner, Bundorf, Singer, & Baker, 2005; Whaley, 2004). The digital divide is especially significant in rural communities, whose residents often lack access to information technologies and have little experience using them (Dick, Manson, Hansen, Huggins, & Trullinger, 2007; Wang, Bennett, & Probst, 2011). Nevertheless, healthcare communication technologies (HCT), including email, text messaging, social networking platforms, videoconferencing, and the Internet, are vital to rural health. Promoting their use is a priority for both the Department of Health and Human Services and the Indian Health Service (Carroll et al., 2011; Sequist, Cullen, & Acton, 2011; Sequist, Cullen, & Ayanian, 2005; Sequist et al., 2007).

HCT has been successful in engaging community members with regard to health and well-being in a variety of resource-limited healthcare settings (Armstrong et al., 2009; Buchanan, Morris, & Kauley, 1999; Burdette, Herchline, & Oehler, 2008; Cegala, Street, & Clinch, 2007; Geraghty, Glynn, Amin, & Kinsella, 2008; Hammel, 2003; Jackson et al., 2008; Kleiboer et al., 2010; Lorence, Park, & Fox, 2006; Menachemi, Prickett, & Brooks, 2011; Salovey et al., 2009; Sequist et al., 2005; Street, Gordon, & Haidet, 2007; Street, Makoul, Arora, & Epstein, 2009; Street, O'Malley, Cooper, & Haidet, 2008; Zolfo et al., 2010). The SCF uses a wide array of technologies to deliver and support healthcare services. Initiatives for meaningful use of HCT include developing web-based tools (Doorenbos, Demiris, et al., 2011; Doorenbos et al., 2010; Doorenbos, Kundu, et al., 2011; Garritano & Goldenberg, 2011; Hild, 2004; Kokesh, Ferguson, & Patricoski, 2004, 2011; Patricoski, 2004) and collaborative initiatives to improve the delivery and quality of healthcare services for our patients, including physical, mental, social, and spiritual care (Carroll et al., 2011). However, these tools and initiatives are effective only if they reach their target population and are tailored to meet that population's health needs.

Colorectal cancer (CRC) is one of the leading causes of cancer-related deaths among all US residents, including AN/AI people (Jemal, Siegel, Xu, & Ward, 2010; Kelly, Alberts, Sacco,

& Lanier, 2012; Swan & Edwards, 2003). Joint consensus guidelines recommend regular screening for asymptomatic adults 50 years of age and older (Levin, Lieberman, McFarland, Andrews, et al., 2008). CRC screening has been shown to reduce both the incidence of and mortality from CRC, but despite its effectiveness, screening uptake remains suboptimal, and current low-tech interventions such as telephone and postal mail reminders may not address the needs of AN/AIs (Brouwers et al., 2011; Doorenbos, Jacobsen, Corpuz, Forquera, & Buchwald, 2011; Lee, Groessl, Ganiats, & Ho, 2011; Naylor, Ward, & Polite, 2012; Redwood et al., 2011; Sarfaty et al., 2012; Sequist, Zaslavsky, Marshall, Fletcher, & Ayanian, 2009; Yabroff et al., 2011).

Computerized reminders and other HCT have been shown to improve healthcare delivery in a subset of patients with specific health conditions and to assist providers with panel management, such as tracking patient outcomes by provider and monitoring the usage and delivery of screening services. Several studies have shown that automated messaging can improve care and screening for chronic diseases, including cancer (Armstrong et al., 2009; Feldstein et al., 2009; Greaney et al., 2012; Stone et al., 2002; Williams et al., 2011). SCF plans to implement an HCT intervention that includes reminders for chronic disease screening. The intervention will begin with messaging on CRC screening. However, it is unclear whether AN/AI people, especially those older than 50 years, have access to the HCT required to use such a service (Carroll et al., 2011; Dick et al., 2007; Rushing & Stephens, 2011; Sequist et al., 2011; Sequist et al., 2005; Wood et al., 2003). According to the principles of patient-centered care, health system changes should be driven by desired outcomes and align with patient preferences. The outcomes desired by SCF provide an initial framework for developing a formative evaluation, identifying key stakeholders (e.g., patients, providers, and leadership), formulating CRC screening messages, and implementing the intervention.

In 2011, SCF conducted a survey on communication technology to determine whether AN/AI people in SCF's region of Alaska have access to the HCT needed for the planned CRC screening intervention. The results will help to determine the potential reach of the intervention, maximize its success, and lead to improvements in monitoring and managing AN/AI population health.

Methods

Study Design

We performed a nonrandomized, cross-sectional, self-report survey of HCT among AN/AI adults. The survey was administered on paper to SCF patients in the health center's waiting rooms. The survey was written in English, the language predominantly used at the SCF. The survey was also distributed electronically through Survey Monkey to SCF staff. Survey administration to patients began in mid-February 2011; electronic recruitment of staff began on March 4. All surveys were completed by March 18, 2011.

Eligibility criteria included a self-reported age of at least 18 years and current eligibility for care at SCF. As an incentive, all participants were entered in a draw to win one of six \$50 gift cards.

SCF is an Alaska Native owned and managed regional health corporation operating under the tribal authority of Cook Inlet Region, Inc. SCF and Alaska Native Tribal Health Consortium (ANTHC) approved all aspects of the study in accordance with requirements for tribal review of research. Additionally, SCF and ANTHC provided prepublication review and approval of this manuscript prior to peer review. The Alaska Area Institutional Review Board of the Indian Health Service also approved all aspects of the study.

Study Instrument

The study instrument was a 15-item survey developed to ascertain health information, technology access (e.g., Internet, mobile phone), use of specific services (e.g., email, text messaging), anticipated use of HCT (e.g., scheduling appointments, refilling prescriptions), and participant demographics among the study population. The frequency of anticipated HCT use was rated on a 4-point Likert scale (*often, sometimes, rarely, and never*). To obtain community approval and engagement, we gave AN/AI community members and tribal leadership a voice in developing the survey. It was based on a nonvalidated instrument, which was shortened for ease of administration with input from tribal leadership and culturally adapted by AN/AI researchers at SCF. Survey questions were designed at an 81% Flesch Reading Ease Readability score and a 3.9 Flesch–Kincaid Grade Level.

Analyses

We used SAS (Version 9.2) for data manipulation and statistical analysis. Descriptive statistics were calculated and gender associations with availability and use of technology were assessed with chi-square tests. Because CRC screening is recommended on the basis of age, we present results for the entire sample as well as for the subsample of participants aged 50 years and older. Associations were considered significant for p values less than .05.

Results

Our recruitment methods resulted in a convenience sample of 673 participants (Table 1). Ninety-six percent reported gender, among whom 79% were women. Most participants were employees at the SCF (84%) and of AN/AI descent (72%). The demographics of the full sample were similar to those of the subsample of participants aged 50 years and older ($n = 168$). In the older subsample, 75% were women, 74% were SCF employees, and 68% were AN/AI. Approximately 98% of respondents reported having computer access (with or without Internet capability), 97% reported email access, and 94% reported mobile phone use (Table 2). Among mobile phone users, 60% had access to the Internet on their phones. Sixty-one percent of respondents spent 5 to 9 hours per day using a computer, a mobile phone, or the Internet. Ninety percent of participants with text messaging capabilities reported sending text messages, with 65% sending at least one message per day and 40% sending 10 or more messages per day. As shown in Table 2, rates of computer access and email use were higher among women than among men, both in the full sample ($p = .011$ and $p = .005$, respectively) and in the older subsample ($p = .016$ and $p = .011$, respectively). The rate of mobile phone access was also significantly higher for women in the full sample ($p < .001$), but we found no gender difference in the older subsample ($p = .198$). Nor did we find any significant gender differences in Internet access by mobile phone (full sample, $p = .332$; older

subsample, $p = .480$) or number of text messages sent daily (full sample, $p = .074$; older subsample, $p = .202$).

Most participants in the full sample reported that they found the following options useful and would probably use them at least sometimes: email communication with their healthcare providers (78%), online appointment scheduling (76%), online medication refills (75%), and text or email screening reminders (70%). Within the older subsample, slightly smaller majorities reported the same preferences: email communication with their healthcare providers (71%), online medication refills (68%), online appointment scheduling (62%), and text or email screening reminders (61%). SCF employees were more likely than non-employees to anticipate use of HCT to schedule appointments, refill medications, receive text or email screening reminders, and communicate by email with healthcare providers ($p < .001$, $p < .001$, $p = .002$, and $p < .001$, respectively; data not shown). The preferred method for receipt of health information did not appear to differ between the full sample and the older subsample, except that a smaller percentage of the older subsample preferred text messages ($p < .001$). When we examined age distributions at a more granular level (16–25, 26–45, 46–64, and 65–80 years), people in the oldest age category (65–80) were significantly less likely to anticipate using the Internet to schedule appointments, refill medications, or communicate with their providers (all $p < .001$).

Limitations

A key limitation of this study is our use of a convenience sample to increase the speed and ease of data collection and reduce administrative cost. Participants may have self-selected on the basis of interest in technology and may therefore overrepresent technology use in the community. In addition, the large proportion of SCF employees in our sample may have elevated results for HCT access and use. Nevertheless, 84% of SCF employees are also SCF beneficiaries, so their participation may not affect the representativeness of our study sample. An additional limitation of the study was that the survey was offered only in English, and AN/AI people who were not as comfortable or fluent in English may have chosen not to participate.

Discussion

Americans depend on technology to receive and communicate information (Denizard-Thompson, Feiereisel, Stevens, Miller, & Wofford, 2011; Miller & West, 2009; Salovey et al., 2009). However, widespread, equitable access to information technology has been limited by a knowledge divide as well as a digital divide (Goodall, Ward, & Newman, 2010; Haughton, Kreuter, Hall, Holt, & Wheatley, 2005; Hsu et al., 2005; Lorence et al., 2006; Salovey et al., 2009). Those most at risk of poor health outcomes are likely to fall farther behind as health systems increasingly rely on HCT for communication (Sarkar et al., 2011). For example, adoption of digital technologies by older adults has increased, but it is still limited and based on the perceived utility of these tools (Heart & Kalderon, 2013; Tran, Buckley, Bertera, & Gonzales, 2009).

Among AN/AI people aged 50 years and older, our study found rates of access and of reported and anticipated use of digital technologies that were higher than we expected

(Goodall et al., 2010; Haughton et al., 2005; Lorence et al., 2006; McNeill, Puleo, Bennett, & Emmons, 2007). Computer access and email use appeared consistent across the entire respondent population, regardless of age. However, as also observed in the literature, a more granular examination of our age data revealed that anticipated Internet use to manage healthcare needs declined with increasing age (Balfour et al., 2009; Heart & Kalderon, 2013; Tran et al., 2009; Womeodu & Bailey, 1996).

Studies of the knowledge and digital divides have returned conflicting results with regard to gender (Jackson et al., 2008; Swartz, Cowan, & Batista, 2004). Unlike other work, our study found significant differences between men and women in computer access, mobile phone access, and email use. However, it is important to note that our results included a large number of women who were employed at the SCF and thus were likely to have access to computers as part of their daily job activities (Denizard-Thompson et al., 2011).

Across the US, disparities in technology use have been noted in underrepresented populations (e.g., African Americans, Hispanics, AN/AIs; Brodie et al., 2000; Hanauer, Dibble, Fortin, & Col, 2004; Hsu et al., 2005; Jackson et al., 2008; Whaley, 2004). A 2007 study found that among eighth-grade students in public schools, AN/AIs were less likely than all other racial and ethnic groups to use computers at home, resulting in poorer performance on standardized tests (Jackson et al., 2006; Kind et al., 2005). However, the AN/AI participants in our study appeared to be familiar with email and text messaging, and were interested in using these technologies for clinical purposes. Despite anticipated variations, overall community access to and interest in HCT were high.

The digital divide has affected rural communities in such a way that residents often lack experience with technological resources (Dick et al., 2007), whereas urban residents typically have Internet access in their homes, schools, and community facilities (Wang et al., 2011). Because a sizeable portion of the AN/AI population resides in remote areas, expanded Internet access can increase their use of health services through HCT. Access to, use of, and optimization of HCT are vital both to patient-centered care and to the expansion of healthcare services to AN/AI communities (Carroll et al., 2011; Sequist et al., 2011; Sequist et al., 2005; Sequist et al., 2007; Wagner et al., 2005; Watson, Bell, Kvedar, & Grant, 2008). However, HCT-based initiatives will be effective only if they reach AN/AIs and are tailored to meet their unique healthcare needs.

Conclusion

Information on technology access and use enables healthcare providers and administrators to deploy HCT more efficiently and cost-effectively to address health concerns and disparities in AN/AI communities. Outcomes of this formative evaluation will drive future qualitative research on HCT. Our results will facilitate the identification and subsequent engagement of key stakeholders in HCT interventions and contribute to the meaningful use of clinical data. In particular, our findings will assist in developing a telecommunication messaging initiative to promote CRC screening and to better manage and monitor AN/AI population health.

Acknowledgements

This research was supported by the National Cancer Institute funded Regional Native American Community Network.

References

- Armstrong AW, Watson AJ, Makredes M, Frangos JE, Kimball AB, Kvedar JC. Text-message reminders to improve sunscreen use: A randomized, controlled trial using electronic monitoring. *Archives of Dermatology*. 2009; 145(11):1230–1236. [PubMed: 19917951]
- Balfour DC III, Evans S, Januska J, Lee HY, Lewis SJ, Nolan SR, Thapar K. Health information technology—results from a roundtable discussion. *Journal of Managed Care Pharmacy*. 2009; 15(1 Suppl S-a):S10–S17.
- Brodie M, Flournoy RE, Altman DE, Blendon RJ, Benson JM, Rosenbaum MD. Health information, the Internet, and the digital divide. *Health Affairs (Millwood)*. 2000; 19(6):255–265.
- Brouwers MC, De Vito C, Bahirathan L, Carol A, Carroll JC, Cotterchio M, Wathen N. What implementation interventions increase cancer screening rates? A systematic review. *Implementation Science*. 2011; 6:111.10.1186/1748-5908-6-111 [PubMed: 21958556]
- Brown L, Copas A, Stephenson J, Gilleran G, Ross JD. Preferred options for receiving sexual health screening results: A population and patient survey. *International Journal of STD & AIDS*. 2008; 19(3):184–187. [PubMed: 18397559]
- Buchanan HS, Morris RC, Kauley TD. Serving Native health needs: Merging technology and traditional information services. *Western Journal of Medicine*. 1999; 171(5–6):373–375. [PubMed: 10639882]
- Burdette SD, Herchline TE, Oehler R. Surfing the web: Practicing medicine in a technological age: Using smartphones in clinical practice. *Clinical Infectious Diseases*. 2008; 47(1):117–122. [PubMed: 18491969]
- Burke D, Menachemi N, Brooks RG. Diffusion of information technology supporting the Institute of Medicine's quality chasm care aims. *Journal for Healthcare Quality*. 2005; 27(1):24–32. 39. [PubMed: 16416889]
- Carroll M, Cullen T, Ferguson S, Hogge N, Horton M, Kokesh J. Innovation in Indian healthcare: Using health information technology to achieve health equity for American Indian and Alaska Native populations. *Perspectives in Health Information Management*. 2011; 8:1d.
- Cegala DJ, Street RL Jr, Clinch CR. The impact of patient participation on physicians' information provision during a primary care medical interview. *Health Communication*. 2007; 21(2):177–185. [PubMed: 17523863]
- Denizard-Thompson NM, Feiereisel KB, Stevens SF, Miller DP, Wofford JL. The digital divide at an urban community health center: Implications for quality improvement and health care access. *Journal of Community Health*. 2011; 36(3):456–460.10.1007/s10900-010-9327-5 [PubMed: 21086028]
- Dick RW, Manson SM, Hansen AL, Huggins A, Trullinger L. The Native Telehealth Outreach and Technical Assistance Program: A community-based approach to the development of multimedia-focused health care information. *American Indian and Alaska Native Mental Health Research*. 2007; 14(2):49–66. [PubMed: 17874365]
- Doorenbos AZ, Demiris G, Towle C, Kundu A, Revels L, Colven R, Buchwald D. Developing the Native People for Cancer Control Telehealth Network. *Telemedicine and e-Health*. 2011; 17(1): 30–34.10.1089/tmj.2010.0101 [PubMed: 21214371]
- Doorenbos AZ, Eaton LH, Haozous E, Towle C, Revels L, Buchwald D. Satisfaction with telehealth for cancer support groups in rural American Indian and Alaska Native communities. *Clinical Journal of Oncology Nursing*. 2010; 14(6):765–770.10.1188/10.CJON.765-770 [PubMed: 21112853]
- Doorenbos AZ, Jacobsen C, Corpuz R, Forquera R, Buchwald D. A randomized controlled calendar mail-out to increase cancer screening among urban American Indian and Alaska Native patients.

- Journal of Cancer Education. 2011; 26(3):549–554.10.1007/s13187-011-0217-z [PubMed: 21472495]
- Doorenbos AZ, Kundu A, Eaton LH, Demiris G, Haozous EA, Towle C, Buchwald D. Enhancing access to cancer education for rural healthcare providers via telehealth. *Journal of Cancer Education*. 2011; 26(4):682–686.10.1007/s13187-011-0204-4 [PubMed: 21336979]
- Eby D. Integrated primary care. *International Journal of Circumpolar Health*. 1998; 57(Suppl 1):665–667. [PubMed: 10093362]
- Eby DK. Primary care at the Alaska Native Medical Center: A fully deployed “new model” of primary care. *International Journal of Circumpolar Health*. 2007; 66(Suppl 1):4–13. [PubMed: 18154227]
- Feldstein AC, Perrin N, Rosales AG, Schneider J, Rix MM, Keels K, Glasgow RE. Effect of a multimodal reminder program on repeat mammogram screening. *American Journal of Preventive Medicine*. 2009; 37(2):94–101. [PubMed: 19589447]
- Garritano FG, Goldenberg D. Successful telemedicine programs in otolaryngology. *Otolaryngologic Clinics of North America*. 2011; 44(6):1259–1274. vii.10.1016/j.otc.2011.08.003 [PubMed: 22032480]
- Geraghty M, Glynn F, Amin M, Kinsella J. Patient mobile telephone “text” reminder: A novel way to reduce non-attendance at the ENT out-patient clinic. *Journal of Laryngology & Otology*. 2008; 122(3):296–298. [PubMed: 17470313]
- Goodall K, Ward P, Newman L. Use of information and communication technology to provide health information: What do older migrants know, and what do they need to know? *Quality in Primary Care*. 2010; 18(1):27–32.
- Greaney ML, Puleo E, Sprunck-Harrild K, Bennett GG, Cunningham MA, Gillman MW, Emmons KM. Electronic reminders for cancer prevention: Factors associated with preference for automated voice reminders or text messages. *Preventive Medicine*. 2012; 55(2):151–154.10.1016/j.ypmed.2012.05.014 [PubMed: 22659227]
- Hammel J. Technology and the environment: Supportive resource or barrier for people with developmental disabilities? *Nursing Clinics of North America*. 2003; 38(2):331–349. [PubMed: 12914311]
- Hanauer D, Dibble E, Fortin J, Col NF. Internet use among community college students: Implications in designing healthcare interventions. *Journal of American College Health*. 2004; 52(5):197–202. [PubMed: 15029941]
- Haughton LT, Kreuter M, Hall J, Holt CL, Wheetley E. Digital divide and stability of access in African American women visiting urban public health centers. *Journal of Health Care for the Poor and Underserved*. 2005; 16(2):362–374. [PubMed: 15937398]
- Heart T, Kalderon E. Older adults: Are they ready to adopt health-related ICT? *International Journal of Medical Informatics*. 2013; 82(11):e209–231.10.1016/j.ijmedinf.2011.03.002 [PubMed: 21481631]
- Hild C. Arctic telehealth: North to the future. *International Journal of Circumpolar Health*. 2004; 63(Suppl 2):63–70. [PubMed: 15736624]
- Hsu J, Huang J, Kinsman J, Fireman B, Miller R, Selby J, Ortiz E. Use of e-Health services between 1999 and 2002: A growing digital divide. *Journal of the American Medical Informatics Association*. 2005; 12(2):164–171. [PubMed: 15561786]
- Jackson LA, von Eye A, Biocca FA, Barbatsis G, Zhao Y, Fitzgerald HE. Does home Internet use influence the academic performance of low-income children? *Developmental Psychology*. 2006; 42(3):429–435. [PubMed: 16756435]
- Jackson LA, Zhao Y, Kolenic A III, Fitzgerald HE, Harold R, Von Eye A. Race, gender, and information technology use: The new digital divide. *CyberPsychology & Behavior*. 2008; 11(4):437–442. [PubMed: 18721092]
- Jemal A, Siegel R, Xu J, Ward E. Cancer statistics, 2010. *CA: A Cancer Journal for Clinicians*. 2010; 60(5):277–300.10.3322/caac.20073 [PubMed: 20610543]
- Kelly JJ, Alberts SR, Sacco F, Lanier AP. Colorectal cancer in Alaska Native people, 2005–2009. *Gastrointestinal Cancer Research*. 2012; 5(5):149–154. [PubMed: 23112882]
- Kind T, Huang ZJ, Farr D, Pomerantz KL. Internet and computer access and use for health information in an underserved community. *Ambulatory Pediatrics*. 2005; 5(2):117–121. [PubMed: 15780014]

- Kleiboer A, Gowing K, Holm Hansen C, Hibberd C, Hodges L, Walker J, Sharpe M. Monitoring symptoms at home: What methods would cancer patients be comfortable using? *Quality of Life Research*. 2010; 19(7):965–968.10.1007/s11136-010-9662-0 [PubMed: 20446045]
- Kokesh J, Ferguson AS, Patricoski C. Telehealth in Alaska: Delivery of health care services from a specialist's perspective. *International Journal of Circumpolar Health*. 2004; 63(4):387–400. [PubMed: 15709314]
- Kokesh J, Ferguson AS, Patricoski C. The Alaska experience using store-and-forward telemedicine for ENT care in Alaska. *Otolaryngologic Clinics of North America*. 2011; 44(6):1359–1374. ix. 10.1016/j.otc.2011.08.010 [PubMed: 22032488]
- Lee JK, Grossell EJ, Ganiats TG, Ho SB. Cost-effectiveness of a mailed educational reminder to increase colorectal cancer screening. *BMC Gastroenterology*. 2011; 11:93.10.1186/1471-230X-11-93 [PubMed: 21867492]
- Levin B, Lieberman DA, McFarland B, Andrews KS, Brooks D, Bond J, Winawer SJ. Screening and surveillance for the early detection of colorectal cancer and adenomatous polyps, 2008: A joint guideline from the American Cancer Society, the US Multi-Society Task Force on Colorectal Cancer, and the American College of Radiology. *Gastroenterology*. 2008; 134(5):1570–1595. [PubMed: 18384785]
- Lorence DP, Park H, Fox S. Racial disparities in health information access: Resilience of the digital divide. *Journal of Medical Systems*. 2006; 30(4):241–249. [PubMed: 16978003]
- McNeill LH, Puleo E, Bennett GG, Emmons KM. Exploring social contextual correlates of computer ownership and frequency of use among urban, low-income, public housing adult residents. *Journal of Medical Internet Research*. 2007; 9(4):e35. [PubMed: 18093903]
- Menachemi N, Prickett CT, Brooks RG. The use of physician–patient email: A follow-up examination of adoption and best-practice adherence 2005–2008. *Journal of Medical Internet Research*. 2011; 13(1):e23.10.2196/jmir.1578 [PubMed: 21447468]
- Miller EA, West DM. Where's the revolution? Digital technology and health care in the Internet age. *Journal of Health Politics, Policy and Law*. 2009; 34(2):261–284.
- Naylor K, Ward J, Polite BN. Interventions to improve care related to colorectal cancer among racial and ethnic minorities: A systematic review. *Journal of General Internal Medicine*. 2012; 27(8):1033–1046.10.1007/s11606-012-2044-2 [PubMed: 22798214]
- Patricoski C. Alaska telemedicine: Growth through collaboration. *International Journal of Circumpolar Health*. 2004; 63(4):365–386. [PubMed: 15709313]
- Redwood D, Holman L, Zandman-Zeman S, Hunt T, Besh L, Katinszky W. Collaboration to increase colorectal cancer screening among low-income uninsured patients. *Preventing Chronic Disease*. 2011; 8(3):A69. [PubMed: 21477509]
- Rushing SC, Stephens D. Use of media technologies by Native American teens and young adults in the Pacific Northwest: Exploring their utility for designing culturally appropriate technology-based health interventions. *Journal of Primary Prevention*. 2011; 32(3–4):135–145.10.1007/s10935-011-0242-z [PubMed: 21805055]
- Sahm L, MacCurtain A, Hayden J, Roche C, Richards HL. Electronic reminders to improve medication adherence—are they acceptable to the patient? *Pharmacy World & Science*. 2009; 31(6):627–629. [PubMed: 19731069]
- Salovey P, Williams-Piehota P, Mowad L, Moret ME, Edlund D, Andersen J. Bridging the digital divide by increasing computer and cancer literacy: Community technology centers for Head-Start parents and families. *Journal of Health Communication*. 2009; 14(3):228–245. [PubMed: 19440907]
- Sarfaty M, Myers RE, Harris DM, Borsky AE, Sifri R, Cocroft J, Johnson M. Variation in colorectal cancer screening steps in primary care: Basis for practice improvement. *American Journal of Medical Quality*. 2012; 27(6):458–466. [PubMed: 22267818]
- Sarkar U, Karter AJ, Liu JY, Adler NE, Nguyen R, Lopez A, Schillinger D. Social disparities in Internet patient portal use in diabetes: Evidence that the digital divide extends beyond access. *Journal of the American Medical Informatics Association*. 2011; 18(3):318–321.10.1136/jamia.2010.006015 [PubMed: 21262921]

- Sepucha KR, Levin CA, Uzogara EE, Barry MJ, O'Connor AM, Mulley AG. Developing instruments to measure the quality of decisions: Early results for a set of symptom-driven decisions. *Patient Education and Counseling*. 2008; 73(3):504–510. [PubMed: 18718734]
- Sequist TD, Cullen T, Acton KJ. Indian health service innovations have helped reduce health disparities affecting American Indian and Alaska Native people. *Health Affairs (Millwood)*. 2011; 30(10):1965–1973.10.1377/hlthaff.2011.0630
- Sequist TD, Cullen T, Ayanian JZ. Information technology as a tool to improve the quality of American Indian health care. *American Journal of Public Health*. 2005; 95(12):2173–2179. [PubMed: 16257947]
- Sequist TD, Cullen T, Hays H, Taulii MM, Simon SR, Bates DW. Implementation and use of an electronic health record within the Indian Health Service. *Journal of the American Medical Informatics Association*. 2007; 14(2):191–197. [PubMed: 17213495]
- Sequist TD, Zaslavsky AM, Marshall R, Fletcher RH, Ayanian JZ. Patient and physician reminders to promote colorectal cancer screening: A randomized controlled trial. *Archives of Internal Medicine*. 2009; 169(4):364–371. [PubMed: 19237720]
- Stone EG, Morton SC, Hulscher ME, Maglione MA, Roth EA, Grimshaw JM, Shekelle PG. Interventions that increase use of adult immunization and cancer screening services: A meta-analysis. *Annals of Internal Medicine*. 2002; 136(9):641–651. [PubMed: 11992299]
- Street RL Jr, Gordon H, Haidet P. Physicians' communication and perceptions of patients: Is it how they look, how they talk, or is it just the doctor? *Social Science & Medicine*. 2007; 65(3):586–598. [PubMed: 17462801]
- Street RL Jr, Makoul G, Arora NK, Epstein RM. How does communication heal? Pathways linking clinician–patient communication to health outcomes. *Patient Education and Counseling*. 2009; 74(3):295–301. [PubMed: 19150199]
- Street RL Jr, O'Malley KJ, Cooper LA, Haidet P. Understanding concordance in patient–physician relationships: Personal and ethnic dimensions of shared identity. *Annals of Family Medicine*. 2008; 6(3):198–205. [PubMed: 18474881]
- Swan J, Edwards BK. Cancer rates among American Indians and Alaska Natives: Is there a national perspective? *Cancer*. 2003; 98(6):1262–1272. [PubMed: 12973851]
- Swartz SH, Cowan TM, Batista IA. Using claims data to examine patients using practice-based Internet communication: Is there a clinical digital divide? *Journal of Medical Internet Research*. 2004; 6(1):e1. [PubMed: 15111267]
- Thompson E. Patient vs. “customer-owner.”. *Family Practice Management*. 2008; 15(4):11. [PubMed: 18446931]
- Tran, BQ.; Buckley, KM.; Bertera, EM.; Gonzales, PL. Benefits and barriers to adoption of health IT in an elderly low-income, minority community-based environment. *Conference Proceedings: Annual International Conference of the IEEE Engineering in Medicine and Biology Society*, 2009; 2009. p. 6441–6443.
- Wagner TH, Bundorf MK, Singer SJ, Baker LC. Free Internet access, the digital divide, and health information. *Medical Care*. 2005; 43(4):415–420. [PubMed: 15778645]
- Wang JY, Bennett K, Probst J. Subdividing the digital divide: Differences in Internet access and use among rural residents with medical limitations. *Journal of Medical Internet Research*. 2011; 13(1):e25.10.2196/jmir.1534 [PubMed: 21371989]
- Watson AJ, Bell AG, Kvedar JC, Grant RW. Reevaluating the digital divide: Current lack of Internet use is not a barrier to adoption of novel health information technology. *Diabetes Care*. 2008; 31(3):433–435. [PubMed: 18056885]
- Whaley KC. America's digital divide: 2000–2003 trends. *Journal of Medical Systems*. 2004; 28(2):183–195. [PubMed: 15195849]
- Williams LS, Ofner S, Yu Z, Beyth RJ, Plue L, Damush T. Pre-post evaluation of automated reminders may improve detection and management of post-stroke depression. *Journal of General Internal Medicine*. 2011; 26(8):852–857.10.1007/s11606-011-1709-6 [PubMed: 21499827]
- Womeodu RJ, Bailey JE. Barriers to cancer screening. *Medical Clinics of North America*. 1996; 80(1):115–133. [PubMed: 8569291]

- Wood FB, Sahali R, Press N, Burroughs C, Mala TA, Siegel ER, Fuller SS. Tribal Connections health information outreach: Results, evaluation, and challenges. *Journal of the Medical Library Association*. 2003; 91(1):57–66. [PubMed: 12568158]
- Yabroff KR, Zapka J, Klabunde CN, Yuan G, Buckman DW, Haggstrom D, Taplin SH. Systems strategies to support cancer screening in U.S. primary care practice. *Cancer Epidemiology, Biomarkers & Prevention*. 2011; 20(12):2471–2479.10.1158/1055-9965.EPI-11-0783
- Zolfo M, Iglesias D, Kiyon C, Echevarria J, Fucay L, Llacsahuanga E, Lynen L. Mobile learning for HIV/AIDS healthcare worker training in resource-limited settings. *AIDS Research and Therapy*. 2010; 7:35.10.1186/1742-6405-7-35 [PubMed: 20825677]

Table 1

Demographic Descriptive Statistics

Characteristic	Entire sample (<i>N</i> = 673)		Sample 50 years (<i>n</i> = 168)	
	<i>n</i>	% ^a	<i>n</i>	% ^a
Gender				
Female	511	79%	107	75%
Male	135	21%	36	25%
Employment at SCF				
Yes	540	84%	104	74%
No	102	16%	36	26%
Ethnicity				
Alaska Native/American Indian	460	72%	97	68%
Not Alaska Native/American Indian	183	28%	46	32%

^aMissing count and percentage are not presented, thus *n* will not add up to *N*. The percentage is of those who specified gender, employment at SCF, or ethnicity.

Table 2

Reported Technology Access and Use by Age and Gender

	Entire sample (<i>N</i> = 673)			Sample 50 years (<i>n</i> = 168)		
	Female <i>n</i> (%) ^{<i>a</i>}	Male <i>n</i> (%) ^{<i>a</i>}	<i>p</i> ^{<i>b</i>}	Female <i>n</i> (%) ^{<i>a</i>}	Male <i>n</i> (%) ^{<i>a</i>}	<i>p</i> ^{<i>b</i>}
Computer availability^{<i>c</i>}						
Yes, every day	440 (86)	115 (85)		89 (84)	26 (72)	
Yes, but not every day	61 (12)	12 (9)	.011	15 (14)	5 (14)	.016
No	8 (2)	8 (6)		2 (2)	5 (14)	
Email use						
Yes, every day	397 (78)	102 (76)		76 (72)	22 (61)	
Yes, but not every day	102 (20)	23 (17)	.005	25 (24)	7 (19)	.011
No	10 (2)	10 (7)		4 (4)	7 (19)	
Mobile phone access						
Yes, I have a monthly plan	459 (90)	105 (78)		89 (84)	26 (72)	
Yes, I have a prepaid plan	25 (5)	16 (12)	< .001	6 (6)	5 (14)	.198
No	25 (5)	14 (10)		11 (10)	5 (14)	
Mobile phone Internet access	296 (61)	67 (56)	.332	45 (47)	12 (40)	.480
Number of text messages sent						
10 or more daily	201 (42)	40 (33)		9 (9)	2 (6)	
1 to 9 daily	119 (25)	31 (26)		16 (17)	6 (19)	
Some messages but not daily	125 (26)	32 (26)	.074	50 (53)	11 (35)	.202
I never send texts	38 (8)	18 (15)		20 (21)	12 (39)	

^{*a*} Missing count and percentage are not presented, thus *n* will not add up to *N*. The percentage is of people who responded to the question of interest and who specified gender.

^{*b*} *p*-value calculated using the χ^2 test of independence.

^{*c*} Any computer availability with or without Internet access.