Rapid Whole-Cell Assay of Antitubercular Drugs Using Second-Generation Fluoromycobacteriophages

Estefanía Urdániz,^a Liliana Rondón,^a Marcelo A. Martí,^a Graham F. Hatfull,^b Mariana Piuri^a

Departamento de Química Biológica, Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires, IQUIBICEN-CONICET, Buenos Aires, Argentina^a; Department of Biological Sciences and Pittsburgh Bacteriophage Institute, University of Pittsburgh, Pittsburgh, Pennsylvania, USA^b

Tuberculosis (TB) is a major human health concern and kills about 3,500 people each day (1). The emergence of *Mycobacterium tuberculosis*-resistant strains has become a serious public health problem worldwide, complicating treatment and control of the disease. There is a need for new and efficient antitubercular drugs that shorten the time of treatment and frequency of administration, have less toxicity, and require less patient surveillance (2–5). A novel, rapid, and sensitive assay to be used in activity testing and screening of new compounds could be part of the solution to the TB health challenges.

We previously described the development of fluoromycobacteriophages as a new class of reporter phages that provide a simple

means of indicating the metabolic state of *M. tuberculosis* cells and thus their response to antibiotics (6) (7). Mycobacterial cells can

Accepted manuscript posted online 14 March 2016

Citation Urdániz E, Rondón L, Martí MA, Hatfull GF, Piuri M. 2016. Rapid whole-cell assay of antitubercular drugs using second-generation fluoromycobacteriophages. *Antimicrob Agents Chemother* 60:3253–3256. doi:10.1128/AAC.03016-15.

Address correspondence to Mariana Piuri, mpiuri@qb.fcen.uba.ar.

Copyright © 2016, American Society for Microbiology. All Rights Reserved.

FIG 1 Construction of new *mCherry_{bomb}* ϕ . (A and B) *M. smegmatis* mc²155 colonies after transformation with pJL37::hsp60_{prom}-mCherry_{bomb} (A) or pJL37::hsp60_{prom}-gp9RBS-mCherry_{bomb} (B). (C) Flow cytometry analysis of *M. smegmatis* mc²155 control (gray) transformed as described for panel A (pink) or panel B (purple). The intensity of fluorescence is plotted against the number of events normalized to the mode. PE-A, phycoerythrin A. (D and E) Infection of *M. tuberculosis* mc²6230 with the new *mCherry_{bomb}* ϕ for 5 h at 37°C. Phase-contrast images (D) and fluorescence micrograph images (E) are shown. Magnification, $\times 1,000$.

FIG 2 Monitoring of fluorescence after infection of *M. tuberculosis* with fluoromycobacteriophages using a fluorimeter. (A) Fluorescence units (FU) versus time after infection of *M. tuberculosis* mc²6230 with phAE87::hsp60-EGFP (circles) or mCherry_{bomb}Φ (squares). (B) Schematic representation of the multiwell plate setup and output fluorescence reading for activity testing of drugs. Black represents maximal FU and white absence of signal. Different shades of gray are used for intermediate values. Control (CTRL) wells without drug are shown. EMB and EMB(24), EMB without and with 24 h of preincubation; INH and INH(24), INH without and with 24 h of preincubation. (C to J) Relative fluorescence units (RFU) versus time after infection of *M. tuberculosis* mc²6230 Kan^r with second-generation fluoromycobacteriophages in the presence of increasing concentrations of the following antibiotics: rifampin (RIF) (C); streptomycin (STR) (D); ofloxacin (OFX) (E); kanamycin (KAN) (F); ethambutol (EMB) (G and I); isoniazid (INH) (H and J). Antibiotics were added simultaneously with mCherry_{bomb}Φ (C to H) or preincubated with cells for 24 h before infection (I and J).

easily be detected by fluorescence microscopy or flow cytometry after infection with the reporter phage.

To construct second-generation fluoromycobacteriophages with greater sensitivity and shorter time to *M. tuberculosis* detection, we used a modified *mCherry_{bomb}* version with codon usage optimized for mycobacteria under the control of the *hsp60* promoter and with the predicted ribosome binding site (RBS) of gp9 of mycobacteriophage TM4, the major capsid protein (8, 9). Panels A to C of Fig. 1 show a comparison between mycobacterial cells transformed with the expression plasmid before and after substitution of the RBS associated with the *hsp60* promoter. The original vector in shuttle phasmid pAE159 (10) was replaced with pYUB854 carrying the *hsp60*prom-gp9RBS-*mCherry_{bomb}* cassette. Phage particles were obtained as previously described (6), and the newly constructed phage was called *mCherry_{bomb}*Φ.

After infection of *M. tuberculosis* mc²6230 (Δ RD1 Δ panCD) (11), bright red cells were observed by fluorescence microscopy after only 5 h (Fig. 1D and E), showing a notable reduction of the time required to detect the signal compared to previous pAE87::*hsp60*-EGFP (enhanced green fluorescent protein) (6) and other fluorescent reporter phages (12). We then evaluated the detection of *M. tuberculosis*-infected cells by monitoring the expression kinetics of *mCherry_{bomb}* using an automated multiwell plate reader fluorimeter. After infection with *mCherry_{bomb}*Φ, the maximal readout was reached at 4 to 6 h versus 10 h with the previous GFP phage (6). Moreover, the signal/background ratio (total number of fluorescence units/number of fluorescence units of cells plus phage at time zero) was higher for the new reporter phage, allowing better discrimination between noninfected and infected cells (Fig. 2A).

We further measured the fluorescent signal after infection in the presence of 2-fold dilutions of the drugs most commonly used for TB treatment. A schematic representation of the 96-well plate setup and the output signal for this assay is presented in Fig. 2B. To avoid background differences among *mCherry_{bomb}*Φ stocks, we calculated relative fluorescence units (RFU) as follows:

$$\text{RFU} = \frac{\text{mean fluorescence units} - \text{background}}{\text{maximal fluorescence reading units} - \text{background}}$$

A decrease in the fluorescent signal was observed for increasing concentrations of the tested compounds (Fig. 2), allowing MIC determinations. With this assay, we were able to corroborate the antimicrobial resistance phenotype of the *M. tuberculosis* mc²6230 kanamycin-resistant (Kan^r) derivative strain used in this work (Fig. 2F).

As we previously reported (6), when the target of the tested drug is gene expression (transcription or translation), phage and antibiotics could effectively be added simultaneously (Fig. 2C to F). When the antibiotic had a different target (e.g., cell wall synthesis), a preincubation of 24 h was necessary prior to addition of *mCherry_{bomb}*Φ (Fig. 2G to J). Based on this requirement, in testing new compounds, a preliminary mechanism of action of the drug can be inferred.

In comparison with other whole-cell assays for activity testing of compounds (13–18), results using fluoromycobacteriophages can be obtained in hours in contrast to days; no addition of substrate is needed (13, 14, 19), and, since mycobacterial cultures can be maintained constantly growing, they are readily available when needed. All these features make *mCherry_{bomb}*Φ in

combination with automated fluorimetric detection a convenient tool for activity testing of new antitubercular drugs and a potential rapid drug susceptibility testing (DST) assay of *M. tuberculosis* clinical isolates.

ACKNOWLEDGMENTS

We thank Eric Rubin (Harvard School of Public Health) for providing the codon-optimized *mCherry_{bomb}* and William Jacobs, Jr. (Albert Einstein College of Medicine), for the *M. tuberculosis* mc²6230 Kan^r strain.

FUNDING INFORMATION

This work, including the efforts of Mariana Piuri, was funded by Fundación Bunge y Born (FBBE16/12). This work, including the efforts of Graham F. Hatfull and Mariana Piuri, was funded by FIRCA-BB | National Institutes of Health (NIH) (R03TW008926). This work, including the efforts of Marcelo Martí, was funded by MINCYT | Agencia Nacional de Promoción Científica y Tecnológica (ANPCyT) (PICT-O 0057).

This work was funded by U.S. National Institutes of Health (FIRCA-BB) R03TW008926 to G.F.H. and M.P. (LMICC); Bunge & Born Foundation FBBE16/12 to M.P. and ANPCyT PICT-O 0057 to M.M. E.U. and L.R. are doctoral fellows of CONICET (Consejo Nacional de Investigaciones Científicas y Tecnológicas, Argentina).

REFERENCES

1. WHO. 2014. Global tuberculosis report. World Health Organization, Geneva, Switzerland.
2. Schito M, Migliori GB, Fletcher HA, McNerney R, Centis R, D'Ambrosio L, Bates M, Kibiki G, Kapata N, Corrah T, Bomanji J, Vilaplana C, Johnson D, Mwaba P, Maeurer M, Zumla A. 2015. Perspectives on advances in tuberculosis diagnostics, drugs, and vaccines. Clin Infect Dis 61(Suppl 3):S102–S118. <http://dx.doi.org/10.1093/cid/civ609>.
3. Zumla A, Chakaya J, Centis R, D'Ambrosio L, Mwaba P, Bates M, Kapata N, Nyirenda T, Chanda D, Mfinanga S, Hoelscher M, Maeurer M, Migliori GB. 2015. Tuberculosis treatment and management—an update on treatment regimens, trials, new drugs, and adjunct therapies. Lancet Respir Med 3:220–234. [http://dx.doi.org/10.1016/S2213-2600\(15\)00063-6](http://dx.doi.org/10.1016/S2213-2600(15)00063-6).
4. Zuniga ES, Early J, Parish T. 2015. The future for early-stage tuberculosis drug discovery. Future Microbiol 10:217–229. <http://dx.doi.org/10.2217/fmb.14.125>.
5. Sotgiu G, Centis R, D'Ambrosio L, Migliori GB. 2015. Tuberculosis treatment and drug regimens. Cold Spring Harb Perspect Med 5:a017822. <http://dx.doi.org/10.1101/cshperspect.a017822>.
6. Piuri M, Jacobs WR, Jr, Hatfull GF. 2009. Fluoromycobacteriophages for rapid, specific, and sensitive antibiotic susceptibility testing of Mycobacterium tuberculosis. PLoS One 4:e4870. <http://dx.doi.org/10.1371/journal.pone.0004870>.
7. Rondón L, Piuri M, Jacobs WR, Jr, de Waard J, Hatfull GF, Takiff HE. 2011. Evaluation of fluoromycobacteriophages for detecting drug resistance in Mycobacterium tuberculosis. J Clin Microbiol 49:1838–1842. <http://dx.doi.org/10.1128/JCM.02476-10>.
8. Ford ME, Stenstrom C, Hendrix RW, Hatfull GF. 1998. Mycobacteriophage TM4: genome structure and gene expression. Tuber Lung Dis 79: 63–73. <http://dx.doi.org/10.1054/tuld.1998.0007>.
9. Piuri M, Rondón L, Urdaniz E, Hatfull GF. 2013. Generation of affinity-tagged fluoromycobacteriophages by mixed assembly of phage capsids. Appl Environ Microbiol 79:5608–5615. <http://dx.doi.org/10.1128/AEM.01016-13>.
10. Bardarov S, Jr, Pavelka MS, Jr, Sambandamurthy V, Larsen M, Tufariello J, Chan J, Hatfull G, Jacobs WR, Jr. 2002. Specialized transduction: an efficient method for generating marked and unmarked targeted gene disruptions in Mycobacterium tuberculosis, M. bovis BCG and M. smegmatis. Microbiology 148:3007–3017. <http://dx.doi.org/10.1099/00221287-148-10-3007>.
11. Sambandamurthy VK, Jacobs WR, Jr. 2005. Live attenuated mutants of Mycobacterium tuberculosis as candidate vaccines against tuberculosis. Microbes Infect 7:955–961. <http://dx.doi.org/10.1016/j.micinf.2005.04.001>.
12. Jain P, Hartman TE, Eisenberg N, O'Donnell MR, Kriakov J, Govender

- K, Makume M, Thaler DS, Hatfull GF, Sturm AW, Larsen MH, Moodley P, Jacobs WR, Jr. 2012. phi(2)GFP10, a high-intensity fluorophage, enables detection and rapid drug susceptibility testing of *Mycobacterium tuberculosis* directly from sputum samples. *J Clin Microbiol* 50:1362–1369. <http://dx.doi.org/10.1128/JCM.06192-11>.
13. Riska PF, Jacobs WR, Jr, Bloom BR, McKittrick J, Chan J. 1997. Specific identification of *Mycobacterium tuberculosis* with the luciferase reporter mycobacteriophage: use of p-nitro-alpha-acetylamino-beta-hydroxy propiophenone. *J Clin Microbiol* 35:3225–3231.
14. Shawar RM, Humble DJ, Van Dalsen JM, Stover CK, Hickey MJ, Steele S, Mitscher LA, Baker W. 1997. Rapid screening of natural products for antimycobacterial activity by using luciferase-expressing strains of *Mycobacterium bovis* BCG and *Mycobacterium intracellulare*. *Antimicrob Agents Chemother* 41:570–574.
15. Andreu N, Fletcher T, Krishnan N, Wiles S, Robertson BD. 2012. Rapid measurement of antituberculosis drug activity in vitro and in macrophages using bioluminescence. *J Antimicrob Chemother* 67:404–414. <http://dx.doi.org/10.1093/jac/dkr472>.
16. Ollinger J, Bailey MA, Moraski GC, Casey A, Florio S, Alling T, Miller MJ, Parish T. 2013. A dual read-out assay to evaluate the potency of compounds active against *Mycobacterium tuberculosis*. *PLoS One* 8:e60531. <http://dx.doi.org/10.1371/journal.pone.0060531>.
17. Franzblau SG, DeGroote MA, Cho SH, Andries K, Nuermberger E, Orme IM, Mdluli K, Angulo-Barturen I, Dick T, Dartois V, Lenaerts AJ. 2012. Comprehensive analysis of methods used for the evaluation of compounds against *Mycobacterium tuberculosis*. *Tuberculosis (Edinb)* 92: 453–488. <http://dx.doi.org/10.1016/j.tube.2012.07.003>.
18. Collins LA, Torrero MN, Franzblau SG. 1998. Green fluorescent protein reporter microplate assay for high-throughput screening of compounds against *Mycobacterium tuberculosis*. *Antimicrob Agents Chemother* 42: 344–347.
19. Banaiee N, Bobadilla-del-Valle M, Riska PF, Bardarov S, Jr, Small PM, Ponce-de-Leon A, Jacobs WR, Jr, Hatfull GF, Sifuentes-Osornio J. 2003. Rapid identification and susceptibility testing of *Mycobacterium tuberculosis* from MGIT cultures with luciferase reporter mycobacteriophages. *J Med Microbiol* 52:557–561. <http://dx.doi.org/10.1099/jmm.0.05149-0>.