Published in final edited form as:

J Am Chem Soc. 2016 March 23; 138(11): 3659–3662. doi:10.1021/jacs.6b01187.

Direct Catalytic Asymmetric Mannich Reactions for the Construction of Quaternary Carbon Stereocenters

Barry M. Trost*, Tanguy Saget, and Chao-I (Joey) Hung

Department of Chemistry, Stanford University, Stanford, California 94305-5080, United States

Abstract

Herein, we report a Zn-ProPhenol catalyzed Mannich reaction using α -branched ketones as nucleophilic partners for the direct enantio- and diastereoselective construction of quaternary carbon stereocenters. The reaction can be run on a gram-scale with a low catalyst loading without impacting its efficiency. Moreover, the Mannich adducts can be further elaborated with complete diastereocontrol to access molecules possessing complex stereotriads.

Graphical abstract

One of the modern challenges for the synthetic chemist is the fast, selective and atom-economic access to molecular complexity starting from simple and widely available starting materials.¹ The Mannich reaction is an important tool for C-C bond formation.² Moreover, it is also one of the most robust ways to produce nitrogen-containing compounds which are ubiquitous in nature. All-carbon quaternary stereocenters are also a common feature of natural products and their construction still represents a synthetic challenge, especially in a catalytic enantioselective fashion.³ In this context, we report herein the first direct asymmetric Mannich reaction that allow the use of α -branched ketone donors. This atom-economic transformation provides an efficient enantio- and diastereoselective access to chiral β -amino ketones decorated with a quaternary carbon stereocenter.

Corresponding Author: bmtrost@stanford.edu.

The authors declare no competing financial interest.

Supporting Information

Experiment details, compound characterization data, and spectra. This material is available free of charge via the Internet at <http://pubs.acs.org>.

The Mannich reaction was discovered in the early 20th century and has received a lot of attention, especially in the context of the total synthesis of alkaloids.⁴ Despite its synthetic importance, the first examples of catalytic enantioselective Mannich reaction only appeared in the late 90's.⁵ These reactions require preformed enolates such as silyl enol ethers that are reacted with an imine activated by a chiral Lewis acid catalyst. One serious drawback of this approach is the preparation and the instability of the preformed enolates. A more effective and atom-economic approach was later introduced by directly using an unmodified carbonyl donor.⁶ This strategy really became popular with the advent of organocatalysis.⁷ Bimetallic catalysis has also emerged as a powerful way to perform direct enantioselective Mannich reactions.⁸ However, the use of ketones as donors for such reactions is so far limited to methyl ketones, cyclohexanones or α -hydroxyketones.⁶⁻⁹

The formation of α -quaternary carbonyl compounds using a direct Mannich reaction has been limited to the use of easily enolizable ketones¹⁰ such as 1,3-dicarbonyl compounds¹¹ or 3-substituted oxindoles¹² (Scheme 1a). To the best of our knowledge, the lone exception is a proline-catalyzed reaction using branched aldehydes developed by Barbas and coworkers¹³ (Scheme 1b). However, this reaction proceeds with poor to moderate diastereoselectivities and is restricted to the use of *N*-PMP-protected glyoxylate ethyl ester as acceptor. In this respect, we wished to develop the first direct catalytic enantioselective Mannich reaction allowing the use of a broad array of α -branched aromatic ketones together with a broad range of imine acceptors.

The ProPhenol ligand is part of the aza-hemicrown family of ligands and forms dinuclear main group metal catalysts when treated with alkyl metal reagents such as Et₂Zn. The Zn-ProPhenol catalytic system has proven to be particularly efficient for a range of enantioselective aldol and Mannich reactions.^{8c} However, the scope of donors that is compatible with this catalytic system is mostly limited to methyl ketones (acetophenones, acetone, methyl vinyl ketone, methyl ynones) or unsubstituted α -hydroxy carbonyl compounds. Very recently, we reported the use of more substituted ketones¹⁴ for Mannich-type reactions but so far the formation of quaternary stereogenic centers using this strategy has remained elusive.

Our recent work with fluorinated aromatic ketones^{14a} led us to wonder about the special importance of the fluorine atom concerning the reactivity/selectivity for this Mannich reaction. Could α -branched ketones substituted with simple alkyl substituents be suitable for related processes using our Zn-ProPhenol catalytic system? A priori, such alkyl substituents should deactivate the ketone partner for both steric and electronic reasons thus significantly impacting the outcome of the reaction. With these challenges in mind, we initiated our studies with 2-methyl tetralone **1a** to explore the feasibility of such a transformation (Table 1). Pleasingly, when **1a** was reacted with Boc-protected aldimine **3a** in the presence of 20 mol% of the Zn-ProPhenol catalyst in THF at 60°C, the desired Mannich product **4aa** was obtained with very promising selectivities albeit in low yield due to a poor conversion (entry 1). When the reaction temperature was raised to 80°C, a better conversion was observed and remarkably, the selectivities were not affected (entry 2). Increasing the concentration of the reaction was beneficial too (entry 3) and the catalyst loading could be reduced to 10 mol% to afford **4aa** with only a slightly reduced yield (entry 4). Under similar reaction conditions, 2-

methyl indanone **2a** proved to be more reactive than **1a** (entry 5–7) and **5aa** was also obtained with excellent selectivities. Pleasingly, the catalyst loading could even be lowered to 5 mol% without impacting the efficiency of the reaction (entry 8).

We selected **2a** to evaluate the generality of the reaction concerning the imine partner using our optimized conditions. A variety of aromatic Boc-aldimines were successfully reacted under our optimized reaction conditions to afford highly functionalized β -amino ketones **5ab–5ag** (Scheme 2). Both electron-withdrawing and electron-donating groups were tolerated on the aromatic ring as well as heteroaromatic imines. The reaction is not restricted to aromatic imines; vinyl imine **3f** and alkyl imine **3g** gave the Mannich products **5af** and **5ag** in high yields and selectivities. Finally, Cbz-protected imines can be used with similar efficiency which allows the use of orthogonal protecting group strategies.

We then investigated several 5-membered aromatic ketones for this direct Mannich reaction (Scheme 3). Indanones **2a–e** which are substituted with linear alkyl groups are competent partners and the Mannich products **5aa–5eh** were obtained in good yields together with perfect selectivities. Notably, an ethyl ester and a free terminal alkyne are well tolerated under these reaction conditions. Substitution with a branched alkyl group did not impact the efficiency of the reaction as shown with **5fa**. Importantly, even a *tert*-butyl substituent could be used to afford **5ga** possessing two adjacent quaternary carbons. 3-Coumaranones are important heterocyclic compounds that are found in numerous natural products and bioactive compound.¹⁵ In this context, we were pleased to see that 3-coumaranone **2h** could be used for this direct Mannich reaction.

Moreover, this method was extended to other important aromatic ketones (Scheme 4) such as tetralone **1a–b** and 4-chromanone **1c** which afforded the Mannich products **4aa–4cf** with similar levels of efficiency.¹⁶ Thiochromanone **4d**, possessing a coordinating sulfur atom that could potentially poison the Zn-catalyst, was also successfully used but in this case the innate diastereoselectivity of the reaction was 2.6:1. However, the two diastereoisomers could be easily separated by chromatography and the major diastereoisomer **4da** was isolated in 72% yield together with excellent selectivities. Benzosuberone **4e** was also a competent partner for this Mannich reaction. Finally, challenging acyclic substrates were attempted for this reaction. Pleasingly, isobutyrophenone afforded β -amino ketone **4fa** in good yield and high enantioselectivity. However, only traces of the Mannich product were detected when the highly hindered phenone **4g** was reacted under these reaction conditions, the remainder being unreacted starting material.

To showcase the scalability and the practicality of the process, we performed a gram-scale reaction using **2a** and 1.05 equivalent of imine **3a** (Scheme 5). Pleasingly, the catalyst loading could be reduced to only 2 mol% without impacting the outcome of the reaction and **5aa** was obtained with comparable excellent yield and selectivities.

We then attempted to further elaborate the Mannich products **5** to demonstrate their synthetic utility (Scheme 6).

First, the Boc protecting group was quantitatively removed with trifluoroacetic acid in dichloromethane at room temperature. Of note, this type of reaction is usually plagued by retro-Mannich processes that were not observed in this case. Then, primary amine **6** can be further functionalized and a protecting group switch to a *para*-nosyl group afforded compound **7** from which crystals were grown. The relative and absolute configuration of **7** derived from **5aa** was unambiguously established by X-ray crystallographic analysis and by analogy the same configuration was assigned to all compounds **4–5**. To account for the stereochemical outcome of the reaction, we propose a transition state where the Boc-imine is engaged *via* two-point binding with the dinuclear catalyst as shown in scheme 7. When **5aa** was reacted with MeLi at -78°C in THF, a diastereoselective addition of the methyl group was achieved to form **8** possessing three contiguous stereocenters and featuring two vicinal tetrasubstituted carbons. The use of MeMgCl instead of MeLi led to a sluggish reaction under similar conditions and to decomposition when the reaction temperature was increased. Additionally, a diastereoselective reduction of **5aa** with NaBH_4 /methanol at 0°C was also developed. The seemingly hindered neopentyl secondary alcohol **9** could then be acylated under standard conditions using DMAP as a catalyst. A diastereoselective reduction of ketone **5eg** was also performed using our optimized reaction conditions and **11** was used for the next step without chromatography purification. Removal of the Cbz protecting group under hydrogenolysis in methanol directly followed by the addition of K_2CO_3 led to the complex spirolactam **12** in excellent yield.

In summary, we have developed the first direct asymmetric Mannich reaction using α -branched ketones. This strategy allows an enantio- and diastereoselective access to a range of functionalized β -amino ketones featuring an all-carbon quaternary stereocenter. The reaction exhibits many interesting features: the pre-activation of the reaction partners is not required; the reaction is highly atom-economic and can be run on a gram-scale with a low catalyst loading. Moreover, two convenient and orthogonal protecting groups can be used on the imines with similar efficiency. Finally, the Mannich adducts can be further elaborated to complex molecules possessing three contiguous stereogenic centers with complete control of diastereoselectivity. The extension of this strategy to the synthesis of quaternary carbon in acyclic systems is a major goal of our future efforts and will be reported in due course.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank the National Science Foundation (CHE-1360634) and the National Institute of health (GM-033049) for their generous support of our programs. We thank Dr. Jana Maclaren (Stanford University) for X-ray crystallographic analysis and Jacob S. Tracy (Stanford University) for conducting NOE experiments. T.S. is extremely grateful to the Swiss National Science Foundation for a fellowship.

References

1. Trost BM. Science. 1991; 254:1471. [PubMed: 1962206]
2. Tramontini M, Angiolini L. Tetrahedron. 1990; 46:1791.

3. (a) Corey EJ, Guzman-Perez A. *Angew Chem Int Ed.* 1998; 37:388.(b) Douglas CJ, Overman LE. *Proc Nat Acad Sci USA.* 2004; 101:5363. [PubMed: 14724294] (c) Das JP, Marek I. *Chem Commun.* 2011; 47:4593.(d) Quasdorf KW, Overman LE. *Nature.* 2014; 516:181. [PubMed: 25503231]
4. Arend M, Westermann B, Rish N. *Angew Chem Int Ed.* 1998; 37:1044.
5. Ishitani H, Ueno M, Kobayashi S. *J Am Chem Soc.* 1997; 119:7153. For an early review, see: Kobayashi S, Ishitani H. *Chem Rev.* 1999; 99:1069. [PubMed: 11749440]
6. (a) Cordova A. *Acc Chem Res.* 2004; 37:102. [PubMed: 14967057] (b) Arrayas RG, Carretero JC. *Chem Soc Rev.* 2009; 38:1940. [PubMed: 19551174]
7. For the first organocatalytic direct asymmetric Mannich reaction, see: List B. *J Am Chem Soc.* 2000; 122:9336. For general reviews, see: Notz W, Tanaka F, Barbas CF III. *Acc Chem Res.* 2004; 37:580. [PubMed: 15311957] Mukherjee S, Yang JW, List B. *Chem Rev.* 2007; 107:5471. [PubMed: 18072803] Verkade JMM, Van Hemert LJC, Quaedflieg PJLM, Rutjes FPJT. *Chem Soc Rev.* 2008; 37:29. [PubMed: 18197331]
8. (a) Shibasaki M, Yoshikawa N. *Chem Rev.* 2002; 102:2187. [PubMed: 12059266] (b) Shibasaki M, Kanai M, Matsunaga S, Kumagai N. *Acc Chem Res.* 2009; 42:1117. [PubMed: 19435320] (c) Trost BM, Bartlett MJ. *Acc Chem Res.* 2015; 48:688. [PubMed: 25650587]
9. (a) Matsunaga S, Yoshida T, Morimoto H, Kumagai N, Shibasaki M. *J Am Chem Soc.* 2004; 126:8777. [PubMed: 15250731] (b) Trost BM, Jaratjaroonphong J, Reutrakul V. *J Am Chem Soc.* 2006; 128:2778. [PubMed: 16506738]
10. For a general review, see: Kobayashi S, Mori Y, Fossey JS, Salter MM. *Chem Rev.* 2011; 111:2626. [PubMed: 21405021]
11. For selected examples, see: Hamashima Y, Sasamoto N, Hotta D, Somei H, Umebayashi N, Sodeoka M. *Angew Chem Int Ed.* 2005; 44:1525. Ting A, Lou S, Schaus SE. *Org Lett.* 2006; 8:2003. [PubMed: 16671767] Kang YK, Kim DY. *J Org Chem.* 2009; 74:5734. [PubMed: 19555053] Hatano M, Horibe T, Ishihara K. *J Am Chem Soc.* 2010; 132:56. [PubMed: 20000707]
12. (a) Tian X, Jiang K, Peng J, Du W, Chen YC. *Org Lett.* 2008; 10:3583. [PubMed: 18642826] (b) Cheng L, Liu L, Jia H, Wang D, Chen YJ. *J Org Chem.* 2009; 74:4650. [PubMed: 19518155] (c) He R, Ding C, Maruoka K. *Angew Chem Int Ed.* 2009; 48:4559.(d) Shimizu S, Tsubogo T, Xu P, Kobayashi S. *Org Lett.* 2015; 17:2006. [PubMed: 25849712]
13. Chowdari NS, Suri JT, Barbas CF III. *Org Lett.* 2004; 6:2507. [PubMed: 15255677] . For a single example using a branched ketone, see: Yang X, Toste FD. *J Am Chem Soc.* 2015; 137:3205. [PubMed: 25719604]
14. (a) Trost BM, Saget T, Lerchen A, Hung C-I. *Angew Chem Int Ed.* 2016; 55:781.(b) Trost BM, Hung C-I. *J Am Chem Soc.* 2015; 137:15940. [PubMed: 26630114]
15. (a) Boumendjel A. *Curr Med Chem.* 2003; 10:2621. [PubMed: 14529476] (b) Seabra RM, Andrade PB, Ferreres F, Moreira MM. *Phytochemistry.* 1997; 45:839.(c) Lee CY, Chew EH, Go ML. *Eur J Med Chem.* 2010; 45:2957. [PubMed: 20392544]
16. The utilization of saturated alkyl imines with **4c** and **4e** has proven to be problematic due to competitive isomerization to the corresponding enecarbamates and will be a focus for future efforts.

a) Stabilized enolates

b) Non-stabilized enolates

Only reported example, Barbas and coworkers:

This work:

Scheme 1.

Direct Mannich reactions forming quaternary carbon stereocenters

Scheme 2.

Scope of the reaction^a

^aReaction conditions: 0.20 mmol **2a**, 0.24 mmol **3**, 5 mol% (*R,R*)-ProPhenol, 10 mol% Et₂Zn (1M in hexanes), 3 Å molecular sieves (5 mg), in THF (0.4 M) at 80°C for 16 h. ^bReaction run at 70°C.

Scheme 3.

Scope of the reaction^a^aReaction conditions: 0.20 mmol **2**, 0.24 mmol **3**, 10 mol% (*R,R*)-ProPhenol, 20 mol% Et₂Zn (1M in hexanes), 3 Å molecular sieves (5 mg), in THF (0.4 M) at 80°C for 16 h.^bReaction run in THF (1.0 M).

Scheme 4.

Scope of the reaction^a

^aReaction conditions: 0.20 mmol **1**, 0.24 mmol **3**, 10 mol% (*R,R*)-ProPhenol, 20 mol% Et₂Zn (1M in hexanes), 3 Å molecular sieves (5 mg), in THF (1.0 M) at 80°C for 16 h. ^b8:1 d.r. before chromatography. ^c2.6:1 d.r. before chromatography. ^cReaction run in Et₂O (1.0 M).

Scheme 5.
Gram-scale reaction

Scheme 6.
Synthetic applications

Scheme 7.
Proposed transition-state for the formation of 5aa

Table 1

Optimization of the reaction conditions^a

entry	1	x	[conc.], T	yield ^b	d.r. ^c	ee ^d
1	1a	20	(0.4 M), 60°C	28%	>20:1	97%
2	1a	20	(0.4 M), 80°C	76%	>20:1	96%
3 ^e	1a	20	(1.0 M), 80°C	93%	>20:1	99%
4 ^e	1a	10	(1.0 M), 80°C	75%	>20:1	95%
5	2a	20	(0.4 M), 60°C	99%	>20:1	99%
6	2a	10	(0.4 M), 60°C	91%	>20:1	99%
7 ^e	2a	10	(0.4 M), 80°C	99%	>20:1	99%
8 ^e	2a	5	(0.4 M), 80°C	95%	>20:1	99%

^aReaction conditions: 0.20 mmol **1a** or **2a**, 0.24 mmol **3a**, x mol% (R,R)-ProPhenol, 2x mol% Et₂Zn (1M in hexanes), 3 Å molecular sieves (5 mg), in THF for 40 h at the indicated temperature and concentration.

^bIsolated yield.

^cDetermined by ¹H-NMR analysis.

^dDetermined by HPLC on a chiral stationary phase.

^eReaction time is 16h.