Published in final edited form as:

ACS Chem Biol. 2015 December 18; 10(12): 2733–2742. doi:10.1021/acscchembio.5b00648.

Muscular Dystrophy Mutations Impair the Nuclear Envelope Emerin Self-assembly Properties

Isaline Herrada[†], Camille Samson[†], Christophe Velours[‡], Louis Renault[‡], Cecilia Östlund[§], Pierre Chervy[†], Dmytro Puchkov^{||}, Howard J Worman[§], Brigitte Buendia[⊥], and Sophie Zinn-Justin^{†,*}

[†]Laboratoire de Biologie Structurale et Radiobiologie, Institute for Integrative Biology of the Cell (I2BC), CEA Saclay Bât. 144, 91191 Gif-sur-Yvette Cedex, France

[‡]Laboratoire d'Enzymologie et Biochimie Structurales, Institute for Integrative Biology of the Cell (I2BC), CNRS Bât.34, 1 avenue de la terrasse, 91190 Gif-sur-Yvette, France

[§]Department of Medicine and Department of Pathology and Cell Biology, College of Physicians and Surgeons, Columbia University, 630 West 168th Street, New York, New York 10032, United States

^{||}Department of Molecular Pharmacology and Cell Biology, Leibniz-Institut für Molekulare Pharmakologie (FMP), 13125 Berlin, Germany

[⊥]Laboratoire de Physiologie du Muscle Strié, Université Paris Diderot-Paris 7, CNRS, UMR 8251, Institut de Biologie Fonctionnelle et Adaptative, 4 rue M.A. Lagroua Weill Halle, 75205 Paris Cedex 13, France

Abstract

More than 100 genetic mutations causing X-linked Emery–Dreifuss muscular dystrophy have been identified in the gene encoding the integral inner nuclear membrane protein emerin. Most mutations are nonsense or frameshift mutations that lead to the absence of emerin in cells. Only very few cases are due to missense or short in-frame deletions. Molecular mechanisms explaining the corresponding emerin variants' loss of function are particularly difficult to identify because of the mostly intrinsically disordered state of the emerin nucleoplasmic region. We now demonstrate that this EmN region can be produced as a disordered monomer, as revealed by nuclear magnetic resonance, but rapidly self-assembles *in vitro*. Increases in concentration and temperature favor the formation of long curvilinear filaments with diameters of approximately 10 nm, as observed by electron microscopy. Assembly of these filaments can be followed by fluorescence through Thioflavin-T binding and by Fourier-transform Infrared spectrometry through formation of β -

*Corresponding Author. Tel.: +33(0)169083026. Fax: +33(0)169084712. sophie.zinn@cea.fr.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acscchembio.5b00648.

Alignment of emerin sequences (Supporting Figure 1), analytical ultracentrifugation plots (Supporting Figure 2), gel filtration profiles of EmN and its mutants (Supporting Figure 3), and fluorescence emission spectra recorded in the presence of ThT on EmN mutants at pH 6.8 (Supporting Figure 4) (PDF)

The authors declare no competing financial interest.

structures. Analysis of the assembly properties of five EmN variants reveals that del95–99 and Q133H impact filament assembly capacities. In cells, these variants are located at the nuclear envelope, but the corresponding quantities of emerin–emerin and emerin–lamin proximities are decreased compared to wild-type protein. Furthermore, variant P183H favors EmN aggregation *in vitro*, and variant P183T provokes emerin accumulation in cytoplasmic foci in cells. Substitution of residue Pro183 might systematically favor oligomerization, leading to emerin aggregation and mislocalization in cells. Our results suggest that emerin self-assembly is necessary for its proper function and that a loss of either the protein itself or its ability to self-assemble causes muscular dystrophy.

Graphical abstract

In the metazoan nucleus, the genome is surrounded by the nuclear envelope, containing a double lipid membrane and a large number of proteins. A filamentous protein meshwork called the nuclear lamina lines the inner nuclear membrane. It is primarily composed of intermediate filament proteins called A-type (lamin A/C) and B-type lamins. It is anchored to the nuclear membrane by an extensive array of inner nuclear membrane proteins. Lamins and inner nuclear membrane proteins contribute to the architecture of the nuclear envelope. Emerin is one of these inner nuclear membrane proteins. Its gene was discovered in 1994 because it is mutated in patients with X-linked Emery–Dreifuss muscular dystrophy (EDMD¹). Its mRNA shows ubiquitous tissue distribution with the highest expression in skeletal and cardiac muscles. In 1996, the availability of antiemerin antibodies enabled localization of the protein at the nuclear membrane.^{2,3} More precisely, emerin is anchored at the inner nuclear membrane via its hydrophobic C-terminal domain from amino acid 223 to amino acid 243.⁴ The fragment of emerin from amino acid 117 to 170 is sufficient for nuclear accumulation, but only the fragment from amino acid 3 to 170 is capable of targeting an integral membrane protein to the inner nuclear membrane.⁵ In the heart and cultured cardiomyocytes, emerin might additionally be associated with intercalated discs.⁶ Thus, it could play a ubiquitous role in association of the nuclear membrane with the lamins and a tissue-specific role in the heart in membrane anchorage to the cytoskeleton.

Emerin interacts with several proteins essential for cytoplasmic and nuclear structure and chromatin organization. Binding partners include lamin A/C, B-type lamins, actin, BAF, and lamina-associated polypeptide 1.^{7–11} Emerin interaction with lamin A/C was revealed by several teams and is supported by the fact that a large number of mutations in lamin A/C also cause EDMD.¹² Emerin generally colocalizes with lamins, being mainly at the nuclear periphery in the interphase.¹³ Furthermore, lamin A is essential for anchorage of emerin to

the inner nuclear membrane.¹⁴ However, emerin and lamins do not remain colocalized during mitosis, demonstrating that their interaction is regulated in a cell cycle-dependent manner. Emerin also binds the integral inner nuclear membrane protein lamina-associated polypeptide 1, which similarly associates with lamins and causes muscular dystrophy and cardiomyopathy when deleted from striated muscle.¹¹ Emerin and lamin A/C further bind actin at the late stages of myotube differentiation and in mature muscle, and these interactions are regulated by phosphorylation.¹⁵ Wilson and co-workers proposed that the emerin region between residues 1 and 221 caps the pointed end of actin filaments.¹⁶ Lammerding and co-workers showed that emerin regulates the activity of the mechanosensitive transcription factor megakaryoblastic leukemia 1, a myocardin family member that is pivotal in cardiac development and function, through modulation of nuclear and cytoskeletal actin polymerization.¹⁷ Furthermore, emerin organizes actin flow for centrosome orientation and nuclear movement in migrating cells.¹⁸ Emerin is also tyrosine phosphorylated after force is applied on isolated nuclei, and this event mediates the nuclear mechanical response to tension.¹⁹ As emerin-deficient mouse embryo fibroblasts have apparently normal nuclear mechanics but impaired expression of mechanosensitive genes in response to strain, it has been suggested that disease-causing loss of function mutations do not act by increasing nuclear fragility but through altered transcriptional regulation.²⁰ Interaction of emerin with the DNA-bridging protein BAF is important for nuclear membrane targeting to chromatin and chromatin decondensation during nuclear assembly²¹ and is regulated through mitotic phosphorylation.^{22,23}

Most mutations in the gene encoding emerin in patients with X-linked EDMD are nonsense or out-of-frame mutations that lead to the absence of emerin in patient cells.²⁴ Only very few EDMD cases are due to missense or short in-frame deletions. Attempts to describe the impact of these mutations on emerin structure and binding properties have been hampered by difficulties in obtaining pure and soluble fragments of the emerin nucleoplasmic region from amino acid 1 to 221.²⁵ The three-dimensional structure of the emerin N-terminal LEM domain, from amino acid 1 to 45, has been solved,²⁶ and a model of its structure in complex with a BAF dimer has been proposed.²⁷ However, it was demonstrated in HeLa cells that the LEM domain and distal tyrosines (a.a. 59, 74, 95, 161) contribute to BAF binding.²⁸ The region of emerin from amino acid 46 to 221 is predicted to be mainly intrinsically disordered.²⁹ It contains the lamin and actin binding regions of emerin. It also comprises four positions (a.a 54, 95–99, 133, 183) whose substitution or deletion do not hinder emerin expression but cause EDMD.^{7,30–32} The structural consequences of these changes in the protein are yet unknown. They could preclude specific emerin modification events or locally modify the three-dimensional structure and conformational plasticity of the protein. They could also affect assembly of protein complexes involving emerin, either by altering emerin oligomerization and/or hindering partner recognition.

We now present a combined biochemical and cellular analysis of the nucleoplasmic region of emerin, describing its self-assembly properties and providing novel data on the architecture of emerin oligomers. Wilson and co-workers recently proposed that residues 187–220, at the C-terminus of the nucleoplasmic region, are essential for emerin intermolecular association both *in vitro* and at the nuclear envelope.²⁹ We further show that emerin fragment EmN, from residue 1 to residue 187, forms filaments *in vitro*, and that two

variants with alterations in this fragment modify the organization of emerin at the nuclear envelope. From the comparison between results obtained *in vitro* and in cells on the full-length emerin protein, we discuss mechanisms that may lead to self-assembly defect and loss of function of emerin variants with amino acid substitutions or a small deletion associated with X-linked EDMD.

RESULTS AND DISCUSSION

The Region of Emerin from Amino Acid 1 to 187 Is Able to Self-associate Either through Disulfide Bridge Formation or *via* Hydrophobic Interactions

We produced two recombinant fragments of emerin in *Escherichia coli*: the whole nucleoplasmic region preceding the transmembrane domain from amino acid 1 to 221 (EmN0) and a smaller nucleoplasmic fragment from amino acid 1 to 187 (EmN; Figure 1A,B). Bioinformatics analysis using the Disopred3 Web server predicted that, outside of the LEM domain, these fragments, which include amino acid substitutions and deletions in expressed disease-associated emerin variants, are essentially intrinsically disordered (Figure 1C). However, three regions show a tendency for local order (Metadisorder score lower than 0.7); they comprise residues 74 to 106, residues 159 to 186, and residues 203 to 208, respectively. The two first regions are rich in conserved hydrophobic residues (red squared in Supporting Figure 1). One of these regions contains the residues deleted in the EDMD-associated variant del95–99 (blue squared in Supporting Figure 1). The capacities of these regions to locally adopt preferred 3D structures are currently unknown.

As EmN0 and EmN are mainly present in inclusion bodies when expressed in bacterial cells, we purified these fragments by affinity chromatography in 8 M urea. The two fragments were then refolded when dialyzed in Tris 50 mM (pH 6.8) and NaCl 150 mM. EmN0 was essentially only obtained as aggregates in all tested conditions (including pH from 5.0 to 9.0 and NaCl concentrations from 30 to 500 mM), as previously observed.²⁵ Therefore, we decided to use the more soluble EmN for the rest of our study. Analysis of the NMR ¹H–¹⁵N HSQC spectrum of refolded EmN showed that this emerin fragment is only partially structured. Indeed, about 50 HSQC signals show a nonrandom coil ¹H chemical shift and/or correspond to positive ¹H → ¹⁵N NOE values (Figure 1D). Most of these signals can be assigned to the emerin LEM domain based on previous ¹H chemical shift assignments (BMRB entry 5074). The remaining HSQC signals have a ¹H chemical shift comprised between 7.8 and 8.4 ppm (Figure 1D). Such low dispersion of the backbone ¹H resonances suggests that they correspond to EmN unstructured residues. Moreover, analysis of the NMR ¹H → ¹⁵N NOE experiment shows that the corresponding ¹H–¹⁵N correlations are absent from the saturated experiment spectrum when compared to the ¹H–¹⁵N HSQC spectrum, which confirmed that these EmN residues are highly flexible in solution (Figure 1D).

We next tested the impact of the refolding buffer on emerin solubility and oligomerization by analytical ultracentrifugation (Figure 2A). In Tris 20 mM (pH 8.0), EmN at 20 μM exists mainly as two different species in solution, as observed in sedimentation velocity experiments. These species are characterized by sedimentation coefficients of 2.2 and 3.3 S, as well as form factors f/f_0 of 1.52 and 1.58, respectively. They correspond to elongated

monomeric and dimeric EmN, as confirmed by sedimentation equilibrium experiments (Supporting Figure 2). Nonreducing SDS-PAGE analysis of the EmN samples in the presence of increasing β -mercaptoethanol concentrations showed that the unique cysteine of EmN (Cys147) is responsible for dimerization through disulfide bridge formation (Figure 2B).

Increasing the salt concentration close to physiological ionic strength favored the formation of larger molecular mass oligomers and/or aggregates of emerlin (Table in Figure 2A) that were too large to be observed by NMR. The presence of these large species at a high salt concentration suggested that hydrophobic interactions play a role in emerlin self-assembly and/or aggregation. Similar experiments carried out after dialysis at lower pH's (5.0 and 6.8), and thus closer to the isoelectric point (5.0) of EmN, led to a massive precipitation of the samples (data not shown). This further illustrated that increasing the weight of hydrophobic interactions compared to electrostatic interactions favored emerlin self-assembly and/or aggregation. Finally, gel filtration experiments were performed in Tris 20 mM (pH 8.0) at two different protein concentrations (40 and 80 μ M, corresponding to 1 and 3 mg mL⁻¹, respectively). Increasing EmN concentration led to the formation of high molecular oligomers under these conditions (Figure 2C).

We concluded that EmN was capable of forming different oligomers at physiological ionic strength, and the addition of a reducing agent and/or salt regulated EmN oligomerization/aggregation state. As the nucleus is characterized by a relatively reducing environment³³ and as the emerlin unique cysteine is not conserved between species (residue 147; Supporting Figure 1), the biological relevance of the disulfide-mediated dimer seemed limited. We choose to control the EmN oligomeric state by adding a reducing agent (10 mM β -mercaptoethanol) to the urea buffer and analyze EmN after an overnight dialysis in Tris 20 mM (pH 8.0) and NaCl 30 mM. In these reducing and low salt conditions, EmN is initially mainly monomeric (Supporting Figure 3). Moreover, it forms high molecular weight oligomers/aggregates with time, as revealed for example by following, using NMR, a sample of EmN at 600 μ M in Tris 20 mM (pH 7.2), NaCl 30 mM, and 50 mM β -mercaptoethanol: the EmN NMR signals disappeared within a few days at 10 °C because of the formation of species too large to be observed by NMR.

EmN Forms Filaments Rich in β -Structure *in Vitro*

We studied oligomerization of EmN over time using negative staining electron microscopy (EM). After dialysis, the protein sample was split into four aliquots which were concentrated to either 80 or 400 μ M, and either kept at RT or heated at 65 °C for 1 h; all aliquots were finally kept at RT over 3 days. Observation of these aliquots using EM (Figure 3) revealed that at 20 °C, EmN forms either amorphous aggregates and a few irregular filament-like structures (80 μ M; Figure 3A) or spherical particles of a diameter of about 10 nm as well as filament-like structures (400 μ M; Figure 3B). After heating the sample at 65 °C for 1 h, EmN forms spherical particles as well as short filaments already at 80 μ M (Figure 3C). At a high concentration (400 μ M), heating promotes the formation of long and curvilinear filaments with diameters of about 10 nm (Figure 3D). These data show that an increase in both the temperature (20 to 65 °C) and the concentration (80 and 400 μ M) favors the *in vitro*

formation of long and well-separated filaments. In order to test the impact of oxidation on filament assembly, either H₂O₂ or β -mercaptoethanol was added to the EmN sample just before heating over 1 h at 65 °C. EM analyses showed that only amorphous aggregates form in the presence of H₂O₂ (Figure 3E). Thus, although oxidative conditions favor dimer formation (as shown in Figure 2B), they inhibit filament assembly. Instead, reducing conditions allow filament assembly (Figure 3F).

We monitored first structural events associated with oligomerization of EmN by the fluorescent dye thioflavin T (ThT). This benzothiol dye has been used for decades in the diagnosis of protein-misfolding diseases and in kinetic studies of self-assembly. It interacts with β -sheet structures, in particular by docking onto surfaces formed by a single tyrosine ladder.³⁴ This interaction is revealed by a significant increase in fluorescence intensity at 480 nm. We observed a 10-fold increase in fluorescence intensity at 480 nm when EmN at 300 μ M is incubated for either 1 h at 65 °C or 3 h at 37 °C (Figure 4A). This suggested that EmN with time forms species characterized by the presence of an amyloid-like β -structure. Furthermore, the emerlin samples were characterized by ATR-FTIR spectroscopy. The amide I spectral region (1600–1700 cm⁻¹) is informative about the stretching mode vibrations of the backbone carbonyl groups and is affected by the secondary structure of the polypeptide chain. In the spectrum of monomeric emerlin, amide I vibrations evolved at 1650 cm⁻¹ (black in Figure 4B), which suggested the presence of a random coil-like structure. After incubation for 1 h at 65 °C, an extra shoulder at 1615 cm⁻¹ (green in Figure 4B) was observed, which revealed the formation of a β -sheet structure related to emerlin peptide self-assembly. Analysis using SDS-PAGE revealed that incubation for 4 h at 65 °C led to the formation of high molecular weight oligomers unable to enter the gel (Figure 4C). The presence of these oligomers yielded to a substantial decrease in total protein staining. EM images of EmN incubated for 4 h at 65 °C confirmed that these oligomers were either spherical particles or filaments with a diameter of about 10 nm (Figure 4D). Similar analyses carried out after incubation at 37 °C showed that after 4 h the EmN sample can still be observed by SDS-PAGE (Figure 4C) and that well-structured filaments can only be observed after several days (Figure 4D). Thus, both the formation of a β -structure and assembly of filaments were facilitated when increasing the temperature from 37 to 65 °C. However, we also observed the EmN filaments after incubation at 37 °C.

EmN Variants del95–99 and Q133H Occurring in EDMD Are Impaired in Their Capacity to Self-assemble

We produced five EmN variants, corresponding to emerlin variants that occur in EDMD: S54F, del95–99 (the deleted sequence being YEESY), Q133H, P183T, and P183H. These variants are similarly mainly monomeric after dialysis (Supporting Figure 3). We further tested their capacity to self-assemble.

Following EmN self-assembly kinetics at pH 8.0 and 37 °C using ThT revealed that variants S54F and P183T behave as wild-type EmN (Figure 5A). In contrast, Q133H showed a reduced ability to form β -rich structures, and del95–99 did not form any type of β -rich assembly within 1 day. Further analysis of del95–99 after incubation for 4 h at 65 °C showed that even under these conditions, the variant could be observed on SDS-PAGE gel, whereas

the wild-type EmN was already assembled into high molecular weight oligomers (Figure 5B). Observation by negative staining EM of the different variants after incubation at a concentration higher than 300 μ M and a temperature of 65 °C for 1 h confirmed that, if del95–99 was unable to form filaments under these conditions (data not shown), Q133H and P183T self-assembled into filaments (Figure 6). Filaments obtained from variant P183T were indistinguishable from wild-type filaments, whereas filaments formed by Q133H at a similar time point were systematically shorter and less regular. In the case of P183H, the self-assembly kinetics were not significantly different from those measured for wild-type EmN. However, after 1 day, further self-assembly or aggregation systematically led to a decrease in detectable fluorescent signal. Such an event was also observed in the case of wild-type EmN and variants S54F and P183T but only after several days. When wild-type EmN and the variants were dialyzed against a buffer at pH 6.8, closer to the EmN isoelectric point (5.0), most of them could not be concentrated to 300 μ M because of aggregation. Only Q133H and del95–99 could yield this concentration. Characterization of their capacity to form a β -structure after 1 day using the ThT assay showed that, at pH 6.8, Q133H had an improved capacity to self-assemble as compared to pH 8.0, whereas del95–99 still did not form filament-like structures (Supporting Figure 4). Thus, lowering the pH from 8.0 to 6.8 favored either aggregation or self-assembly but did not change the relative capacities of the variants to self-assemble. Altogether, these experiments suggest that P183H has the strongest capacity to self-assemble or aggregate, whereas Q133H and then del95–99 have the lowest capacities to form filament-like structures.

Proximity Ligation Assays Suggest the Presence of Emerin Oligomers in Cells and Reveal the Impact of EDMD-associated Variants on Emerin–Emerin and Emerin–Lamin Proximities at the Nuclear Periphery

We carried out experiments in HeLa cells in order to detect intermolecular emerin–emerin and emerin–lamin proximities. Cells were first transfected with plasmids coding for both wildtype green fluorescent protein (GFP)-full length emerin and FLAG-full length emerin (Figure 7A). GFP-FLAG proximities were then assessed using proximity ligation assays (PLA; Figure 7B). These experiments revealed emerin–emerin proximities mostly located in the nucleus, and particularly enriched at the nuclear periphery. Additional proximities were observed in the cytoplasm that could correspond to emerin located in the endoplasmic reticulum. The effects of three emerin variants with different filament assembly properties *in vitro*, del95–99, Q133H, and P183T, were then tested on emerin architecture in cells. Cells were transfected with a couple of plasmids coding for GFP and FLAG fusions of each variant (Figure 7A). Although HeLa cells express a pool of endogenous wild-type emerin, the design of our PLA assay allowed quantification of emerin–emerin proximity events occurring only between ectopic GFP and FLAG-tagged emerins. Quantification of PLA signals within the nuclear compartment demonstrated that variants del95–99 and Q133H created less emerin–emerin proximities than wild-type emerin at the nuclear periphery of transfected cells (Figure 7C). Variant P183T uniquely accumulated into foci in the endoplasmic reticulum. Depending on the experiment and due to variability in the fraction of this variant to localize in the inner nuclear membrane, the amount of emerin–emerin proximities visualized by PLA was not reproducible.

Cells were also transfected with only one plasmid encoding GFP-emerin (Figure 8A). In these cells, proximities between GFP-emerin and endogenous lamin A/C were observed at the intranuclear periphery (Figure 8B). We further measured the proximities between the emerin del95–99, Q133H, P183T, and lamin A/C by PLA. Variants del95–99 and Q133H were significantly less frequently close to lamin A/C in HeLa cells (Figure 8C). In the case of emerin P183T, which is not correctly targeted to the nuclear envelope, the amount of emerin–lamin A/C proximities visualized by PLA was not reproducible. However, intense PLA signals could be observed at the nuclear envelope, suggesting that variant P183T is still capable of interacting with lamin A/C.

A Specific Role for Proline 183 in Regulating Emerin Oligomerization

Proline is a residue playing an important structural role in IDRs by hindering their folding into aggregation prone structures. Mutation of this residue is frequently described as favoring the formation of amyloid-like structures. In the case of EmN, amino acid substitutions P183T and P183H do not significantly modify EmN filament assembly properties *in vitro* at pH 8.0. The molecular impact of these mutations is limited to a lower solubility of variant P183H at pH 8.0. However, we show in HeLa cells and in the context of the whole emerin protein that GFP-emerin P183T shows a significant tendency to form cytoplasmic foci close to the nuclear membrane. Ellis et al. previously demonstrated that in patient cells, emerin variants P183T and P183H were no longer confined to the nuclear fraction but were also distributed in other membranous fractions, with the majority being associated with “heavier” membrane fractions (lysosomes, plasma membrane, mitochondria), most particularly in the case of P183H.³¹ Wilson and co-workers demonstrated that deletion of the region from amino acid 168 to 186 favored assembly of emerin oligomers stabilized by intermolecular interactions involving the region from amino acid 187 to 220 *in vitro* and in cells.²⁹ They showed that it caused a relocalization of emerin into perinuclear aggregates. These data suggest that the region from amino acid 168 to 186 and in particular Pro183 limits oligomerization through the region from amino acid 187 to 220 in full-length emerin. We propose that emerin can form two types of intermolecular interactions in the reducing environment of the nucleus, one resulting from EmN self-assembly and the other resulting from dimerization of the region from amino acid 187 to 220. In this frame, Pro183 would play a critical role in regulating emerin self-assembly, not by directly influencing EmN filament formation but by limiting the formation of oligomers stabilized by self-assembly of the region from amino acid 187 to 220 in cells. Variants P183T and P183H would consistently show a stronger tendency to form dimers through the region from amino acid 187 to 220, which would lead to aggregation and mislocalization in the cytoplasm.

Relationship between Emerin Oligomeric State and Lamin Binding Capacity

Binding of emerin to lamin A, which has been described using several techniques including yeast two-hybrid assays and coimmunoprecipitation experiments, involves the fragment from amino acid 70 to 164 of emerin.³⁵ Within this region, deletion del95–99 caused emerin to be present but significantly less associated with membranes and to the nuclear lamina components in patient cells³⁰ and hindered binding to lamin A *in vitro* as judged by blot overlay assays.¹⁰ From our results, we now propose a model in which a loss of function of

emerin variant del95–99 (weaker lamin A/C binding by PLA) relies on its defective organization at the nuclear envelope (weaker emerin–emerin associations by PLA) and is related to its self-assembly defects (absence of oligomerization *in vitro*). For variant Q133H, no qualitative or quantitative differences in nuclear targeting were observed between mutant and normal emerin, and quantitative BIAcore analysis showed no significant change in lamin A binding to emerin when the mutation was present.³² However, our results showed that the amount of emerin–lamin A/C proximities is decreased, suggesting that emerin self-assembly defects impact lamin recognition in cells. It is also possible that emerin self-assembly defects affects its interaction with other binding proteins, such as lamina-associated polypeptide 1.

Variants P83T and P183H present a mutation outside of the lamin A/C binding region of emerin. In contrast to mutations del95–99 and Q133H, we observed that mutation P183T does not prevent binding to lamin A/C (as judged by PLA). Based on an *in vitro* blot overlay assay,¹⁰ Lee et al. proposed that variant P183H has an even increased lamin binding capacity as compared to wild-type emerin. We could not see such a clear effect for variant P183T in our cell assay. We propose that the observed mistargeting of P183T emerin within cytoplasmic aggregates might be favored upon specific stimuli in the muscular tissue and consequently impact (in an indirect fashion) the formation of lamin A/C–emerin complexes at the nuclear envelope. Further work is needed to determine to what extent amino acid substitutions P183T and P183H affect lamin A/C binding in various cellular contexts and to determine whether this can be related to the proposed increased dimerization capacity through region 170–220 of that variant (see above).

Conclusion

Our study provides a first structural description of the region from amino acid 1 to amino acid 187 of emerin and reveals the structural consequences of several mutations that occur in EDMD on the self-assembly capacity of this EmN fragment *in vitro* and the amount of emerin–emerin proximities in cells. Full-length emerin self-associates either through the EmN fragment, as shown in this study, or through the region between residues 187 and 220, as shown by others.²⁹ Oxidation of EmN hinders its self-assembly into high molecular weight oligomers. Mutations at positions 95–99 and 133 affect EmN self-assembly and impact emerin binding to lamin A/C in cells, whereas mutations of Pro183 affect emerin self-assembly probably mainly through the region between residues 187 and 220 and impact emerin localization in cells. Virtually all of the known EDMD-causing mutations lead to an absence of emerin in cells, suggesting that S54F, del95–99, Q133T, P183T, and P183H are loss-of-function mutations. Defects in oligomerization properties essential for emerin localization and function might explain the decreased functional capacities of the variants del95–99, Q133T, P183T, and P183H. Variant S54F does not impact EmN self-assembly *in vitro*, and it is still correctly localized in cells.³⁶ However, Ser54 is either phosphorylated or *O*-GlcNAc-modified in wild-type emerin, and mutation of Ser54 into phenylalanine impairs post-translational modifications at several other positions in full-length emerin.³⁶ As the emerin regions proposed to mediate EmN self-assembly are hotspots for phosphorylation and *O*-GlcNAcylation in cells, we suggest that S54F might interfere with emerin–emerin interactions by perturbing the modification state of emerin *in vivo*. Overall, our results

suggest that emerin self-assembly is necessary for its proper function and that a loss of either the protein itself or its ability to self-assemble causes muscular dystrophy.

METHODS

Protein Expression and Purification

Human wild-type emerin fragments from amino acid 1 to amino acid 187 (EmN) and from amino acid 1 to amino acid 221 (EmN0) were expressed using a pETM13 vector as N-terminal octa-histidine fusions in *Escherichia coli* BL21 DE3 Star (Novagen). The emerin cDNAs were optimized for expression in *Escherichia coli* (GenScript). Mutations encoding emerin variants S54F, Q133H, 95–99, P183H, and P183T were inserted in the EmN cDNA using a standard QuikChange Site-Directed Mutagenesis kit (Stratagene). The EmGCN4 protein was also expressed using pETM13. Its cDNA was obtained by synthesizing an optimized sequence coding for the N-terminal octa-histidine tagged EmN, a TEV cleavage site, and a GCN4-derived peptide of 33 amino acids that efficiently dimerizes.³⁷ Bacteria were cultured in ¹⁵N-labeled minimum medium, induced at an optical density of 1 with 0.1 mM isopropyl β-D-1-thiogalactopyranoside, grown overnight at 20 °C, and lysed in 50 mM Tris-HCl (pH 8.0), 300 mM NaCl, 40 mM imidazole, 5% glycerol, 1% Triton X-100, and 1 mM phenylmethanesulfonylfluoride. After centrifugation at 20 000g for 20 min at 4 °C, the pellet was resuspended in buffer C8 (50 mM Tris-HCl pH 8.0, 150 mM NaCl, 40 mM imidazole, 8 M urea). A second centrifugation step was performed at 20 000g for 20 min at 4 °C. The soluble extract was then filtered and loaded onto a Ni-NTA column (GE-Healthcare) equilibrated with buffer C8. Proteins were eluted directly with buffer E8 (50 mM Tris-HCl pH 8, 150 mM NaCl, 1 M imidazole, 8 M urea). Proteins were refolded by dialysis in the desired buffer. The final yield was typically 10 mg of purified protein per liter of bacterial culture. We verified that purification either from the bacterial soluble fraction or from inclusion bodies yielded a protein with the same conformation by recording NMR ¹H–¹⁵N HSQC spectra on both preparations. These two NMR spectra were superimposable (data not shown), which validates the use of a protocol including denaturation and refolding steps to obtain a pure, soluble, and concentrated preparation. Oligomerization studies were performed using proteins that were reduced after affinity purification in urea by adding 10 mM β-mercaptoethanol and refolded in a buffer containing 30 mM NaCl and 10 mM dithiothreitol (DTT) to observe oligomer formation from a monomeric preparation.

NMR Spectroscopy

NMR experiments were performed on a 700 MHz Bruker Avance spectrometer equipped with a cryogenic probe. Two-dimensional ¹H–¹⁵N correlation spectra were acquired using a HSQC pulse sequence at 20 °C, on a 3-mm-diameter NMR sample tube containing 100 μM uniformly ¹⁵N-labeled EmN in 50 mM Tris (pH 6.8), 150 mM NaCl, 10 mM DTT, and 90:10%, H₂O/D₂O.

Analytical Ultracentrifugation

The 360 μL protein samples at a concentration of 20 to 30 μM were dialyzed against 20 mM Tris (pH 8.0) and 0 to 150 mM NaCl. They were subjected to sedimentation velocity runs at

20 °C and 45 000 rpm (147 280*g*) on a XLA70 analytical ultracentrifuge (Beckman Coulter, Palo Alto, USA) using a An-60Ti rotor. Optical density scans recorded every 5 min at 280 nm were analyzed using the Sedfit software³⁸ to determine the sedimentation coefficients. The EmN sample was also subjected to a sedimentation equilibrium experiment in 50 mM Tris (pH 8.0) at two successive speeds (11 500 and 13 800 rpm/9619 and 13 851*g*) and 20 °C. Data analysis was performed on the scans acquired on the equilibrated system using the Sedphat software³⁸ to determine the molecular masses of the species.

Analytical Gel Filtration

The proteins were first dialyzed in the gel-filtration buffer (20 mM Tris-HCl pH 8.0, 30 mM NaCl) before being diluted to 1 mg mL⁻¹ and loaded onto the column. Analytical gel-filtration experiments were carried out at 4 °C on a Superdex 75 10/30 column (GE Healthcare).

EM

To follow large oligomer assembly, EmN, its variants, or EmGCN4 in buffer E8 and 10 mM β -mercaptoethanol were dialyzed against 20 mM Tris (pH 8.0), 30 mM NaCl, and 10 mM DTT. They were then incubated at 20, 37, or 65 °C. Samples were taken at regular time points and observed by negative-staining transmission electron microscopy. For experiments at 20 and 65 °C, His-tagged wild-type EmN was used. For experiments at 37 °C, the His-Tag from the wild-type and mutated EmN constructs was cleaved using Tobacco Etch Virus (TEV) protease (see below), and proteins without Tag were incubated in 8 M urea and redialyzed against 20 mM Tris (pH 8.0), 30 mM NaCl and 10 mM DTT. For EM experiments, about 2–5 μ L of the sample solution was deposited on a carbon-coated Formvar copper grid. After 30 s, the sample droplet was blotted with filter paper. It was placed in contact with 10 μ L of water and then uranyl acetate 0.5%. Samples were imaged using a JEOL MET 1400 (120 keV) or a Tecnai G2 (200 keV) transmission electron microscope at 10 000 $\times$ to 30 000 $\times$ magnification.

Thioflavin Kinetics

For these experiments, the His-tag of the wild-type and mutated EmN constructs was cleaved using Tobacco Etch Virus (TEV) protease for 2 h at RT. EmN mixed to the His-tagged TEV was then incubated with Ni-NTA beads for 1 h in buffer C0 (50 mM Tris-HCl pH 8.0, 150 mM NaCl, 40 mM imidazole). It was collected by washing with C0, incubated in 8 M urea, and redialyzed in 20 mM Tris (pH 8.0), 30 mM NaCl, and 10 mM DTT. Filament assembly was followed with time by incubating EmN at the targeted concentration and temperature and by regularly taking protein aliquots to be analyzed by fluorescence. These aliquots were diluted in a thioflavin (ThT) containing buffer so as to obtain 20 μ M protein and 2.5 μ M ThT in 20 mM Tris (pH 8.0), 30 mM NaCl, and 10 mM β -mercaptoethanol. The fluorescence measurements were carried out in a 60 μ L cuvette at 20 °C using a fluorimeter JASCO ADP-303T. ThT fluorescence was monitored using an excitation wavelength of 450 nm. Fluorescence emission was read at 480 nm.

Attenuated Total Reflection–Fourier-Transform Infrared (ATR-FTIR) Spectrometry

ATR-FTIR spectra were recorded at 4 cm⁻¹ resolution with a Bruker IFS 66 spectrophotometer equipped with a 45° N ZnSe attenuated total reflection attachment. For the FTIR experiments, EmN samples were concentrated to 800 μM. The concentrator flow-through was used as the reference buffer sample for measuring buffer signal. FTIR spectra of the EmN sample and its corresponding buffer solution were initially recorded just after concentration. Thereafter, the EmN sample was incubated for 1 h at 65 °C, and additional spectra of both protein and buffer samples were recorded. Spectra of both protein and buffer samples were recorded with an average of 30 scans. The buffer spectra were subsequently subtracted from the protein spectra.

Transfection

HeLa cells were obtained from American Type Culture Collection and cultured in Minimum Essential Medium containing Glutamax (Gibco), 1% nonessential amino acids, and 10% fetal bovine serum. HeLa cells were transfected using XtremeGene 9 (Roche). After 24 h, cells were processed for immunoblotting, immunofluorescence, or Proximity Ligation Assay (PLA).

Immunoblotting

Whole cell protein extracts were suspended in Laemmli sample buffer, separated by SDS-PAGE, and transferred to nitrocellulose membranes. Membranes were blocked for 1.5 h in TBS (10 mM Tris, pH 8.0, 150 mM NaCl, 0.05% Tween-20) containing 5% dry milk, incubated with mouse antiemerin antibody (Leica; 1:300) and rabbit antilamin A/C (see ref 39 1:5000) for 1 h in TBS with 1% milk, washed four times and incubated with HRP-conjugated secondary antibodies. After four washes in TBS, proteins were detected by enhanced chemiluminescence.

Immunofluorescence Microscopy

Cells were fixed with 3% paraformaldehyde for 12 min at RT, permeabilized with phosphate-buffered saline containing 0.5% Triton for 5 min at RT, and quenched with 2% bovine serum albumin diluted in phosphate-buffered saline containing 0.1% Triton X-100. Primary antibodies were rabbit antilamin A/C (1:500;⁴⁰), rabbit anti-FLAG (Sigma, 1:150), and mouse anti-GFP (Roche, 1:150). Fluorescent labeled secondary antibodies (donkey antimouse Cy2 1:60 and donkey antirabbit Cy3 1:200) were from Jackson ImmunoResearch. DNA was stained with Hoechst 33258 (1 μg/mL).

Proximity Ligation Assay (PLA)

PLA was used to detect GFP-emerin–lamin A/C interactions and GFP-emerin–FLAG-emerin interactions based on proximity (<40 nm) of two secondary antibodies directed against these proteins. After cell fixation, cell permeabilization, and quenching (as above), pairs of primary antibodies, mouse anti-GFP 1:150 and rabbit antilamin A/C 1:500 or mouse anti-GFP 1:150 and rabbit anti-FLAG 1:150, were added to HeLa cells expressing GFP-emerin either alone or together with FLAG-emerin, respectively, for 30 min at RT. Next, Duolink PLA probe antirabbit plus, Duolink PLA probe antimouse minus, and Duolink

detection reagents orange (detected with a Cy3 filter) were used according to manufacturer's instructions (Olink, Bioscience). Confocal microscopy image acquisition was performed using a LSM 700 Laser scanning microscope (Zeiss). Quantitative analysis of PLA signals was done on images using ImageJ. Data were then analyzed by comparing median values for integrated densities of signals (Cy3) per nucleus, and statistical analyses were performed using Kruskal–Wallis tests.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Funding

This work was supported by CNRS and University Paris 7 and by funding from the French Association against Myopathies (AFM; research grant no. 17243 to S.Z.-J. and Ph.D. fellowship no. 18159 to C.S.) and from the Foundation for Medical Research (FRM; grant FDT20140931008 to I.H.). H.J.W. was supported by a grant (R01AR048997) from NIH/NIAMS.

The authors thank E. Kasotakis and M. Paternostre for constant help in carrying out ATR-FTIR and EM experiments and F.X. Theillet and P. Selenko for fruitful discussions. They acknowledge the French national infrastructure FRISBI program for access to the NMR spectrometers at CEA Saclay (Gif-sur-Yvette, France), the Functional Biology and Adaptive (BFA) institute for access to the imaging facility at University Paris 7, and the Leibniz-Institut für Molekulare Pharmakologie (FMP Berlin, Germany) for access to the fluorescence and electron microscopy facilities of the institute.

REFERENCES

1. Bione S, Maestrini E, Rivella S, Mancini M, Regis S, Romeo G, Toniolo D. Identification of a novel X-linked gene responsible for Emery-Dreifuss muscular dystrophy. *Nat. Genet.* 1994; 8:323–327. [PubMed: 7894480]
2. Manilal S, Nguyen TM, Sewry CA, Morris GE. The Emery-Dreifuss muscular dystrophy protein, emerin, is a nuclear membrane protein. *Hum. Mol. Genet.* 1996; 5:801–808. [PubMed: 8776595]
3. Nagano A, Koga R, Ogawa M, Kurano Y, Kawada J, Okada R, Hayashi YK, Tsukahara T, Arahata K. Emerin deficiency at the nuclear membrane in patients with Emery-Dreifuss muscular dystrophy. *Nat. Genet.* 1996; 12:254–259. [PubMed: 8589715]
4. Yorifuji H, Tadano Y, Tsuchiya Y, Ogawa M, Goto K, Umetani A, Asaka Y, Arahata K. Emerin, deficiency of which causes Emery-Dreifuss muscular dystrophy, is localized at the inner nuclear membrane. *Neurogenetics.* 1997; 1:135–140. [PubMed: 10732816]
5. Ostlund C, Ellenberg J, Hallberg E, Lippincott-Schwartz J, Worman HJ. Intracellular trafficking of emerin, the Emery-Dreifuss muscular dystrophy protein. *J. Cell Sci.* 1999; 112(Pt 11):1709–1719. [PubMed: 10318763]
6. Cartegni L, di Barletta MR, Barresi R, Squarzoni S, Sabatelli P, Maraldi N, Mora M, Di Blasi C, Cornelio F, Merlini L, Villa A, Cobiauchi F, Toniolo D. Heart-specific localization of emerin: new insights into Emery-Dreifuss muscular dystrophy. *Hum. Mol. Genet.* 1997; 6:2257–2264. [PubMed: 9361031]
7. Fairley EA, Kendrick-Jones J, Ellis JA. The Emery-Dreifuss muscular dystrophy phenotype arises from aberrant targeting and binding of emerin at the inner nuclear membrane. *J. Cell Sci.* 1999; 112(Pt 15):2571–2582. [PubMed: 10393813]
8. Clements L, Manilal S, Love DR, Morris GE. Direct interaction between emerin and lamin A. *Biochem. Biophys. Res. Commun.* 2000; 267:709–714. [PubMed: 10673356]
9. Sakaki M, Koike H, Takahashi N, Sasagawa N, Tomioka S, Arahata K, Ishiura S. Interaction between emerin and nuclear lamins. *J. Biochem.* 2001; 129:321–327. [PubMed: 11173535]

10. Lee KK, Haraguchi T, Lee RS, Koujin T, Hiraoka Y, Wilson KL. Distinct functional domains in emerin bind lamin A and DNA-bridging protein BAF. *J. Cell Sci.* 2001; 114:4567–4573. [PubMed: 11792821]
11. Shin JY, Mendez-Lopez I, Wang Y, Hays AP, Tanji K, Lefkowitz JH, Schulze PC, Worman HJ, Dauer WT. Lamina-associated polypeptide-1 interacts with the muscular dystrophy protein emerin and is essential for skeletal muscle maintenance. *Dev. Cell.* 2013; 26:591–603. [PubMed: 24055652]
12. Bonne G, Di Barletta MR, Varnous S, Becane HM, Hammouda EH, Merlini L, Muntoni F, Greenberg CR, Gary F, Urtizberea JA, Duboc D, Fardeau M, Toniolo D, Schwartz K. Mutations in the gene encoding lamin A/C cause autosomal dominant Emery-Dreifuss muscular dystrophy. *Nat. Genet.* 1999; 21:285–288. [PubMed: 10080180]
13. Manilal S, Nguyen TM, Morris GE. Colocalization of emerin and lamins in interphase nuclei and changes during mitosis. *Biochem. Biophys. Res. Commun.* 1998; 249:643–647. [PubMed: 9731189]
14. Vaughan A, Alvarez-Reyes M, Bridger JM, Broers JL, Ramaekers FC, Wehnert M, Morris GE, Whitfield WGF, Hutchison CJ. Both emerin and lamin C depend on lamin A for localization at the nuclear envelope. *J. Cell Sci.* 2001; 114:2577–2590. [PubMed: 11683386]
15. Lattanzi G, Cenni V, Marmiroli S, Capanni C, Mattioli E, Merlini L, Squarzoni S, Maraldi NM. Association of emerin with nuclear and cytoplasmic actin is regulated in differentiating myoblasts. *Biochem. Biophys. Res. Commun.* 2003; 303:764–770. [PubMed: 12670476]
16. Holaska JM, Kowalski AK, Wilson KL. Emerin caps the pointed end of actin filaments: evidence for an actin cortical network at the nuclear inner membrane. *PLoS Biol.* 2004; 2:E231. [PubMed: 15328537]
17. Ho CY, Jaalouk DE, Vartiainen MK, Lammerding J. Lamin A/C and emerin regulate MKL1-SRF activity by modulating actin dynamics. *Nature.* 2013; 497:507–511. [PubMed: 23644458]
18. Chang W, Folker ES, Worman HJ, Gundersen GG. Emerin organizes actin flow for nuclear movement and centrosome orientation in migrating fibroblasts. *Mol. Biol. Cell.* 2013; 24:3869–3880. [PubMed: 24152738]
19. Guilluy C, Osborne LD, Van Landeghem L, Sharek L, Superfine R, Garcia-Mata R, BurrIDGE K. Isolated nuclei adapt to force and reveal a mechanotransduction pathway in the nucleus. *Nat. Cell Biol.* 2014; 16:376–381. [PubMed: 24609268]
20. Lammerding J, Hsiao J, Schulze PC, Kozlov S, Stewart CL, Lee RT. Abnormal nuclear shape and impaired mechanotransduction in emerin-deficient cells. *J. Cell Biol.* 2005; 170:781–791. [PubMed: 16115958]
21. Segura-Totten M, Kowalski AK, Craigie R, Wilson KL. Barrier-to-autointegration factor: major roles in chromatin decondensation and nuclear assembly. *J. Cell Biol.* 2002; 158:475–485. [PubMed: 12163470]
22. Hirano Y, Segawa M, Ouchi FS, Yamakawa Y, Furukawa K, Takeyasu K, Horigome T. Dissociation of emerin from barrier-to-autointegration factor is regulated through mitotic phosphorylation of emerin in a xenopus egg cell-free system. *J. Biol. Chem.* 2005; 280:39925–39933. [PubMed: 16204256]
23. Bengtsson L, Wilson KL. Barrier-to-autointegration factor phosphorylation on Ser-4 regulates emerin binding to lamin A in vitro and emerin localization in vivo. *Mol. Biol. Cell.* 2006; 17:1154–1163. [PubMed: 16371512]
24. Yates JR, Wehnert M. The Emery-Dreifuss Muscular Dystrophy Mutation Database. *Neuromuscul. Disord.* 1999; 9:199. [PubMed: 10382916]
25. Roberts RC, Sutherland-Smith AJ, Wheeler MA, Jensen ON, Emerson LJ, Spiliotis II, Tate CG, Kendrick-Jones J, Ellis JA. The Emery-Dreifuss muscular dystrophy associated-protein emerin is phosphorylated on serine 49 by protein kinase A. *FEBS J.* 2006; 273:4562–4575. [PubMed: 16972941]
26. Wolff N, Gilquin B, Courchay K, Callebaut I, Worman HJ, Zinn-Justin S. Structural analysis of emerin, an inner nuclear membrane protein mutated in X-linked Emery-Dreifuss muscular dystrophy. *FEBS Lett.* 2001; 501:171–176. [PubMed: 11470279]

27. Cai M, Huang Y, Suh JY, Louis JM, Ghirlando R, Craigie R, Clore GM. Solution NMR structure of the barrier-to-autointegration factor-Emerin complex. *J. Biol. Chem.* 2007; 282:14525–14535. [PubMed: 17355960]
28. Tifft KE, Bradbury KA, Wilson KL. Tyrosine phosphorylation of nuclear-membrane protein emerin by Src, Abl and other kinases. *J. Cell Sci.* 2009; 122:3780–3790. [PubMed: 19789182]
29. Berk JM, Simon DN, Jenkins-Houk CR, Westerbeck JW, Gronning-Wang LM, Carlson CR, Wilson KL. The molecular basis of emerin-emerin and emerin-BAF interactions. *J. Cell Sci.* 2014; 127:3956–3969. [PubMed: 25052089]
30. Ellis JA, Craxton M, Yates JR, Kendrick-Jones J. Aberrant intracellular targeting and cell cycle-dependent phosphorylation of emerin contribute to the Emery-Dreifuss muscular dystrophy phenotype. *J. Cell Sci.* 1998; 111(Pt 6):781–792. [PubMed: 9472006]
31. Ellis JA, Yates JR, Kendrick-Jones J, Brown CA. Changes at P183 of emerin weaken its protein-protein interactions resulting in X-linked Emery-Dreifuss muscular dystrophy. *Hum. Genet.* 1999; 104:262–268. [PubMed: 10323252]
32. Holt I, Clements L, Manilal S, Morris GE. How does a g993t mutation in the emerin gene cause Emery-Dreifuss muscular dystrophy? *Biochem. Biophys. Res. Commun.* 2001; 287:1129–1133. [PubMed: 11587540]
33. Go YM, Jones DP. Redox control systems in the nucleus: mechanisms and functions. *Antioxid. Redox Signaling.* 2010; 13:489–509.
34. Biancalana M, Makabe K, Koide A, Koide S. Molecular mechanism of thioflavin-T binding to the surface of beta-rich peptide self-assemblies. *J. Mol. Biol.* 2009; 385:1052–1063. [PubMed: 19038267]
35. Berk JM, Tifft KE, Wilson KL. The nuclear envelope LEM-domain protein emerin. *Nucleus.* 2013; 4:298–314. [PubMed: 23873439]
36. Berk JM, Maitra S, Dawdy AW, Shabanowitz J, Hunt DF, Wilson KL. O-Linked beta-N-acetylglucosamine (O-GlcNAc) regulates emerin binding to barrier to autointegration factor (BAF) in a chromatin- and lamin B-enriched "niche". *J. Biol. Chem.* 2013; 288:30192–30209. [PubMed: 24014020]
37. Ciani B, Bjelic S, Honnappa S, Jawhari H, Jaussi R, Payapilly A, Jowitt T, Steinmetz MO, Kammerer RA. Molecular basis of coiled-coil oligomerization-state specificity. *Proc. Natl. Acad. Sci. U. S. A.* 2010; 107:19850–19855. [PubMed: 21045134]
38. Schuck P. Size-distribution analysis of macromolecules by sedimentation velocity ultracentrifugation and lamm equation modeling. *Biophys. J.* 2000; 78:1606–1619. [PubMed: 10692345]
39. Duband-Goulet I, Woerner S, Gasparini S, Attanda W, Konde E, Tellier-Lebegue C, Craescu CT, Gombault A, Roussel P, Vadrot N, Vicart P, Ostlund C, Worman HJ, Zinn-Justin S, Buendia B. Subcellular localization of SREBP1 depends on its interaction with the C-terminal region of wild-type and disease related A-type lamins. *Exp. Cell Res.* 2011; 317:2800–2813. [PubMed: 21993218]
40. Vadrot N, Duband-Goulet I, Cabet E, Attanda W, Barateau A, Vicart P, Gerbal F, Briand N, Vigouroux C, Oldenburg AR, Lund EG, Collas P, Buendia B. The p.R482W substitution in A-type lamins deregulates SREBP1 activity in Dunnigan-type familial partial lipodystrophy. *Hum. Mol. Genet.* 2015; 24:2096–2109. [PubMed: 25524705]
41. Kozlowski LP, Bujnicki JM. MetaDisorder: a meta-server for the prediction of intrinsic disorder in proteins. *BMC Bioinf.* 2012; 13:111.

Figure 1.

Structural analysis of emerlin region from residue 1 to residue 187 (EmN). This region is predicted and observed to be mainly unstructured in solution. (A) Schematic representation of emerlin. The two structured domains are represented in orange for the LEM domain (LEM 1–45) and in green for the transmembrane domain (TM 223–243). (B) Emerlin fragments expressed for this study. (C) Metadisorder profile of emerlin:⁴¹ apart from the two structured domains (LEM and TM) that have a disorder probability lower than 0.5, most of the other residues are predicted as disordered. Nevertheless, three regions, between residues 74 and 106, residues 159 and 186, and residues 203 and 208, show intermediate scores between 0.5 and 0.7. The red dots mark positions mutated in EDMD causing missense or microdeletion variants. (D) 2D NMR ^1H - ^{15}N spectra recorded on a EmN sample at 100 μM in 50 mM Tris-HCl pH6.7, 150 mM NaCl, 10 mM DTT at 20 $^{\circ}\text{C}$, and 700 MHz. Superposition of the ^1H - ^{15}N HSQC spectrum in black (each peak corresponds to one residue; peaks corresponding to poorly structured residues have a ^1H chemical shift comprised between 7.8 and 8.4 ppm) and the saturated $^1\text{H} \rightarrow ^{15}\text{N}$ NOE spectrum in orange (positive peaks corresponding to structured residues) and blue (negative peaks corresponding to fully disordered residues). Residues with intermediate dynamical behavior have NOE values close to zero and thus are not visible on the saturated ^1H to ^{15}N NOE spectrum.

Figure 2.

Identification of EmN dimers and higher molecular weight oligomers/aggregates formed in solution. (A) AUC sedimentation velocity results obtained on samples of EmN at 20 μ M in 20 mM Tris-HCl pH 8, under three different salt conditions (0 mM, 30 mM, and 150 mM NaCl). Three species are present in these conditions, which might correspond to an elongated monomer (sedimentation coefficient of 2.1S and frictional ratio of 1.52), an elongated dimer (sedimentation coefficient of 3.2S and frictional ratio of 1.58), and high molecular weight oligomers and/or aggregates. (B) Nonreducing SDS-PAGE of EmN in 20 mM Tris (pH 8.0), 30 mM NaCl, and different concentrations of β -mercaptoethanol showing a disappearance of EmN dimers in the presence of a reducing agent. (C) Gel filtration profile obtained after dialysis of EmN: samples 1 (black full lines) and 2 (black dashed lines) were injected on the column at 1 and 3 mg mL^{-1} , respectively. Clearly, sample 2 contains oligomers and/or aggregates that could not pass the column prefilter. Both samples also contain monomers (elution volume corresponding to 33 kDa, the protein being a 25 kDa elongated protein), dimers (elution volume corresponding to 66 kDa), and oligomers (elution at the void volume).

Figure 3.

EmN filaments formed in a concentration/temperature/reducing conditions dependent manner. (A–D) EM images acquired on an EmN sample concentrated at either 80 or 400 μM in a low salt concentration buffer (20 mM Tris-HCl [pH 8.0], 30 mM NaCl, 10 mM β-mercaptoethanol) and not heated or heated for 1 h at 65 °C. (E,F) EM images acquired on a EmN sample concentrated at 400 μM with either (E) 10 mM H₂O₂ or (F) 10 mM β-mercaptoethanol and heated 1 h at 65 °C in a low salt concentration buffer (20 mM Tris-HCl [pH 8.0], 30 mM NaCl). All samples were observed by negative staining EM using uranyl acetate after 3 days at 20 °C. The black bars correspond to 200 nm.

Figure 4.

EmN filament assembly at either 37 or 65 °C associated with an increase of β -structure.

EmN self-assembly kinetics followed by (A) ThT fluorescence on a EmN sample at 300 μ M in 20 mM Tris-HCl (pH8.0), 30 mM NaCl, and 10 mM β -mercaptoethanol, heated at either 37 °C (orange) or 65 °C (green) for 4 h, (B) ATR-FTIR on a EmN sample at 1.2 mM before (black) and after (green) heating for 1 h at 65 °C, (C) SDS-PAGE on a EmN sample at 300 μ M (S and P correspond to Supernatant and Pellet, respectively; the arrow corresponds to the emerim monomer), and (D) EM on a EmN sample at 700 μ M (the black bar corresponds to 200 nm; negative staining was achieved using uranyl acetate).

Figure 5.

Oligomerization capacities of EmN EDMD-causing variants. Del95–99 and Q133H have defective assembly properties. (A) Fluorescence signals measured at 480 nm after the addition of ThT. The wild-type and mutated EmN samples were concentrated up to 300 μ M in 20 mM Tris-HCl pH 8, 30 mM NaCl, and 10 mM β -mercaptoethanol and heated for 1 day at 37 °C. (B) SDS-PAGE of wild-type and del95–99 EmN after heating the samples at 65 °C for either 1 or 4 h (S and P correspond to Supernatant and Pellet, respectively; the arrows correspond to the protein monomers).

Figure 6.

Filament assembly of variants Q133H and P183T. These variants form filaments after heating for 1 h at 65 °C. Filaments formed by Q133H were systematically shorter and less regular than filaments obtained from variant P183T and wild-type EmN at a similar time point. EM images were acquired on samples of wild-type EmN (left) and variants Q133H (center) and P183T (right), concentrated above 300 μ M and heated for 1 h at 65 °C. The black bars correspond to 100 nm. Negative staining was achieved using uranyl acetate.

Figure 7.

Impact of emerlin variants expressed in EDMD on nuclear emerlin–emerlin proximities as detected by PLA. (A) Whole cell protein extracts prepared from HeLa cells coexpressing either wild-type, 95–99, Q133H, or P183T GFP and FLAG-emerlins were analyzed by Western blot using antiemerlin antibodies. The expression of endogenous A-type lamins revealed by anti-LA/C antibodies is also shown. (B) HeLa cells coexpressing either wild-type (WT), del95–99, Q133H, or P183T GFP and FLAG-tagged emerlins were fixed, labeled with anti-GFP and anti-FLAG antibodies, and processed for PLA before analysis at the confocal microscope. The PLA signals are shown alone (upper row) and merged with DNA staining and GFP fluorescence (lower row). (C) Quantification of PLA signals per nucleus for cells as shown in B. Shown are the median values for the total signal intensity per nucleus for three independent experiments (EXP1, EXP2, EXP 3; $n = 150\text{--}330$ nuclei per

sample). Boxes show first and third quartiles (** $p < 0.001$, * $p < 0.05$ with Kruskal and Wallis test).

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Figure 8.

Impact of emerlin variants expressed in EDMD on nuclear emerlin–lamin A/C proximities as detected by PLA. (A) Whole cell protein extracts prepared from HeLa cells expressing either WT, 95–99, Q133H, or P183T GFP–emerlin were analyzed by Western blot using anti-emerlin antibodies. Shown is also the expression of endogenous A-type lamins revealed by anti-LA/C antibodies, and processed for PLA. (B) HeLa cells overexpressing either WT or mutant del95–99, Q133H, and P183T GFP–emerlin were fixed, labeled with anti-GFP and anti-LA/C antibodies, and processed for PLA before analysis at the confocal microscope. The PLA signals are shown alone (upper row) and merged with DNA staining and GFP

fluorescence (lower row). (C) Quantification of PLA signals per nucleus for cells as shown in B. Shown are the median values for the total signal intensity per nucleus for three independent experiments (EXP1, EXP2, EXP 3; $n = 180\text{--}340$ nuclei per sample). Boxes show first and third quartiles (** $p < 0.001$, * $p < 0.05$ with Kruskal and Wallis test).