Published in final edited form as:

Adv Exp Med Biol. 2015 ; 887: 1–14. doi:10.1007/978-3-319-22380-3_1.

A Fleeting Glimpse Inside microRNA, Epigenetics, and Micropeptidomics

Gaetano Santulli, M.D., Ph.D.

Columbia University Medical Center, New York, NY, USA, gsantulli001@gmail.com

Abstract

MicroRNAs (miRs) are important regulators of gene expression in numerous biological processes. Their maturation process is herein described, including the most updated insights from the current literature. Circa 2000 miR sequences have been identified in the human genome, with over 50,000 miR-target interactions, including enzymes involved in epigenetic modulation of gene expression. Moreover, some “pieces of RNA” previously annotated as noncoding have been recently found to encode micropeptides that carry out critical mechanistic functions in the cell. Advanced techniques now available will certainly allow a precise scanning of the genome looking for micropeptides hidden within the “noncoding” RNA.

Keywords

miRNA; Micropeptides; ORF; Micropeptidome; Mitochondria Myogenin; Humanin; SERCA; MOTS-c; Micropeptidomics; RISC; Pharmacogenomics; Drosha; Dicer; METTL3; Exportin; TargetScan; miRWalk; miRBase; EpimiR; Transcriptome; Precision medicine

Introduction

MicroRNAs (miRs) are an evolutionarily conserved family of small (~22 nucleotides) generally [1–4] noncoding RNAs, first discovered in *Caenorhabditis elegans* [5–8]. They represent a vital component of genetic regulation, existing in virtually all organisms, suggesting thereby a pivotal role in biological processes. Undeniably, miRs are important regulators of gene expression in a plethora of biological processes including cellular proliferation, differentiation, and tumorigenesis [9–26]. Other examples of noncoding RNAs are reported in Table 1.1.

Biogenesis

Classically, miRs are regarded as negative regulators of gene expression that inhibit translation and/or promote mRNA degradation by base pairing to complementary sequences within the 3'-untranslated region (3'-UTR) of protein-coding mRNA transcripts [27, 28]—mRNA degradation accounts for the majority of miR activity [29]. By altering levels of key

regulators within complex genetic pathways, miRs provide a posttranscriptional level of control of homeostatic and developmental events [30–32].

Specific structural aspects of miRs are discussed in detail in Chap. 2 of this book. Briefly, maturation of miRs involves a multi-step process [33–35] that starts from the transcription (mainly operated by RNA polymerase II) of single-stranded nonprotein-coding RNAs, which are either transcribed as stand-alone transcripts (*intergenic* miRs), often encoding various miRs, or generated by the processing of introns of protein-coding genes (*intragenic* or *intronic* miRs). Transcription of intergenic miRs leads to the formation of primary miRs (pri-miRs) with a characteristic hairpin or stem–loop structure [36], which are subsequently processed by the nuclear RNase III, Drosha [37], and its partner proteins, including the DiGeorge Syndrome Critical Region 8 (DGCR8, known as Pasha in invertebrates), named for its association with DiGeorge Syndrome [38, 39], to become precursor miRs (pre-miRs). On the other hand, intronic miRs are obtained by the regular transcription of their host genes and then spliced to form looped pre-miRs, bypassing thereby the Drosha pathway [33, 40]. Recently, Claudio Alarcón and colleagues discovered that the addition of an m6A mark to primary miRs by methyltransferase-like 3 (METTL3) is required for their recognition by DGCR8 [41]. They also proved that METTL3 is sufficient to enhance miR maturation in a global and non-cell-type-specific manner, acting as a strategic posttranscriptional modification that promotes the initiation of miR biogenesis.

Pre-miRs are exported from the nucleus in the cytoplasm in a process involving the Ran-GTP-dependent shuttle Exportin-5 [42]. Once in the cytosol, the pre-miR hairpin is cleaved by the RNase III enzyme Dicer [43, 44], yielding a mature miR:miR* duplex about 22 nucleotides in length, which is subsequently incorporated into the protein complex called RNA-induced silencing complex (RISC) to form miRISC [45, 46]. At this point, one of the double strands, the guide strand, is selected by the argonaute protein [47], the catalytically active RNase in the RISC complex, on the basis of the thermodynamic stability of the 5' end. In particular, the strand with a less thermodynamically stable 5' end is commonly chosen and loaded into the RISC complex [48], serving as a guide for miRISC to find its complementary motifs in the 3'-UTR of the target mRNA(s). Although either strand of the mature duplex may potentially act as a functional miR, only one strand is usually incorporated into the RISC where the miR and its mRNA target interact [49, 50]. Such a binding inhibits the translation of the protein that the target mRNA encodes or promotes gene silencing via mRNA degradation [51, 52].

Nearly 2000 miR sequences have been heretofore identified in the human genome, with over 50,000 miR-target interactions. Several algorithms and bioinformatics websites, including TargetScan and miRWalk [53, 54], have been developed to predict specific mRNA/miR interactions. However, miR binding rules are quite complex and not fully understood, resulting in a lack of consensus in the literature.

Given all these crucial features, miRs could represent an important way for the cell to establish intercellular (with other cells, via secreted miRs) and intracellular (among its own genes) communication. Determining direct cause-and-effect links between miRs and mRNA

targets is essential to understanding the molecular mechanisms underlying disease and the subsequent development of targeted therapies [55, 56].

Walking through an Apparently Complicated Nomenclature: miR, *miR*, miR-Xa, miR*

Nomenclature of miRs may appear confusing to the naïve readers. Briefly, mature miRs are named using the non-italicized prefix “miR-” followed by a roman number (with the exception of a few early miRNAs including the let family); stem-loop precursor miRs are all named using the italicized prefix “*mir*-”.

Similar miR sequences are distinguished by a lettered suffix, for example, miR-200a, miR-200b, and miR-200c, without implying shared targets or functions. Identical miR sequences are distinguished by a numerical suffix: for instance, mir-7-1 (located on chromosome 9), mir-7-2 (located on chromosome 15), and mir-7-3 (located on chromosome 19) can all produce identical mature miRNAs. Mature miRs can be formed from either arm of the stem-loop precursor miRNA (pre-miR). In the majority of cases, one arm is more commonly formed than the other (guide strand). Previous convention was to name these strands according to their relative abundance, with the less common form (“passenger strand”) taking the name miR-X*. However, the latest convention is to name mature miRs by the arm of the pre-miR from which they are derived, regardless of their abundance: those from the 5' arm are named miR-X-5p and those from the 3' arm as miR-X-3p. Therefore, miR-181a is now known as miR-181a-5p and miR-181a* is now known as miR-181a-3p, avoiding problems with the previous system if the abundance of each arm changes between tissues, developmental stages, or species. All of the above naming conventions can be preceded by a three-letter code which identifies the species the miRNA is from: hsa = *homo sapiens* (human); rno = *rattus norvegicus* (rat). Therefore, miR-181a-5p found in humans could be represented as hsa-miR-181a-5p. Of note, identical miRNAs are given the same number, regardless of species.

Epigenetics and miRs: An Intricate Affair

Epigenetics is the study of heritable changes in gene expression caused by mechanisms other than changes in the underlying DNA sequences [57], including DNA methylation [58] and posttranslational modifications of chromatin proteins [59, 60]. The main enzymes involved in this process are DNA methyltransferases (DNMT), histone demethylases (HDM), histone acetylases (HAT), and histone deacetylases (HDAC). Mounting evidence demonstrates that epigenetics and miRs can affect each other in an intricate connection [61–64]. Indeed, miRs play a key role in regulating DNA methylation or histone modifications through means of directly targeting epigenetic enzymes or functional protein complexes. For instance, a global DNA hypomethylation is induced by miR-29b leading to marked reduction of the expression of DNMT1, DNMT3A, and DNMT3B and subsequent reactivation of tumor suppressor genes p15 (INK4b) and ESR1 [65, 66]. Another example is given by miR-200a, which upregulates histone H3 acetylation via direct targeting of the 3' untranslated region of the HDAC4 mRNA [67].

On the other hand, epigenetic control is involved in the regulation of miR expression. DNA methylation of promoter-associated CpG dinucleotides generally correlates with reduced transcription levels of corresponding miRs [68–70], thereby inducing the expression of miR target genes. A novel miR-148a/DNMT1 regulatory circuit has been identified in hepatocellular carcinogenesis: a member of the miR-148/152 family, miR-148a is a tumor suppressor that can be silenced by hypermethylation and interacts with DNMT1 [71].

A comprehensive database, EpimiR, in which experimentally validated mutual interactions between epigenetics and miRs are described, has been recently published [72].

The Emerging Functional Role of Micropeptidomics

Intriguingly, some so-called “noncoding” pieces of RNA may actually encode short proteins (micropeptides) that carry out critical mechanistic functions within the cell(s). A conserved micropeptide (46 amino acids), named myoregulin, encoded by RNA that had been previously misannotated as noncoding, has been recently identified by Olson’s group [2]. Myoregulin is a skeletal muscle-specific micropeptide that forms a transmembrane alpha helix within the membrane of the sarcoplasmic reticulum (SR), where it modulates Ca^{2+} handling interacting with the SR Ca^{2+} ATPase (SERCA).

Such a micropeptide displays a structural resemblance to phospholamban and sarcolipin, which inhibit SERCA activity in the heart and in slow-type and developing skeletal muscle [73]. The fact that putative long noncoding RNA may harbor hidden micropeptides had been suggested by recent genome-wide analyses [74]. However, heretofore the microproteome has largely been overlooked in gene annotations [75, 76].

Due to their small size, micropeptides could not be identified by genome annotation or by protein prediction algorithms whose threshold of detection is relatively high: indeed, in scans of the genome, a DNA sequence is usually not considered potentially protein-coding unless it can encode a string of more than 100 amino acids [77]. Of note, albeit some short peptides have crucial biological functions, these peptides are generally fragments chipped off larger proteins [78]. More of these “mysterious” RNA molecules could produce peptides too small to be considered true proteins but which nonetheless carry out important functions (Fig. 1.1).

Recently, other nonclassical peptides—encoded by small open reading frames (ORF)—have been discovered. These micropeptides are translated from ORF shorter than 100 amino acids. In contrast to other bioactive peptides, micropeptides are not cleaved from a larger precursor protein and lack an amino-terminal signaling sequence [79]. An estimated 40 % of mRNAs in the fruit fly *Drosophila melanogaster*, in which the first micropeptides were identified [80–82], might contain upstream ORFs in 5′-regions and some show signs of evolutionary conservation [83].

Exploring the mitochondrial genome, a compact circular genetic system that encodes for 13 proteins essentially dedicated to energy production, Pinchas Cohen and colleagues have identified a short ORF encoded within the mitochondrial 12S rRNA that yields a bioactive peptide, named MOTS-c (mitochondrial ORF of the 12S rRNA type-c) involved in the

regulation of metabolic homeostasis. The Cohen's laboratory was among the three groups [84–86] that independently discovered another important mitochondrial peptide, humanin, encoded in the mitochondrial genome by the 16S ribosomal RNA gene, MT-RNR2, which displays fundamental cytoprotective effects.

Similar short peptides could be hiding in several places in the genome, including in transcripts of unknown function. Hence, exploiting state-of-the-art techniques [1, 87], a major exciting field of research in the next years will be represented by scanning the microproteome embedded in the (previously annotated) “noncoding” RNA.

Is this the way toward precision medicine? We'll see.

Acknowledgments

Dr. Gaetano Santulli is supported by the National Institutes of Health (K99DK107895) and by the American Heart Association (AHA 15SDG25300007).

References

1. Bazzini AA, Johnstone TG, Christiano R, Mackowiak SD, Obermayer B, Fleming ES, Vejnar CE, Lee MT, Rajewsky N, Walther TC, Giraldez AJ. Identification of small ORFs in vertebrates using ribosome footprinting and evolutionary conservation. *EMBO J.* 2014; 33:981–993. [PubMed: 24705786]
2. Anderson DM, Anderson KM, Chang CL, Makarewich CA, Nelson BR, McAnally JR, Kasaragod P, Shelton JM, Liou J, Bassel-Duby R, Olson EN. A micropeptide encoded by a putative long noncoding RNA regulates muscle performance. *Cell.* 2015; 160:595–606. [PubMed: 25640239]
3. Laressergues D, Couzigou JM, Clemente HS, Martinez Y, Dunand C, Becard G, Combier JP. Primary transcripts of microRNAs encode regulatory peptides. *Nature.* 2015; 520:90–93. [PubMed: 25807486]
4. Wronska A, Kurkowska-Jastrzebska I, Santulli G. Application of microRNAs in diagnosis and treatment of cardiovascular disease. *Acta Physiol.* 2015; 213:60–83.
5. Ambros V. The functions of animal microRNAs. *Nature.* 2004; 431:350–355. [PubMed: 15372042]
6. Gan Z, Rumsey J, Hazen BC, Lai L, Leone TC, Vega RB, Xie H, Conley KE, Auwerx J, Smith SR, Olson EN, Kralli A, Kelly DP. Nuclear receptor/microRNA circuitry links muscle fiber type to energy metabolism. *J Clin Invest.* 2013; 123:2564–2575. [PubMed: 23676496]
7. Ruvkun G, Giusto J. The *Caenorhabditis elegans* heterochronic gene *lin-14* encodes a nuclear protein that forms a temporal developmental switch. *Nature.* 1989; 338:313–319. [PubMed: 2922060]
8. Ruvkun G, Ambros V, Coulson A, Waterston R, Sulston J, Horvitz HR. Molecular genetics of the *Caenorhabditis elegans* heterochronic gene *lin-14*. *Genetics.* 1989; 121:501–516. [PubMed: 2565854]
9. Dvinge H, Git A, Graf S, Salmon-Divon M, Curtis C, Sottoriva A, Zhao Y, Hirst M, Armisen J, Miska EA, Chin SF, Provenzano E, Turashvili G, Green A, Ellis I, Aparicio S, Caldas C. The shaping and functional consequences of the microRNA landscape in breast cancer. *Nature.* 2013; 497:378–382. [PubMed: 23644459]
10. Shen J, Xia W, Khotskaya YB, Huo L, Nakanishi K, Lim SO, Du Y, Wang Y, Chang WC, Chen CH, Hsu JL, Wu Y, Lam YC, James BP, Liu X, Liu CG, Patel DJ, Hung MC. EGFR modulates microRNA maturation in response to hypoxia through phosphorylation of AGO2. *Nature.* 2013; 497:383–387. [PubMed: 23636329]
11. Song SJ, Poliseno L, Song MS, Ala U, Webster K, Ng C, Beringer G, Brikbak NJ, Yuan X, Cantley LC, Richardson AL, Pandolfi PP. MicroRNA-antagonism regulates breast cancer stemness and metastasis via TET-family-dependent chromatin remodeling. *Cell.* 2013; 154:311–324. [PubMed: 23830207]

12. Care A, Catalucci D, Felicetti F, Bonci D, Addario A, Gallo P, Bang ML, Segnalini P, Gu Y, Dalton ND, Elia L, Latronico MV, Hoydal M, Autore C, Russo MA, Dorn GW 2nd, Ellingsen O, Ruiz-Lozano P, Peterson KL, Croce CM, Peschle C, Condorelli G. MicroRNA-133 controls cardiac hypertrophy. *Nat Med*. 2007; 13:613–618. [PubMed: 17468766]
13. Gomez IG, MacKenna DA, Johnson BG, Kaimal V, Roach AM, Ren S, Nakagawa N, Xin C, Newitt R, Pandya S, Xia TH, Liu X, Borza DB, Grafals M, Shankland SJ, Himmelfarb J, Portilla D, Liu S, Chau BN, Duffield JS. Anti-microRNA-21 oligonucleotides prevent Alport nephropathy progression by stimulating metabolic pathways. *J Clin Invest*. 2015; 125:141–156. [PubMed: 25415439]
14. Epstein DM. Special delivery: microRNA-200-containing extracellular vesicles provide metastatic message to distal tumor cells. *J Clin Invest*. 2014; 124:5107–5108. [PubMed: 25401465]
15. Sachdeva M, Mito JK, Lee CL, Zhang M, Li Z, Dodd RD, Cason D, Luo L, Ma Y, Van Mater D, Gladdy R, Lev DC, Cardona DM, Kirsch DG. MicroRNA-182 drives metastasis of primary sarcomas by targeting multiple genes. *J Clin Invest*. 2014; 124:4305–4319. [PubMed: 25180607]
16. Santulli G, Wronska A, Uryu K, Diacovo TG, Gao M, Marx SO, Kitajewski J, Chilton JM, Akat KM, Tuschl T, Marks AR, Totary-Jain H. A selective microRNA-based strategy inhibits restenosis while preserving endothelial function. *J Clin Invest*. 2014; 124:4102–4114. [PubMed: 25133430]
17. Bertero T, Lu Y, Annis S, Hale A, Bhat B, Saggari R, Saggari R, Wallace WD, Ross DJ, Vargas SO, Graham BB, Kumar R, Black SM, Fratz S, Fineman JR, West JD, Haley KJ, Waxman AB, Chau BN, Cottrell KA, Chan SY. Systems-level regulation of microRNA networks by miR-130/301 promotes pulmonary hypertension. *J Clin Invest*. 2014; 124:3514–3528. [PubMed: 24960162]
18. Chao CH, Chang CC, Wu MJ, Ko HW, Wang D, Hung MC, Yang JY, Chang CJ. MicroRNA-205 signaling regulates mammary stem cell fate and tumorigenesis. *J Clin Invest*. 2014; 124:3093–3106. [PubMed: 24911147]
19. Alexander MS, Casar JC, Motohashi N, Vieira NM, Eisenberg I, Marshall JL, Gasperini MJ, Lek A, Myers JA, Estrella EA, Kang PB, Shapiro F, Rahimov F, Kawahara G, Widrick JJ, Kunkel LM. MicroRNA-486-dependent modulation of DOCK3/PTEN/AKT signaling pathways improves muscular dystrophy-associated symptoms. *J Clin Invest*. 2014; 124:2651–2667. [PubMed: 24789910]
20. Latreille M, Hausser J, Stutzer I, Zhang Q, Hastoy B, Gargani S, Kerr-Conte J, Pattou F, Zavolan M, Esguerra JL, Eliasson L, Rulicke T, Rorsman P, Stoffel M. MicroRNA-7a regulates pancreatic beta cell function. *J Clin Invest*. 2014; 124:2722–2735. [PubMed: 24789908]
21. Bang C, Batkai S, Dangwal S, Gupta SK, Foinquinos A, Holzmann A, Just A, Remke J, Zimmer K, Zeug A, Ponimaskin E, Schmiedl A, Yin X, Mayr M, Halder R, Fischer A, Engelhardt S, Wei Y, Schober A, Fiedler J, Thum T. Cardiac fibroblast-derived microRNA passenger strand-enriched exosomes mediate cardiomyocyte hypertrophy. *J Clin Invest*. 2014; 124:2136–2146. [PubMed: 24743145]
22. Li CJ, Cheng P, Liang MK, Chen YS, Lu Q, Wang JY, Xia ZY, Zhou HD, Cao X, Xie H, Liao EY, Luo XH. MicroRNA-188 regulates age-related switch between osteoblast and adipocyte differentiation. *J Clin Invest*. 2015; 125:1509–1522. [PubMed: 25751060]
23. Murugaiyan G, da Cunha AP, Ajay AK, Joller N, Garo LP, Kumaradevan S, Yosef N, Vaidya VS, Weiner HL. MicroRNA-21 promotes Th17 differentiation and mediates experimental autoimmune encephalomyelitis. *J Clin Invest*. 2015; 125:1069–1080. [PubMed: 25642768]
24. Santa-Maria I, Alaniz ME, Renwick N, Cela C, Fulga TA, Van Vactor D, Tuschl T, Clark LN, Shelanski ML, McCabe BD, Crary JF. Dysregulation of microRNA-219 promotes neurodegeneration through post-transcriptional regulation of tau. *J Clin Invest*. 2015; 125:681–686. [PubMed: 25574843]
25. Santulli G, Iaccarino G, De Luca N, Trimarco B, Condorelli G. Atrial fibrillation and microRNAs. *Front Physiol*. 2014; 5:15. [PubMed: 24478726]
26. Sardu C, Marfella R, Santulli G, Paolisso G. Functional role of miRNA in cardiac resynchronization therapy. *Pharmacogenomics*. 2014; 15:1159–1168. [PubMed: 25084208]
27. van Rooij E, Olson EN. MicroRNA therapeutics for cardiovascular disease: opportunities and obstacles. *Nat Rev Drug Discov*. 2012; 11:860–872. [PubMed: 23080337]

28. Meijer HA, Kong YW, Lu WT, Wilczynska A, Spriggs RV, Robinson SW, Godfrey JD, Willis AE, Bushell M. Translational repression and eIF4A2 activity are critical for microRNA-mediated gene regulation. *Science*. 2013; 340:82–85. [PubMed: 23559250]
29. Guo H, Ingolia NT, Weissman JS, Bartel DP. Mammalian microRNAs predominantly act to decrease target mRNA levels. *Nature*. 2010; 466:835–840. [PubMed: 20703300]
30. Callis TE, Pandya K, Seok HY, Tang RH, Tatsuguchi M, Huang ZP, Chen JF, Deng Z, Gunn B, Shumate J, Willis MS, Selzman CH, Wang DZ. MicroRNA-208a is a regulator of cardiac hypertrophy and conduction in mice. *J Clin Invest*. 2009; 119:2772–2786. [PubMed: 19726871]
31. Yates LA, Norbury CJ, Gilbert RJ. The long and short of microRNA. *Cell*. 2013; 153:516–519. [PubMed: 23622238]
32. Yamamoto S, Niida S, Azuma E, Yanagibashi T, Muramatsu M, Huang TT, Sagara H, Higaki S, Ikutani M, Nagai Y, Takatsu K, Miyazaki K, Hamashima T, Mori H, Matsuda N, Ishii Y, Sasahara M. Inflammation-induced endothelial cell-derived extracellular vesicles modulate the cellular status of pericytes. *Sci Rep*. 2015; 5:8505. [PubMed: 25687367]
33. Bartel DP. MicroRNAs: genomics, biogenesis, mechanism, and function. *Cell*. 2004; 116:281–297. [PubMed: 14744438]
34. Cullen BR. Transcription and processing of human microRNA precursors. *Mol Cell*. 2004; 16:861–865. [PubMed: 15610730]
35. Roberts TC. The microRNA biology of the mammalian nucleus. *Mol Ther Nucleic Acids*. 2014; 3:e188. [PubMed: 25137140]
36. Denli AM, Tops BB, Plasterk RH, Ketting RF, Hannon GJ. Processing of primary microRNAs by the microprocessor complex. *Nature*. 2004; 432:231–235. [PubMed: 15531879]
37. Zeng Y, Yi R, Cullen BR. Recognition and cleavage of primary microRNA precursors by the nuclear processing enzyme Drosha. *EMBO J*. 2005; 24:138–148. [PubMed: 15565168]
38. Shiohama A, Sasaki T, Noda S, Minoshima S, Shimizu N. Molecular cloning and expression analysis of a novel gene DGCR8 located in the DiGeorge syndrome chromosomal region. *Biochem Biophys Res Commun*. 2003; 304:184–190. [PubMed: 12705904]
39. Roth BM, Ishimaru D, Hennig M. The core microprocessor component DiGeorge syndrome critical region 8 (DGCR8) is a nonspecific RNA-binding protein. *J Biol Chem*. 2013; 288:26785–26799. [PubMed: 23893406]
40. Ruby JG, Jan CH, Bartel DP. Intronic microRNA precursors that bypass Drosha processing. *Nature*. 2007; 448:83–86. [PubMed: 17589500]
41. Alarcon CR, Lee H, Goodarzi H, Halberg N, Tavazoie SF. N6-methyladenosine marks primary microRNAs for processing. *Nature*. 2015; 519:482–485. [PubMed: 25799998]
42. Lund E, Guttinger S, Calado A, Dahlberg JE, Kutay U. Nuclear export of microRNA precursors. *Science*. 2004; 303:95–98. [PubMed: 14631048]
43. Saxena A, Tabin CJ. miRNA-processing enzyme Dicer is necessary for cardiac outflow tract alignment and chamber septation. *Proc Natl Acad Sci U S A*. 2010; 107:87–91. [PubMed: 20018673]
44. Marasovic M, Zocco M, Halic M. Argonaute and Triman generate Dicer-independent priRNAs and mature siRNAs to initiate heterochromatin formation. *Mol Cell*. 2013; 52(2):173–183. [PubMed: 24095277]
45. Filipowicz W, Bhattacharyya SN, Sonenberg N. Mechanisms of post-transcriptional regulation by microRNAs: are the answers in sight? *Nat Rev Genet*. 2008; 9:102–114. [PubMed: 18197166]
46. Wu PH, Isaji M, Carthew RW. Functionally diverse microRNA effector complexes are regulated by extracellular signaling. *Mol Cell*. 2013; 52:113–123. [PubMed: 24055343]
47. Pfaff J, Hennig J, Herzog F, Aebersold R, Sattler M, Niessing D, Meister G. Structural features of Argonaute-GW182 protein interactions. *Proc Natl Acad Sci U S A*. 2013; 110:E3770–E3779. [PubMed: 24043833]
48. Siomi H, Siomi MC. On the road to reading the RNA-interference code. *Nature*. 2009; 457:396–404. [PubMed: 19158785]
49. von Brandenstein M, Richter C, Fries JW. MicroRNAs: small but amazing, and their association with endothelin. *Life Sci*. 2012; 91:475–489. [PubMed: 22771694]

50. Fabian MR, Sonenberg N. The mechanics of miRNA-mediated gene silencing: a look under the hood of miRISC. *Nat Struct Mol Biol.* 2012; 19:586–593. [PubMed: 22664986]
51. Kallen AN, Ma J, Huang Y. Does Lin28 antagonize miRNA-mediated repression by displacing miRISC from target mRNAs? *Front Genet.* 2012; 3:240. [PubMed: 23162570]
52. Papait R, Kunderfranco P, Stirparo GG, Latronico MV, Condorelli G. Long noncoding RNA: a new player of heart failure? *J Cardiovasc Transl Res.* 2013; 6:876–883. [PubMed: 23835777]
53. Lewis BP, Burge CB, Bartel DP. Conserved seed pairing, often flanked by adenosines, indicates that thousands of human genes are microRNA targets. *Cell.* 2005; 120:15–20. [PubMed: 15652477]
54. Dweep H, Sticht C, Pandey P, Gretz N. miRWalk--database: prediction of possible miRNA binding sites by “walking” the genes of three genomes. *J Biomed Inform.* 2011; 44:839–847. [PubMed: 21605702]
55. Zacharewicz E, Lamon S, Russell AP. MicroRNAs in skeletal muscle and their regulation with exercise, ageing, and disease. *Front Physiol.* 2013; 4:266. [PubMed: 24137130]
56. Ambros V, Bartel B, Bartel DP, Burge CB, Carrington JC, Chen X, Dreyfuss G, Eddy SR, Griffiths-Jones S, Marshall M, Matzke M, Ruvkun G, Tuschl T. A uniform system for microRNA annotation. *RNA.* 2003; 9:277–279. [PubMed: 12592000]
57. Tycko B, Ashkenas J. Epigenetics and its role in disease. *J Clin Invest.* 2000; 105:245–246. [PubMed: 10675348]
58. Boulard M, Edwards JR, Bestor TH. FBXL10 protects Polycomb-bound genes from hypermethylation. *Nat Genet.* 2015; 47:479–485. [PubMed: 25848754]
59. Hohl M, Wagner M, Reil JC, Muller SA, Tauchnitz M, Zimmer AM, Lehmann LH, Thiel G, Bohm M, Backs J, Maack C. HDAC4 controls histone methylation in response to elevated cardiac load. *J Clin Invest.* 2013; 123:1359–1370. [PubMed: 23434587]
60. Kraft M, Cirstea IC, Voss AK, Thomas T, Goehring I, Sheikh BN, Gordon L, Scott H, Smyth GK, Ahmadian MR, Trautmann U, Zenker M, Tartaglia M, Ekici A, Reis A, Dorr HG, Rauch A, Thiel CT. Disruption of the histone acetyltransferase MYST4 leads to a Noonan syndrome-like phenotype and hyperactivated MAPK signaling in humans and mice. *J Clin Invest.* 2011; 121:3479–3491. [PubMed: 21804188]
61. Leucci E, Patella F, Waage J, Holmstrom K, Lindow M, Porse B, Kauppinen S, Lund AH. MicroRNA-9 targets the long non-coding RNA MALAT1 for degradation in the nucleus. *Sci Rep.* 2013; 3:2535. [PubMed: 23985560]
62. Swierczynski S, Klieser E, Illig R, Alinger-Scharinger B, Kiesslich T, Neureiter D. Histone deacetylation meets miRNA: epigenetics and post-transcriptional regulation in cancer and chronic diseases. *Expert Opin Biol Ther.* 2015; 15:651–664. [PubMed: 25766312]
63. Tu J, Liao JJ, Luk AC, Tang NL, Chan WY, Lee TL. MicroRNAs mediated targeting on the Yin-yang dynamics of DNA methylation in disease and development. *Int J Biochem Cell Biol.* 2015
64. Greco CM, Condorelli G. Epigenetic modifications and noncoding RNAs in cardiac hypertrophy and failure. *Nat Rev Cardiol.* 2015; 12:488–497. [PubMed: 25962978]
65. Garzon R, Liu S, Fabbri M, Liu Z, Heaphy CE, Callegari E, Schwind S, Pang J, Yu J, Muthusamy N, Havelange V, Volinia S, Blum W, Rush LJ, Perrotti D, Andreeff M, Bloomfield CD, Byrd JC, Chan K, Wu LC, Croce CM, Marcucci G. MicroRNA-29b induces global DNA hypomethylation and tumor suppressor gene reexpression in acute myeloid leukemia by targeting directly DNMT3A and 3B and indirectly DNMT1. *Blood.* 2009; 113:6411–6418. [PubMed: 19211935]
66. Robaina MC, Mazzocchi L, Arruda VO, Reis FR, Apa AG, de Rezende LM, Klumb CE. Deregulation of DNMT1, DNMT3B and miR-29s in Burkitt lymphoma suggests novel contribution for disease pathogenesis. *Exp Mol Pathol.* 2015; 98:200–207. [PubMed: 25746661]
67. Yuan JH, Yang F, Chen BF, Lu Z, Huo XS, Zhou WP, Wang F, Sun SH. The histone deacetylase 4/SPI1/microRNA-200a regulatory network contributes to aberrant histone acetylation in hepatocellular carcinoma. *Hepatology.* 2011; 54:2025–2035. [PubMed: 21837748]
68. Lujambio A, Calin GA, Villanueva A, Ropero S, Sanchez-Cespedes M, Blanco D, Montuenga LM, Rossi S, Nicoloso MS, Faller WJ, Gallagher WM, Eccles SA, Croce CM, Esteller M. A microRNA DNA methylation signature for human cancer metastasis. *Proc Natl Acad Sci U S A.* 2008; 105:13556–13561. [PubMed: 18768788]

69. Bueno MJ, Perez de Castro I, Gomez de Cedron M, Santos J, Calin GA, Cigudosa JC, Croce CM, Fernandez-Piqueras J, Malumbres M. Genetic and epigenetic silencing of microRNA-203 enhances ABL1 and BCR-ABL1 oncogene expression. *Cancer Cell*. 2008; 13:496–506. [PubMed: 18538733]
70. Hu W, Wang M, Yin H, Yao C, He Q, Yin L, Zhang C, Li W, Chang G, Wang S. MicroRNA-1298 is regulated by DNA methylation and affects vascular smooth muscle cell function by targeting connexin 43. *Cardiovasc Res*. 2015
71. Long XR, He Y, Huang C, Li J. MicroRNA-148a is silenced by hypermethylation and interacts with DNA methyltransferase 1 in hepatocellular carcinogenesis. *Int J Oncol*. 2014; 44:1915–1922. [PubMed: 24714841]
72. Dai E, Yu X, Zhang Y, Meng F, Wang S, Liu X, Liu D, Wang J, Li X, Jiang W. EpimiR: a database of curated mutual regulation between miRNAs and epigenetic modifications. *Database (Oxford)*. 2014; 2014:bau023. [PubMed: 24682734]
73. Santulli G, Marks AR. Essential roles of intracellular calcium release channels in muscle, brain, metabolism, and aging. *Curr Mol Pharmacol*. 2015; 8:206–222. [PubMed: 25966694]
74. Andrews SJ, Rothnagel JA. Emerging evidence for functional peptides encoded by short open reading frames. *Nat Rev Genet*. 2014; 15:193–204. [PubMed: 24514441]
75. Carninci P, Kasukawa T, Katayama S, Gough J, Frith MC, Maeda N, Oyama R, Ravasi T, Lenhard B, Wells C, Kodzius R, Shimokawa K, Bajic VB, Brenner SE, Batalov S, Forrest AR, Zavolan M, Davis MJ, Wilming LG, Aidinis V, Allen JE, Ambesi-Impiomato A, Apweiler R, Aturaliya RN, Bailey TL, Bansal M, Baxter L, Beisel KW, Bersano T, Bono H, Chalk AM, Chiu KP, Choudhary V, Christoffels A, Clutterbuck DR, Crowe ML, Dalla E, Dalrymple BP, de Bono B, Della Gatta G, di Bernardo D, Down T, Engstrom P, Fagiolini M, Faulkner G, Fletcher CF, Fukushima T, Furuno M, Futaki S, Gariboldi M, Georgii-Hemming P, Gingeras TR, Gojobori T, Green RE, Gustincich S, Harbers M, Hayashi Y, Hensch TK, Hirokawa N, Hill D, Huminiecki L, Iacono M, Ikeo K, Iwama A, Ishikawa T, Jakt M, Kanapin A, Katoh M, Kawasawa Y, Kelso J, Kitamura H, Kitano H, Kollias G, Krishnan SP, Kruger A, Kummerfeld SK, Kurochkin IV, Lareau LF, Lazarevic D, Lipovich L, Liu J, Liuni S, McWilliam S, Madan Babu M, Madera M, Marchionni L, Matsuda H, Matsuzawa S, Miki H, Mignone F, Miyake S, Morris K, Mottagui-Tabar S, Mulder N, Nakano N, Nakauchi H, Ng P, Nilsson R, Nishiguchi S, Nishikawa S, Nori F, Ohara O, Okazaki Y, Orlando V, Pang KC, Pavan WJ, Pavesi G, Pesole G, Petrovsky N, Piazza S, Reed J, Reid JF, Ring BZ, Ringwald M, Rost B, Ruan Y, Salzberg SL, Sandelin A, Schneider C, Schonbach C, Sekiguchi K, Semple CA, Seno S, Sessa L, Sheng Y, Shibata Y, Shimada H, Shimada K, Silva D, Sinclair B, Sperling S, Stupka E, Sugiura K, Sultana R, Takenaka Y, Taki K, Tammoya K, Tan SL, Tang S, Taylor MS, Tegner J, Teichmann SA, Ueda HR, van Nimwegen E, Verardo R, Wei CL, Yagi K, Yamanishi H, Zabarovsky E, Zhu S, Zimmer A, Hide W, Bult C, Grimmond SM, Teasdale RD, Liu ET, Brusic V, Quackenbush J, Wahlestedt C, Mattick JS, Hume DA, Kai C, Sasaki D, Tomaru Y, Fukuda S, Kanamori-Katayama M, Suzuki M, Aoki J, Arakawa T, Iida J, Imamura K, Itoh M, Kato T, Kawaji H, Kawagashira N, Kawashima T, Kojima M, Kondo S, Konno H, Nakano K, Ninomiya N, Nishio T, Okada M, Plessy C, Shibata K, Shiraki T, Suzuki S, Tagami M, Waki K, Watahiki A, Okamura-Oho Y, Suzuki H, Kawai J, Hayashizaki Y. FANTOM Consortium, RIKEN Genome Exploration Research Group, Genome Science Group. The transcriptional landscape of the mammalian genome. *Science*. 2005; 309:1559–1563. [PubMed: 16141072]
76. Lee LJ, Hughes TR, Frey BJ. How many new genes are there? *Science*. 2006; 311:1709–1711. author reply 1709–11. [PubMed: 16556823]
77. Troy CS, MacHugh DE, Bailey JF, Magee DA, Loftus RT, Cunningham P, Chamberlain AT, Sykes BC, Bradley DG. Genetic evidence for Near-Eastern origins of European cattle. *Nature*. 2001; 410:1088–1091. [PubMed: 11323670]
78. Rosenberg MI, Desplan C. Molecular biology. Hiding in plain sight. *Science*. 2010; 329:284–285. [PubMed: 20647453]
79. Crappe J, Ndah E, Koch A, Steyaert S, Gawron D, De Keulenaer S, De Meester E, De Meyer T, Van Crielinge W, Van Damme P, Menschaert G. PROTEOFORMER: deep proteome coverage through ribosome profiling and MS integration. *Nucleic Acids Res*. 2015; 43:e29. [PubMed: 25510491]

80. Kondo T, Plaza S, Zanet J, Benrabah E, Valenti P, Hashimoto Y, Kobayashi S, Payre F, Kageyama Y. Small peptides switch the transcriptional activity of Shavenbaby during *Drosophila* embryogenesis. *Science*. 2010; 329:336–339. [PubMed: 20647469]
81. Martinho RG, Kunwar PS, Casanova J, Lehmann R. A noncoding RNA is required for the repression of RNAPolIII-dependent transcription in primordial germ cells. *Curr Biol*. 2004; 14:159–165. [PubMed: 14738740]
82. Galindo MI, Pueyo JI, Fouix S, Bishop SA, Couso JP. Peptides encoded by short ORFs control development and define a new eukaryotic gene family. *PLoS Biol*. 2007; 5:e106. [PubMed: 17439302]
83. Hayden CA, Bosco G. Comparative genomic analysis of novel conserved peptide upstream open reading frames in *Drosophila melanogaster* and other dipteran species. *BMC Genomics*. 2008; 9:61. [PubMed: 18237443]
84. Hashimoto Y, Niikura T, Tajima H, Yasukawa T, Sudo H, Ito Y, Kita Y, Kawasumi M, Kouyama K, Doyu M, Sobue G, Koide T, Tsuji S, Lang J, Kurokawa K, Nishimoto I. A rescue factor abolishing neuronal cell death by a wide spectrum of familial Alzheimer's disease genes and Abeta. *Proc Natl Acad Sci U S A*. 2001; 98:6336–6341. [PubMed: 11371646]
85. Guo B, Zhai D, Cabezas E, Welsh K, Nouraini S, Satterthwait AC, Reed JC. Humanin peptide suppresses apoptosis by interfering with Bax activation. *Nature*. 2003; 423:456–461. [PubMed: 12732850]
86. Ikonen M, Liu B, Hashimoto Y, Ma L, Lee KW, Niikura T, Nishimoto I, Cohen P. Interaction between the Alzheimer's survival peptide humanin and insulin-like growth factor-binding protein 3 regulates cell survival and apoptosis. *Proc Natl Acad Sci U S A*. 2003; 100:13042–13047. [PubMed: 14561895]
87. Washietl S, Findeiss S, Muller SA, Kalkhof S, von Bergen M, Hofacker IL, Stadler PF, Goldman N. RNaCode: robust discrimination of coding and noncoding regions in comparative sequence data. *RNA*. 2011; 17:578–594. [PubMed: 21357752]
88. Prokhorova IV, Osterman IA, Burakovsky DE, Serebryakova MV, Galyamina MA, Pobeguts OV, Altukhov I, Kovalchuk S, Alexeev DG, Govorun VM, Bogdanov AA, Sergiev PV, Dontsova OA. Modified nucleotides m(2)G966/m(5)C967 of *Escherichia coli* 16S rRNA are required for attenuation of tryptophan operon. *Sci Rep*. 2013; 3:3236. [PubMed: 24241179]
89. Selitsky SR, Baran-Gale J, Honda M, Yamane D, Masaki T, Fannin EE, Guerra B, Shirasaki T, Shimakami T, Kaneko S, Lanford RE, Lemon SM, Sethupathy P. Small tRNA-derived RNAs are increased and more abundant than microRNAs in chronic hepatitis B and C. *Sci Rep*. 2015; 5:7675. [PubMed: 25567797]
90. Katayama S, Tomaru Y, Kasukawa T, Waki K, Nakanishi M, Nakamura M, Nishida H, Yap CC, Suzuki M, Kawai J, Suzuki H, Carninci P, Hayashizaki Y, Wells C, Frith M, Ravasi T, Pang KC, Hallinan J, Mattick J, Hume DA, Lipovich L, Batalov S, Engstrom PG, Mizuno Y, Faghihi MA, Sandelin A, Chalk AM, Mottagui-Tabar S, Liang Z, Lenhard B, Wahlestedt C, Group RGER, Genome Science G, Consortium F. Antisense transcription in the mammalian transcriptome. *Science*. 2005; 309:1564–1566. [PubMed: 16141073]
91. Mizuno T, Chou MY, Inouye M. A unique mechanism regulating gene expression: translational inhibition by a complementary RNA transcript (micRNA). *Proc Natl Acad Sci U S A*. 1984; 81:1966–1970. [PubMed: 6201848]
92. Kim TK, Hemberg M, Gray JM, Costa AM, Bear DM, Wu J, Harmin DA, Laptewicz M, Barbara-Haley K, Kuersten S, Markenscoff-Papadimitriou E, Kuhl D, Bito H, Worley PF, Kreiman G, Greenberg ME. Widespread transcription at neuronal activity-regulated enhancers. *Nature*. 2010; 465:182–187. [PubMed: 20393465]
93. Orom UA, Derrien T, Beringer M, Gumireddy K, Gardini A, Bussotti G, Lai F, Zytnicki M, Notredame C, Huang Q, Guigo R, Shiekhattar R. Long noncoding RNAs with enhancer-like function in human cells. *Cell*. 2010; 143:46–58. [PubMed: 20887892]
94. Ranzani V, Rossetti G, Panzeri I, Arrigoni A, Bonnal RJ, Curti S, Gruarin P, Provati E, Sugliano E, Marconi M, De Francesco R, Geginat J, Bodega B, Abrignani S, Pagani M. The long intergenic noncoding RNA landscape of human lymphocytes highlights the regulation of T cell differentiation by linc-MAF-4. *Nat Immunol*. 2015; 16:318–325. [PubMed: 25621826]

95. Han BW, Wang W, Li C, Weng Z, Noncoding ZPD. RNA. piRNA-guided transposon cleavage initiates Zucchini-dependent, phased piRNA production. *Science*. 2015; 348:817–821. [PubMed: 25977554]
96. Kang S, Kang D, Bakhtiar Ul Islam SM, Je S, Kim JH, Song JJ. Silencing Daxx increases the anti-tumor activity of a TRAIL/shRNA Bcl-xL-expressing oncolytic adenovirus through enhanced viral replication and cellular arrest. *Cell Signal*. 2015; 27:1214–1224. [PubMed: 25748050]
97. Herrmann A, Priceman SJ, Kujawski M, Xin H, Cherryholmes GA, Zhang W, Zhang C, Lahtz C, Kowolik C, Forman SJ, Kortylewski M, Yu H. CTLA4 aptamer delivers STAT3 siRNA to tumor-associated and malignant T cells. *J Clin Invest*. 2014; 124:2977–2987. [PubMed: 24892807]
98. Chu L, Su MY, Maggi LB Jr, Lu L, Mullins C, Crosby S, Huang G, Chng WJ, Vij R, Tomasson MH. Multiple myeloma-associated chromosomal translocation activates orphan snoRNA ACA11 to suppress oxidative stress. *J Clin Invest*. 2012; 122:2793–2806. [PubMed: 22751105]
99. Taft RJ, Hawkins PG, Mattick JS, Morris KV. The relationship between transcription initiation RNAs and CCCTC-binding factor (CTCF) localization. *Epigenetics Chromatin*. 2011; 4:13. [PubMed: 21813016]
100. Lerner MR, Boyle JA, Hardin JA, Steitz JA. Two novel classes of small ribonucleoproteins detected by antibodies associated with lupus erythematosus. *Science*. 1981; 211:400–402. [PubMed: 6164096]
101. Wang I, Kowalski MP, Langley AR, Rodriguez R, Balasubramanian S, Hsu ST, Krude T. Nucleotide contributions to the structural integrity and DNA replication initiation activity of noncoding y RNA. *Biochemistry*. 2014; 53:5848–5863. [PubMed: 25151917]
102. Davis CA, Ares M Jr. Accumulation of unstable promoter-associated transcripts upon loss of the nuclear exosome subunit Rrp6p in *Saccharomyces cerevisiae*. *Proc Natl Acad Sci U S A*. 2006; 103:3262–3267. [PubMed: 16484372]
103. Zhang Y, Liu XS, Liu QR, Wei L. Genome-wide in silico identification and analysis of cis natural antisense transcripts (cis-NATs) in ten species. *Nucleic Acids Res*. 2006; 34:3465–3475. [PubMed: 16849434]
104. Li S, Liberman LM, Mukherjee N, Benfey PN, Ohler U. Integrated detection of natural antisense transcripts using strand-specific RNA sequencing data. *Genome Res*. 2013; 23:1730–1739. [PubMed: 23816784]
105. Kapranov P, Cheng J, Dike S, Nix DA, Duttagupta R, Willingham AT, Stadler PF, Hertel J, Hackermuller J, Hofacker IL, Bell I, Cheung E, Drenkow J, Dumais E, Patel S, Helt G, Ganesh M, Ghosh S, Piccolboni A, Sementchenko V, Tammanna H, Gingeras TR. RNA maps reveal new RNA classes and a possible function for pervasive transcription. *Science*. 2007; 316:1484–1488. [PubMed: 17510325]
106. Affymetrix ENCODE Transcriptome Project, Cold Spring Harbor Laboratory ENCODE Transcriptome Project. Post-transcriptional processing generates a diversity of 5'-modified long and short RNAs. *Nature*. 2009; 457:1028–1032. [PubMed: 19169241]
107. Kurokawa R. Promoter-associated long noncoding RNAs repress transcription through a RNA binding protein TLS. *Adv Exp Med Biol*. 2011; 722:196–208. [PubMed: 21915790]
108. Preker R, Almvig K, Christensen MS, Valen E, Mapendano CK, Sandelin A, Jensen TH. PROMoter uPstream Transcripts share characteristics with mRNAs and are produced upstream of all three major types of mammalian promoters. *Nucleic Acids Res*. 2011; 39:7179–7193. [PubMed: 21596787]
109. Guerrier-Takada C, Gardiner K, Marsh T, Pace N, Altman S. The RNA moiety of ribonuclease P is the catalytic subunit of the enzyme. *Cell*. 1983; 35:849–857. [PubMed: 6197186]
110. Howard MJ, Klemm BP, Fierke CA. Mechanistic studies reveal similar catalytic strategies for phosphodiester bond hydrolysis by protein-only and RNA-dependent ribonuclease P. *J Biol Chem*. 2015; 290:13454–13464. [PubMed: 25817998]
111. Gold HA, Topper JN, Clayton DA, Craft J. The RNA processing enzyme RNase MRP is identical to the Th RNP and related to RNase P. *Science*. 1989; 245:1377–1380. [PubMed: 2476849]
112. Mattijssen S, Hinson ER, Onnekink C, Hermanns P, Zabel B, Cresswell P, Pruijn GJ. Viperin mRNA is a novel target for the human RNase MRP/RNase P endoribonuclease. *Cell Mol Life Sci*. 2011; 68:2469–2480. [PubMed: 21053045]

113. Varshney D, Vavrova-Anderson J, Oler AJ, Cowling VH, Cairns BR, White RJ. SINE transcription by RNA polymerase III is suppressed by histone methylation but not by DNA methylation. *Nat Commun.* 2015; 6:6569. [PubMed: 25798578]
114. Ohshima K, Koishi R, Matsuo M, Okada N. Several short interspersed repetitive elements (SINEs) in distant species may have originated from a common ancestral retrovirus: characterization of a squid SINE and a possible mechanism for generation of tRNA-derived retroposons. *Proc Natl Acad Sci U S A.* 1993; 90:6260–6264. [PubMed: 8327507]
115. Feng J, Funk WD, Wang SS, Weinrich SL, Avilion AA, Chiu CP, Adams RR, Chang E, Allsopp RC, Yu J, et al. The RNA component of human telomerase. *Science.* 1995; 269:1236–1241. [PubMed: 7544491]
116. Kannan R, Helston RM, Dannebaum RO, Baumann P. Diverse mechanisms for spliceosome-mediated 3' end processing of telomerase RNA. *Nat Commun.* 2015; 6:6104. [PubMed: 25598145]
117. Fassan M, Dall'Olmo L, Galasso M, Braconi C, Pizzi M, Realdon S, Volinia S, Valeri N, Gasparini P, Baffa R, Souza RF, Vicentini C, D'Angelo E, Bornschein J, Nuovo GJ, Zaninotto G, Croce CM, Rugge M. Transcribed ultraconserved noncoding RNAs (T-UCR) are involved in Barrett's esophagus carcinogenesis. *Oncotarget.* 2014; 5:7162–7171. [PubMed: 25216530]
118. Calin GA, Liu CG, Ferracin M, Hyslop T, Spizzo R, Sevignani C, Fabbri M, Cimmino A, Lee EJ, Wojcik SE, Shimizu M, Tili E, Rossi S, Taccioli C, Pichiorri F, Liu X, Zupo S, Herlea V, Gramantieri L, Lanza G, Alder H, Rassenti L, Volinia S, Schmittgen TD, Kipps TJ, Negrini M, Croce CM. Ultraconserved regions encoding ncRNAs are altered in human leukemias and carcinomas. *Cancer Cell.* 2007; 12:215–229. [PubMed: 17785203]
119. St Laurent G, Shtokalo D, Dong B, Tackett MR, Fan X, Lazorthes S, Nicolas E, Sang N, Triche TJ, McCaffrey TA, Xiao W, Kapranov P. VlnRNAs controlled by retroviral elements are a hallmark of pluripotency and cancer. *Genome Biol.* 2013; 14:R73. [PubMed: 23876380]
120. Ferracin M, Gautheret D, Hubé D, Mani SA, Mattick JS, Andersson Ørom U, Santulli G, Slotkin RK, Szweykowska-Kulinska Z, Taube JH, Vazquez F, Yang JH. The Non-Coding RNA Journal Club: Highlights on Recent Papers. *Non-Coding RNA.* 2015; 1:87–93.

Fig. 1.1.
Micropeptidomics: *multa paucis* or *hic sunt leones*?

Table 1.1

Characteristics of noncoding RNAs within the cell

Abbreviation	Complete name	Main functions	Length (nt)	Ref.
miRNA	MicroRNA	Gene silencing	21–25	[4]
rRNA	Ribosomal RNA	Translation	120–4700	[88]
tRNA	Transfer RNA	Translation	70–100	[89]
aRNA	Antisense RNA	Transcriptional attenuation	>30	[90, 91]
eRNA	Enhancer-like ncRNA (1D/2D)	Transcriptional enhancers	50–2000	[92, 93]
lincRNA	Long intergenic RNA	Transcriptional and posttranscriptional regulation	<50 kb	[94]
piRNA	PIWI-interacting RNA	Genome stabilization	24–30	[95]
shRNA	Short hairpin RNA	Gene silencing	19–29	[96]
siRNA	Short interfering RNA	Gene silencing	21–25	[97]
snRNA	Small nuclear RNA	Splicing		
snoRNA	Small nucleolar RNA	Methylation (C/D box), pseudouridylation (H/ACA box)	~20–24	[98]
SRP-RNA	Signal recognition particle RNA	Translocation of proteins across the endoplasmic reticulum		
tiRNA	Transcription initiation RNA	Transcriptional regulation	18	[99]
Y RNA	Y RNA	DNA replication and RNA processing (repressor of Ro60)	83–112	[100, 101]
CUT	Cryptic unstable transcript	Gene regulation	200–800	[102]
NAT	Natural antisense transcript	RNA interference	Variable	[103, 104]
PALR	Promoter-associated long RNA	Transcriptional regulation	200–1000	[105–107]
PROMPT	Promoter upstream transcript	Gene transcription	long	[108]
RNase P	Ribonuclease P	Endonucleolytic 5' cleavage of tRNA precursors (ribozyme)	354–417	[109, 110]
RNase MRP	Mitochondrial RNA processing ribonuclease	Mitochondrial DNA replication and rRNA maturation (ribozyme)	265–340	[111, 112]

Abbreviation	Complete name	Main functions	Length (nt)	Ref.
SINE	Short interspersed repetitive elements	Transcriptional suppressor (e.g. Alu element)	<500	[113, 114]
TERC	Telomerase RNA component	Telomere synthesis	451	[115, 116]
T-UCR	Transcribed ultra-conserved region	Transcriptional enhancer	>200	[117, 118]
vlincRNAs	Very long intergenic RNA	Transcriptional and posttranscriptional regulation	>50 kb	[119, 120]