Published in final edited form as:

Psychiatry Res. 2015 March 30; 231(3): 333–340. doi:10.1016/j.psychres.2015.01.017.

Frontal and striatal alterations associated with psychopathic traits in adolescents

Yaling Yang^{a,*}, Katherine L. Narr^b, Laura A. Baker^c, Shantanu H. Joshi^b, Neda Jahanshad^d, Adrian Raine^e, and Paul M. Thompson^d

^aDepartment of Pediatrics, Children's Hospital Los Angeles, University of Southern California, 4650 Sunset Blvd, Smith Research Tower Room 400B, Los Angeles, CA 90027, USA

^bDepartment of Neurology, David Geffen School of Medicine at UCLA, Los Angeles, CA, USA

^cDepartment of Psychology, University of Southern California, Los Angeles, CA, USA

^dInstitute for Neuroimaging and Informatics, University of Southern California, Los Angeles, CA, USA

^eDepartments of Criminology, Psychiatry, and Psychology, University of Pennsylvania, Philadelphia, PA, USA

Abstract

Neuroimaging research has demonstrated a range of structural deficits in adults with psychopathy, but little is known about structural correlates of psychopathic tendencies in adolescents. Here we examined structural magnetic resonance imaging (sMRI) data obtained from 14-year-old adolescents ($n=108$) using tensor-based morphometry (TBM) to isolate global and localized differences in brain tissue volumes associated with psychopathic traits in this otherwise healthy developmental population. We found that greater levels of psychopathic traits were correlated with increased brain tissue volumes in the left putamen, left ansa peduncularis, right superiomedial prefrontal cortex, left inferior frontal cortex, right orbitofrontal cortex, and right medial temporal regions and reduced brain tissues volumes in the right middle frontal cortex, left superior parietal lobule, and left inferior parietal lobule. Post hoc analyses of parcellated regional volumes also showed putamen enlargements to correlate with increased psychopathic traits. Consistent with earlier studies, findings suggest poor decision-making and emotional dysregulation associated with psychopathy may be due, in part, to structural anomalies in frontal and temporal regions whereas striatal structural variations may contribute to sensation-seeking and reward-driven behavior in psychopathic individuals. Future studies will help clarify how disturbances in brain maturational

*Corresponding Author: Yaling Yang, Ph.D. Department of Pediatrics, Children's Hospital Los Angeles, University of Southern California, 4650 Sunset Blvd, Smith Research Tower Room 400B, Los Angeles, CA 90027, USA. Tel.: +1 323-361-6398. yayang@chla.usc.edu.

Financial disclosures

All authors report no potential conflicts of interest, financial or otherwise.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

processes might lead to the developmental trajectory from psychopathic tendencies in adolescents to adult psychopathy.

Keywords

Developmental; MRI; Tensor-based morphometry; Putamen

1. Introduction

Psychopathy is a clinical condition defined by a combination of persistent antisocial behavior, marked sensation-seeking, impulsivity, shallow emotion, blunted empathy, and punishment insensitivity that emerges early in life (Hare, 2003). Although a diagnostic level of psychopathy is present in only 1–2% of the general population, it is widely present on a spectrum in otherwise healthy populations. Research on the neural bases of psychopathy has focused largely on the profound emotional deficits and antisocial behavior observed in psychopathic adults, emphasizing the possible contributions of a disturbed frontolimbic circuitry (Gao et al., 2009; Yang and Raine, 2009). To date, findings from several structural imaging studies have supported this argument by showing frontolimbic deficits in adults with high psychopathy scores, with the most robust findings being reduced volumes in frontal and temporal regions including the amygdala (de Oliveira-Souza et al., 2008; Muller et al., 2008; Yang et al., 2009a; Yang et al., 2009b; Ermer et al., 2012; Bertsch et al., 2013). However, conflicting findings have also been presented, particularly for the hippocampus, insula, and the anterior cingulate cortex for psychopathic adults (Laakso et al., 2001; Boccardi et al., 2010; Glenn et al., 2010b; Cope et al., 2012).

More recently, initial evidence has begun to emerge suggesting that deficits in regions densely connected with the frontolimbic circuitry, particularly the striatum, may also contribute to traits associated with psychopathy (Glenn and Yang, 2012; Blair, 2013). The striatum is comprised of the caudate nucleus and the putamen and has been linked to traits such as reward-seeking, stimulus-reinforcement learning, and impulsivity (Barros-Loscertales et al., 2006; Cohen et al., 2009). Studies have linked psychopathy to increased sensitivity to reward and decreased sensitivity to punishment, suggesting potential abnormalities in the striatal regions that contribute to overly focus on the prospect of reward despite signals of potential later punishment (Newman and Kosson, 1986; van Honk and Schutter, 2006; Glenn et al., 2009). To date, *increased* volumes have been found in the striatal in adults with psychopathy (Glenn et al., 2009) and antisocial personality disorder (Barkataki et al., 2006), findings that are in line with the functional imaging findings of abnormal activation or connectivity with the striatum (Osumi et al., 2012; Carre et al., 2013). Therefore, it may be suggested that an overactive reward system (e.g. striatum) may further compromise the weakened regulation system (e.g., prefrontal cortex), leading to heightened psychopathic tendencies.

Despite the accumulating knowledge in the neural basis of psychopathy, studies of psychopathic traits have focused mainly on adults and the examination of psychopathic traits in children and adolescence remains scarce. By revealing structural abnormalities in regions

overlap with those found impaired in adults with psychopathy, findings from studies of conduct disorder (some with comorbid callous-unemotional traits) seem to suggest a neurodevelopmental basis to psychopathy (Blair, 2006a; Gao et al., 2009). For example, reduced volumes in the amygdala, dorsolateral prefrontal cortex and orbitofrontal cortex (Kruesi et al., 2004; Sterzer et al., 2007; Boes et al., 2008; Huebner et al., 2008; Dalwani et al., 2011; Fairchild et al., 2011; Hyatt et al., 2012) have been reported in adolescents with conduct problems. In addition, callous-unemotional traits were found to correlate positively with increased volumes in the orbitofrontal cortex (De Brito et al., 2009; Fairchild et al., 2012), suggesting potential delays in brain maturation in association with certain psychopathic traits. In one study, callous-unemotional traits were also found to correlate negatively with reduced volume in the striatum (Fairchild et al., 2012), but the effect did not survive when controlling for conduct disorder symptoms.

Although findings to date have provided initial evidence suggesting abnormal brain structures associated with psychopathic traits in children and adolescents, there are notable variations among findings. One potential contribution to discrepancy among findings may be that most existing studies have been conducted on relatively small, heterogeneous samples. The aggregation of participants with a wide age range may be particularly alarming because it has been demonstrated that gray matter volumes increase in early childhood and decline in adolescence, but white matter volumes tend to increase well into adulthood (Sowell et al., 2003). The pattern of age-related changes in gray and white matter has also been shown to vary across brain regions during brain maturation (Sowell, 2002). The involvement of substance use in older participants, often co-occurring with antisocial behavior and psychopathic traits, may also complicate the matter by introducing substance-related brain changes. Furthermore, the majority of the findings are from samples with comorbid psychiatric disorders (e.g. conduct disorders, attention deficits hyperactivity disorder), making it difficult to determine to what extent previous findings can be applied to psychopathic traits in the general population.

To address these limitations in the literature, the present study included a community sample of homogeneously aged, healthy adolescents (all aged 14 years old at the time of the scanning) to examine the relationship between psychopathic tendencies and regional brain tissue volumes. We used a recently developed method of tensor-based morphometry (TBM), which allows the illustration of statistical effects on regional volumes of gray matter, white matter, and cerebrospinal fluid (CSF) across the entire brain. Specifically, global and regional differences in brain tissue volume are estimated by applying localized deformations to adjust the anatomy of each individual to match a sample-specific template. By correlating these deformation fields with psychopathic traits across the sample, very subtle morphological changes associated with psychopathic traits in adolescents can be identified, with high accuracy and sensitivity. To date, TBM has been validated and applied to study several disorders (Leow et al., 2009; Yang et al., 2011); however, it has yet to be applied to explore patterns of brain tissue alterations associated with psychopathic traits in adolescents. Based on prior reports, we predicted that psychopathic tendencies would correlate with reduced brain tissue volumes in the fronto-limbic circuitry and increased brain tissue volumes in the striatal regions in this adolescent sample.

2. Methods

2.1. Subjects

The 108 adolescent twins (54 males and 54 females, age 14 years) included in this study were drawn from participants in the University of Southern California (USC) Risk Factors for Antisocial Behavior Twin Study (Baker et al., 2006; Baker et al., 2013), and comprised 27 monozygotic twin pairs (16 male and 11 female twin pairs) and 27 dizygotic twin pairs (5 male, 10 female, and 12 opposite-sex twin pairs). In brief, the USC Twin Study is a longitudinal study assessing the development of antisocial behaviors from childhood to young adulthood, which began in 2000 and now is approaching the end of fifth wave data collection. The adolescents and their families were recruited from Los Angeles County through advertisements, schools, and mothers of twins clubs. The sample is representative of the ethnic and socioeconomic diversity of the Great Los Angeles areas. The present study used data obtained from the third wave of assessment, when the twins were 14 years old, as this was the time point when magnetic resonance imaging (MRI) scans were collected. Participants were excluded if they had a history of significant head injury, major neurological, psychiatric illness, substance abuse, or contraindication to scanning (Yang et al., 2012; Baker et al., 2013). The adolescents and their primary caregivers participated in 6–8 h of laboratory assessment at USC including a 1-h scan. Assessment of psychopathic traits was provided by caregivers, who were predominantly biological mothers ($n=50$). The remaining were biological father ($n=1$), grandmother ($n=1$), foster mother ($n=1$) and adoptive mother ($n=1$). Each child's ethnicity was determined by the ethnicity of his/her biological mother and father, as reported by the primary caregiver. The ethnicity breakdown of the sample was as follows: 36.7% Hispanic, 27.4% Caucasian, 14% Black, 4.4% Asian, 0.16% Native American, and 17.3% mixed. Both caregivers and children gave written informed consent/assent prior to the study. The study was approved by the CHLA/USC Institutional Review Boards.

2.2. Behavioral measurements

Psychopathic traits were measured using a slightly extended version of the Child Psychopathy Scale (CPS) – Revised Extended (Lynam, 1997). The CPS is a well-validated instrument for measuring psychopathic traits in children and adolescents and is composed of 14 subscales including glibness, untruthfulness, lack of guilt, callousness, impulsiveness, boredom susceptibility, manipulation, poverty of affect, parasitic lifestyle, behavioral dyscontrol, lack of planning, unreliability, failure to accept responsibility, and grandiosity. The CPS was administered to the caregivers of the adolescents in an interview form. Scores for each item were added to create a total CPS score for each individual. The internal reliabilities for the composite score have been established in previous reports (Baker et al., 2007; Bezdjian et al., 2011), and the frequency of the total CPS scores of this sample is reported in Fig. S1 (see Supplementary Material). To summarize, the parent-report CPS total scores of our sample ranged from 2 to 33 (mean = 12.18, SD = 7.09), with males (mean = 13.96, SD = 7.92, range: 3–33) scoring higher than females (mean = 10.39, SD = 5.68, range: 2–31) ($p = 0.008$). As a complementary analysis, we also categorized the participants into High-CPS (with CPS total scores in the top 20%) and Low-CPS (with CPS total scores in the low 20%) group, and the TBM results were included in the Supplementary Material

(Fig. S2). For a subset of the sample ($n = 62$), intelligence quotient (IQ) was obtained using Wechsler Intelligence Scale for Children (Wechsler, 2004). For the subsample, the intelligence mean was 100.1 for Full-Scale IQ (range: 77–129), 98.44 for Performance IQ (range 67–124) and 102.15 for Verbal IQ (range: 65–128). Further analysis showed that parent-report CPS total scores did not correlate with Full-Scale, Performance, or Verbal IQ scores (all $p > 0.3$). Post hoc analyses showed that TBM results were unchanged when controlling for IQ, therefore only results controlling for whole brain volume, sex, and subject relatedness were included in this report.

2.3. MRI acquisition and preprocessing

All participants were scanned using a 3 Tesla Siemens Magnetom Trio whole-body scanner at the USC Dornsife Cognitive Neuroscience Imaging Center. Three-dimensional high-resolution T1-weighted images were acquired with a magnetization prepared rapid gradient echo (MP-RAGE) protocol as follows: inversion time (TI)/ repetition time (TR)/echo time (TE) = 800 / 2530/3.09 ms, slice thickness = 1 mm without gap, matrix = 256×256 , and field of view (FOV) = $256 \text{ mm} \times 256 \text{ mm}$. Image preprocessing (e.g. skull-stripping, correction of signal intensity and inhomogeneity artifacts) was conducted using Freesurfer processing streams where outputs were visually inspected and manually corrected for accuracy (Yang et al., 2011).

TBM relies on matching structures with similar intensity patterns, and measures volumetric differences in a population by analyzing the gradients of the non-linear deformation fields required to align individual images to an anatomical template specific to the population studied (Yang et al., 2011). To detect local differences in brain tissue structure associated with psychopathic traits, TBM processing streams were implemented in the LONI Pipeline environment (Rex et al., 2003) using methods similar to those described in earlier investigations (Leow et al., 2009; Yang et al., 2011). Specifically, TBM processing included optimized non-rigid registration models that quantify the symmetric Kullback-Leibler (KL)-distance between the anatomical template and the transformed individual subject images. Processing steps were as follows: (1) Each preprocessed image volume was first registered to a single image using a nine-parameter registration to adjust for global brain scale, head tilt and alignment. (2) The images were then used to create an anatomical template or minimal deformation target (MDT). This step matches each 3D volume to all other volumes using a mutual information-based inverse-consistent algorithm, followed by applying the inverse of the mean displacement field from all subjects to the MDT. (3) Image volumes from all subjects were each subsequently aligned to the MDT by nonlinearly deforming the anatomy of each individual image to match the anatomical template. The Jacobian operator was then applied to the deformation fields to produce univariate Jacobian determinants (i.e., Jacobian maps) that index the extent of local expansion or contraction required to non-linearly warp each brain to match each subject's anatomy to the MDT. These 3D Jacobian maps represent relative tissue volume differences between each individual and the MDT, and were used to characterize local differences in brain tissue structure associated with the targeted behavioral measure across the sample.

2.4. Statistical analysis

The R statistical package (version 2.9.2; <http://www.r-project.org/>) was used to compute differences in the deformation fields associated with CPS total scores across the entire brain using the mixed effects model while controlling for sex, whole brain volume, and subject relatedness. Random intercepts were included for each family to account for relatedness within families. The analysis was implemented using the ‘nlme’ library in the R statistical package (Pinheiro and Bates, 2000). Uncorrected probability values and the corresponding beta (regression coefficient) values from the analyses were mapped onto the MDT. Since comparisons were made at thousands of voxels, results were further thresholded using an enhanced False Discovery Rate (FDR)-control method (Langers et al., 2007). By setting the FDR to 5%, this study was able to control the expected proportion of incorrectly rejected null hypotheses (Benjamini and Hochberg, 1995). In the FDR-corrected maps, 95% of the reported findings reported are expected to be true positives, irrespective of how many contrasts were conducted. FDR-thresholded probability values from each comparison were mapped onto the MDT atlas and color was used to encode the regions and the directions of the FDR-controlled significant correlations between psychopathic traits and brain tissue volume respectively.

For descriptive purposes and to report the location of the observed effects, we extracted MNI atlas coordinate locations for regions showing significant structural volume differences using the Anatomy Toolbox V1.5 (Eickhoff et al., 2005) of the Statistical Parametric Mapping software (SPM8; <http://www.fil.ion.ucl.ac.uk/spm/software/spm8>) executed in MATLAB (Mathworks, Sherborn, Massachusetts) (Yang et al., 2011). Anatomic locations of clusters with > 2000 voxels are provided in Table S1 (see the Supplementary Material).

2.5. Post hoc analyses

Finally, we applied an independent and widely-used volume quantification method to confirm that local striatum expansions, specifically in the caudate nucleus, putamen and the pallidum, reflect volumetric differences associated with psychopathic traits, as observed in the TBM analyses. The caudate nucleus, putamen and the pallidum volumes were measured by employing the completely independent image segmentation methods of Freesurfer v5.1.0 (<http://surfer.nmr.mgh.harvard.edu>; (Fischl et al., 2002)). For these procedures, any topographical errors in segmentation were corrected manually on a case-by-case basis (Yang et al., 2011).

3. Results

3.1. TBM analysis across the whole brain

As shown in Fig. 1, parent-reported CPS total scores were significantly correlated with brain tissue volume differences in several frontal, temporal and striatal regions after FDR thresholding. Specifically, higher CPS total scores were correlated with increased brain tissue volumes in the left putamen, including the ansa peduncularis, right superiomedial frontal cortex, right orbitofrontal cortex and right medial temporal cortex across the subjects while controlling for sex, whole brain volume, and subject relatedness (see Supplemental Material, Figs. S3–S4 and Table S1). In addition, higher CPS total scores were found to

correlate with reduced brain tissue volumes in the right middle frontal gyrus, bilateral inferior frontal gyrus, right superior parietal lobule, left inferior parietal gyrus, left supplementary motor area and the occipital cortex (see Supplemental Material, Figs. S3–S4 and Table S1).

As an exploratory measure, we also conducted the analysis separately for males and females and found similar patterns for both sexes. For males, higher CPS total scores were found to correlate with increased volumes in the left putamen, left ansa peduncularis, and left superiomedial frontal gyrus, and reduced volumes in the right middle frontal, right superior parietal lobule, left inferior parietal lobule, left supplementary motor area and the occipital cortex (Fig. 1, b). For females, higher CPS total scores were found to correlate with increased volumes in the left putamen, bilateral gyrus rectus, left inferior frontal gyrus, and left insula and reduced volume in the occipital cortex (Fig. 2, c).

3.2. Post hoc analysis on striatal regions

Post hoc analyses performed to compare subcortical volume estimates obtained independently using Freesurfer confirmed volumetric increases of the left putamen to be correlated with increased parent-reported CPS scores ($p = 0.01$) across the sample of 108 adolescents (Fig. 2), adjusting for whole brain volume and sex. Even so, the associations for the left caudate nucleus and the left pallidum were below the threshold of significance ($p = 0.22, 0.88$, respectively). In addition, Freesurfer showed the volume of the right putamen was also significantly increased ($p = 0.015$). No significant correlation was found for right caudate or pallidum ($p = 0.81, 0.75$, respectively).

For male adolescents, we found increased parent-reported CPS scores to correlate with increased volume in the left putamen ($p = 0.016$) while adjusting for whole brain volume and subject relatedness, but not other striatal structures (all $p > 0.1$). For female adolescents, we found increased parent-reported CPS scores to correlate with increased volume in the right putamen ($p = 0.019$), but below the threshold of significance for the left putamen ($p = 0.08$) and left caudate ($p = 0.065$). No significant correlation was found for other striatal structures for females (all $p > 0.1$).

4. Discussion

Using a homogeneously aged sample of healthy young adolescents, we captured unique patterns of global and local brain dysmorphology associated with psychopathic scores. The principal findings of this study are that abnormal brain tissue volumes in several regions, particularly increased volumes in the putamen, ansa peduncularis, superiomedial prefrontal cortex, inferior frontal cortex, medial temporal regions, and reduced volumes in the middle frontal cortex, superior parietal lobule, and inferior parietal lobule, were significantly correlated with increased psychopathic tendency in adolescents. Interestingly, we observed sex-specific abnormalities associated with psychopathy traits. Particularly, higher psychopathy scores correlate more strongly with enlarged left putamen, reduced right middle frontal volume, and increased right superiomedial frontal volume in males. For females, higher psychopathy scores correlate more prominently with increased volumes in the bilateral gyrus rectus, left inferior frontal cortex, left insula, and right putamen. These

findings support the possibility that psychopathy has a neurodevelopmental basis, with disorder-related brain morphological characteristics detectable in younger populations. In addition, findings of this study further highlight the complexity of brain maturational processes that occur during adolescence with regard to their possible contributions to symptom development such as elevated psychopathic traits.

In line with previous sMRI studies of adults with high psychopathy scores (Yang et al., 2005; Raine et al., 2009; Yang et al., 2010; Ermer et al., 2012), psychopathic tendency in adolescents were found to be associated with brain tissue abnormalities in several frontal regions including reduced middle frontal volume, increased superiomedial and inferior frontal volumes, and increased volume in the gyrus rectus. Findings are in line with those from studies of boys with conduct disorders. For example, De Brito and colleagues showed increased grey matter concentrations in several frontal regions including the medial orbitofrontal cortex in boys with callous-unemotional traits and conduct problems (De Brito et al., 2009). Using the same sample, they also found abnormal white matter concentrations in several frontal regions including decreased white matter concentrations in the right superior frontal cortex and increased white matter concentrations in the bilateral middle frontal gyrus in boys with psychopathic tendency (De Brito et al., 2011). The frontal cortex is centrally involved in functions frequently associated with key impairments of psychopathic individuals, including response inhibition, fear conditioning, and delay of gratification (Patrick, 1994; Dolan and Fullam, 2006; Raine and Yang, 2006). Findings of morphological alterations in these frontal regions may also be supported by previous reports of abnormal activation in these frontal regions in psychopathic individuals (Birbaumer et al., 2005; Rilling et al., 2007; Finger et al., 2008; Veit et al., 2010). For example, psychopathy scores were associated with increased activation in middle frontal gyrus during judgments of fear-evoking moral scenarios in healthy adults (Marsh and Cardinale, 2012). Although any interpretation can only be highly speculative, stronger correlations observed in the middle and superiomedial frontal cortex in males may suggest that poor impulsivity and behavioral control plays a center role in the development of psychopathy in males. On the other hand, stronger correlations observed in the gyrus rectus in females may suggest that impaired emotional regulation and moral cognition is crucial in the development of psychopathy in females. Nonetheless, findings here provide strong support for morphological abnormalities in the frontal cortex to be potential biomarkers for psychopathic tendencies in adolescents.

Another key finding of this study is the correlation between brain tissue contractions in parietal lobe, specifically superior and inferior parietal lobules, and greater psychopathic tendencies in adolescents. Although these brain regions have not received much attention so far, they both play central roles in functions that are closely associated with psychopathic traits. One of the most critical contributions of the parietal lobules is that they are part of the mirror neuron system (Cattaneo and Rizzolatti, 2009). This system is composed of neural circuits activated during the observation of actions, emotions, and sensations - *and* during execution and experiencing of the same actions, emotions and sensations (Gallese, 2003). Therefore, the mirror neuron system is thought to play a role in key aspects of social behavior, such as mentalizing and empathy (Iacoboni et al., 2005; Gazzola et al., 2006). One core feature of psychopathy is the lack of empathy, including impaired recognition and response to sadness, observational learning, and lack of concern about the impact of one's

actions on others (Blair, 2005; Dolan and Fullam, 2006). This is supported by a previous report showing psychopathic traits, specifically coldheartedness, were positively correlated with the level of corticospinal excitability in the motor cortex while watching short videos known to activate the sensorimotor mirror neuron system for pain (Fecteau et al., 2008).

Findings here provide initial evidence suggesting that neuroanatomical disturbances in regions within the mirror neuron network may also be linked to psychopathy. Finding of abnormal insula associated with psychopathic traits in females may be of particular significance. The insula has long been implicated in empathy, the capability to understand and resonate with the affective experience of another, and is crucial in the inhibition of antisocial, aggressive behavior. Abnormal activation in the insula during empathy-related tasks has been linked to higher levels of psychopathy in several adult studies (Decety et al., 2013; Decety et al., 2014; Molenberghs et al., 2014). For developmental populations, prior studies on children with conduct problems have also demonstrated abnormal neural responses to others in pain in the insula (Sterzer et al., 2007). For example, Lockwood and colleagues found children with conduct problems to show reduced activation in the bilateral anterior insula while viewing pictures of others in pain (Lockwood et al., 2013). Furthermore, higher levels of callous traits were associated with reduced responses to others' pain in the anterior insula. Our findings provide further support for the argument that insula plays an important role in the developmental vulnerability to psychopathy, and is a potentially sensitive biomarker for psychopathic tendencies in female adolescents.

In addition to brain tissue abnormalities in the frontal and parietal regions, we also found brain tissue expansions in several subcortical regions, most prominently in the putamen, to be associated with heightened psychopathic tendencies in adolescents. Although the enlargement was more prominent in the left putamen for the males, and right putamen for the females, both sexes show similar positive relationships between increased volumes in the putamen and increase psychopathy scores. Findings are in line with some previous studies (Glenn et al., 2010a; Schiffer et al., 2011), but not the others (Fairchild et al., 2012), which could be due to differences in sample size, age of the cohort studied, and whether the focus was on psychopathy in general or more specific traits (i.e. callous-unemotional).

The striatum, including the putamen, is one of the structures most crucially involved in reward prediction and guiding action selection towards immediate over delayed reward (O'Doherty, 2004; Hariri et al., 2006). Structural brain abnormalities in the putamen may thus contribute to some of the most distinct features of psychopathy such as increased stimulation seeking, perseverative response, and reward-driven and impulsive behavior (Newman et al., 1987; Newman et al., 1992; Hare, 2003). The striatum is also a critical part of the neural circuitry underlying stimulus-reinforcement learning, i.e., learning appropriate behaviors for responding to rewarding stimuli and refraining from responding to those that lead to punishment (Newman and Kosson, 1986; Blair, 2006b). Stimulus-reinforcement learning is a critical component of the passive avoidance in psychopathy, which has been studied extensively (Newman and Kosson, 1986; Blair, 2006b), and is closely linked to decision-making, socialization, and empathy-based learning. Enlarged striatum as observed here may provide neurobiological evidence supporting the theoretical explanation of psychopathy that overactivation by reward leads to impaired ability to modulate the

dominant response for reward in psychopathic individuals (O'Brien and Frick, 1996). Sufficient evidence from animal studies showed that the striatum, particularly the lateral dorsal striatum (i.e. putamen) appears to be involved in habitual stimulus-response associations (Liljeholm and O'Doherty, 2012), which may further explain the response perseveration that is often observed in psychopathic individuals. Furthermore, enlarged ansa peduncularis, which is a main division of the ventral amygdalofugal pathway, was also found to be associated with higher psychopathy scores, particularly in males. The ventral amygdalofugal pathway contains fibers leaving the amygdaloid nucleus to several targets including the hypothalamus, thalamus, septal nuclei and nucleus accumbens, thus enlargement observed in this area may reflect abnormal connection between amygdala and associated structures. This is supported by findings of reduced connectivity with the amygdala (Cohen et al., 2009; Finger et al., 2012; Passamonti et al., 2012; Contreras-Rodriguez et al., 2014) and structural abnormalities in the amygdala (Yang et al., 2009c; Yang et al., 2010; Pardini et al., 2014) in psychopathic individuals.

An important question remains concerning how increased striatal volumes may predispose a person to psychopathic traits. Enlarged putamen has been observed in several other populations including individuals with methamphetamine-dependent (Chang et al., 2005), cocaine-dependent (Jacobsen et al., 2001) and schizophrenia patients with neuroleptic treatment (Gur et al., 1998; Taylor et al., 2005), and was argued to be the result of inflammation and reactive gliosis in the region, possibly assisted by glia-mediated neurotrophic effects to increase striatal sprouting (Song and Haber, 2000). Therefore, one explanation for the association between enlargement of the putamen and psychopathic traits as observed here would be that it represents a compensatory response to maintain function of the striatum. Another explanation for the enlarged putamen and sex-specific changes observed here might come from a developmental perspective. One recent study showed putamen declined in volume with age, and the decreases were more for females than males and greater in the left hemisphere (Dennison et al., 2013). This may be due to sex hormone changes through puberty that induce sex-specific synaptogenesis and synaptic pruning during adolescence. Nonetheless, our findings provide initial evidence suggesting that a delay or disruption in the development of putamen may contribute to the development of psychopathic tendency in adolescents, particularly heightened sensation-seeking and reward-dominated behavior.

Limitations of this study include the use of a twin sample, which may limit the generalizability of findings to the general population. However, our findings are mostly consistent with previous studies of singletons, suggesting structural alterations in frontal, temporal, parietal and striatal regions as critical components underlying early onset psychopathic tendency. Unlike other whole-brain volumetric methods, TBM does not require a segmentation step, which avoids potential partial volume errors associated with tissue classification. As such, results from this current study that point to differences in both gray and white matter structure may be less directly comparable to previous findings concentrating mainly on gray matter. Notably, for subcortical regions, results from the independent FreeSurfer method are complementary to those of TBM, however bilateral instead of unilateral enlargement in putamen and sex-related hemispheric effect was also observed. Nonetheless, both methods are supportive of the argument that striatal

abnormalities may contribute to the development of psychopathic traits in adolescents and future studies are needed to validate the sex- and hemisphere-specific results observed here. Last, there has been some evidence suggesting that pubertal stage may be a better predictor than chronological age for examining developmental changes in brain morphology, particularly subcortical structures (Blanton et al., 2012). Although the potential age-related brain variations among our subjects were limited by using a homogeneously-aged sample, future studies could benefit from further controlling for pubertal stage.

Overall, this study reveals valuable knowledge regarding the neural mechanisms underlying psychopathic traits in adolescents. Future studies examining how genetic predispositions interact with environmental influences in shaping the brain structure and function could help understand the development of psychopathic tendencies throughout the lifespan.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was supported by grants from National Institute of Mental Health (K99 MH093388 to Y.Y., R01MH58354 to L.A.B., R01MH092301 K.L.N., R01 AG040060 and P41 RR013642 to P.M.T.), and by a grant from the Brain and Creativity Institute, University of Southern California.

References

- Baker LA, Barton M, Lozano DI, Raine A, Fowler JH. The Southern California Twin Register at the University of Southern California. II. Twin Research and Human Genetics. 2006; 9:933–940. [PubMed: 17254433]
- Baker LA, Jacobson KC, Raine A, Lozano DI, Bezdjian S. Genetic and environmental bases of childhood antisocial behavior: a multi-informant twin study. *Journal of Abnormal Psychology*. 2007; 116:219–235. [PubMed: 17516756]
- Baker LA, Tuvblad C, Wang P, Gomez K, Bezdjian S, Niv S, Raine A. The Southern California Twin Register at the University of Southern California: III. Twin Research and Human Genetics. 2013; 16:336–343. [PubMed: 23394193]
- Barkataki I, Kumari V, Das M, Taylor P, Sharma T. Volumetric structural brain abnormalities in men with schizophrenia or antisocial personality disorder. *Behavioural Brain Research*. 2006; 169:239–247. [PubMed: 16466814]
- Barros-Loscertales A, Meseguer V, Sanjuan A, Belloch V, Parcet MA, Torrubia R, Avila C. Striatum gray matter reduction in males with an overactive behavioral activation system. *European Journal of Neuroscience*. 2006; 24:2071–2074. [PubMed: 17040475]
- Benjamini Y, Hochberg Y. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society. Series B*. 1995; 57:289–300.
- Bertsch K, Grothe M, Prehn K, Vohs K, Berger C, Hauenstein K, Keiper P, Domes G, Teipel S, Herpertz SC. Brain volumes differ between diagnostic groups of violent criminal offenders. *European Archives of Psychiatry and Clinical Neuroscience*. 2013; 263:593–606. [PubMed: 23381548]
- Bezdjian S, Baker LA, Tuvblad C. Genetic and environmental influences on impulsivity: a meta-analysis of twin, family and adoption studies. *Clinical Psychology Review*. 2011; 31:1209–1223. [PubMed: 21889436]
- Birbaumer N, Veit R, Lotze M, Erb M, Hermann C, Grodd W, Flor H. Deficient fear conditioning in psychopathy: a functional magnetic resonance imaging study. *Archives of General Psychiatry*. 2005; 62:799–805. [PubMed: 15997022]

- Blair RJ. Applying a cognitive neuroscience perspective to the disorder of psychopathy. *Developmental Psychopathology*. 2005; 17:865–891.
- Blair RJ. The emergence of psychopathy: implications for the neuropsychological approach to developmental disorders. *Cognition*. 2006a; 101:414–442. [PubMed: 16904094]
- Blair RJ. The neurobiology of psychopathic traits in youths. *Nature Reviews Neuroscience*. 2013; 14:786–799. [PubMed: 24105343]
- Blair, RJR. Subcortical brain systems in psychopathy. In: Patrick, CJ., editor. *Handbook of Psychopathy*. Guilford Press; New York: 2006b. p. 296–312.
- Blanton RE, Cooney RE, Joormann J, Eugene F, Glover GH, Gotlib IH. Pubertal stage and brain anatomy in girls. *Neuroscience*. 2012; 217:105–112. [PubMed: 22569152]
- Boccardi M, Ganzola R, Rossi R, Sabatoli F, Laakso MP, Repo-Tiihonen E, Vaurio O, Kononen M, Aronen HJ, Thompson PM, Frisoni GB, Tiihonen J. Abnormal hippocampal shape in offenders with psychopathy. *Human Brain Mapping*. 2010; 31:438–447. [PubMed: 19718651]
- Boes AD, Tranel D, Anderson SW, Nopoulos P. Right anterior cingulate: a neuroanatomical correlate of aggression and defiance in boys. *Behavioural Neuroscience*. 2008; 122:677–684.
- Carre JM, Hyde LW, Neumann CS, Viding E, Hariri AR. The neural signatures of distinct psychopathic traits. *Social Neuroscience*. 2013; 8:122–135. [PubMed: 22775289]
- Cattaneo L, Rizzolatti G. The mirror neuron system. *Archives of Neurology*. 2009; 66:557–560. [PubMed: 19433654]
- Chang L, Cloak C, Patterson K, Grob C, Miller EN, Ernst T. Enlarged striatum in abstinent methamphetamine abusers: a possible compensatory response. *Biological Psychiatry*. 2005; 57:967–974. [PubMed: 15860336]
- Cohen MX, Schoene-Bake JC, Elger CE, Weber B. Connectivity-based segregation of the human striatum predicts personality characteristics. *Nature Neuroscience*. 2009; 12:32–34. [PubMed: 19029888]
- Contreras-Rodriguez O, Pujol J, Batalla I, Harrison BJ, Soriano-Mas C, Deus J, Lopez-Sola M, Macia D, Pera V, Hernandez-Ribas R, Pifarre J, Menchon JM, Cardoner N. Functional connectivity bias in the prefrontal cortex of psychopaths. *Biological Psychiatry*. 2014 electronic publication 4 March. 10.1016/j.biopsych.2014.03.007
- Cope LM, Shane MS, Segall JM, Nyalakanti PK, Stevens MC, Pearlson GD, Calhoun VD, Kiehl KA. Examining the effect of psychopathic traits on gray matter volume in a community substance abuse sample. *Psychiatry Research: Neuroimaging*. 2012; 204:91–100. [PubMed: 23217577]
- Dalwani M, Sakai JT, Mikulich-Gilbertson SK, Tanabe J, Raymond K, McWilliams SK, Thompson LL, Banich MT, Crowley TJ. Reduced cortical gray matter volume in male adolescents with substance and conduct problems. *Drug and Alcohol Dependence*. 2011; 118:295–305. [PubMed: 21592680]
- De Brito SA, McCrory EJ, Mechelli A, Wilke M, Jones AP, Hodgins S, Viding E. Small, but not perfectly formed: decreased white matter concentration in boys with psychopathic tendencies. *Molecular Psychiatry*. 2011; 16:476–477. [PubMed: 20548295]
- De Brito SA, Mechelli A, Wilke M, Laurens KR, Jones AP, Barker GJ, Hodgins S, Viding E. Size matters: increased grey matter in boys with conduct problems and callous-unemotional traits. *Brain*. 2009; 132:843–852. [PubMed: 19293245]
- de Oliveira-Souza R, Hare RD, Bramati IE, Garrido GJ, Azevedo Ignacio F, Tovar-Moll F, Moll J. Psychopathy as a disorder of the moral brain: fronto-temporo-limbic grey matter reductions demonstrated by voxel-based morphometry. *Neuroimage*. 2008; 40:1202–1213. [PubMed: 18289882]
- Decety J, Chen C, Harenski C, Kiehl KA. An fMRI study of affective perspective taking in individuals with psychopathy: imagining another in pain does not evoke empathy. *Frontiers in Human Neuroscience*. 2013; 7:489. [PubMed: 24093010]
- Decety J, Skelly L, Yoder KJ, Kiehl KA. Neural processing of dynamic emotional facial expressions in psychopaths. *Social Neuroscience*. 2014; 9:36–49. [PubMed: 24359488]
- Dennison M, Whittle S, Yucel M, Vijayakumar N, Kline A, Simmons J, Allen NB. Mapping subcortical brain maturation during adolescence: evidence of hemisphere- and sex-specific longitudinal changes. *Developmental Sciences*. 2013; 16:772–791.

- Dolan M, Fullam R. Face affect recognition deficits in personality-disordered offenders: association with psychopathy. *Psychological Medicine*. 2006; 36:1563–1569. [PubMed: 16893483]
- Eickhoff SB, Stephan KE, Mohlberg H, Grefkes C, Fink GR, Amunts K, Zilles K. A new SPM toolbox for combining probabilistic cytoarchitectonic maps and functional imaging data. *Neuroimage*. 2005; 25:1325–1335. [PubMed: 15850749]
- Ermer E, Cope LM, Nyalakanti PK, Calhoun VD, Kiehl KA. Aberrant paralimbic gray matter in criminal psychopathy. *Journal of Abnormal Psychology*. 2012; 121:649–658. [PubMed: 22149911]
- Fairchild G, Hagan CC, Walsh ND, Passamonti L, Calder AJ, Goodyer IM. Brain structure abnormalities in adolescent girls with conduct disorder. *Journal of Child Psychology and Psychiatry*. 2012; 54:86–95. [PubMed: 23082797]
- Fairchild G, Passamonti L, Hurford G, Hagan CC, von dem Hagen EA, van Goozen SH, Goodyer IM, Calder AJ. Brain structure abnormalities in early-onset and adolescent-onset conduct disorder. *American Journal of Psychiatry*. 2011; 168:624–633. [PubMed: 21454920]
- Fecteau S, Pascual-Leone A, Theoret H. Psychopathy and the mirror neuron system: preliminary findings from a non-psychiatric sample. *Psychiatry Research*. 2008; 160:137–144. [PubMed: 18599127]
- Finger EC, Marsh A, Blair KS, Majestic C, Evangelou I, Gupta K, Schneider MR, Sims C, Pope K, Fowler K, Sinclair S, Tovar-Moll F, Pine D, Blair RJ. Impaired functional but preserved structural connectivity in limbic white matter tracts in youth with conduct disorder or oppositional defiant disorder plus psychopathic traits. *Psychiatry Research*. 2012; 202:239–244. [PubMed: 22819939]
- Finger EC, Marsh AA, Mitchell DG, Reid ME, Sims C, Budhani S, Kosson DS, Chen G, Towbin KE, Leibenluft E, Pine DS, Blair JR. Abnormal ventromedial prefrontal cortex function in children with psychopathic traits during reversal learning. *Archives of General Psychiatry*. 2008; 65:586–594. [PubMed: 18458210]
- Fischl B, Salat DH, Busa E, Albert M, Dieterich M, Haselgrove C, van der Kouwe A, Killiany R, Kennedy D, Klaveness S, Montillo A, Makris N, Rosen B, Dale AM. Whole brain segmentation: automated labeling of neuroanatomical structures in the human brain. *Neuron*. 2002; 33:341–355. [PubMed: 11832223]
- Gallese V. The roots of empathy: the shared manifold hypothesis and the neural basis of intersubjectivity. *Psychopathology*. 2003; 36:171–180. [PubMed: 14504450]
- Gao Y, Glenn AL, Schug RA, Yang Y, Raine A. The neurobiology of psychopathy: A neurodevelopmental perspective. *Canadian Journal of Psychiatry*. 2009; 54:813–823. [PubMed: 20047720]
- Gazzola V, Aziz-Zadeh L, Keysers C. Empathy and the somatotopic auditory mirror system in humans. *Current Biology*. 2006; 16:1824–1829. [PubMed: 16979560]
- Glenn AL, Raine A, Yaralian PS, Yang Y. Increased volume of the striatum in psychopathic individuals. *Biological Psychiatry*. 2009; 67:52–58. [PubMed: 19683706]
- Glenn AL, Yang Y. The potential role of the striatum in antisocial behavior and psychopathy. *Biological Psychiatry*. 2012; 72:817–822. [PubMed: 22672927]
- Glenn AL, Yang Y, Raine A, Colletti P. No volumetric differences in the anterior cingulate of psychopathic individuals. *Psychiatry Research: Neuroimaging*. 2010b; 183:140–143. [PubMed: 20630717]
- Gur RE, Maany V, Mozley PD, Swanson C, Bilker W, Gur RC. Subcortical MRI volumes in neuroleptic-naïve and treated patients with schizophrenia. *American Journal of Psychiatry*. 1998; 155:1711–1717. [PubMed: 9842780]
- Hare, RD. *Manual for the Revised Psychopathy Checklist. 2*. Multi-Health Systems; Toronto, ON, Canada: 2003.
- Hariri AR, Brown SM, Williamson DE, Flory JD, de Wit H, Manuck SB. Preference for immediate over delayed rewards is associated with magnitude of ventral striatal activity. *Journal of Neuroscience*. 2006; 26:13213–13217. [PubMed: 17182771]
- Huebner T, Vloet TD, Marx I, Konrad K, Fink GR, Herpertz SC, Herpertz-Dahlmann B. Morphometric brain abnormalities in boys with conduct disorder. *Journal of the American Academy of Child and Adolescent Psychiatry*. 2008; 47:540–547. [PubMed: 18356764]

- Hyatt CJ, Haney-Caron E, Stevens MC. Cortical thickness and folding deficits in conduct-disordered adolescents. *Biological Psychiatry*. 2012; 72:207–214. [PubMed: 22209639]
- Iacoboni M, Molnar-Szakacs I, Gallese V, Buccino G, Mazziotta JC, Rizzolatti G. Grasping the intentions of others with one's own mirror neuron system. *PLoS Biology*. 2005; 3:e79. [PubMed: 15736981]
- Jacobsen LK, Giedd JN, Gottschalk C, Kosten TR, Krystal JH. Quantitative morphology of the caudate and putamen in patients with cocaine dependence. *American Journal of Psychiatry*. 2001; 158:486–489. [PubMed: 11229995]
- Kruesi MJP, Casanova MV, Mannheim G, Jonson-Bilder A. Reduced temporal lobe volume in early-onset conduct disorder. *Psychiatry Research: Neuroimaging*. 2004; 132:1–11. [PubMed: 15546698]
- Laakso MP, Vaurio O, Koivisto E, Savolainen L, Eronen M, Aronen HJ, Hakola P, Repo E, Soininen H, Tiihonen J. Psychopathy and the posterior hippocampus. *Behavioral Brain Research*. 2001; 118:187–193.
- Langers DR, Jansen JF, Backes WH. Enhanced signal detection in neuroimaging by means of regional control of the global false discovery rate. *Neuroimage*. 2007; 38:43–56. [PubMed: 17825583]
- Leow AD, Yanovsky I, Parikshak N, Hua X, Lee S, Toga AW, Jack CR Jr, Bernstein MA, Britson PJ, Gunter JL, Ward CP, Borowski B, Shaw LM, Trojanowski JQ, Fleisher AS, Harvey D, Kornak J, Schuff N, Alexander GE, Weiner MW, Thompson PM. Alzheimer's disease neuroimaging initiative: a one-year follow up study using tensor-based morphometry correlating degenerative rates, biomarkers and cognition. *Neuroimage*. 2009; 45:645–655. [PubMed: 19280686]
- Liljeholm M, O'Doherty JP. Contributions of the striatum to learning, motivation, and performance: an associative account. *Trends in Cognitive Sciences*. 2012; 16:467–475. [PubMed: 22890090]
- Lockwood PL, Sebastian CL, McCrory EJ, Hyde ZH, Gu X, De Brito SA, Viding E. Association of callous traits with reduced neural response to others' pain in children with conduct problems. *Current Biology*. 2013; 23:901–905. [PubMed: 23643836]
- Lynam DR. Pursuing the psychopath: capturing the fledgling psychopath in a nomological net. *Journal of Abnormal Psychology*. 1997; 106:425–438. [PubMed: 9241944]
- Marsh AA, Cardinale EM. When psychopathy impairs moral judgments: Neural responses during judgments about causing fear. *Social Cognitive & Affective Neuroscience*. 2012; 9:3–11. [PubMed: 22956667]
- Molenberghs P, Bosworth R, Nott Z, Louis WR, Smith JR, Amiot CE, Vohs KD, Decety J. The influence of group membership and individual differences in psychopathy and perspective taking on neural responses when punishing and rewarding others. *Human Brain Mapping*. 2014
- Muller JL, Ganssbauer S, Sommer M, Dohnel K, Weber T, Schmidt-Wilcke T, Hajak G. Gray matter changes in right superior temporal gyrus in criminal psychopaths. Evidence from voxel-based morphometry. *Psychiatry Research: Neuroimaging*. 2008; 163:213–222. [PubMed: 18662867]
- Newman JP, Kosson DS. Passive avoidance learning in psychopathic and nonpsychopathic offenders. *Journal of Abnormal Psychology*. 1986; 95:252–256. [PubMed: 3745647]
- Newman JP, Kosson DS, Patterson CM. Delay of gratification in psychopathic and nonpsychopathic offenders. *Journal of Abnormal Psychology*. 1992; 101:630–636. [PubMed: 1430601]
- Newman JP, Patterson CM, Kosson DS. Response perseveration in psychopaths. *Journal of Abnormal Psychology*. 1987; 96:145–148. [PubMed: 3584663]
- O'Brien BS, Frick PJ. Reward dominance: associations with anxiety, conduct problems, and psychopathy in children. *Journal of Abnormal Child Psychology*. 1996; 24:223–240. [PubMed: 8743246]
- O'Doherty JP. Reward representations and reward-related learning in the human brain: insights from neuroimaging. *Current Opinion in Neurobiology*. 2004; 14:769–776. [PubMed: 15582382]
- Osumi T, Nakao T, Kasuya Y, Shinoda J, Yamada J, Ohira H. Amygdala dysfunction attenuates frustration-induced aggression in psychopathic individuals in a non-criminal population. *Journal of Affective Disorders*. 2012; 142:331–338. [PubMed: 22840629]
- Pardini DA, Raine A, Erickson K, Loeber R. Lower amygdala volume in men is associated with childhood aggression, early psychopathic traits, and future violence. *Biological Psychiatry*. 2014; 75:73–80. [PubMed: 23647988]

- Passamonti L, Fairchild G, Fornito A, Goodyer IM, Nimmo-Smith I, Hagan CC, Calder AJ. Abnormal anatomical connectivity between the amygdala and orbitofrontal cortex in conduct disorder. *PLoS One*. 2012; 7:e48789. [PubMed: 23144970]
- Patrick CJ. Emotion and psychopathy: startling new insights. *Psychophysiology*. 1994; 31:319–330. [PubMed: 10690912]
- Pinheiro, JC.; Bates, DM. *Mixed-Effects Models in S and S-PLUS*. Springer; New York: 2000.
- Raine A, Yang Y. Neural foundations to moral reasoning and antisocial behavior. *Social Cognitive & Affective Neuroscience*. 2006; 1:203–213. [PubMed: 18985107]
- Raine A, Yang Y, Narr KL, Toga AW. Sex differences in orbitofrontal gray as a partial explanation for sex difference in antisocial personality. *Molecular Psychiatry*. 2009; 16:227–236. [PubMed: 20029391]
- Rex DE, Ma JQ, Toga AW. The LONI Pipeline Processing Environment. *Neuroimage*. 2003; 19:1033–1048. [PubMed: 12880830]
- Rilling JK, Glenn AL, Jairam MR, Pagnoni G, Goldsmith DR, Elfenbein HA, Lilienfeld SO. Neural correlates of social cooperation and non-cooperation as a function of psychopathy. *Biological Psychiatry*. 2007; 61:1260–1271. [PubMed: 17046722]
- Schiffer B, Muller BW, Scherbaum N, Hodgins S, Forsting M, Wiltfang J, Gizewski ER, Leygraf N. Disentangling structural brain alterations associated with violent behavior from those associated with substance use disorders. *Archives of General Psychiatry*. 2011; 68:1039–1049. [PubMed: 21646569]
- Song DD, Haber SN. Striatal responses to partial dopaminergic lesion: evidence for compensatory sprouting. *Journal of Neuroscience*. 2000; 20:5102–5114. [PubMed: 10864967]
- Sowell E. Mapping cortical gray matter asymmetry patterns in adolescents with heavy prenatal alcohol exposure. *Neuroimage*. 2002; 17:1807–1819. [PubMed: 12498754]
- Sowell ER, Peterson BS, Thompson PM, Welcome SE, Henkenius AL, Toga AW. Mapping cortical change across the human life span. *Nature Neuroscience*. 2003; 6:309–315. [PubMed: 12548289]
- Sterzer P, Stadler C, Poustka F, Kleinschmidt A. A structural neural deficit in adolescents with conduct disorder and its association with lack of empathy. *Neuroimage*. 2007; 37:335–342. [PubMed: 17553706]
- Taylor S, Christensen JD, Holcomb JM, Garver DL. Volume increases in striatum associated with positive symptom reduction in schizophrenia: a preliminary observation. *Psychiatry Research: Neuroimaging*. 2005; 140:85–89. [PubMed: 16194599]
- van Honk J, Schutter DJ. Unmasking feigned sanity: a neurobiological model of emotion processing in primary psychopathy. *Cognitive Neuropsychiatry*. 2006; 11:285–306. [PubMed: 17354073]
- Veit R, Lotze M, Sewing S, Missenhardt H, Gaber T, Birbaumer N. Aberrant social and cerebral responding in a competitive reaction time paradigm in criminal psychopaths. *Neuroimage*. 2010; 49:3365–3372. [PubMed: 19948223]
- Wechsler, D. *The Wechsler Intelligence Scale for Children*. 4. Pearson Assessment; London: 2004.
- Yang Y, Joshi AA, Joshi SH, Baker LA, Narr KL, Raine A, Thompson PM, Damasio H. Genetic and environmental influences on cortical thickness among 14-year-old twins. *Neuroreport*. 2012; 23:702–706. [PubMed: 22713927]
- Yang Y, Nuechterlein KH, Phillips OR, Gutman B, Kurth F, Dinov I, Thompson PM, Asarnow RF, Toga AW, Narr KL. Disease and genetic contributions toward local tissue volume disturbances in schizophrenia: a tensor-based morphometry study. *Human Brain Mapping*. 2012; 33:2081–2091. [PubMed: 22241649]
- Yang Y, Raine A. Prefrontal structural and functional brain imaging findings in antisocial, violent, and psychopathic individuals: a meta-analysis. *Psychiatry Research: Neuroimaging*. 2009; 174:81–88. [PubMed: 19833485]
- Yang Y, Raine A, Colletti P, Toga AW, Narr KL. Abnormal temporal and prefrontal cortical gray matter thinning in psychopaths. *Molecular Psychiatry*. 2009a; 14:561–562. 555. [PubMed: 19455172]
- Yang Y, Raine A, Colletti P, Toga AW, Narr KL. Morphological alterations in the prefrontal cortex and the amygdala in unsuccessful psychopaths. *Journal of Abnormal Psychology*. 2010; 119:546–554. [PubMed: 20677843]

- Yang Y, Raine A, Lencz T, Bihle S, LaCasse L, Colletti P. Volume reduction in prefrontal gray matter in unsuccessful criminal psychopaths. *Biological Psychiatry*. 2005; 57:1103–1108. [PubMed: 15866549]
- Yang Y, Raine A, Narr KL, Colletti P, Toga AW. Localization of deformities within the amygdala in individuals with psychopathy. *Archives of General Psychiatry*. 2009b; 66:986–994. [PubMed: 19736355]

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

- Structural magnetic resonance imaging data were obtained from 108 adolescents using tensor-based morphometry.
- Higher levels of psychopathic traits correlated with increased volumes in the left putamen, left ansa peduncularis, right superiomedial prefrontal cortex, left inferior frontal cortex, right orbitofrontal cortex, and right medial temporal regions.
- Higher levels of psychopathic traits also correlated with reduced volumes in the right middle frontal cortex, left superior parietal lobule, and left inferior parietal lobule.
- Poor decision-making and emotional dysregulation associated with psychopathy may reflect structural anomalies in frontal and temporal regions whereas striatal structural variations may contribute to sensation-seeking and reward-driven behavior in psychopathic individuals.

Fig. 1.

The left panel illustrates FDR-corrected ($q = 0.05$) probability maps for (a) total sample, (b) males, and (c) females showing significant correlations between psychopathic tendencies and brain tissue volumes, and the right panel shows the corresponding beta maps indicating the direction of the effects. Cold colors indicate negative and hot colors positive correlations.

Fig. 2.
Correlations between total Childhood Psychopathy Scale (CPS) scores and gray matter volumes of the left and right putamen, corrected for total brain volumes and subject relatedness.