Adjustments of lipid pathways in plant adaptation to temperature stress

Qiang Li¹, Wenyun Shen², Qian Zheng³, D. Brian Fowler¹, and Jitao Zou^{2,*}

¹Department of Plant Science; University of Saskatchewan; Saskatoon, Saskatchewan, Canada; ²National Research Council Canada; Saskatoon, Saskatchewan, Canada;

³Department of Food and Bioproduct Science; University of Saskatchewan; Saskatoon, Saskatchewan, Canada

Modulation of membrane lipid composition under varying environmental conditions is an important part of plant stress adaptation. Most notably, proportional changes of lipid composition in response to temperature changes are a major cellular response to requirements of membrane fluidity adjustment. In higher plants, synthesis of glycerolipids is accomplished by 2 major pathways, the prokaryotic and eukaryotic pathway, located in the chloroplast and the endoplasmic reticulum (ER), respectively. Recently, we systematically investigated the re-adjustments of glycerolipid pathways under temperature stress at the metabolite and transcript levels using 3 plant species with distinct lipid profiles. The relative contributions of 2 pathways and lipid channeling from the ER and chloroplast were both observed in plants under temperature stress. Potential factors controlling the lipid flux were identified through transcriptome analysis.

Keywords: fatty acid metabolism, glycerolipid pathways, metabolic regulation, RNA sequencing, temperature adaptation

Abbreviations: PG, phosphatidylglycerol; MGDG, monogalactosyldiacylglycerol; DGDG, digalactosyldiacylglycerol; C16, chain with 16 carbon; C18, chain with 18 carbon; DAG, diacylglycerol; TAG, triacylglycerol; ER, endoplasmic reticulum.

*Correspondence to: Jitao Zou; Email: jitao.zou@nrc-cnrc.gc.ca

Submitted: 05/22/2015

Accepted: 06/01/2015

<http://dx.doi.org/10.1080/15592324.2015.1058461>

Addendum to: Li Q, Zheng Q, Shen W, Cram D, Fowler DB, Wei Y, Zou J. Understanding the biochemical basis of temperature-induced lipid pathway adjustments in plants. *Plant Cell* 2015; 27:86-103. To whom correspondence should be sent. Email: Jitao.Zou@nrc-cnrc.gc.ca

of membrane lipid attributes pertinent to temperature stress adaptation.

In plant cells, the cytosolic and chloroplast compartment each possesses its own set of glycerolipid assembly pathway and generates glycerolipid species of defined molecular characteristics.⁷⁻⁹ The lipid pathway located in the chloroplast, which traces its origin to symbiogenesis, is called the prokaryotic pathway. The cytosolic endoplasmic reticulum (ER) lipid pathway, on the other hand, is known as the eukaryotic pathway. The two pathways are separated by barriers of the chloroplast envelope, but they are by no means insulated from each other. In fact, a significant portion of the glycerolipid moieties required for the biogenesis of photosynthetic membrane systems in the chloroplast is supplied by the eukaryotic pathway. The relative contributions of the 2 pathways, however, are very different among plant species.^{10,11} In species like *Arabidopsis*, about half of the glycerolipid moieties in chloroplast come from the eukaryotic pathway.⁸ In species, such as wheat, construction of photosynthesis apparatus relies almost entirely on the eukaryotic pathway, while the contribution of prokaryotic pathway is limited to the production of phosphatidylglycerol (PG) only.^{11,12} Despite the divergent pattern of relevant inputs from the 2 pathways, at the core is a basic requirement of glycerolipid channeling from the ER to chloroplasts in all plant species.

Changes in the proportion of glycerolipid classes under temperature stress have been previously reported in different plant species. The most common change is the adjustment between monogalactosyldiacylglycerol (MGDG) and digalactosyldiacylglycerol (DGDG), and it has been proposed that this metabolic change aids

A key aspect of plant adaptation to temperature stress concerns modulation of membrane fluidity.¹⁻³ Membrane fluidity is affected by 2 key aspects of lipid composition: (1) the degree of unsaturation in the fatty acyl groups of glycerolipid molecules and (2) the relative proportions of various lipid classes in the lipid bilayer.²⁻⁵ We recently conducted a comprehensive survey on glycerolipid metabolite as well as enzyme gene transcript changes under temperature stress in 3 plant species, *Arabidopsis thaliana* (*Arabidopsis*), *Atriplex lentiformis* (saltbush) and *Triticum aestivum* L. (wheat).⁶ Our results show that glycerolipid pathway adjustment between the chloroplast and cytosolic compartments are fundamental to the modulation

in survival at sub-optimal temperatures.¹³ A greater proportion of DGDG has been observed at high temperatures in *Arabidopsis*, while low temperature induced the synthesis of MGDG in *Brassica napus*.^{13,14} In addition to the alterations in the proportion of glycerolipid species, changes were also observed in the same lipid species with different fatty acid combinations. In *Brassica napus*, the major molecular species of MGDG is a mixture of *sn*-118:3/*sn*-2 16:3 and *sn*-1 18:3/*sn*-2 18:3 molecule species.^{8,11} An increase in MGDG (18:3/16:3) relative to MGDG (18:3/18:3) molecule species were observed in *Brassica napus* leaves grown at low temperature.¹⁴ In contrast, a decrease in MGDG (18:3/16:3) relative to MGDG (18:3/18:3) was observed in *Arabidopsis* grown at high temperatures.¹³

Rebalancing of the 2 glycerolipid pathways influences both the distribution of glycerolipid classes of different head groups and the degree of fatty acid desaturation in glycerolipid molecules. Due to substrate specificity of the plastidic lysophosphatidic acyltransferase, the prokaryotic pathway produces glycerolipids with C16 fatty acyl moiety at the *sn*-2 position of the glycerol backbone, and hence a molecular configuration of C34 (C18/C16, *sn*-1/*sn*-2).^{8,11} The *sn*-2 C16 fatty acids in the MGDG originating from the prokaryotic pathway are subsequently desaturated to form C16:3 fatty acids, leading to the formation of the highly unsaturated C34:6 (18:3/16:3) MGDG. The eukaryotic pathway generates glycerolipids with only C18 fatty acid at the *sn*-2, and produces C36 (C18/C18) and C34 (C16:0/C18). Since the *sn*-1 16:0 does not undergo any further desaturation, the eukaryotic C34:3 (16:0/18:3) has a level of desaturation much lower than that of the prokaryotic C34:6 (18:3/16:3). Through comprehensive lipid profiling and lipidomics analysis, we have shown that high temperatures repress the input of the prokaryotic pathway and promote the channeling of eukaryotic lipids to the chloroplast, thereby reducing potential of fatty acid desaturation in the photosynthesis apparatus. Significantly, we also observed that there was a differential channeling of eukaryotic glycerolipid species from the ER to the chloroplast: C34

diacylglycerol (DAG) moieties of 16:0/C18 (*sn*-1/*sn*-2) from the ER were preferentially transported to the chloroplast over that of C36 (C18/C18) under high temperature. These two layers of metabolic adjustments collectively reduce the overall degree of fatty acid desaturation in chloroplast lipids, particularly that of MGDG, at higher temperature. In wheat, where chloroplast lipid synthesis is dependent completely on imports from the ER, a differential channeling of C34 (C16/C18) DAG versus C36 (C18/C18) DAG was also observed. Under low temperature, there was a reduction of C34 DAG with a corresponding increase of C36 DAG. This metabolic alteration increased the overall potential of glycerolipid unsaturation in the chloroplast, i.e. increased level of C36 (18:3/18:3). The California desert plant *A. lentiformis* presents an exceptional case, which shuts off its prokaryotic pathways at high temperature, and falls back on the eukaryotic pathway completely for chloroplast lipid synthesis.¹⁵ This metabolic rebalancing allows the ER to deliver more C34 DAG (16:0/C18), thereby reducing the degree of lipid desaturation under extreme heat. These three species demonstrate the essential role glycerolipid pathway adjustment during adaptation to temperature stress.

Transcript analysis of our study and previous reports showed that, consistent with changes observed at the metabolite level, glycerolipid pathway coordination was reflected at the level of gene transcripts.^{16,17} The prokaryotic glycerol-3-phosphate acyltransferase *ACT1*^{18,19} and the chloroplast desaturase *FATTY ACID DESATURASE 5 (FAD5)*^{20,21} are key components influencing the prokaryotic glycerolipid pathway, while the *FATTY ACID DESATURASE 2 (FAD2)*²²⁻²⁴ in the ER was tightly associated with activities of the eukaryotic pathway. In addition to polar lipid biosynthesis pathways, we also detected a large number of differentially expressed genes involved in other branches of lipid metabolism (Figure 1).^{6,25} Triacylglycerols (TAGs) in *Arabidopsis* leaf tissues has been implicated in short-term glycerolipid intermediate provision during membrane lipid remodeling.^{26,27} Accumulation of TAGs in plant leaves

has been reported in *Arabidopsis* under freezing temperature.²⁸ In wheat, about 50 genes related to TAG synthesis were induced at 4°C. This was contrasted by only a few genes affected under heat at 43°C in *A. lentiformis*. We also detected up-regulation of a large number of genes involved in fatty acid elongation and wax biosynthesis at 4°C in wheat. In both *A. lentiformis* and wheat, significant numbers of phospholipid signaling genes were perturbed, suggesting active lipid signaling and lipid transfer events in this process.

Our work draws attention to the importance of glycerolipid pathway coordination during adaptation to temperature stress. Differential channeling of DAG moieties with various combinations of fatty acid moieties from the ER to chloroplast presents another layer of regulatory fine tuning for membrane unsaturation in chloroplast. A challenging question is why plants possess so many desaturase genes but still depend on glycerolipid pathway adjustment to modulate membrane desaturation. The simplest explanation would be that different intracellular compartments as well as developmental and tissues specific regulations demand the participation of multiple members of desaturase. Members of the desaturase family generally have defined glycerolipid substrate and/or fatty acid chain length specificity.^{29,30} Hence, their roles in mediating the desaturation of non-glycerolipids components of membrane systems during stress adaptation are also important.^{30,31} We suggest that specification of desaturases and pathway integration are 2 sides of one regulatory regime; together it allows the membrane system responding to temperature changes, but not in a fashion of a one way extreme, because in nature temperature fluctuates constantly. Adjustment of the 2 pathways provide a framework for desaturases to impart membrane compositional changes, but at the same time, stabilizes the lipid metabolic network against drastic changes to ensure biological robustness. Questions remain as to how lipids are selectively transported between chloroplast and the ER. In addition, the roles of other lipid species such as triacylglycerol, wax, oxylipin are also worthy topics for further investigation.

Figure 1. Number of lipid metabolism genes with significant changes in plants grown at low or high temperatures. *A. lentiformis* plants were grown at 43°C (high temperature) and 23°C (control) while wheat plants were grown at 4°C (low temperature) and 23°C (control). RNA sequencing was performed for *A. lentiformis* and wheat at high or low temperature respectively. Differentially expression genes (fold change >2 or <-2, p value <0.05) in *A. lentiformis* (43°C vs 23°C) and wheat (4°C vs 23°C) were identified.⁶ Genes involved in lipid metabolism were compiled from the Arabidopsis Lipid Gene Database (<http://aralip.plantbiology.msu.edu/>)²⁵ and categorized based on different pathways.

Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

Acknowledgments

The authors would like to thank Drs Mark Smith, Faouzi Bekkaoui of National Research Council Canada-Saskatoon, and Professor Melike Bor of Ege University, Turkey, for helpful discussions.

Funding

This research was supported in part by Canadian Wheat Alliance (CWA) program. This research is a National Research Council Canada Publication #56136.

References

- Murata N and Los DA. Membrane fluidity and temperature perception. *Plant Physiol.* 1997; 115:875-9; PMID:12223851
- Harwood J. Strategies for coping with low environmental temperatures. *Trends Biochem Sci.* 1991; 16:126-7; PMID:1877086; [http://dx.doi.org/10.1016/0968-0004\(91\)90052-W](http://dx.doi.org/10.1016/0968-0004(91)90052-W)
- Iba K. Acclimative response to temperature stress in higher plants: approaches of gene engineering for temperature tolerance. *Ann. Rev. Plant. Biol.* 2002; 53:225-45; <http://dx.doi.org/10.1146/annurev.arplant.53.100201.160729>
- Murakami Y, Tsuyama M, Kobayashi Y, Kodama H, Iba K. Trienoic fatty acids and plant tolerance of high temperature. *Science* 2000; 287:476-9; PMID:10642547; <http://dx.doi.org/10.1126/science.287.5452.476>
- Murata N, Ishizaki-Nishizawa O, Higashi S, Hayashi H, Tasaka Y, Nishida I. Genetically engineered alteration in the chilling sensitivity of plants. *Nature* 1992; 356:710-3; <http://dx.doi.org/10.1038/356710a0>
- Li Q, Zheng Q, Shen W, Cram D, Fowler DB, Wei Y, Zou J. Understanding the biochemical basis of temperature-induced lipid pathway adjustments in plants. *Plant Cell* 2015; 27:86-103; PMID:25564555; <http://dx.doi.org/10.1105/tpc.114.134338>
- Browse J and Somerville C. Glycerolipids synthesis: biochemistry and regulation. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 1991; 42:467-506; <http://dx.doi.org/10.1146/annurev.pp.42.060191.002343>
- Ohlrogge J and Browse J. Lipid biosynthesis. *Plant Cell* 1995; 7: 957-70; PMID:7640528; <http://dx.doi.org/10.1105/tpc.7.7.957>
- Hölzl G, Witt S, Gaude N, Melzer M, Schöttler MA, and Dörmann P. The role of diglycosyl lipids in photosynthesis and membrane lipid homeostasis in *Arabidopsis*. *Plant Physiol* 2009; 150:1147-59; PMID:19403724; <http://dx.doi.org/10.1104/pp.109.139758>
- Mongrand S, Bessoule J, Cabantous F and Cassagne C. The C_{16:3}/C_{18:3} fatty acid balance in photosynthetic tissues from 468 plant species. *Phytochemistry* 1998; 49:1049-64; [http://dx.doi.org/10.1016/S0031-9422\(98\)00243-X](http://dx.doi.org/10.1016/S0031-9422(98)00243-X)
- Heinz E and Roughan PG. Similarities and differences in lipid metabolism of chloroplasts isolated from 18:3 and 16:3 Plants. *Plant Physiol.* 1983; 72:273-9; PMID:16662992; <http://dx.doi.org/10.1104/pp.72.2.273>
- Arunga RO and Morrison WR. The structural analysis of wheat flour glycerolipids. *Lipids* 1971; 6:768-76; <http://dx.doi.org/10.1007/BF02531305>
- Chen J, Burke JJ, Xin Z, Xu C and Velten J. Characterization of the *Arabidopsis* thermosensitive mutant atts02 reveals an important role for galactolipids in thermotolerance. *Plant, Cell and Environment* 2006; 29:1437-48; <http://dx.doi.org/10.1111/j.1365-3040.2006.01527.x>
- Johnson G, Williams JP. Effect of growth temperature on the biosynthesis of chloroplastic galactosyl diacylglycerol molecular species in *Brassica napus* leaves. *Plant Physiol.* 1989; 91:924-9; PMID:16667158; <http://dx.doi.org/10.1104/pp.91.3.924>
- Pearcy RW. Effects of growth temperature on the fatty acid composition of the leaf lipids in *Atriplex lentiformis* (Torr.) Wats. *Plant Physiol.* 1978; 61:484-86; PMID:16660321; <http://dx.doi.org/10.1104/pp.61.4.484>
- Shen W, Li JQ, Dauk M, Huang Y, Periappuram C, Wei Y and Zou J. Metabolic and transcriptional responses of glycerolipid pathways to a perturbation of glycerol-3-phosphate metabolism in *Arabidopsis*. *J. Biol. Chem.* 2010; 285:22957-65; PMID:20304913; <http://dx.doi.org/10.1074/jbc.M109.097758>
- Szymanski J, Brotman Y, Willmiter L, Cuadros-Inostroza A. Linking gene expression and membrane lipid composition of *Arabidopsis*. *Plant Cell* 2014; 26:915-

- 928; PMID:24642935; <http://dx.doi.org/10.1105/tpc.113.118919>
18. Kunst L, Browse J and Somerville CR. Altered regulation of lipid biosynthesis in a mutant of *Arabidopsis* deficient in chloroplast glycerol-3-phosphate acyltransferase activity. *Proc. Natl. Acad. Sci. USA* 1988; 85:4143-7; <http://dx.doi.org/10.1073/pnas.85.12.4143>
19. Kunst L, Browse J and Somerville C. Enhanced thermal tolerance in a mutant of *Arabidopsis* deficient in palmitic Acid unsaturation. *Plant Physiol.* 1989; 91:401-8; PMID:16667033; <http://dx.doi.org/10.1104/pp.91.1.401>
20. Heilmann I, Mekhedov S, King B, Browse J and Shinklin J. Identification of the *Arabidopsis* palmitoylmonogalactosyldiacylglycerol delta7-desaturase gene FAD5, and effects of plastidial retargeting of *Arabidopsis* desaturases on the *fad5* mutant phenotype. *Plant Physiol.* 2004; 136:4237-45; PMID:15579662; <http://dx.doi.org/10.1104/pp.104.052951>
21. Hugly S, Kunst L, Browse J and Somerville C. Enhanced thermal tolerance of photosynthesis and altered chloroplast ultrastructure in a mutant of *Arabidopsis* deficient in lipid desaturation. *Plant Physiol.* 1989; 90:1134-42; PMID:16666863; <http://dx.doi.org/10.1104/pp.90.3.1134>
22. Hugly S and Somerville C. A role for membrane lipid polyunsaturation in chloroplast biogenesis at low temperature. *Plant Physiol.* 1992; 99:197-202; PMID:16668849; <http://dx.doi.org/10.1104/pp.99.1.197>
23. Miquel M, James D Jr, Dooner H and Browse J. *Arabidopsis* requires polyunsaturated lipids for low temperature survival. *Proc. Natl. Acad. Sci. USA* 1993; 90:6208-12
24. Miquel M, and Browse J. *Arabidopsis* mutants deficient in polyunsaturated fatty acid synthesis. Biochemical and genetic characterization of a plant oleoyl-phosphatidylcholine desaturase. *J. Biol. Chem.* 1992; 267:1502-9; PMID:1730697
25. Beisson F, Koo AJK, Ruuska S, Schwender J, Pollard M, Thelen JJ, Paddock T, Salas JJ, Savage L, Milcamps A, Mhaske VB, Cho YH, and Ohlrogge JB. *Arabidopsis* Genes Involved in Acyl Lipid Metabolism. A 2003 Census of the Candidates, a Study of the Distribution of Expressed Sequence Tags in Organs, and a Web-Based Database. *Plant Physiol.* 2003; 132:681-697; PMID:12805597; <http://dx.doi.org/10.1104/pp.103.022988>
26. Fan J, Yan C, Zhang X and Xu C. Dual role for phospholipid:diacylglycerol acyltransferase: enhancing fatty acid synthesis and diverting fatty acids from membrane lipids to triacylglycerol in *Arabidopsis* leaves. *Plant Cell* 2013; 25:3506-18; PMID:24076979; <http://dx.doi.org/10.1105/tpc.113.117358>
27. Winichayakul S, Scott RW, Roldan M, Hatier JH, Livingston S, Cookson R, Curran AC and Roberts NJ. *In vivo* packaging of triacylglycerols enhances *Arabidopsis* leaf biomass and energy density. *Plant Physiol.* 2013; 162:626-39; PMID:23616604; <http://dx.doi.org/10.1104/pp.113.216820>
28. Moellering ER, Muthan B and Benning C. Freezing tolerance in plants requires lipid remodeling at the outer chloroplast membrane. *Science* 2010; 330:226-228; PMID:20798281; <http://dx.doi.org/10.1126/science.1191803>
29. Smith MA, Dauk M, Ramadan H, Yang H, Seamons LE, Haslam RP, Beaudoin F, Ramirez-Erosa I, Forseille L. Involvement of *Arabidopsis* ACYL-COENZYME A DESATURASE-LIKE2 (At2g31360) in the biosynthesis of the very-long-chain monounsaturated fatty acid components of membrane lipids. *Plant Physiol.* 2013 161:81-96; PMID:23175755; <http://dx.doi.org/10.1104/pp.112.202325>
30. Chen M, Thelen JJ. ACYL-LIPID DESATURASE2 is required for chilling and freezing tolerance in *Arabidopsis*. *Plant Cell.* 2013 25:1430-44; PMID:23585650; <http://dx.doi.org/10.1105/tpc.113.111179>
31. Chen M, Markham JE, Cahoon EB. Sphingolipid Δ8 unsaturation is important for glucosylceramide biosynthesis and low-temperature performance in *Arabidopsis*. *Plant J.* 2012 69:769-81; PMID:22023480; <http://dx.doi.org/10.1111/j.1365-3113X.2011.04829.x>