Firm foundation for senior house officers

Five years is long enough

EDITOR—The editorial by Gallen and Peile signals anything but a firm foundation for training junior doctors.¹ The first foundation year will essentially be an extension of undergraduate training, its doctors allowed only to observe medical practice. And doctors in foundation year 2 will be “career tourists,” spending too little time in each specialty to be able to offer much service in return.

Medical educationalists should realise that nurses and other paramedical groups are encroaching on traditional medical practice and do not seem to need the sort of extended training we demand of medical students. For example, a six week course apparently enables paramedics to administer safely almost all drugs used in emergency practice. By contrast we seem intent on postponing the time at which our bright and enthusiastic young doctors can start treating patients.

Five years is quite long enough to learn the theory of medicine. After that, the sooner young doctors are allowed to start treating patients, in a supervised setting, the better.

Alan Leaman *consultant in emergency medicine*
Princess Royal Hospital, Telford TF1 6TF
caleaman@doctors.org.uk

Competing interests: None declared.

1 Gallen D, Peile E. A firm foundation for senior house officers. *BMJ* 2004;328:1390-1. (12 June.)

Junior doctors should become progressively more enabled

EDITOR—Gallen and Peile discuss establishing a firm foundation for senior house officers.¹ Concerns about trends in junior doctor training are growing in acute medical specialties. The proposed reforms may exacerbate a deteriorating situation.

In the past decade the competencies of junior grade doctors have collapsed. Previously junior house officers would rapidly acquire emergency medicine skills by being supervised participants in on-call ward cover and acute takes, consolidating these skills and acquiring considerable responsibilities as senior house officers.

Thus on-site medical competency has become extremely thin on the ground out of hours and in emergencies. Worse, prospective registrars are ill equipped to deal with emergencies owing to lack of exposure as senior house officers. Intensive care out-

reach, increased tertiary or subspecialty referrals, and the appointment of acute medicine consultants may absorb some of this experience deficit, but it still represents a worrying threat to patients' safety in the face of increasing workloads.

The foundation year is likely to become a two year junior house officer year. This poverty of expectation is creating a cadre of underused, frustrated, demoralised junior doctors and inexperienced middle graders. Junior doctors should be becoming progressively more competent and enabled, not the reverse.

Chris M Laing *specialist registrar nephrology and general and internal medicine*
Hammersmith Hospital, London W12 0HS
claing@mac.com

Competing interests: None declared.

1 Gallen D, Peile E. A firm foundation for senior house officers. *BMJ* 2004;328:1390-1. (12 June.)

Quality assurance programme needs to be in place

EDITOR—The development of a medical workforce with a wider range of general medical competencies is a highly desirable target. However, proposing a shorter specialist training period raises the fear of inappropriate development of specialist competencies at the end of the “run through” period. Gallen and Peile's article about incomplete work on the development of tasks, such as curriculum development, competency framework, and assessment methods, highlights another worrying development.¹

Learning opportunities for work based learning depend on the way in which work is organised and allocated, and that in turn depends on prevailing assumptions about the competence of the people concerned.² To establish a competency based training system, core competencies of educational supervisors also need to be established and supported.

The end point of improved patient care can be realised only if competent supervisors and a robust quality assurance system support the foundation programme.

Rafey A Faruqui *specialist registrar*
Charing Cross Higher Training Scheme, West London Mental Health NHS Trust, Uxbridge, Middlesex UB1 3EU
rafey.faruqui@imperial.ac.uk

Alia Faruqui *student in epidemiology (principles and practice)*
External Programme, University of London, London WC1E 7HU

Competing interests: AF is hoping to find a placement in the foundation training programme.

1 Gallen D, Peile E. A firm foundation for senior house officers. *BMJ* 2004;328:1390-1. (12 June.)

2 Eraut M. Concepts of competence and their limitations. In: *Developing professional knowledge and competence*. London: Falmer, 1994:163-81.

A firm foundation for general practice

EDITOR—Gallen and Peile discuss the initiative to modernise medical careers, a concept that could improve the recruitment and retention crisis in primary care by exposing young doctors to general practice.¹ This component may, however, flounder owing to a lack of support from deaneries, primary care trusts, and workforce development confederations.

The development of educational supervisors' and associate trainers' roles needs to be encouraged so that doctors in foundation year 2 can have an appropriate and well supported experience in primary care that can enable attainment and proper assessment of the foundation competencies.

Although training and non-training practices are enthusiastic, workforce development confederations and deaneries are not committed to fund training programmes for educational supervisors and associate trainers properly. For many practices, space and accommodation for doctors in foundation year 2 are also issues that need to be dealt with urgently.

Primary care trusts may not yet fully understand the importance of enthusing doctors in foundation year 2 to promote future recruitment to general practice. Unless these issues are tackled properly, the general practice component of modernising medical careers is unlikely to become established.

Robin Christie *general practitioner*
Portmill Surgery, Hitchin, Hertfordshire SG4 9TH
robanmel@ntlworld.com

Competing interests: None declared.

1 Gallen D, Peile E. A firm foundation for senior house officers. *BMJ* 2004;328:1390-1. (12 June.)

Bone scanning in lung cancer

Evidence is not sufficient to justify routine bone scanning

EDITOR—Hetzl et al claim that patients with lung cancer may be undergoing “futile” surgery owing to incomplete preoperative staging.¹ Their high incidence of bone

metastases may be explained by the unusually high proportion of small cell lung cancer (30%). For potentially resectable early stage tumours, metastases are unlikely without clinical signs.² Were the bone metastases in the field scanned by staging computed tomography and, if so, were they detected? Without TNM (tumour, node, and metastases) staging for the study population the authors cannot claim that a positive bone scan would have altered clinical management.

Positron emission tomography may be preferable to isotope bone scanning if concern exists about occult skeletal metastases in patients with potentially operable lung cancer. Studies have shown that positron emission tomography improves the detection rate of occult distant metastases, including bony lesions, compared with standard methods of staging³ and is cost effective in preventing futile operations.⁴

The gold standard used by Hetzel et al is magnetic resonance imaging of the vertebral column and patients' subsequent clinical course. However, they do not mention the length of clinical follow up. A previous pilot study showed that magnetic resonance imaging may have a role in detecting occult metastases in patients with potentially resectable cancer, but with a false positive rate.⁵ Hetzel et al do not comment on the accuracy of magnetic resonance imaging or what happened to cases with discrepant results on magnetic resonance imaging and bone scanning.

This paper does not provide sufficient evidence to recommend the routine use of bone scans in the staging of lung cancer.

Rachel E Benamore *specialist registrar radiology*
rachelbenamore@doctors.org.uk

James J Entwisle *consultant radiologist*
Department of Radiology, Glenfield Hospital,
University Hospitals of Leicester NHS Trust,
Leicester LE3 9QP

Mick D Peake *consultant chest physician*
Department of Respiratory Medicine, Glenfield
Hospital

Competing interests: None declared.

1 Hetzel M, Hetzel J, Arslanemir C, Nussle K, Schirrmeister H. Reliability of symptoms to determine use of bone scans to identify bone metastases in lung cancer: prospective study. *BMJ* 2004;328:1051-2. (1 May.)

2 Tanaka K, Kubota K, Kodama T, Nagai K, Nishiwaki Y. Extrathoracic staging is not necessary for non-small-cell lung cancer with clinical stage T1-2N0. *Ann Thoracic Surg* 1999;68:1039-42

3 Pieterman RM, van Putten JW, Meuzelaar JJ, Mooyaart EL, Vaalburg W, Koeter GH, et al. Preoperative staging of non-small-cell lung cancer with positron-emission tomography. *New Engl J Med* 2000;343:254-262

4 Verboom P, Van Tinteren H, Hoekstra OS, Smit EF, Van Den Bergh JH, Schreurs AJ, et al. Cost-effectiveness of FDG-PET in staging non-small cell lung cancer: the PLUS study. *Eur J Nucl Med Mol Imaging* 2003;30:1444-9.

5 Earnest F 4th, Ryu JH, Miller GM, Luetmer PH, Forstrom LA, Burnett OL, et al. Suspected non-small cell lung cancer: incidence of occult brain and skeletal metastases and effectiveness of imaging for detection—pilot study. *Radiology* 1999;211:137-45.

Pretest probability is of value

EDITOR—Hetzel et al discuss the reliability of symptoms in indicating the use of bone scanning in patients with lung cancer.¹ They reported a low sensitivity of 53% in detecting bone metastases when bone scanning is restricted to patients with skeletal complaints.

This value seems unacceptably low and is in contrast with the guidelines of leading professional societies that recommend bone scanning only in patients with clinical symptoms indicating bone involvement. However, Hetzel et al do not state the stage of disease at the time of scanning, which is crucial to understand the value of their results. Prevalence (or pretest probability) of bone metastases and reliability of symptoms in driving use of bone scanning is expected to vary according to disease stage.²

Fabio Puglisi *medical oncologist*
University of Udine, I-33100

Udine, Italy
fabio.puglisi@med.uniud.it

Competing interests: None declared.

1 Hetzel M, Hetzel J, Arslanemir C, Nussle K, Schirrmeister H. Reliability of symptoms to determine use of bone scans to identify bone metastases in lung cancer: prospective study. *BMJ* 2004;328:1051-2. (1 May.)

2 Jaeschke R, Guyatt GH, Sackett DL. Users' guides to the medical literature. III. How to use an article about a diagnostic test. B. What are the results and will they help me in caring for my patients? The Evidence-Based Medicine Working Group. *JAMA* 1994;271:703-7.

Authors' reply

EDITOR—Benamore et al assume that our incidence of bone metastases was high because of the unusually high proportion of patients with small cell lung cancer. The incidence of distant metastases is 30-50% at initial presentation of non-small cell lung cancer.¹ The incidence of bone metastases was highest in the small cell variety at initial presentation but was identical at necropsy in the four main types.² Many bone metastases are therefore likely to be missed at initial diagnosis of non-small cell lung cancer.

We used an extensive imaging algorithm to exclude bone metastases. In contrast to earlier clinical data and in agreement with results at necropsy,² we found an almost identical incidence of bone metastases of 32.4% in non-small cell lung cancer and 33.3% in small cell lung cancer (table).

We assessed the reliability of symptoms in detecting bone metastases and hence did not evaluate the number of metastases incidentally detected on computed tomography. To exclude small metastases Benamore et al suggest using fluorodesoxyglucose positron emission tomography. This method, however, is currently not recommended by German and European professional societies for routine use in lung cancer.

Stage dependent (according to computed tomography criteria) distribution of patients with skeletal metastases. Values are numbers of patients (numbers of patients with skeletal metastases)

	N0	N1	N2	N3
Non-small cell lung cancer				
T1	5(1)*	3(1)*	1(0)	0(0)
T2	15(5)*	7(3)*	10(4)	4(2)
T3	4(0)*	1(1)	3(0)	5(1)
T4	0(0)	0(0)	10(4)	16(6)
Small cell lung cancer				
T1	1(1)*	1 (0)	0(0)	1(0)
T2	2(0)*	0 (0)	4(1)	3(1)
T3	1(0)	0 (0)	2(0)	8(4)
T4	0(0)	1 (0)	3(1)	10(4)

*Tumour node stages amenable to primary surgery.

We used magnetic resonance imaging to exclude bone metastases probably missed on bone scanning or computed tomography. We used data about the clinical course or at necropsy to decide finally whether a patient had bone metastases.

All patients in our series were questioned and examined by the doctors before bone scanning and the reference methods. This guaranteed that they were blinded to the stage of disease (table).

The study by Tanaka et al quoted by Benamore et al was a retrospective analysis and included patients between 1982 and 1996 but did not report whether the localisation of pain was identical with the site of bone metastases. Chronic back pain is reported by 80% of people aged 50-80,³ so the pretest probability for bone pain is generally high.

Conventional radiography was the standard technique used to confirm focal lesions identified in bone scans in early studies. Skeletal scintigraphy detects metastases several months before they become visible in conventional radiographs. Metastases not confirmed by conventional radiography were therefore regarded as false positive findings, giving a presumed low specificity of bone scintigraphy. Furthermore, only very large, potentially symptomatic metastases were considered true positives.

Martin Hetzel *consultant physician*
martin.hetzel@medizin.uni-ulm.de

Coskun Arslanemir *physician*
Department of Internal Medicine II, University of
Ulm, D-89081 Ulm, Germany

Holger Schirrmeister *consultant in nuclear medicine*
Department of Nuclear Medicine, University of
Ulm

Competing interests: None declared.

1 Vaporciyan AA, Nesbitt JC, Lee JS. Cancer of the lung. In: Holland JF, Frei E, eds. *Cancer medicine*. 5th ed. London: BC Decker, Hamilton, 2000:1297-92.

2 Muggia FM, Chervu LR. Lung cancer: diagnosis in metastatic sites. *Semin Oncol* 1974;1:217-28.

3 Deyo RA, Tsui-Wu Y-J. Descriptive epidemiology of low-back pain and its related medical care in the United States. *Spine* 1987;12:264-8.

The full version of this reply is available on
bmj.com

Low back pain

Sacroiliac joint pain may be myth

EDITOR—I disagree with only one bit of Speed's review of low back pain¹: except in inflammatory disease or infection, I am not certain that the sacroiliac joint is responsible for much of what is called sacroiliac pain. The rigidity of the joint is such that mechanical disruption is almost impossible: the pelvis will fracture first. Cures of pain reported by injection of the joint are, I think, not because the joint has been injected but because the injection has gone into part of the gluteal muscle origin.

Many years ago a colleague tried to assess the accuracy of joint injection. Injecting corpses, with Indian ink, using classical localisation and x ray screening, he dissected the region and found no evidence of ink in the joint on any of eight attempts. The ink was all in the muscle, or in some cases had tracked into the vertebral venous plexuses. (I imagine such an experiment would nowadays be impossible to arrange.)

Although the finding of a "stress reaction" on radiography (sclerosis without erosion on the ilial side only, without erosion) might represent some sort of mechanical instability that could cause pain, I suspect that most cases of "sacroiliac strain" are really injuries of the gluteal origin—and injections work because that, and not the joint, is what has been injected. Unilateral pain spreading across the iliac crest, worse when the patient bends to the opposite side, might also, in theory, arise from the iliolumbar ligament rather than the disc.

Andrew N Bamji consultant in rheumatology and rehabilitation

Queen Mary's Hospital, Sidcup, Kent DA14 6LT
andrewbamji@lineone.net

Competing interests: None declared.

¹ Speed C. ABC of rheumatology. Low back pain. *BMJ* 2004;328:1119-1121. (8 May)

Author's reply

EDITOR—Bamji's views reflect the ongoing active discussion on the contribution of the sacroiliac joint to pain in this region. Although, as he points out, the sacroiliac joint has extreme mechanical strength, it exists within the pelvic ring; instability at the symphysis (common in women and active men) may result in pain further along the kinetic chain—including the innervated and partly synovial sacroiliac joint. Sacroiliac pain may arise through dysfunction through minute laxity or instability, or pain due to ligamentous irritation without any instability.

Any further comments are based on differences in clinical perspective in the face of a vacuum of evidence. It is a difficult joint to assess. The history, including potential mechanisms for injury (such as hurdling, or football), and examination, in particular to exclude other sources of pain, are crucial. Approaches to examination of the joint

have been proposed but remain unvalidated. Pain truly arising from the sacroiliac joint should, in my view, be considered as one of the differential diagnoses of pain in the pelvic region, particularly in young and active patients. Nevertheless, "sacroiliac pain" is an overdiagnosed condition, and it should be diagnosed only in the absence of any other cause for pain, although conditions can coexist.

Cathy Speed honorary consultant in rheumatology and sports medicine

Addenbrooke's Hospital, Cambridge CB2 2QQ
Cas50@medschl.cam.ac.uk

Competing interests: None declared.

Monitoring procalcitonin is of value in acute pancreatitis

EDITOR—In their discussion of the important issue of antibiotic prophylaxis in acute pancreatitis, I was surprised that O'Reilly and Kingsnorth made no mention of monitoring the plasma concentration of procalcitonin.¹ A considerable literature now exists, establishing the value of procalcitonin monitoring in acute pancreatitis,²⁻⁴ and it is our practice to use the APACHE II score and the value of this inflammatory marker to guide the requirement for early antibiotic prophylaxis with a carbapenem and fluconazole.

A recent study has shown that the plasma concentration of procalcitonin may reflect the derangement in gut barrier function (rather than the extent of systemic inflammation) and may hence predict those patients in whom the translocation of bacteria and fungi into dead pancreas is more likely.⁵

I agree with the authors that we do not want to give all patients with acute pancreatitis antibiotic prophylaxis. But I have experience of fatal acute pancreatitis from untreated Gram negative sepsis because of too "purist" an approach. The issue in my mind is how to identify the patients at risk early on and then include both bacterial and fungal prophylaxis.

David Bihari associate professor

University of New South Wales Lismore Base Hospital, Lismore, NSW 2480, Australia
biharid@aol.com

Competing interests: None declared.

¹ O'Reilly DA, Kingsnorth AN. Management of acute pancreatitis. *BMJ* 2004;328:968-9. (24 April)

² Rau B, Steinbach G, Gansauge F, Mayer JM, Grunert A, Beger HG. The potential role of procalcitonin and interleukin 8 in the prediction of infected necrosis in acute pancreatitis. *Gut* 1997;41:832-40

³ Kylanpaa-Back M-L, Takala A, Kemppainen EA, Puolakkainen PA, Leppaniemi AK, Karonen S-L, et al. Procalcitonin, soluble interleukin-2 receptor and soluble E-selectin in predicting the severity of acute pancreatitis. *Crit Care Med* 2001;29:63-9.

⁴ Ammori BJ, Becker KL, Kite P, Snider RH, Nylen ES, White JC, Larvin M, et al. Calcitonin precursors in the prediction of severity of acute pancreatitis on the day of admission. *Br J Surg* 2003;90:197-204.

⁵ Ammori BJ, Becker KL, Kite P, Snider RH, Nylen ES, White JC, et al. Calcitonin precursors: early markers of gut barrier dysfunction in patients with acute pancreatitis. *Pancreas* 2003;27:239-43.

Responsibility for ancillary care in clinical trials

Research patrimony and unintended coercion are hazards

EDITOR—I agree with many of the assessments and suggestions made by Belsky and Richardson on medical researchers' responsibilities for ancillary care,¹ but important issues remain to be considered.

The need to explore ancillary care is just as important in developed countries as it is in developing countries. British study participants, for example, are also sometimes found by the research team to have previously unknown medical conditions. More often they develop a good rapport with the research team, which can then be led to assume the role of a general practice with a long term research participant. Although this is understandable and shows the research team's patient centredness, it can cause problems.

The research doctor may be expected to see and diagnose every medical concern of the participant. But researchers with a specialist interest are really not trained to screen and treat many of the day to day ailments of these patients. They might alter the study subjects' expectations without being able to provide real care. This can interfere with subjects' use of normal medical care and be harmful to their interests.

The second problem is that acceptance of ancillary care responsibilities for research subjects from developing countries might coerce them into participating in the study and induce a strong element of volunteer bias. The quid pro quo arrangement of offering medical care in exchange for study participation would be no different to the payment of research subjects (in fact, it could mean greater financial incentives for poorer subjects), and has ethical implications that are not discussed in this paper.

Arvindan Veiraiah research registrar in cardiology
Royal Sussex County Hospital, Brighton BN2 1ES
drveiraiah@hotmail.com

Competing interests: None declared.

¹ Belsky L, Richardson HS. Medical researchers' ancillary clinical care responsibilities. *BMJ* 2004;328:1494-6. (19 June)

Finding out what participants think may be way forward

EDITOR—Belsky and Richardson highlight an important issue of ancillary care in the conduct of research trials and propose a potentially useful framework for dealing with the ethical problems raised.¹ What seems to be missing, however, is an exploration of the issue from the participants' perspective. It may be clear to the researcher that entrustment is partial. Participants may, however, believe (rightly or wrongly) that by

agreeing to participate, they are fully entrusting the researcher with responsibility for their wellbeing.

A need may therefore arise to delineate the limits of entrustment clearly in the process of obtaining consent. Whether this is practical or feasible is uncertain, but it would be useful and interesting to know what participants think they are entrusting to researchers when they agree to participate in a trial.

Ike Anya *specialist registrar in public health medicine*
Bristol North Primary Care Trust, Bristol BS2 8EE
ikechuku.anya@lshim.ac.uk

Competing interests: None declared.

1 Belsky L, Richardson HS. Medical researchers' ancillary clinical care responsibilities. *BMJ* 2004;328:1494-6. (19 June.)

A time to live or a time to die?

Are we losing our humanity?

EDITOR—My sympathy, congratulations, and deep respect to the unnamed author (and her husband) of “A time to live or a time to die?” for having the courage to present her harrowing account of being kept alive by percutaneous endoscopic gastrostomy.¹ Too often we hear of such tragedies and of well meaning but fundamentally misguided examples of medical intervention.

I believe the most challenging aspect of all forms of medical practice is the need for, and the exercise of, judgment. These days, in many walks of life, “judgment” carries connotations of authority and control which are often viewed as being unacceptable.

But judgment is really about making a decision. While, globally, the move within medicine is increasingly towards the objective goal of “evidence based medicine,” this philosophy must inevitably struggle with the conflicts of subjectivity. And there is little that is more subjective than the morality of what is right and what is wrong in such a situation as that presented by your correspondent.

It's easy to be objective. It's easy to base one's decisions on evidential data and, of course, such decisions do not bear the same degree of individual responsibility for their outcomes which more subjective decisions might entail.

Having qualified in 1975 I have seen numerous such cases and, regrettably, more so in recent years. Clinicians have always faced such ethical and moral dilemmas, but I fear that our increasing commitment to “evidence” may be undermining our need to exercise compassion. In our drive for objectivity, are doctors losing that compassion? Has the intellectual status quo of

medicine shifted so much that subjective considerations can be discounted? Or, worse, can it be an excuse not to exercise compassion?

Perhaps as we continue down the path of increasingly scientific objectivity, it may be time for us to remember the humanities as well as the science.

Stephen R Taylor *general practitioner*
Auckland, New Zealand 1311
steve@dcox.co.nz

Competing interests: None declared.

1 A time to live or a time to die? *BMJ* 2004;328:1445. (12 June.)

Let those who have ears hear

EDITOR—Perhaps one reason for the limited response to “A time to live or a time to die?” is that it is so powerful, and perhaps so close to home, that we feel some of the impotence that the writer feels.¹ I wonder if it was her choice to remain anonymous and what that may say about her trust towards the medical profession. Or perhaps she was simply trying to give her husband the dignity of anonymity at a time when so much of his dignity has been stripped away.

As someone who teaches medical ethics

I recognised too many areas of contention—the nature of capacity and consent; the relationships between doctor, patient, and relative; feeding through a percutaneous endoscopic gastrostomy (PEG) tube; professional boundaries; living wills; quality of life issues, etc. However, the millions of words written in guidelines and books (including the BMA's own *Medical Ethics Today—Handbook of Ethics and Law*, 2004) only take us so far—they do not make decisions for us, simply equip us with some of the tools.

Sadly there is no algorithm for these dilemmas, but what seems to stand out in this case is that the family do not believe that their views were heard or considered at the critical decision making moments. Their views may ultimately have been superseded but they may have at least felt that they were being heard.

Barry A Clark *hospital chaplain*
Selly Oak Hospital, Birmingham B29 6JD
barry.clark@uhb.nhs.uk

Competing interests: None declared.

1 A time to live or a time to die? *BMJ* 2004;328:1445. (12 June.)

Summary of responses

EDITOR—The personal view entitled “A time to live or a time to die?”—a reader's tale of how her dying husband has been kept alive by percutaneous endoscopic gastrostomy—sparked a small but impassioned debate on

what one correspondent highlights as the right of patients and their families to influence medical treatment.¹

Donald Couper, a general practitioner in Manchester, is surprised at the lack of commentary on a theme that must be familiar to doctors working in all types of clinical practice but thinks that fear of legal and disciplinary proceedings may have a bearing on clinical decisions of this kind. Another general practitioner, J Hopkins from Newton Aycliffe, is also certain that the restraint in the case is not medical but legal and advises mature debate about the ethical and legal issues to balance the rights of families against the legal and regulatory framework within which NHS clinical teams are expected to work.

Some correspondents suggest how to improve matters. Herbert Nehrllich, a private practitioner in Australia, for whom being in hospital means to leave one's identity and dignity at reception, only half humorously recommends assigning a “companion” to each hospital inpatient “to reassure, interpret, make conversation, and in general treat you like royalty.” Clive Barker, a general practitioner in Chorley, mentions advance directives as a possible way forward.

The dilemma clearly has no easy solution. One step in the right direction could come from clear and honest communication, say several correspondents. Nigin Bagla and Ritika Mangal from India use personal experience to illustrate what a doctor—in this case a family member—is faced with when treating a patient who is at death's door. And Chris Hudson, consultant geriatrician in Swansea, reminds us that there are two sides to every story and that the problem must have been a failure in communication between family and health-care team.

The consensus is with Naseem Qureshi from Saudia Arabia, that life and death should take their destined journeys. But the matter is obviously not simple. Doctors should never forget that they are human first, then doctors, surgeons, and specialists in sequence—argues Erik Walbeehm, specialist registrar in the Netherlands—and everything else should follow on from that.

Birte Twisselmann *technical editor*
BMJ

1 Electronic responses. A time to live or a time to die? *bmj.com*2004. <http://bmj.bmjournals.com/cgi/eletters/328/7453/1445> (accessed 15 July).

Ecclesiastes: A time to be born, and a time to die

bmj.com

bmj.com

Letters appearing here are an edited selection of rapid responses originally posted on bmj.com

We ask for all letters to the editor to be submitted as rapid responses via bmj.com

For advice see: bmj.com/rapidresponses