Published in final edited form as:

Org Lett. 2015 April 3; 17(7): 1624–1627. doi:10.1021/ol503765n.

5'- β,γ -CHF-ATP Diastereomers: Synthesis and Fluorine-Mediated Selective Binding by c-Src Protein Kinase

Candy S. Hwang[†], Alvin Kung^{†,‡}, Boris A. Kashemirov[†], Chao Zhang^{†,‡}, and Charles E. McKenna^{†,*}

[†]Department of Chemistry, University of Southern California, Los Angeles, California 90089, United States

[‡]Loker Hydrocarbon Research Institute, University of Southern California, Los Angeles, California 90089, United States

Abstract

The first preparation of the individual β,γ -CHF-ATP stereoisomers **12a** and **12b** is reported. Configurationally differing solely by the orientation of the C–F fluorine, **12a** and **12b** have discrete ³¹P (202 MHz, pH 10.9, $\delta_{\text{P}\alpha}$ 6 Hz, $\delta_{\text{P}\beta}$ 4 Hz) and ¹⁹F NMR (470 MHz, pH 9.8, δ_{F} 25 Hz) spectral signatures and exhibit a 6-fold difference in IC₅₀ values for c-Src kinase, attributed to a unique interaction of the (*S*)-fluorine of bound **12b** with R388 in the active site.

Graphical abstract

ATP is ubiquitous in biological systems where it plays a central role in energy storage, transduction, and utilization. Modification of the triphosphate moiety by replacing the oxygen between the β - and γ -phosphate with a methylene (CH_2)¹ creates nonhydrolyzable inhibitors for enzymes that cleave the γ -phosphate, such as ATPases and protein kinases.^{2,3} It was proposed many years ago that α -fluorination of the bridging methylene affords pCHFp analogues (CXY, X, Y = HF) that are more isoacidic with ATP than the

*Corresponding Author. ; Email: mckenna@usc.edu

ASSOCIATED CONTENT

Supporting Information

Detailed information on the synthesis of **4–12** and **14–17** including, ¹H, ¹⁹F, and ³¹P NMR and MS spectra and HPLC analytical and preparative traces. Experimental conditions and data for enzyme and characterization assays. Details of molecular docking studies. This material is available free of charge via the Internet at <http://pubs.acs.org>.

The authors declare no competing financial interest.

corresponding pCH₂p analogues.^{4,6} Besides adjusting the electronegativity of the bridging carbon to more closely resemble that of the bridging oxygen atom with minimal steric perturbation, the general absence of the fluorine atom from natural biological systems allows it to function as a bioorthogonal and readily detectable probe for ¹⁹F NMR studies. At the same time, replacement of the pOp oxygen by CHF introduces a new chiral center into the nucleotide, resulting in two diastereomeric forms of β,γ-CHF-ATP that offer the potential advantage of comparing biochemical properties of two enzyme inhibitors essentially identical in structure except with respect to the orientation of the C–F fluorine within a protein binding site. By switching a defined fluorine orientation in an otherwise similarly bound ATP analogue, a unique platform for investigating weak F bonding interactions^{7–11} with proteins is thus made available.

Beginning in the late 1990s, reports began to appear^{12–15} suggesting that the CHF stereochemistry of α-fluorinated monophosphonates can affect their binding to enzymes, and interest in this phenomenon has continued to grow.^{9,16–18} However, the preparation of the individual diastereomers of β,γ-CHF-ATP itself has remained an unmet challenge, preventing this unique set of enzyme probes from realizing their original potential. Standard approaches to the synthesis of β,γ-CXY ATPs such as conjugation of the appropriate methylenebis(phosphonic acid) salt with a 5'-activated AMP¹⁹ (Figure 1) result in a mixture of both diastereomers when X ≠ Y.

β,γ-CXY and α,β-CXY *desoxy*NTP analogues have recently demonstrated their utility as probes of the ground and transition states of DNA polymerase β (pol β).^{20–24} As part of these studies, the individual synthesis of all four β,γ-CHF and β,γ-CHCl dGTP diastereomers was achieved, with the absolute configurations assigned by X-ray crystallography of their ternary complex with DNA and pol β.²⁵ These analogues were found to exhibit significantly different *K_d* and *k_{pol}* values with the enzyme, modulated only by the position of the halogen atom substituent at the CXY carbon.²⁴ Furthermore, the ¹⁹F NMR spectra of these CHF diastereomers were found to be resolvable and useful to monitor their stereoselective turnover catalyzed by the enzyme.^{20,24} Early work on synthetic β,γ-CHF-ATP mixtures²⁶ and a recent paper on β,γ-CHF-UTPs¹⁸ reported inability to distinguish the ribosynucleotide CHF-stereoisomers by ¹⁹F NMR, although the former observation was not confirmed when the fluorinated ATP derivative was prepared by an alternate method.²⁷

These results prompted us to attempt the long-awaited synthesis of the individual β,γ-CHF-ATPs. Our strategy was based on the key chiral bisphosphonate synthons **9a** and **9b** (separable by preparative HPLC) prepared in eight steps from tetraisopropyl methylenebis(phosphonate) **1**;²⁵ however, the synthesis of **9a** and **9b** was modified at several steps, improving the yield significantly (Scheme 1). In the first modification, monodemethylation of **4** (previously prepared with NaI²⁵) with triethylamine^{28,29} under reflux raised the isolated yield of **5** to 90% with a substantial reduction in reaction time to 10–20 min.

Compound **5** was then transformed into the stereoisomer ester mixture **6a/6b**, purified on a silica gel column replacing the original preparative TLC procedure for convenient scalability.²⁵ In the second important modification, after conversion of **6a/6b** to the acids

8a/8b,²⁵ the latter were derivatized to the chiral dimorpholidates **9a/9b** using a nonaqueous solvent system, shortening the reaction time from several hours to <1 h after addition of DCC and also modestly increasing the yield from 87% to 96%, additionally aided by a more performant reversed-phase HPLC column.

Prior to separation and purification of the individual isomers, reaction mixture **9a/9b** taken up in D₂O showed discrete, but overlapping, multiplets in the ¹⁹F (470 or 564 MHz, pH 8.8–9.3, Figures S7C,D and S11, Supporting Information) and ³¹P NMR spectra (202 MHz, pH 8.8–8.9, Figures S9 and S13, Supporting Information). Interestingly, the same sample dissolved in CD₃OD gave a clean separation between the isomer multiplets in ¹⁹F NMR (470 or 564 MHz, Figures S7A,B and S10, Supporting Information), but the relative position of the two resonances was inverted, which was confirmed at higher spectrometer frequency (564 MHz) and by spiking with **9a** (Figure S10–11, Supporting Information).

The separated individual isomers in acid form (**10a, 10b**)²⁵ were coupled conventionally with AMP-morpholidate¹⁹ (AMP 5'-M, **14**) and then treated to remove the protecting groups as described previously²⁵ (Scheme 2). Gratifyingly, coupling of **14** to the individual synthons did not require additional protection of the 2',3'-hydroxyl groups of the ribose sugar. The (*R*)/(*S*)-CHF chirality of the separated and purified dimorpholidated chiral synthons (**9a** and **9b**) was assigned by HPLC and NMR analysis in comparison with the literature values.²⁵ ³¹P NMR spectral splitting patterns in D₂O (Figures S20 and S24, Supporting Information), where **9a** exhibits second order splitting, additionally distinguish the two isomers. (The original assignments of the absolute configurations of **9a** and **9b** were established retrosynthetically from the X-ray structure of (*R*)-β,γ-CHF-dGTP complexed to DNA pol.²⁵)

β,γ-Methylene (CH₂)-, mixed β,γ-CHF-, and β,γ-difluoromethylene (CF₂)-ATPs were synthesized by conjugating **14** with the appropriate bisphosphonate (Figure 1).^{1,6,19} A discrete ¹⁹F and ³¹P NMR spectrum for each diastereomer **12a/12b** in a synthetic mixture (**12**) was observed (Figures S44 and S45, Supporting Information), contrary to early observations²⁶ but consistent with prior observations for β,γ-CHF-dGTP^{20,22,25} and -UTP analogues.³⁰ The discrepancy may be attributed to the crucial role of the sample pH³¹ (10) and, possibly, the nature of the counteranion (e.g., NH₄⁺ gives broadened line widths).^{30,32} A 3:1 mixture of the separated diastereomers was made, and its ¹⁹F and ³¹P NMR spectra were measured to confirm (Figures S55 and S56, Supporting Information) that **12a** (the (*S*)-CHF-isomer) resonates more downfield in the ¹⁹F NMR and has P_α and P_β, more downfield and upfield, respectively, in the ³¹P NMR.²⁵ To further validate the assignment of each pairwise resonance²⁵ to a specific diastereomer, the ¹⁹F NMR spectra were simulated³³ and compared to that of the experimental mixture of isomers (Figure S57, Supporting Information).^{22,30} The ability to discern each diastereomer by ¹⁹F and ³¹P NMR provides a convenient method for determining selective substrate consumption by a selective enzyme.^{18,24}

In search of an enzyme interaction that might be stereospecific for **12a** vs **12b**, we examined a protein kinase. Protein kinases catalyze the transfer of a phosphoryl group from ATP to substrate proteins. As a major mode of post-translational modification, the phosphorylation

event catalyzed by protein kinases regulates fundamental cellular processes such as cell proliferation, differentiation and motility. Aberrant expression or activation of these enzymes can often lead to a variety of human diseases.^{34–38} For example, overexpression of Src tyrosine kinases is frequently found in cancerous tumors.^{3,34,39–41} Understanding the catalytic mechanism of Src and its role in oncogenesis is critical for design of therapeutic agents to target the diseases associated with its dysregulation.^{3,37,40}

We tested the β,γ -CXY-ATP analogues **12**, **15**, and **16**, (Figure 1) as inhibitors against the kinase domain of Src (c-Src, E.C. 2.7.10.2) and determined their IC₅₀ values using the radioactive disk assay.⁴² The data from the inhibition assays show that fluorination at the β,γ -position increases inhibitor potency (Table 1). β,γ -CH₂-ATP (**15**) was already shown to be a competitive inhibitor of c-Src kinase with respect to ATP and a noncompetitive inhibitor versus the peptide substrate.² The K_m for ATP and c-Src kinase is reported at around 80 μ M.⁴³

The Src kinase inhibition assay IC₅₀ data (Table 2) indicate a 6-fold selectivity for the (*S*)-CHF-ATP diastereomer **12a**. The relative potency of **16** appears to exclude a steric origin for the selectivity exhibited by the enzyme. We hypothesize that the specific orientation of the CHF-fluorine in **12a** makes possible a specific dipolar interaction, adding additional stabilization energy to the complex, which is reflected in the IC₅₀ difference. To examine this possibility, in silico molecular docking simulations using AutoDock Vina 1.1.2⁴⁴ were performed to explore potential protein–ligand interactions (Figure 2). Although our kinase assays were performed using chicken Src kinase, we chose a published X-ray crystal structure of human Src (2SRC) for molecular docking calculations due to the higher resolution available (1.8 Å vs 2.3 Å for chicken Src (3DQX)). The human and chicken Src proteins share 99% sequence identity (using BLASTP 2.2.30+⁴⁵) and are identical within the active site.⁴⁶

The calculations suggested interactions of both the (*S*)-fluorine of **16** and the γ -phosphate with the active site Arg388 (the latter observation is consistent with the crystal structure of c-Src kinase⁴⁷): the (*S*)-fluorine atom of **16** is located 3.0 Å from a guanidinium N of Arg388 (Figure 2A). Additionally, Lys295 stabilizes two phosphates, and Asp404 interacts with a structural water and Mg²⁺. In contrast, interactions of the (*R*)-fluorine with residues in the active site were not identified: the nearest group was the amido NH of Cys277 at 3.6 Å, too distant from the (*R*)-fluorine to have a significant interaction (Figure 2A). The (*R*)-fluorine was found to be 3.8 Å from the carbonyl carbon of Gly276 and 4.6 Å from the nearer N of Arg388.

Fluorine atoms proximal to Arg side chain nitrogens have been frequently documented in the literature,^{14,48} and it has been proposed that the guanidinium moiety of Arg is fluorophilic.^{49–51} We previously observed an apparent interaction of the CHF-fluorine of β,γ -CHF-dGTP with Arg183 (PDB entry 4DO9) in the nucleotide-binding site of pol β .^{20,22,25} The IC₅₀ ratio of 6:1 observed for **12b** vs **12a** corresponds to a bonding interaction on the order of 1.1 kcal/mol, which is consistent with literature values for weak noncovalent fluorine bonds.^{49,52}

In summary, for the first time, the individual diastereomers of β,γ -CHF-ATP have been obtained using an improved method for preparing the separable (*S*)-mandelic acid morpholinamide (*R*)/(*S*)-monoesters **9a** and **9b** as synthons. The two diastereomers have discrete ^{19}F and ^{31}P NMR spectra in D_2O and CD_3OD and show significantly different inhibition constants for c-Src protein kinase, associated with the presence or absence of a noncovalent bonding interaction of C–F with R388 in the active site.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was supported by funding from the Dana and David Dornsife College of the University of Southern California (C.E.M. and C.Z.) and NIH U19CA177547 (C.E.M.) and by a National Science Foundation Graduate Research Fellowship (to C.S.H, DGE-0937362).

REFERENCES

1. Myers TC, Nakamura K, Flesher JW. J. Am. Chem. Soc. 1963; 85:3292.
2. Boerner RJ, Barker SC, Knight WB. Biochemistry. 1995; 34:16419. [PubMed: 8845369]
3. Elphick LM, Lee SE, Gouverneur V, Mann DJ. ACS Chem. Biol. 2007; 2:299. [PubMed: 17518431]
4. McKenna CE, Shen PD. J. Org. Chem. 1981; 46:4573.
5. Leswara ND, Shen PD, McKenna CE. Fed. Proc. 1982; 41:860.
6. Blackburn GM, Kent DE, Kolkman F. J. Chem. Soc., Chem. Commun. 1981:1188.
7. Dalvit C, Vulpetti A. Chem Med Chem. 2011; 6:104. [PubMed: 21117131]
8. O'Hagan D. Chem. Soc. Rev. 2008; 37:308. [PubMed: 18197347]
9. Berkowitz DB, Karukurichi KR, de la Salud-Bea R, Nelson DL, McCune CD. J. Fluorine Chem. 2008; 129:731.
10. Romanenko VD, Kukhar VP. Chem. Rev. 2006; 106:3868–3935. [PubMed: 16967924]
11. Bégué J-P, Bonnet-Delpon D. J. Fluorine Chem. 2006; 127:992.
12. Nieschalk J, Batsanov AS, O'Hagan D, Howard J. Tetrahedron. 1996; 52:165.
13. Berkowitz DB, Bose M, Pfannenstiel TJ, Doukov T. J. Org. Chem. 2000; 65:4498. [PubMed: 10959850]
14. Berkowitz DB, Bose M. J. Fluorine Chem. 2001; 112:13.
15. Xu Y, Qian L, Prestwich GD. J. Org. Chem. 2003; 68:5320. [PubMed: 12816494]
16. Cox RJ, Gibson JS, Hadfield AT. Chem Bio Chem. 2005; 6:2255.
17. Forget SM, Bhattasali D, Hart VC, Cameron TS, Syvitski RT, Jakeman DL. Chem. Sci. 2012; 3:1866.
18. Beaton SA, Jiang PM, Melong JC, Loranger MW, Mohamady S, Veinot TI, Jakeman DL. Org. Biomol. Chem. 2013; 11:5473. [PubMed: 23857455]
19. Roseman S, Distler JJ, Moffatt JG, Khorana HG. J. Am. Chem. Soc. 1961; 83:659.
20. McKenna CE, Kashemirov BA, Upton TG, Batra VK, Goodman MF, Pedersen LC, Beard WA, Wilson SH. J. Am. Chem. Soc. 2007; 129:15412. [PubMed: 18031037]
21. Sucato CA, Upton TG, Kashemirov BA, Batra VK, Martinek V, Xiang Y, Beard WA, Pedersen LC, Wilson SH, McKenna CE, Florian J, Warshel A, Goodman MF. Biochemistry. 2007; 46:461. [PubMed: 17209556]
22. Batra VK, Pedersen LC, Beard WA, Wilson SH, Kashemirov BA, Upton TG, Goodman MF, McKenna CE. J. Am. Chem. Soc. 2010; 132:7617. [PubMed: 20465217]
23. Sucato CA, Upton TG, Kashemirov BA, Osuna J, Oertell K, Beard WA, Wilson SH, Florian J, Warshel A, McKenna CE, Goodman MF. Biochemistry. 2008; 47:870. [PubMed: 18161950]

24. Oertell K, Wu Y, Zakharova VM, Kashemirov BA, Shock DD, Beard WA, Wilson SH, McKenna CE, Goodman MF. *Biochemistry*. 2012; 51:8491. [PubMed: 23043620]
25. Wu Y, Zakharova VM, Kashemirov BA, Goodman MF, Batra VK, Wilson SH, McKenna CE. *J. Am. Chem. Soc.* 2012; 134:8734. [PubMed: 22397499]
26. Blackburn GM, Kent DE, Kolkman F. *J. Chem. Soc., Perkin Trans.* 1984; 1:1119.
27. McKenna CE, Harutunian V. *FASEB J.* 1988; 2:6148.
28. Vepsäläinen JJ, Kivikoski J, Ahlgrén M, Nupponen HE, Pohjala EK. *Tetrahedron*. 1995; 51:6805.
29. Ahlmark MJ, Vepsäläinen JJ. *Tetrahedron*. 1997; 53:16153.
30. Hwang CS, Kashemirov BA, McKenna CE. *J. Org. Chem.* 2014; 79:5315. [PubMed: 24819695]
31. The pH values for deuterated water solutions as reported here and by us previously (refs ²⁰, ²², ²⁵, ³⁰) are given as measured with a glass electrode, without correction to pD (ref ³²).
32. Kręgel A, Bal W. *J. Inorg. Biochem.* 2004; 98:161. [PubMed: 14659645]
33. Fluorine NMR were simulated using spin simulation on MestReNova 8.1.2.
34. Schwartz PA, Murray BW. *Bioorg. Chem.* 2011; 39:192. [PubMed: 21872901]
35. Kalesh KA, Liu K, Yao SQ. *Org. Biomol. Chem.* 2009; 7:5129. [PubMed: 20024108]
36. Xie J, Corneau AB, Seto CT. *Org. Lett.* 2003; 6:83. [PubMed: 14703356]
37. Moffat D, Nichols CJ, Riley DA, Simpkins NS. *Org. Biomol. Chem.* 2005; 3:2953. [PubMed: 16186927]
38. Zhang C, Lopez MS, Dar AC, Ladow E, Finkbeiner S, Yun CH, Eck MJ, Shokat KM. *ACS Chem. Biol.* 2013; 8:1931. [PubMed: 23841803]
39. Kwarciński FE, Fox CC, Steffey ME, Soellner MB. *ACS Chem. Biol.* 2012; 7:1910. [PubMed: 22928736]
40. Brandvold KR, Steffey ME, Fox CC, Soellner MB. *ACS Chem. Biol.* 2012; 7:1393. [PubMed: 22594480]
41. Ulrich SM, Kenski DM, Shokat KM. *Biochemistry*. 2003; 42:7915. [PubMed: 12834343]
42. Witt JJ, Roskoski R Jr. *Anal. Biochem.* 1975; 66:253. [PubMed: 1147218]
43. Knight ZA, Shokat KM. *Chem. Biol.* 2005; 12:621. [PubMed: 15975507]
44. Trott O, Olson AJ. *J. Comput. Chem.* 2010; 31:455. [PubMed: 19499576]
45. Altschul SF, Madden TL, Schäffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ. *Nucleic Acids Res.* 1997; 25:3389. [PubMed: 9254694]
46. Via visual inspection of the active site using PyMol, The PyMOL Molecular Graphics System, Version 1.2r3pre, Schrödinger, LLC.
47. Xu W, Harrison SC, Eck MJ. *Nature*. 1997; 385:595. [PubMed: 9024657]
48. Morgenthaler M, Aebi JD, Grüniger F, Mona D, Wagner B, Kansy M, Diederich F. *J. Fluorine Chem.* 2008; 129:852.
49. Zhou P, Zou J, Tian F, Shang Z. *J. Chem. Inf. Model.* 2009; 49:2344. [PubMed: 19788294]
50. Muller K, Faeh C, Diederich F. *Science*. 2007; 317:1881. [PubMed: 17901324]
51. Hagmann WK. *J. Med. Chem.* 2008; 51:4359. [PubMed: 18570365]
52. Dalvit C, Invernizzi C, Vulpetti A. *Chem.–Eur. J.* 2014; 20:11058. [PubMed: 25044441]
53. Accelrys Software, Inc. *Discovery Studio Modeling Environment*, Release 4.1. San Diego: Accelrys Software, Inc.; 2007.

Figure 1.
Standard synthesis of β,γ -CXY-ATPs.

Figure 2. Superimposition for (A) **16** and (B) **12a** and **12b** docked into the active site of human Src kinase using AutoDock Vina 1.1.2 and X-ray crystallographic data for the apoprotein (2SRC) including Mg (shown as green sphere). The (*S*)-fluorine in both compounds is oriented toward Arg388. The (*R*)-fluorine is oriented toward the pocket entrance, with no protein atoms closer than 3.8 Å. The figures were generated using Accelrys Discovery Studio.⁵³

Scheme 1.

Modified Synthesis of (S)-Mandelic Acid Morpholinamide (R)/(S)-Monoesters 9a and 9b

Scheme 2.

Synthesis of 12a and 12b from 9a and 9b, Respectively

Table 1Effect of Fluorination of β,γ -CH₂-ATP on Inhibition of c-Src Kinase

IC ₅₀	X, Y = H ₂ 15 (μ M)	X, Y = HF 12 (μ M)	X, Y = F ₂ 16 (μ M)
mean $\pm$ SE (<i>n</i>) ^a	145 $\pm$ 25 (3)	47 $\pm$ 7 (6)	46 $\pm$ 7 (3)

^aSE = standard error; *n* = number of replicates; variance test revealed no significant deviation prior to running an ANOVA. Differences (ANOVA, *F* (2, 6) = 12, *p* = 0.008 < 0.05) are statistically significant.

Table 2Inhibition of c-Src Kinase by β,γ -CHF-ATP Diastereomers

IC ₅₀	X, Y = H, F 12a (μ M)	X, Y = F, H 12b (μ M)	mixture (~1:1) 12 (μ M)
mean $\pm$ SE (<i>n</i>) ^a	23 $\pm$ 4 (6)	150 $\pm$ 8 (5)	47 $\pm$ 7 (6)

^aSE = standard error; *n* = number of replicates; variance test revealed no significant deviation prior to ANOVA. Differences (ANOVA, $F(2, 12) = 105$, $p < 0.001$) are statistically significant.