Unique Cyst in a 22-Year-Old Man

Alberto Ramos-Garibay, MD, Gisela Navarrete-Franco, MD, Sandra Gutiérrez-Zambrano, MD, and Mónica Olivia Rivera Ramírez, MD

FIGURE 1. Cyst with a normal epidermis surface (H&E $\times 4$).

(*Am J Dermatopathol* 2015;37:908–909)

CASE REPORT

A 22-year-old healthy man presented to our clinic with a 4-month history of a single, asymptomatic, 1-cm subcutaneous cystic nodule on the pubis.

Histopathologic examination showed a totally normal epidermis. The underlying dermis showed a cystic lesion (Fig. 1) containing many eosinophilic bodies (Fig. 2).

WHAT IS YOUR DIAGNOSIS?

(Continued on page 949)

From the Department of Dermatopathology, Centro Dermatológico Pascua, Buenos Aires, México City.

The authors declare no conflicts of interest.

Reprints: Alberto Ramos-Garibay, MD, Department of Dermatopathology, Centro Dermatológico Pascua, Av Dr. Vertiz 464, Col Buenos Aires 06780, México City, DF. México (e-mail: ramosgari4400@yahoo.com.mx).

Copyright © 2014 Wolters Kluwer Health, Inc. All rights reserved.

FIGURE 2. Many eosinophilic bodies in the wall and into the cyst (H&E).