Published in final edited form as:

Int J Eat Disord. 2016 June ; 49(6): 603–612. doi:10.1002/eat.22538.

Extremes of Eating Are Associated With Reduced Neural Taste Discrimination

Guido K.W. Frank, MD^{1,2,*}, Megan E. Shott, BS¹, Carrie Keffler, BS¹, and Marc-Andre Cornier, MD³

¹ Department of Psychiatry, School of Medicine, University of Colorado Anschutz Medical Campus

² Neuroscience Program, University of Colorado Anschutz Medical Campus

³ Division of Endocrinology, Metabolism, and Diabetes, School of Medicine, University of Colorado Anschutz Medical Campus

Abstract

Objective—Eating disorders are severe psychiatric disorders of unknown etiology.

Understanding how neuronal function affects food choices could help personalize treatment based on brain function. Here we wanted to determine whether disordered eating behavior is associated with alterations in the primary taste cortex's ability to classify taste stimuli, which could interfere with taste reward processing.

Method—One-hundred-and-six women, 27 healthy CW (age 26.15±6.95 years), 21 with restricting-type anorexia nervosa (AN; age 23.10±6.14 years), 19 recovered from restricting-type AN (recovered AN; age 26.95±5.31 years), 20 with bulimia nervosa (BN; age 25.15±5.31 years), and 19 with obesity (age 28.16±8.13 years), received sucrose, control solution or no taste stimulation during functional magnetic resonance brain imaging. Multivariate Bayesian pattern analysis (decoding) and cross-validation tested taste classification accuracy (adjusted for comorbidity, medication use, taste perception, interoception and brain activation volume).

Results—For sucrose versus control solution, classification accuracy differed ($F=2.53, p<0.041$). Post-hoc tests indicated higher classification accuracy in CW compared to women with AN ($p<0.016$) or obesity ($p<0.027$), and in recovered AN compared to AN ($p<0.016$) or obesity ($p<0.047$) groups. Taste stimulation resulted in sparse insula voxel activation across all groups.

Discussion—Reduced classification accuracy across stimuli in women with AN or obesity could indicate low brain encoding discrimination of stimulus quality, which could contribute to altered reward activation and eating drive that is not adjusted to nutritional needs. This deficit appears to normalize with recovery from AN, but adjusting food flavor intensity could aid in the treatment of individuals with AN or obesity.

*Correspondence to: Guido K.W. Frank, Departments of Psychiatry and Neuroscience, Developmental Brain Research Program, University of Colorado Denver, The Children's Hospital Colorado, Gary Pavilion A036/B-130, 13123 East 16th Avenue, Aurora, CO 80045. Guido.Frank@ucdenver.edu.

Keywords

Anorexia; Bulimia; Classification Accuracy; Obesity; Taste; Insula; Recovery; Decoding

Introduction

Eating disorders (EDs) are severe psychiatric disorders (1), with anorexia nervosa (AN) characterized by fear of weight gain and underweight, and bulimia nervosa (BN) by binge eating and purging but normal weight (2). Usually associated with more food intake than physiologically needed is obesity, defined by a body mass index (BMI, kg/m²) ≥ 30 . EDs and obesity are associated with increased mortality (3, 4), their underlying causes are considered multifactorial and treatment success is modest (4-8).

Basic science and human in vivo brain imaging research has suggested that food restriction and overeating are associated with alterations in reward circuit function.

Taste is an important driver of food intake (9) and invariably associated with distinct neuronal patterns in the insula, the brain's primary taste cortex (10). The insula connects to ventral striatum, orbitofrontal cortex and amygdala, higher order brain structures that control how much we eat (11, 12). Thus the insula has a “gate-keeper” function for taste information transmission and could have a central role in the pathophysiology of disordered eating (13). The insula is also important for interoceptive awareness (responding to body cues (14)), which tends to be altered in EDs (13).

Insula *size* is altered in EDs (15, 16), and *functional* alterations in the insula have also been reported repeatedly. For instance, in AN after recovery, insula, anterior cingulate and striatal activation were reduced during repetitive sucrose application (17), but insula and cingulate response were increased when applied randomly (18). In BN, sucrose and milkshake activated insula and frontal cortex less than in CW (19, 20). Imaging research in obesity most commonly implicated the insula (21), with increased or decreased activation depending on anticipation or receipt of visual or actual food stimuli (22, 23).

Functional brain imaging typically studies strength of activation (dependent variable) in response to specific stimuli (independent variable) and may inform on specific brain circuits (24, 25). The measured signal is the sum of many neurons coming together, although some neurons may contribute much and others little to the overall signal. In contrast, so called multivariate (multi-voxel) pattern analysis, or decoding, goes the opposite direction and uses brain activation patterns as the independent variable to classify task conditions (26-28). Multivariate pattern analysis is used to investigate how brain regions code stimuli-specific information as distinct patterns of neural activity. This then allows for differentiation of distinct perceptual states by assessing the characteristic distribution of voxel activation. Thus, decoding tries to identify a neuronal “fingerprint” to predict a condition or psychological state (24, 29). However, there may be many equally likely solutions of voxel activation patterns, and to overcome this problem, one can use constraints or priors in a multivariate Bayesian approach, by testing how voxel patterns are distributed within predefined models (26, 27). Cross-validation then can test the brain's “pattern classification

accuracy”, or how well a person's distinct activation pattern (associated with the stimulus or psychological state) can be generalized to the full data set, which can be compared across individuals and groups.

Various studies have applied decoding strategies to brain activation. For instance, decoding has been used in the study of vision, finding that distinct activation patterns could predict discrete objects (30). Decoding was also used to study attention states, memory generation, decision making bias and lie detection (31, 32), as well as interoception of pain, which could be relevant for ED research (13). To the best of our knowledge decoding has not been studied previously in EDs or obesity.

Here we tested the hypothesis that insula taste classification accuracy is reduced in individuals with disordered eating behavior. We sought to 1) determine the best model of pattern activation in this region across participants and 2) compare taste classification pattern accuracy between groups, while correcting for potential confounds such as comorbidity, medication use, brain volume, but also interoceptive or taste perception differences, factors that may fluctuate with anxiety and self-restraint (33, 34). We anticipated two possible scenarios. One, AN could be associated with higher and obesity with lower pattern classification accuracy with the idea that under- and overweight are associated with higher and lower response to food stimuli (23). Alternatively, having in mind that AN and obesity may respond stronger or weaker to *any* type of stimulus, both groups might distinguish taste stimuli poorly as they may code stimulus salience similarly high or low. We studied two additional groups, individuals recovered from AN to test whether such deficits would improve with recovery, and individuals with BN to strengthen the hypothesis that any alterations seen are more dependent on weight or other biologic factors than simply on ED cognitions.

Methods

Study Participants

We recruited 106 women, 27 healthy comparison women (CW), 21 women with restricting-type AN, 19 women recovered from restricting-type AN, 20 women with BN, and 19 women with obesity (Table 1.). The Colorado Multiple Institutional Review Board approved the study; all participants gave written informed consent. Participants received \$160 for the brain imaging session and completion of assessments.

AN (35) was defined as underweight below 85% of weight expected for age and height, severe fear of gaining weight, body image distortion, lack of menstrual cycle, but without binge eating/purging behavior. Individuals with BN had binge eating/purging episodes at least twice per week for at least three months and self-evaluation was unduly influenced by shape and weight. Obesity was defined by BMI ≥ 30 (36). Individuals recovered from AN had a history of restricting-type AN, normal weight, regular menses and normal exercise patterns for ≥ 1 year. Healthy CW had no history of psychiatric or major medical illness, were not taking medication, and were within normal BMI range life long.

Individuals with AN or BN were within their first 1-2 weeks of inpatient or partial hospitalization treatment, and had no electrolyte, blood count or other laboratory abnormalities. A doctoral level interviewer assessed psychiatric diagnostic status using the structured clinical interview (SCID (35)) for DSM-IV diagnoses. CW, women recovered from AN, women with BN or obesity were studied during the first ten days of the menstrual cycle to keep hormonal variation low (37).

Psychological Assessments

Study participants completed self-assessments for: 1) drive for thinness, bulimia, body dissatisfaction and interoceptive deficits (Eating Disorder Inventory-3) (38); 2) harm avoidance (Temperament and Character Inventory) (39); 3) state and trait anxiety (Spielberger State and Trait Anxiety Inventory) (40); 4) depression (Beck Depression Inventory) (41); and 5) reward and punishment sensitivity (Sensitivity to Reward and Punishment Questionnaire, revised) (42).

Brain Imaging Procedures

Prior to breakfast, participants rated randomly presented, unmarked taste stimuli (distilled water; five sucrose solution strengths, Mallinckrodt Chemicals, Phillipsburg, NJ: 2%, 4%, 8%, 16%, 1Molar; and artificial saliva: 25mM KCl, 2mM NaHCO₃) (43) for sweetness, 'absent' (1) to 'extreme' (9), and pleasantness, 'dislike extremely' (1) to 'like extremely' (9), on 9-point Likert scales. Between 7.00 and 8.00AM, AN and BN groups ate breakfast according to their meal plan, comparison, obese, and women recovered from AN had breakfast matched to the average ED program meal plan breakfast. Blood oxygen level dependent functional magnetic resonance brain imaging (fMRI) was performed between 8.00 and 9.00AM (GE Signa 3T scanner, T2* weighted echo-planar imaging, voxel size 3.4 × 3.4 × 2.6mm, TR 2100ms, TE 30ms, angle 70°, 30 slices, interleaved acquisition, and 2.6mm slice thickness with 1.4mm gap).

Taste Task

Individuals received three taste stimuli during fMRI imaging(43): 1 mol/L sucrose solution (100 trials), no solution (100 trials), and artificial saliva (80 trials). Individuals learned to associate each taste stimulus with a paired conditioned visual stimulus (CS) that is probabilistically associated with its US: the no-solution (null) CS was followed in 20% of trials by sucrose (unexpected sucrose receipt, positive-prediction error), and the sucrose CS was followed in 20% of trials by no-solution (unexpected Sucrose omission, negative-prediction error). The first 10 trials were fixed CS shape for sucrose followed by US sucrose delivery to establish an initial stable association between the CS sucrose shape and US sucrose taste (43). Trials began with the CS (2 seconds), followed by US delivery, tongue swish and swallow and awaiting the next trial (4 seconds). Each trial lasted 6 seconds. Every 2.1 seconds a brain image was recorded. All other trials were fully randomized. Taste stimuli were applied using a customized programmable syringe pump (J-Kem Scientific, St. Louis, Missouri) controlled by E-Prime Software (Psychological Software Tools, Pittsburgh, Pennsylvania) and triggered by MRI-scanner radiofrequency pulse (19). Task duration was 28 minutes. We only included trials with matching CS–US association in order to focus on taste classification and reduce effects from the prediction error response.

Brain Imaging Analysis

Brain-imaging data were preprocessed and analyzed using Statistical Parametric Mapping software (<http://www.fil.ion.ucl.ac.uk/spm/software/spm5/>). Images were realigned to the first volume, normalized to the Montreal Neurological Institute template, and smoothed with a 6-mm FWHM Gaussian kernel (44). Image sequences were manually inspected and images with artifacts or movement >1 voxel removed. Data were modeled with a hemodynamic response convolved boxcar function, using the general linear model, including temporal and dispersion derivatives, autoregression, and 128s high-pass filter.

We computed three first level contrasts for each subject: 1) sucrose versus no solution; 2) artificial saliva versus no solution; 3) sucrose versus artificial saliva.

Analysis 1 (group by condition ANCOVA)—To test whether groups differed in insula activation strength, we used a random effects, whole brain analysis ($p < 0.05$ family-wise error corrected (FWE), cluster size = 5 voxels) (23).

Analysis 2 (Multivariate Bayes, MVB, Decoding Analysis)—Step 1. In the MVB approach (27) the first step is to identify the optimal model of activation distribution in the relevant brain region. The prediction is that brain activation is distributed according to a sparse model solution. The area of interest on the functional images and for the contrasts of interest are identified (from the taste activation task). The next step is a “greedy search” procedure. Here the algorithm's goal is to detect the best model of activation distribution in relation to the task condition. The brain region of interest is partitioned in subsets of increasing size and tested for type of activation distribution: In a sparse model few voxels have large variance while most have small variance. In a smooth model, there is a sparse representation of activation that is spatially coherent over the brain anatomy. The support model is a type of distributed model where each pattern is an individual voxel and a large number of distributed patterns are expressed. The sparse, support, and smooth MVB models were tested for each of the three contrasts of interest (1. expected sucrose contrasted against expected no solution; 2. expected sucrose contrasted against expected artificial saliva; 3. expected artificial saliva contrasted against expected no solution) for each participant. Expectation–maximization (EM) uses the highest voxel weights for fitting the model. This method creates the log-evidence value. The solution with the largest log-evidence indicates the optimal set of activation distribution. EM algorithms can be prone to over-fitting with higher order polynomials. However, this problem is reduced with increasing number of data points and the fMRI study and each condition (sucrose, artificial saliva, no-solution) had 80 trials with 228 images across those 80 trials should provide adequate number of data points (45). Step 2. After delineation of the adequate distribution model, the log evidence can then be used for cross validation to identify classification accuracy. Cross-validation partitions data into subsets so that the analysis is performed on one (training) subset, while the other (test) data are retained to confirm and validate the initial analysis. In k -fold cross validation, data is randomly partitioned into k partitions, training the classifier on all but one and evaluates classification performance on that partition. This procedure is repeated for all k partitions (here $k=8$). Cross-validation prediction accounts properly for serial correlations and confounds by ensuring that the cross-validation weights cannot be influenced by test

data and that the prediction is conditionally independent of the training data. Classification accuracy was then compared across groups.

While there are other classification algorithms such as support vector machine or Gaussian models, we chose MVB for various reasons. We were not interested whether brain response can separate patients from CW, which is a typical goal of classification algorithms. Rather MVB tests different distribution models and compares the model evidence for each participant. Other methods base their analysis on the expectation that a sparse model is at hand. In light of different volumetric measures of the insula across eating disorders we wanted to make sure that we test various models in the eating disorder groups and use the most accurate data for cross validation.

Statistical Analysis

Behavioral data (ANOVA) and classification accuracy (univariate general linear model including factors and covariates, ANCOVA) were analyzed with SPSS-22 software (IBM-SPSS, Chicago, IL). Post-hoc group comparisons were assessed with Dunnett's T3, and estimated marginal means were computed for classification accuracy and multiple-comparison corrected using bootstrap. Several variables were hypothesized a priori to be possible confounds and included in the between-group model (5 group ANCOVA, CW, AN, BN, recovered AN, obese individuals): number of activated voxels (adjusting for volume differences), interoceptive deficits, sweetness perception of control solution as well as sucrose solution; in addition comorbid anxiety and depression diagnoses and medication use were included as factors in the model. Pearson correlation analysis tested brain-behavior correlations. Sweetness perception across sucrose concentrations was tested with repeated measures ANOVA.

Results

Demographic Variables (Table 1.)

Age was similar between groups. BMI was higher in obesity and lower in AN and recovered AN groups compared to CW. Interoceptive deficits, harm avoidance and state and trait anxiety were higher in AN and BN groups compared to CW, depression scores were elevated in AN, recovered AN and BN. Drive for thinness and body dissatisfaction were elevated in all groups compared to CW, bulimia scores were elevated in BN and obesity. Sweetness and pleasantness perception were similar between groups, as was slope for sweetness perception across concentrations ($F=0.819$, $p<0.516$). Some individuals with AN or BN had comorbid psychiatric disorders or were on medication.

Analysis 1 (group by condition ANCOVA)

The 5-group by 3-taste condition contrast including covariates depression, anxiety and medication use, with and without interoceptive deficits or sweet taste perception did not result in significantly different insula activation, however within groups, all contrasts showed positive posterior and mid insula activation.

Analysis 2 (Multivariate Bayes, MVB, Decoding Analysis)

Classification accuracy (mean $\pm$ SEM) for sucrose versus no solution (CW :70.7 $\pm$ 1.6; AN: 70.5 $\pm$ 1.3; BN:68.9 $\pm$ 1.4; obesity:68.8 $\pm$ 1.4; Recovered AN:69.7 $\pm$ 1.5) was not significantly different between groups (df=4,F=0.615,p<0.7). Classification accuracy for artificial saliva versus no solution (CW: 70.2 $\pm$ 1.4; AN:69.8 $\pm$ 1.2; BN:70.3 $\pm$ 1.3; obesity:69.9 $\pm$ 1.2; recovered AN:69.5 $\pm$ 1.3) was also not significantly different across groups (df=4,F=0.109,p<0.9).

Sucrose versus artificial saliva classification accuracy differed significantly across groups (df=4,F=2.601, p<0.041), with lower classification accuracy (mean $\pm$ SEM) in AN (56.6 $\pm$ 0.8;p<0.016) and obesity (58.1 $\pm$ 0.9;p<0.027) compared to CW (60.4 $\pm$ 1.0), and lower values (p<0.047) for AN versus recovered AN (60.0 $\pm$ 0.9;p<0.016) and obesity versus recovered AN. BN (58.20 $\pm$ 0.89) did not differ significantly from any group (Figure 2.). The voxel weights in each group were significantly greater for sparse versus smooth or support distribution (Figure 1., Supplemental Figure 1): CW sparse 79.2 $\pm$ 45.4, smooth 13.1 $\pm$ 17.4, support 8.2 $\pm$ 14.4, F=49.5, p<0.001; AN sparse 75.0 $\pm$ 50.6, smooth 17.9 $\pm$ 28.3, support 12.0 $\pm$ 24.1, F=19.4, p<0.001; BN sparse 79.0 $\pm$ 43.8, smooth 17.5 $\pm$ 23.7, support 9.8 $\pm$ 19.2, F=30.3, p<0.001; obesity sparse 75.4 $\pm$ 34.1, smooth 11.5 $\pm$ 11.6, support 5.2 $\pm$ 9.4, F=62.7, p<0.001; recovered AN sparse 74.2 $\pm$ 37.1, smooth 15.0 $\pm$ 20.2, support 7.8 $\pm$ 20.7, F=34.2, p<0.001.

Correlation Analyses (Table 2.)

CW—Classification accuracy for sucrose versus artificial saliva contrasts correlated positively with number of activated voxels (261.9 $\pm$ 28.7;r=0.688,p<0.001) and negatively with sucrose sweetness perception (r=-0.527,p<0.005).

AN—Classification accuracy for sucrose versus artificial saliva was positively correlated with number of activated voxels (257.6 $\pm$ 48.9;r=0.712,p<0.001).

Recovered AN—Classification accuracy for sucrose versus artificial saliva was positively correlated with interoceptive deficits (r=0.550, p<0.015).

BN—Classification accuracy for sucrose versus artificial saliva was positively correlated with number of activated voxels (274.5 $\pm$ 70.1;r=0.679,p<0.001); classification accuracy for sucrose versus no solution correlated negatively with sucrose sweetness (r=-0.690,p<0.001).

Obesity—Classification accuracy for sucrose versus artificial saliva correlated positively with number of activated voxels (247.6 $\pm$ 48.9;r=0.853,p<0.001) with a tendency to positive correlation with interoceptive deficits (r=0.425, p<0.070).

Discussion

This study indicates that AN and obesity are associated with reduced taste classification accuracy in the insula when contrasting caloric sucrose against a control solution. Pattern classification accuracy in recovered AN and BN did not differ significantly from the CW group and suggests that taste classification accuracy alterations could be adaptations to an abnormal eating and weight state.

The overall accuracy of insula pattern classification between 50% and 70% may seem low but is comparable with other studies that investigated brain pattern classification (46, 47). Larger regions of interest are used for decoding and pattern classification and typically result in higher classification accuracy; we selected the bilateral insula as a larger yet anatomically defined region. There was no group difference in classification accuracy for sucrose or control solution tested against no solution. This suggests that the perceptual state to sucrose or control solution applied individually, which includes taste perception but also general sensory information, is encoded adequately in AN and obesity. However, when directly contrasting the two taste stimuli, to remove effects of texture and other sensory stimulation in the mouth, encoding differences more specific to taste quality become apparent. Thus there may be deficits in neuronal encoding of distinct taste qualities in AN and obesity, but this may remit with recovery from AN as in our recovered sample. Those results seem to be in line with previous studies. Perception of individual taste stimuli in EDs was not altered (48), but research indicated lower olfactory or gustatory stimulus *discrimination* in EDs (49). Some studies in obesity have also shown difficulties with taste differentiation (50), and sweet taste sensitivity increases with weight-reduction in obesity (51, 52).

What determines such alterations could occur on a variety of levels. For instance, leptin and other hormones are altered in EDs and obesity and affect taste perception (53-55); reduced insula pattern classification could be due to primary structural changes *within* the insula (15), or alternatively could result from altered taste signal processing in afferent pathways *to* the insula (56).

The insula has repeatedly been implicated in ED and obesity pathophysiology (13, 17-21, 23, 57, 58). The bilateral anterior and middle insula responds to taste stimulation (59) and transmits information to ventral striatal and orbitofrontal reward pathways (11). The anteroventral insula is connected to the amygdala (12) and aids in generating internal emotional states (60). Especially the *right* anterior insula has been associated with self-recognition, “abstract representation of oneself” (61) and interoceptive awareness (62). *Left* anteroventral insula activation is related to gastric distention (14) and self-reported fullness (63). Thus, many intertwined functions are processed and alterations on different levels within the insula could contribute to our findings. Altered insula function could have important clinical implications. If normal taste discrimination is disturbed, then it is possible that normal insula inputs to basal ganglia and higher order taste processing are altered (64). Subsequently, other circuits such as subcortical reward or prefrontal cognitive-emotional circuits could have greater influence on determining the drive to eat or not eat. AN and obesity have been associated with altered dopamine related reward function (23), elevated or reduced cognitive control (65, 66) as well as high anxiety (67), and disturbances in those circuits could increasingly drive eating pathology the more afferents from the insula are altered. In fact, such a “multifaceted” concept for behavior modulation has been suggested previously for the psychopathology of obesity (68). Hypothetically, if individuals with AN and obesity do not receive appropriate insula signals in response to taste stimuli but basal ganglia dopamine hyper- (AN) or hypo- (obesity) activation respectively have a stronger impact on food choices, then it could be possible that reducing flavor intensity during treatment of AN and enhancing flavor for obesity could counteract those subcortical

mechanisms. Normal insula classification accuracy in the recovered AN group provides hope that classification accuracy alterations do improve with recovery.

Limitations

We can only speculate at this point why AN as well as obesity groups showed similarly reduced classification accuracy when contrasting sucrose versus artificial saliva. Response to taste stimuli may be similarly increased or decreased across stimuli in AN or obesity (18, 23) and the result when contrasting stimuli against each other may be for each case reduced classification accuracy. This will require further research. We included medication use and comorbid conditions as covariates in the analysis but cannot exclude entirely their possible effects (69, 70). Brain volumes were different in parts of the insula in individuals with EDs or obesity in past studies, which could have affected the findings; however, we corrected for the number of activated voxels to adjust for such effects. Interoception was altered in EDs, which could have affected the perception of taste but was included as a covariate to adjust for such effects. We included the bilateral insula in the analysis and sub-regions could have shown different results, but we felt it would be important to include the entire insula in the analysis in order not to exclude important aspects of the circuitry (59). The classification accuracy for taste solution versus no stimulation was relatively high with around 70% for all participants, but was lower for sucrose versus control solution. This could be due to contributions of general aspects of the taste stimuli such as water based solution texture and temperature, which were similar for the study stimuli. However, smaller sub-regions of the insula could have provided more refined results, thus future studies will need to investigate insula subdivisions. Classification accuracy in BN was normal, however, a larger sample size may have shown that also the BN group has lower values. Taste pleasantness and disgust also have important influence on internal response to stimuli and brain response (71, 72). In this study we aimed to avoid conditioned emotional response and we chose a highly sweet 1 molar sucrose solution and a neutral control solution as stimuli. What we did not specifically test was disgust experience. Disgust is a complex emotion that can be related to taste stimuli, but even more so comes into play in everyday situations that for instance involve social and moral values. We could not identify commonly used disgust measures in taste fMRI research but the Disgust Scale (73) is a widely used assessment tool. Its dimensions Core Disgust, Animal Reminder Disgust, and Contamination-Based Disgust can not necessarily be related to taste experience in this paradigm, but measures from this scale have been associated with for instance OCD and it is possible that measures from this scale could be predictive of eating disorder diagnosis and this in turn could be related to brain function. Disgust in fact has been associated with anterior insula activation (74) and therefore differing experience of disgust across groups could have had impact on the brain response or classification accuracy measure. This will need further investigation in the future.

In conclusion, pattern classification is relatively new to fMRI brain imaging analysis, but provides information that goes beyond strength of activation and toward understanding of neuronal patterns, which could inform about innervation and provide a more refined approach to brain function. Based on the study results, we propose that very basic coding mechanisms for taste quality are altered in AN and obesity. ED cognitions should therefore not be responsible for this alteration, but under- and over-weight or food deprivation and

overstimulation might drive the results observed. The functional significance of these results will need further study, but normal values in individuals recovered from AN suggest that taste stimuli coding may be state dependent and recover with weight restoration.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

References

1. Crow SJ, Peterson CB, Swanson SA, Raymond NC, Specker S, Eckert ED, et al. Increased mortality in bulimia nervosa and other eating disorders. *Am J Psychiatry*. 2009; 166(12):1342–6. [PubMed: 19833789]
2. American Psychiatric Association. Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition (DSM-5(TM)). 5th ed.. American Psychiatric Publishing; Arlington, Va.: 2013. 5 edition (May 27, 2013)
3. Ogden CL, Carroll MD, Kit BK, Flegal KM. Prevalence of obesity and trends in body mass index among US children and adolescents, 1999–2010. *JAMA*. 2012; 307(5):483–90. [PubMed: 22253364]
4. Flegal KM, Carroll MD, Kit BK, Ogden CL. Prevalence of obesity and trends in the distribution of body mass index among US adults, 1999–2010. *JAMA*. 2012; 307(5):491–7. [PubMed: 22253363]
5. Fichter MM, Quadflieg N. Twelve-year course and outcome of bulimia nervosa. *Psychol Med*. 2004; 34(8):1395–406. [PubMed: 15724871]
6. Fichter MM, Quadflieg N, Hedlund S. Twelve-year course and outcome predictors of anorexia nervosa. *Int J Eat Disord*. 2005
7. Flegal KM, Carroll MD, Ogden CL, Curtin LR. Prevalence and trends in obesity among US adults, 1999–2008. *JAMA*. 2010; 303(3):235–41. [PubMed: 20071471]
8. Lowe B, Zipfel S, Buchholz C, Dupont Y, Reas DL, Herzog W. Long-term outcome of anorexia nervosa in a prospective 21-year follow-up study. *Psychol Med*. 2001; 31(5):881–90. [PubMed: 11459385]
9. Drewnowski A, Henderson SA, Levine A, Hann C. Taste and food preferences as predictors of dietary practices in young women. *Public Health Nutr*. 1999; 2(4):513–9. [PubMed: 10656470]
10. Yaxley S, Rolls ET, Sienkiewicz ZJ. Gustatory responses of single neurons in the insula of the macaque monkey. *J Neurophysiol*. 1990; 63(4):689–700. [PubMed: 2341869]
11. Rolls ET. Functions of the orbitofrontal and pregenual cingulate cortex in taste, olfaction, appetite and emotion. *Acta Physiol Hung*. 2008; 95(2):131–64. [PubMed: 18642756]
12. Fudge JL, Breitbart MA, Danish M, Pannoni V. Insular and gustatory inputs to the caudal ventral striatum in primates. *J Comp Neurol*. 2005; 490(2):101–18. [PubMed: 16052493]
13. Kaye WH, Fudge JL, Paulus M. New insights into symptoms and neurocircuit function of anorexia nervosa. *Nat Rev Neurosci*. 2009; 10(8):573–84. [PubMed: 19603056]
14. Craig AD. How do you feel--now? The anterior insula and human awareness. *Nature reviews Neuroscience*. 2009; 10(1):59–70.
15. Frank GK, Shott ME, Hagman JO, Mittal VA. Alterations in brain structures related to taste reward circuitry in ill and recovered anorexia nervosa and in bulimia nervosa. *Am J Psychiatry*. 2013; 170(10):1152–60. [PubMed: 23680873]
16. Van den Eynde F, Suda M, Broadbent H, Guillaume S, Van den Eynde M, Steiger H, et al. Structural magnetic resonance imaging in eating disorders: a systematic review of voxel-based morphometry studies. *European eating disorders review : the journal of the Eating Disorders Association*. 2012; 20(2):94–105. [PubMed: 22052722]
17. Wagner A, Aizenstein H, Mazurkewicz L, Fudge J, Frank GK, Putnam K, et al. Altered insula response to taste stimuli in individuals recovered from restricting-type anorexia nervosa. *Neuropsychopharmacology*. 2008; 33(3):513–23. [PubMed: 17487228]

18. Cowdrey FA, Park RJ, Harmer CJ, McCabe C. Increased Neural Processing of Rewarding and Aversive Food Stimuli in Recovered Anorexia Nervosa. *Biol Psychiatry*. 2011
19. Frank GK, Reynolds JR, Shott ME, O'Reilly RC. Altered temporal difference learning in bulimia nervosa. *Biol Psychiatry*. 2011; 70(8):728–35. [PubMed: 21718969]
20. Bohon C, Stice E. Reward abnormalities among women with full and subthreshold bulimia nervosa: A functional magnetic resonance imaging study. *Int J Eat Disord*. 2010
21. Huerta CI, Sarkar PR, Duong TQ, Laird AR, Fox PT. Neural bases of food perception: coordinate-based meta-analyses of neuroimaging studies in multiple modalities. *Obesity (Silver Spring)*. 2014; 22(6):1439–46. [PubMed: 24174404]
22. Frank S, Kullmann S, Veit R. Food related processes in the insular cortex. *Front Hum Neurosci*. 2013; 7:499. [PubMed: 23986683]
23. Frank GK, Reynolds JR, Shott ME, Jappe L, Yang TT, Tregellas JR, et al. Anorexia nervosa and obesity are associated with opposite brain reward response. *Neuropsychopharmacology*. 2012; 37(9):2031–46. [PubMed: 22549118]
24. Kriegeskorte N. Pattern-information analysis: from stimulus decoding to computational-model testing. *Neuroimage*. 2011; 56(2):411–21. [PubMed: 21281719]
25. D'Ardenne K, McClure SM, Nystrom LE, Cohen JD. BOLD responses reflecting dopaminergic signals in the human ventral tegmental area. *Science*. 2008; 319(5867):1264–7. [PubMed: 18309087]
26. Brodersen KH, Wiech K, Lomakina EI, Lin CS, Buhmann JM, Bingel U, et al. Decoding the perception of pain from fMRI using multivariate pattern analysis. *Neuroimage*. 2012; 63(3):1162–70. [PubMed: 22922369]
27. Friston K, Chu C, Mourao-Miranda J, Hulme O, Rees G, Penny W, et al. Bayesian decoding of brain images. *Neuroimage*. 2008; 39(1):181–205. [PubMed: 17919928]
28. Naselaris T, Kay KN, Nishimoto S, Gallant JL. Encoding and decoding in fMRI. *Neuroimage*. 2011; 56(2):400–10. [PubMed: 20691790]
29. Chu C, Mourao-Miranda J, Chiu YC, Kriegeskorte N, Tan G, Ashburner J. Utilizing temporal information in fMRI decoding: classifier using kernel regression methods. *Neuroimage*. 2011; 58(2):560–71. [PubMed: 21729756]
30. Macevoy SP, Epstein RA. Decoding the representation of multiple simultaneous objects in human occipitotemporal cortex. *Curr Biol*. 2009; 19(11):943–7. [PubMed: 19446454]
31. Tong F, Pratte MS. Decoding patterns of human brain activity. *Annu Rev Psychol*. 2012; 63:483–509. [PubMed: 21943172]
32. Haynes JD. Decoding visual consciousness from human brain signals. *Trends Cogn Sci*. 2009; 13(5):194–202. [PubMed: 19375378]
33. Herbert BM, Herbert C, Pollatos O, Weimer K, Enck P, Sauer H, et al. Effects of short-term food deprivation on interoceptive awareness, feelings and autonomic cardiac activity. *Biol Psychol*. 2012; 89(1):71–9. [PubMed: 21958594]
34. Dunn BD, Stefanovitch I, Evans D, Oliver C, Hawkins A, Dalgleish T. Can you feel the beat? Interoceptive awareness is an interactive function of anxiety- and depression-specific symptom dimensions. *Behav Res Ther*. 2010; 48(11):1133–8. [PubMed: 20692645]
35. American Psychiatric Association. *Handbook of Psychiatric Measures*. 4 ed.. American Psychiatric Association; Washington DC: 2000. *Diagnostic and Statistical Manual of Mental Disorders - Text Revision (DSM-IV-TR)*..
36. CDC. <http://www.cdc.gov/>
37. Dreher JC, Schmidt PJ, Kohn P, Furman D, Rubinow D, Berman KF. Menstrual cycle phase modulates reward-related neural function in women. *Proc Natl Acad Sci U S A*. 2007; 104(7):2465–70. [PubMed: 17267613]
38. Garner, D. *Eating Disorder Inventory™-3 (EDI™-3)*. Psychological Assessment Resources, Inc.; Lutz, FL: 2004.
39. Cloninger, C.; Przybeck, T.; Svrakic, D.; Wetzell, R. *The temperament and character inventory (TCI): A guide to its development and use*. Center for Psychobiology of Personality, Washington University; St. Louis, MO: 1994.

40. Spielberger, CD. Manual for the State-Trait Anxiety Inventory. Consulting Psychologists Press, Inc.; Palo Alto, CA: 1983.
41. Beck AT, Ward M, Mendelson M, Mock J, Erbaugh J. An Inventory for measuring depression. *Arch Gen Psychiatry*. 1961; 4:53–63.
42. O'Connor R, Colder C, Hawk L. Confirmatory factor analysis of the Sensitivity to Punishment and Sensitivity to Reward Questionnaire. *Personality and Individual Differences*. 2004; 37:985–1002.
43. O'Doherty JP, Dayan P, Friston K, Critchley H, Dolan RJ. Temporal difference models and reward-related learning in the human brain. *Neuron*. 2003; 38(2):329–37. [PubMed: 12718865]
44. Statistical Parametric Mapping 5: Wellcome Trust Centre for Neuroimaging. 2005. [Available from: <http://www.fil.ion.ucl.ac.uk/spm/software/spm5/>]
45. Bishop, C. Pattern Recognition and Machine Learning.. In: Jordan, M.; Kleinberg, J.; Scholkopf, B., editors. *Information Science and Statistics*. Springer; Cambridge, United Kingdom: 2006.
46. Crouzet SM, Busch NA, Ohla K. Taste quality decoding parallels taste sensations. *Curr Biol*. 2015; 25(7):890–6. [PubMed: 25772445]
47. Misaki M, Kim Y, Bandettini PA, Kriegeskorte N. Comparison of multivariate classifiers and response normalizations for pattern-information fMRI. *Neuroimage*. 2010; 53(1):103–18. [PubMed: 20580933]
48. Sunday S, Halmi K. Taste perceptions and hedonics in eating disorders. *Physiol Behav*. 1990; 48:587–94. [PubMed: 2082356]
49. Dazzi F, Nitto SD, Zambetti G, Lorigio C, Ciofalo A. Alterations of the olfactory-gustatory functions in patients with eating disorders. *Eur Eat Disord Rev*. 2013; 21(5):382–5. [PubMed: 23788398]
50. Overberg J, Hummel T, Krude H, Wiegand S. Differences in taste sensitivity between obese and non-obese children and adolescents. *Arch Dis Child*. 2012; 97(12):1048–52. [PubMed: 22995095]
51. Umabiki M, Tsuzaki K, Kotani K, Nagai N, Sano Y, Matsuoka Y, et al. The improvement of sweet taste sensitivity with decrease in serum leptin levels during weight loss in obese females. *Tohoku J Exp Med*. 2010; 220(4):267–71. [PubMed: 20383037]
52. Pepino MY, Bradley D, Eagon JC, Sullivan S, Abumrad NA, Klein S. Changes in taste perception and eating behavior after bariatric surgery-induced weight loss in women. *Obesity (Silver Spring)*. 2013
53. Bailer, UF.; Kaye, WH. Neuroendocrine and neuropeptide dysregulation in anorexia nervosa and bulimia nervosa.. In: Brewerton, T., editor. *Clinical Handbook of Eating Disorders: An Integrated Approach*. Marcell Dekker, Inc.; New York: 2004. p. 283-95.
54. Goldstone AP. The hypothalamus, hormones, and hunger: alterations in human obesity and illness. *Prog Brain Res*. 2006; 153:57–73. [PubMed: 16876568]
55. Jyotaki M, Shigemura N, Ninomiya Y. Modulation of sweet taste sensitivity by orexigenic and anorexigenic factors. *Endocr J*. 2010; 57(6):467–75. [PubMed: 20431269]
56. Jones LM, Fontanini A, Katz DB. Gustatory processing: a dynamic systems approach. *Curr Opin Neurobiol*. 2006; 16(4):420–8. [PubMed: 16842991]
57. Oberndorfer TA, Frank GK, Simmons AN, Wagner A, McCurdy D, Fudge JL, et al. Altered insula response to sweet taste processing after recovery from anorexia and bulimia nervosa. *Am J Psychiatry*. 2013; 170(10):1143–51. [PubMed: 23732817]
58. Frank GK, Wagner A, Achenbach S, McConaha C, Skovira K, Aizenstein H, et al. Altered brain activity in women recovered from bulimic-type eating disorders after a glucose challenge: A pilot study. *Int J Eat Disord*. 2006; 39(1):76–9. [PubMed: 16254868]
59. Kurth F, Zilles K, Fox PT, Laird AR, Eickhoff SB. A link between the systems: functional differentiation and integration within the human insula revealed by meta-analysis. *Brain structure & function*. 2010; 214(5-6):519–34. [PubMed: 20512376]
60. Phillips ML, Drevets WC, Rauch SL, Lane R. Neurobiology of emotion perception II: Implications for major psychiatric disorders. *Biol Psychiatry*. 2003; 54(5):515–28. [PubMed: 12946880]
61. Devue C, Collette F, Baletau E, Degueldre C, Luxen A, Maquet P, et al. Here I am: the cortical correlates of visual self-recognition. *Brain Res*. 2007; 1143:169–82. [PubMed: 17306235]

62. Critchley HD, Wiens S, Rotshtein P, Ohman A, Dolan RJ. Neural systems supporting interoceptive awareness. *Nat Neurosci.* 2004; 7(2):189–95. [PubMed: 14730305]
63. Wang GJ, Tomasi D, Backus W, Wang R, Telang F, Geliebter A, et al. Gastric distention activates satiety circuitry in the human brain. *Neuroimage.* 2008; 39(4):1824–31. [PubMed: 18155924]
64. Maffei A, Haley M, Fontanini A. Neural processing of gustatory information in insular circuits. *Curr Opin Neurobiol.* 2012; 22(4):709–16. [PubMed: 22554880]
65. Tchanturia K, Anderluh MB, Morris RG, Rabe-Hesketh S, Collier DA, Sanchez P, et al. Cognitive flexibility in anorexia nervosa and bulimia nervosa. *J Int Neuropsychol Soc.* 2004; 10(4):513–20. [PubMed: 15327730]
66. Gearhardt AN, Yokum S, Orr PT, Stice E, Corbin WR, Brownell KD. Neural correlates of food addiction. *Arch Gen Psychiatry.* 2011; 68(8):808–16. [PubMed: 21464344]
67. Kaye WH, Wierenga CE, Bailer UF, Simmons AN, Bischoff-Grethe A. Nothing tastes as good as skinny feels: the neurobiology of anorexia nervosa. *Trends Neurosci.* 2013; 36(2):110–20. [PubMed: 23333342]
68. Volkow ND, Wang GJ, Fowler JS, Telang F. Overlapping neuronal circuits in addiction and obesity: evidence of systems pathology. *Philos Trans R Soc Lond B Biol Sci.* 2008; 363(1507): 3191–200. [PubMed: 18640912]
69. Platte P, Herbert C, Pauli P, Breslin PA. Oral perceptions of fat and taste stimuli are modulated by affect and mood induction. *PLoS One.* 2013; 8(6):e65006. [PubMed: 23755167]
70. McCabe C, Mishor Z, Cowen PJ, Harmer CJ. Diminished neural processing of aversive and rewarding stimuli during selective serotonin reuptake inhibitor treatment. *Biol Psychiatry.* 2010; 67(5):439–45. [PubMed: 20034615]
71. Troop NA, Murphy F, Bramon E, Treasure JL. Disgust sensitivity in eating disorders: a preliminary investigation. *Int J Eat Disord.* 2000; 27(4):446–51. [PubMed: 10744851]
72. Watkins TJ, Di Iorio CR, Olatunji BO, Benningfield MM, Blackford JU, Dietrich MS, et al. Disgust proneness and associated neural substrates in obesity. *Soc Cogn Affect Neurosci.* 2015
73. Olatunji BO, Williams NL, Tolin DF, Abramowitz JS, Sawchuk CN, Lohr JM, et al. The Disgust Scale: item analysis, factor structure, and suggestions for refinement. *Psychol Assess.* 2007; 19(3): 281–97. [PubMed: 17845120]
74. Jabbi M, Bastiaansen J, Keysers C. A common anterior insula representation of disgust observation, experience and imagination shows divergent functional connectivity pathways. *PLoS One.* 2008; 3(8):e2939. [PubMed: 18698355]

Figure 1. Distributed sparse coding pattern of insula activation rendered on template brain across exemplary participants from each study group. A) High classification accuracy; B) Low classification accuracy. CA: classification accuracy.

Figure 2.
Classification accuracy across groups.

Table 1

Demographic variables.

	Healthy Comparison Women (A)		Women with Anorexia Nervosa (B)		Women with Bulimia Nervosa (C)		Women with Obesity (D)		Women Recovered from Anorexia Nervosa (E)		F	P-value
	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD		
	n = 27		n = 21		n = 20		n = 19		n = 19			
Age (years)	26.2	7.0	22.9	6.1	25.2	5.3	28.2	8.1	27.0	5.3	1.9	0.115
BMI (kg/m ²)	21.5	1.4	16.0	1.1	22.6	5.7	34.7	4.6	20.2	1.1	89.0	<0.001
Novelty Seeking	18.5	5.3	13.8	6.8	22.1	6.7	21.2	5.5	17.5	5.8	6.1	<0.001
Harm Avoidance	9.9	4.1	22.7	6.6	23.0	5.8	12.6	5.9	15.4	7.2	22.6	<0.001
Depression	1.3	1.0	23.5	9.8	24.5	11.4	4.8	4.6	3.8	3.6	56.1	<0.001
Interoceptive Deficits (EDI-3)	1.0	1.6	16.0	7.7	18.2	6.2	6.1	9.3	3.8	5.1	32.4	<0.001
Drive for Thinness (EDI-3)	2.5	3.3	19.5	5.9	23.1	4.5	13.1	8.4	8.3	6.0	48.8	<0.001
Bulimia (EDI-3)	1.0	1.4	3.5	3.9	22.7	5.3	12.0	11.9	2.2	2.4	50.2	<0.001
Body Dissatisfaction (EDI-3)	4.0	4.2	24.8	8.9	30.7	8.0	26.6	9.0	9.6	6.5	56.4	<0.001
Punishment Sensitivity	4.2	1.8	12.8	4.7	12.4	3.9	6.9	4.5	7.1	4.1	21.6	<0.001
Reward Sensitivity	4.5	2.9	7.5	3.6	8.4	3.7	6.3	4.6	6.4	3.4	3.9	0.006
State Anxiety	26.0	5.4	52.7	11.5	50.1	13.8	31.8	10.9	32.2	13.3	26.0	<0.001
Trait Anxiety	28.3	5.7	54.9	11.7	58.0	11.1	35.3	9.7	34.8	11.0	39.5	<0.001
Sucrose Sweetness	5.2	2.3	4.5	2.5	5.5	2.9	4.1	2.5	5.1	1.9	1.1	0.362
Sucrose Pleasantness	8.4	0.8	8.6	0.6	8.7	0.6	8.3	1.2	9.2	1.0	1.2	0.336
SU - AS - Number Activated Voxels	159.4	74.0	180.0	97.0	165.4	72.8	174.8	75.5	143.5	46.5	0.7	0.587
SU - NO - Number Activated Voxels	261.9	28.7	257.6	48.9	274.5	70.1	247.6	48.9	241.3	59.4	1.2	0.304

	Healthy Comparison Women (A) n = 27		Women with Anorexia Nervosa (B) n = 21		Women with Bulimia Nervosa (C) n = 20		Women with Obesity (D) n=19		Women Recovered from Anorexia Nervosa (E) n = 19		F	P-value
	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD		
AS - NO - Number Activated Voxels	248.3	47.6	238.1	65.0	246.0	63.2	246.3	52.2	256.0	55.2	0.3	0.905
n.s.												
<hr/>												
<u>Medication Use</u>												
SSRI	0		9		9		0		3			
Atypical Antipsychotic	0		2		0		0		0			
SSRI + Atypical Antipsychotic	0		1		4		0		0			
<hr/>												
<u>Comorbid Diagnoses</u>												
Major Depression	0		3		3		0		3			
Anxiety Disorder	0		4		6		0		3			
Major Depression + Anxiety Disorder	0		5		7		0		1			

BMI: body mass index; EDI-3: Eating Disorder Inventory 3; SSRI: selective serotonin reuptake inhibitor; SU-AS: sucrose vs artificial saliva; SU-NO, sucrose vs no solution; AS-NO, artificial saliva vs no solution.

Table 2

Correlation Results.

Population	Pearson Correlation		ED13-ID	PAS	SAS	PIMSU	SIMSU	Significantly Activated Voxels		
	<i>r</i>	<i>p</i>						SU-AS	SU-NO	AS-NO
Healthy CW	SU-AS % Accuracy	<i>r</i>	0.028	0.254	-0.135	0.25	-.527**	.688***		
		<i>p</i>	0.89	0.202	0.502	0.208	0.005	<0.001		
	SU-NO % Accuracy	<i>r</i>	-0.198			-0.163	-0.094		-0.037	
AS-NO % Accuracy		<i>p</i>	0.322			0.417	0.639		0.854	
		<i>r</i>	-0.11	0.021	-0.031					0.075
		<i>p</i>	0.584	0.917	0.878					0.709
Women with AN	SU-AS % Accuracy	<i>r</i>	-0.143	-0.071	0.178	-0.22	-0.186	.712***		
		<i>p</i>	0.536	0.76	0.44	0.337	0.419	<0.001		
	SU-NO % Accuracy	<i>r</i>	0.318			-0.209	0.135		-0.401	
AS-NO % Accuracy		<i>p</i>	0.16			0.363	0.56		0.072	
		<i>r</i>	0.246	-0.208	-0.01					.797***
		<i>p</i>	0.283	0.367	0.964					<0.001
Women with BN	SU-AS % Accuracy	<i>r</i>	-0.193	.561*	1	0.126	0.093	.679***		
		<i>p</i>	0.414	0.01	.	0.596	0.697	<0.001		
	SU-NO % Accuracy	<i>r</i>	-0.212			-0.04	-0.187		0.32	
AS-NO % Accuracy		<i>p</i>	0.37			0.868	0.431		0.17	
		<i>r</i>	-0.403	0.134	1					0.396
		<i>p</i>	0.078	0.572	.					0.084
Women with obesity	SU-AS % Accuracy	<i>r</i>	0.425	-0.168	0.143	0.16	0.259	.853***		
		<i>p</i>	0.07	0.492	0.559	0.512	0.284	<0.001		
	SU-NO % Accuracy	<i>r</i>	0.17			-0.04	-0.238		0.071	
AS-NO % Accuracy		<i>p</i>	0.487			0.871	0.326		0.771	
		<i>r</i>	0.013	0.125	0.175					-0.153
		<i>p</i>	0.958	0.611	0.474					0.531
Women recovered from AN	SU-AS % Accuracy	<i>r</i>	.550*	0.196	0.098	0.088	0.021	0.217		

Population	Pearson Correlation	EDI3-ID	P AS	S AS	P 1M SU	S 1M SU	Significantly Activated Voxels		
							SU-AS	SU-NO	AS-NO
AN: anorexia nervosa; BN: bulimia nervosa; CW: comparison women; EDI3-ID: Eating Disorder Inventory 3 Interceptive Deficits; P AS: Artificial saliva pleasantness rating; S AS: Artificial saliva sweetness rating; P 1 M SU: 1 molar sucrose pleasantness rating; S 1M SU: 1 molar sucrose sweetness rating; SU-AS: sucrose vs artificial saliva; SU-NO: sucrose vs no solution; AS-NO: artificial saliva vs no solution.	P	0.015	0.422	0.691	0.721	0.932	0.372		
		0.016			-0.093	-0.074		0.33	
	r	0.949			0.706	0.763		0.168	
		0.233	0.162	0.023					.799***
	P	0.338	0.506	0.926					<0.001

AN: anorexia nervosa; BN: bulimia nervosa; CW: comparison women; EDI3-ID: Eating Disorder Inventory 3 Interceptive Deficits; P AS: Artificial saliva pleasantness rating; S AS: Artificial saliva sweetness rating; P 1 M SU: 1 molar sucrose pleasantness rating; S 1M SU: 1 molar sucrose sweetness rating; SU-AS: sucrose vs artificial saliva; SU-NO: sucrose vs no solution; AS-NO: artificial saliva vs no solution.