CASE REPORT

Acute disseminated encephalomyelitis in a case of autoimmune haemolytic anaemia: a rare association

Adrija Hajra,¹ Dhrubajyoti Bandyopadhyay²

¹IPGMER, Kolkata, West Bengal, India

²Department of Accident & Emergency, Lady Hardinge Medical College, Hooghly, West Bengal, India

Correspondence to

Dr Dhrubajyoti Bandyopadhyay, drdhrubajyoti87@gmail.com

Accepted 22 May 2016

SUMMARY

Acute disseminated encephalomyelitis (ADEM) is a demyelinating disease that may occur in a postvaccination condition or as a parainfectious encephalomyelitis. It is almost always monophasic. The underlying pathogenesis of ADEM may include perivascular inflammation, oedema and demyelination in the central nervous system. We present a case of a 15-year-old girl who was diagnosed as having ADEM, as well as detected to be a follow-up case of autoimmune haemolytic anaemia on steroid treatment. She presented with progressive weakness of the right lower limb for the past 4 days. MRI showed multiple subcortical lesions of varying size showing hyperintensities in T2 fluid-attenuated inversion recovery (FLAIR). The patient responded well to steroid therapy. No residual lesion was found on follow-up. Very few cases have been found with this rare association in the literature.

BACKGROUND

Acute disseminated encephalomyelitis (ADEM) is an inflammatory demyelinating illness, characterised by a monophasic course frequently associated with infections (postinfectious) or previous immunisation (postvaccination). ADEM may occur after vaccines for measles, mumps, rubella, rabies, diphtheria, tetanus, polio, influenza, smallpox, Japanese B encephalitis, pertussis or hepatitis B. However, only 0.1–0.2 per 100 000 vaccinated individuals suffer from postvaccination ADEM.¹ They are seen secondary to non-specific upper respiratory tract infections.² We present a case of ADEM, though the patient gave a negative history of any particular infection before the neurological illness. History, clinical features, MRI findings and subsequent improvement with high-dose corticosteroids led us to the diagnosis of ADEM. Paraplegia or quadriplegia, partial or complete, and diminished or losses of reflexes as seen in the myelitic form of ADEM, were absent. Regarding pathogenesis, encephalomyelitis is associated with some immunological mechanisms. Autoimmune haemolytic anaemia (AIHA) is a relatively uncommon disorder. Here, autoantibodies cause the destruction of the body's red blood cells (RBCs).³ There are studies postulating the possibility of a combination of several autoimmune diseases. Our patient, who was a follow-up case of AIHA, developed ADEM without any history of preceding infection or immunisation.

CASE PRESENTATION

A 15-year-old Indian girl, initially diagnosed 2 years prior as having a case of AIHA, presented

with gradually progressive weakness of the right lower limb for the past 4 days. It was acute in onset and gradually progressive. There was no history of any other illness, particularly of infectious origin, before this episode. As she developed additional symptoms such as a headache and blurring of vision, she was subsequently transferred to a higher centre. There were exaggerated deep tendon reflexes in the right lower limb and bilateral extensor plantar response. Within 12 hours of admission, the patient became stuporous.

INVESTIGATIONS

After exclusion of papilloedema, lumbar puncture was performed. An urgent MRI was performed, which revealed multiple subcortical well-marginated hyperintense lesions at the grey–white matter interface (figure 1). Incomplete rim enhancements after administration of intravenous gadodiamide was seen on T1-weighted images (figure 2). In diffusion-weighted image MRI, multiple rounded well-marginated peripherally hyperintense lesions were noted in the grey matter and white matter interface, suggestive of peripheral diffusion restriction of the lesions (figures 3 and 4). Cerebrospinal fluid (CSF) study showed cell count of 40 cells/mm³ with 90% mononuclear cells, protein 55 mg/dL, glucose 45 mg/dL and adenosine deaminase 6.3 U/L. Neither Gram stain nor acid-fast bacilli stain showed microorganisms. No oligoclonal band was present in CSF. Tests for enterovirus, herpes simplex virus (type 1 and type 2), cytomegalovirus, hepatitis virus (A, B, C, D and E), *Cryptococcus*,

Figure 1 MRI of the brain (T2 fluid-attenuated inversion recovery (FLAIR), axial view) shows multiple subcortical well-marginated hyperintense lesions at the grey–white matter interface.

CrossMark

To cite: Hajra A, Bandyopadhyay D. *BMJ Case Rep* Published online: [please include Day Month Year] doi:10.1136/bcr-2016-214440

Figure 2 MRI of the brain (T1-weighted image after intravenous gadodiamide, axial view) shows incomplete rim enhancement after administration of intravenous gadodiamide.

HIV, Epstein-Barr virus, *Mycoplasma pneumoniae* and *Chlamydia pneumoniae* were normal. CSF-PCR for the presence of measles virus was negative. As the patient experienced blurring of vision with normal fundus on examination, she underwent a visual evoked potential (VEP) study, and that suggested a bilateral demyelinating type of optic nerve dysfunction (figures 5 and 6). As previously mentioned, the patient was known to have AIHA. Two years prior, she had been admitted to a local hospital due to jaundice and weakness. On evaluation, unconjugated hyperbilirubinaemia was found along with anaemia. Available clinical notes showed that there was no history suggestive of infection before the development of the anaemia. Available relevant investigations were: haemoglobin (Hb) 6.2 g%, reticulocytes 45% (corrected value 9.2%); total leucocyte count and platelet count were normal. Liver function test showed bilirubin 9.46 mg/dL (normal range 0.3–1.3 mg/dL) with an unconjugated fraction of 8.79 mg/dL (normal range 0.2–0.9 mg/dL). Serum lactate dehydrogenase was 887 U/L

(normal range 115–221 U/L). Serum haptoglobin level was 20 ng/mL. High-performance liquid chromatography revealed no abnormality. Urine routine, as well as microscopic examinations, were within normal limits. Peripheral blood smear showed multiple spherocytes, teardrop cells, reticulocytes and evidence of RBC destruction (figure 7). Bone marrow report showed reactive hyperplasia with the presence of both megaloblastic and microcytic maturation. Maturation failure was noted along with depletion of the iron store. Direct antiglobulin (Coombs test) test was positive for IgG and C3d. The patient was treated with a steroid, of which the dosage was being tapered accordingly. She was on regular follow-up. Her autoimmune profile antinuclear antibody (ANA) and ANA profile were also negative. She had been on a 5 mg maintenance dose of prednisolone daily for the past 2 years, due to AIHA. She had developed ADEM while on the maintenance dose of prednisolone. Her complete blood count report and autoimmune profile were repeated, but no abnormality was detected.

DIFFERENTIAL DIAGNOSIS

As per guidelines provided by the Brighton Collaboration Encephalitis Working Group, our patient was diagnosed as having ADEM. She had a reduced level of consciousness with decreased response to environmental stimuli and extensor plantar response. These are suggestive of encephalitis. She also had motor weakness in the right lower limb and the MRI findings were suggestive of ADEM. All these points guided us to diagnose the case as ADEM.⁴ Multiple sclerosis (MS) was a close differential in this case. Such a presentation may be a clinically isolated syndrome (CIS) that will progress to MS in the future. But 90% of individuals affected with MS and 68% patients with CIS are found to have a positive oligoclonal band in the CSF.⁵ So, a negative oligoclonal band does not rule out these diseases. For this, strict follow-up is necessary in our case.

TREATMENT

After diagnosis, intravenous methylprednisolone was given at 30 mg/kg daily for 5 days. After 72 hours, there was a significant improvement in the patient's clinical condition. A regimen of oral prednisolone 10 mg/day for 2 weeks, and then to 5 mg/day for 2 weeks, was advised after intravenous therapy with methylprednisolone.

Figure 3 (Diffusion-weighted image) Multiple rounded well-margined peripherally hyperintense lesions noted in the grey matter and white matter interface, suggestive of peripheral diffusion restriction of the lesions.

Figure 4 Apparent diffusion coefficient map for figure 3.

OUTCOME AND FOLLOW-UP

After she had recovered from the ADEM, the patient was encouraged to follow-up with ongoing 5 mg oral prednisolone therapy. MRI of the brain (figure 8) and VEP study, performed after 6 months, were normal. The patient continues to be monitored for stabilisation of her Hb level, with the tapering steroid dose for treatment of her AIHA.

DISCUSSION

The incidence of ADEM after viral infections varies with different types of viruses. Postinfection or postvaccination inflammatory reactions in the perivenular region in the central nervous system (CNS) may be explained by a molecular mimicry mechanism.⁶ But an antecedent infection could be identified in 72–77% of paediatric patients.⁷ In the case of adult patients, also, in 50–75% of instances there is a history of preceding infection or vaccination. ADEM may follow a non-specific upper respiratory or gastrointestinal illness.⁸ In these cases, we are unable to detect a particular history of infection. However,

regarding pathogenesis, the structure of some peptides of viral protein or vaccine simulate the structures of myelin basic protein or lipid-protein in CNS. The T lymphocytes sensitised by the peptides attach to the vascular endothelial cells with the help of an adherence factor through blood circulation.

The permeability of the blood–brain barrier increases because of released inflammatory factors, allowing the sensitised T lymphocytes to pass through the blood–brain barrier more quickly. Chemotactic factors cause recruitment of many kinds of lymphocytes to the CNS. Other hypotheses assume that the virus may result in activation or suppression of some clones of T-cells. Previous occult bacterial infection or organ transplantation may be responsible for the pathogenesis of ADEM by changing the immunological status of the affected individual. Association with the human leukocyte antigen (HLA) types has also been found with the occurrence of ADEM.⁶ In summary, the combination of altered vascular permeability and inflammation may cause breakdown of the blood–brain barrier. It allows the release of previously sequestered antigens, and results in a

Figure 5 Visual evoked potential study showing a bilateral demyelinating type of optic nerve dysfunction.

Left Eye	Montage	N75 mS	P100 mS	N135 mS	N75-P100 μ V
1	Oz-Fz	149.3	186.8	263.3	13.6
2	Oz-Fz	154.5	185.3	220.5	14.4
3	Oz-Fz	144.8	176.3	228.0	10.9

Right Eye	Montage	N75 mS	P100 mS	N135 mS	N75-P100 μ V
4	Oz-Fz	139.5	184.5	227.3	14.8
5	Oz-Fz	141.0	186.0	247.5	9.9
6	Oz-Fz	139.5	179.3	244.5	14.2

Figure 6 Visual evoked potential study result.

cell-mediated immune response. The perivenous sleeves of inflammation and demyelination have been postulated as the predominant lesions of ADEM.⁸ The pathology behind these neurological complications is still a matter of debate. Studies suggest that, while some complications are the result of direct viral infection, others may be due to autoimmune mechanisms. Acute disseminated encephalitis and postinfectious encephalomyelitis associated with some viral infections such as measles are immune-mediated phenomena resulting in multifocal demyelination of perivenous white matter. Methylprednisolone pulse therapy seems to be beneficial. In non-responsive cases, intravenous immunoglobulin can be given. Clinicians should follow-up such patients stringently. Proper treatment will indeed reduce possible morbidity or even mortality.⁹ AIHA is a relatively uncommon disorder caused by autoantibodies directed against RBCs. It can be classified as idiopathic or secondary. Other classifications are of warm or cold (cold) haemagglutinin disease (CAD) and paroxysmal cold haemoglobinuria, or a combination thereof, depending on the thermal range of the autoantibody. The occurrence of a mixed type of AIHA is rare, and exceedingly so in children.¹⁰ Therapy for warm AIHA is the use of corticosteroids. They are effective in the majority of cases. The dose should be well monitored and tapered according to the response.³ There are associations of an infective aetiology in the case of secondary AIHA. Apart from common infective aetiologies, we have case reports where AIHA has occurred after *Plasmodium* infection.¹¹ Cases have been reported of AIHA from some secondary causes such as systemic lupus erythematosus, rheumatoid arthritis, scleroderma and ulcerative colitis. Owing to these secondary causes, increased incidence of warm AIHA has been noticed in patients around 40 years of age. Children have a peak incidence in the first 4 years of life. There

is a female predilection (2:1) of AIHA, partially due to its association with autoimmune disease. There may be an association with HLA-A1, B7 and B8.¹² One case report has documented antiphospholipid syndrome with AIHA.¹³ It favours the association of an autoimmune pathology in the case of AIHA. The direct antiglobulin test (DAT) is found to be positive with anti-IgG antisera (and anti-C3d) in the case of warm AIHA, whereas IgM antibodies are found in cold AIHA. False-negative DAT results may be due to IgA autoantibodies. More sensitive techniques can overcome the problem.³ Through this case report, we are trying to focus on whether the occurrence of one autoimmune pathology can be affected by another. Several studies have shown that presence of one autoimmune disease may in future herald another autoimmune disease affecting a different organ system. In about 25% of patients, this can be found. The combination of at least three autoimmune diseases in one patient has been defined as multiple autoimmune syndromes. The pathology of multiple autoimmune disorders is still not very clear. Environmental and genetic factors may be involved. Abnormal humoral or cell-mediated immunity have been described as playing a role. The presence of one autoimmune disease should alert the clinician to watch for another. Continued surveillance for the development of new autoimmune disease should be conducted in predisposed patients. The presence of one autoimmune disease, or a family history of autoimmune diseases, increases the chances of developing additional autoimmune disorders.¹⁴ As there is the association of

Figure 7 Peripheral blood smear showing multiple spherocytes (yellow arrow), teardrop cells (green arrow), reticulocytes (brown arrow) and evidence of red blood cell (RBC) destruction (black arrow). Blue arrow indicates normal RBCs.

Figure 8 Repeat MRI after 6 months (T2 fluid-attenuated inversion recovery (FLAIR) axial view), showing no residual lesion.

familial factors and genetic factors in cases of ADEM as well as AIHA, more studies are required to identify whether there is any relationship between these two autoimmune disorders. We reiterate that our patient was also on 5 mg oral steroid when she presented with this neurological disease. In spite of receiving immunosuppressive therapy, she developed ADEM. Although a history of preceding infection or vaccination was absent, there might be a possibility of her having had a non-specific viral infection. We have to keep close observation on this patient to detect the further course of the disease. There is information in the literature about ADEM after different kinds of infection, for example, Epstein-Barr virus infection, streptococcal infection and *M. pneumoniae* infection.^{6 9 15} But an association of autoimmune disorder is a relatively less-discussed topic. Within 5 years of initial presentation of ADEM, MS has been reported in ~25% of patients. The majority of individuals do not progress beyond 3 months.¹⁶ Our patient showed a normal brain MRI after 6 months; also, her VEP study was normal after 6 months. She has to be followed up further for the development of MS as well as for development of autoimmune disorders affecting other systems.

Learning points

- It is critical to recognise immune-mediated disorders of the central nervous system
- (CNS). These disorders are highly responsive to immunotherapy. Early diagnosis will lead to a better outcome.
- Autoimmune CNS involvement may be associated with other autoimmune systemic diseases, such as autoimmune haemolytic anaemia.
- Follow-up of these patients is essential for detecting the development of other autoimmune disease in such cases.

Acknowledgements The authors acknowledge Dr Deb Kumar Biswas, Postgraduate trainee, Department of Radiology, Calcutta Medical College and Hospital, Kolkata.

Contributors AH treated the patient; and DB prepared the manuscript.

Competing interests None declared.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Machicado JD, Bhagya-Rao B, Davogusto G, *et al.* Acute disseminated encephalomyelitis following seasonal influenza vaccination in an elderly patient. *Clin Vaccine Immunol* 2013;20:1485–6.
- 2 Chowdhary J, Ashraf SM, Khajuria K. Measles with acute disseminated encephalomyelitis (ADEM). *Indian Pediatr* 2009;46:72–4.
- 3 Zanella A, Barcellini W. Treatment of autoimmune hemolytic anemias. *Haematologica* 2014;99:1547–54.
- 4 Sejvar J, Kohl K S, Bilynsky R, *et al.* The Brighton Collaboration Encephalitis Working Group. Encephalitis, myelitis, and acute disseminated encephalomyelitis (ADEM): case definitions and guidelines for collection, analysis, and presentation of immunization safety data. *Vaccine* 2007;25:771–92.
- 5 Dobson R, Ramagopalan S, Davis A, *et al.* Cerebrospinal fluid oligoclonal bands in multiple sclerosis and clinically isolated syndromes. A meta-analysis of prevalence, prognosis and effect of latitude. *J Neurol Neurosurg Psychiatry* 2013;84:909–14.
- 6 Yu YL, Li HF, Xia ZZ, *et al.* Acute disseminated encephalomyelitis following streptococcus pneumoniae meningitis. *HK J Paediatr* 2013;18:37–41.
- 7 Lotze TE, Chadwick DJ. Acute disseminated encephalomyelitis in children: pathogenesis, clinical features and diagnosis. In: Patterson MC, ed. *UpToDate*. 2015. <http://www.uptodate.com/contents/acute-disseminated-encephalomyelitis-in-children-pathogenesis-clinical-features-and-diagnosis> (accessed 27 Dec 2015).
- 8 Waldman AT, Jacobs D. Acute disseminated encephalomyelitis in adults. In: Gonzalez-Scarano F, ed. *UpToDate*. 2015. <http://www.uptodate.com/contents/acute-disseminated-encephalomyelitis-in-adults> (accessed 27 Dec 2015).
- 9 Saeed M, Dabbagh O, Al-Muhaizae M, *et al.* Acute disseminated encephalomyelitis and thrombocytopenia following Epstein-Barr virus infection. *J Coll Physicians Surg Pak* 2014;24:S216–18.
- 10 Reddy V R S, Samayam P, Ravichander B, *et al.* Autoimmune hemolytic anemia: mixed type-a case report. *Indian J Hematol Blood Transfus* 2011;27:107–10.
- 11 Sonani R, Bhatnagar N, Maitrey G. Autoimmune hemolytic anemia in a patient with malaria. *Asian J Transfus Sci* 2013;7:151–2.
- 12 Gehrs B C, Friedberg R C. Autoimmune hemolytic anemia. *Am J Hematol* 2002;69:258–71.
- 13 Karasawa N, Taniguchi Y, Hidaka T, *et al.* Antiphospholipid syndrome with autoimmune hemolytic anemia which mimics thrombotic thrombocytopenic purpura. *Rinsho Ketsueki* 2010;51:275–80.
- 14 Cojocaru M, Cojocaru IM, Silosi I. Multiple autoimmune syndrome. *Maedica (Buchar)* 2010;5:132–4.
- 15 Kong MX, Newman K, Goldenberg R, *et al.* Fatal *Mycoplasma pneumoniae* infection: case report and review of the literature. *North Am J Med Sci* 2012;5:126–30.
- 16 Mahdi N, Abdelmalik PA, Curtis M, *et al.* A case of acute disseminated encephalomyelitis in a middle aged adult. *Case Rep Neurol Med* 2015;2015:601706.

Copyright 2016 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- Submit as many cases as you like
- Enjoy fast sympathetic peer review and rapid publication of accepted articles
- Access all the published articles
- Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow