CORRESPONDENCE

The Treatment of Non-Traumatic Meniscus Lesions: A Systematic Review Comparing Arthroscopic Partial Meniscectomy With Non-Surgical Treatment

by Prof. Dr. med. Wolf Petersen, Dr. med. Andrea Achtnich, Christian Lattermann MD, and Dr. med. Sebastian Kopf in issue 42/2015

PD Dr. med. Ralf Müller-Rath
Berufsverband für Arthroskopie (BVASK e. V.)
Orthopädische Praxisklinik Neuss
mueller-rath@opn-neuss.de

Conflict of interest statement

The author has received speaking honoraria from the companies Arthrex and Aesculap.

Outcome Equivalence

We would like to thank the authors for their comprehensive article (1). The following key aspects should be added:

1. When reporting „300 000 meniscus surgeries“ based on the DRG hospital statistics of the German Federal Statistical Office, it should be taken into consideration that the Federal Statistical Office counts procedures, rather than cases. Since an arthroscopic operation often requires several procedures, each individual case/patient appears many times in the procedure count. Thus, it can be assumed that the actual number of patients operated on primarily due to a meniscus injury is significantly lower.
2. The study by Sihvonen et al. (2) contains a massive selection bias. Over a study period of six years and with the participation of five centers, only 145 patients were enrolled—thus representing a highly-selected patient population. These results therefore cannot be generalized to the entire population of patients with degenerative meniscus damage.
3. The alleged equivalence between arthroscopy and physiotherapy in the study by Katz et al. (3) is obtained only when using the intention-to-treat method, with a crossover rate of 35% from the conservative to the operative group. In an appendix to their work, the authors themselves clarified that using an alternative statistical analysis revealed arthroscopy to be superior to conservative treatment.

Overall, it is incorrect to state that arthroscopy and physiotherapy have outcome equivalence for treating degenerative meniscus damage. Indeed, the authors correctly concluded that arthroscopic partial meniscectomy will still have an important role in the future, and that a significant proportion of patients do not benefit from conservative treatment.

DOI: 10.3238/arztebl.2016.0360a

REFERENCES

1. Petersen W, Achtnich A, Lattermann C, Kopf S: The treatment of non-traumatic meniscus lesions—a systematic review comparing arthroscopic partial meniscectomy with non-surgical treatment. *Dtsch Arztebl Int* 2015; 112: 705–13.
2. Sihvonen R, Paavola M, Malmivaara A, et al.: Arthroscopic partial meniscectomy versus sham surgery for a degenerative meniscal tear. *N Engl J Med* 2013; 369: 2515–24.

Unfounded Concerns

The core statement of this review article (1) can be accepted: the benefit of arthroscopic surgery for degenerative meniscus lesions is questionable unless locking symptoms clearly indicate a mechanical problem in the knee. Therefore, this operation should no longer be considered as standard practice but rather as an exception (2). This conclusion is valid however for osteoarthritis in general, irrespective whether degenerative lesions are present in the meniscus, the cartilage, or both structures (3).

The problem presented by some patients changing treatment during the study (especially if it is in one direction—from the control group to the intervention group) could lead to consideration of treatment strategies, rather than treatment options, in the extreme case.

In the case presented here, having an immediate meniscus resection is compared to undergoing an initial (continued) conservative management strategy, which is followed by a subsequent operation only if necessary.

It does not seem plausible, however, to conclude that arthroscopy is beneficial based on the crossover of some study patients from conservative treatment to arthroscopically resective surgery. Rather, such a treatment crossover firstly shows how deeply rooted the belief in the efficacy of arthroscopy is, and how much effective treatment options are lacking momentarily. Even the statement that „patients with flap tears benefit from arthroscopic partial meniscectomy“ is not based on the available data.

Additionally, the Herrlin study reported that 8 of 13, and not “3 of 13,” switchers had flap tears. (Editor’s note: A correction for this has been published in issue 46/2015 of the German print version; the error had been spotted before the publication of the English language version of the article which is therefore correct).

We believe the concern that a low number of studies “may introduce a systematic bias” is unfounded. More importantly, the strong placebo effects of surgery should be considered. Therefore, if feasible, future studies should perform a sham operation in the comparison group.

DOI: 10.3238/arztebl.2016.0360b

REFERENCES

1. Petersen W, Achtnich A, Lattermann C, Kopf S: The treatment of non-traumatic meniscus lesions: a systematic review comparing arthroscopic partial meniscectomy with non-surgical treatment. *Dtsch Arztebl Int* 2015; 112: 705–13.

2. Institut für Qualität und Wirtschaftlichkeit im Gesundheitswesen (IQWiG): Arthroskopie des Kniegelenks bei Gonarthrose: Abschlussbericht; Auftrag N11-01. (IQWiG-Berichte; Band 211). Köln: IQWiG. www.iqwig.de/download/N11-01_Arthroskopie-des-Kniegelenks-bei-Gonarthrose_Abschlussbericht.pdf [last accessed 26 October 2015].
3. Thorlund JB, Juhl CB, Roos EM, Lohmander LS: Arthroscopic surgery for degenerative knee: systematic review and meta-analysis of benefits and harms. *BMJ* 2015; 350: h2747.

PD Dr. med. Stefan Sauerland, MPH

Sandra Molnar, MSc

PD Dr. med. Stefan Lange

Institut für Qualität und Wirtschaftlichkeit im Gesundheitswesen (IQWiG), Köln
stefan.sauerland@iqwig.de

Conflict of interest statement

The authors declare that no conflict of interest exists.

In Reply:

We thank the authors for their interest in our article (1) and we are happy to comment on the points put forward for discussion.

In our article, we have already mentioned the problem of selection bias in prospectively randomized studies. This problem of course strongly applies—as correctly indicated by Dr. Müller-Rath—to the Sihvonen study (2). We were unable to find a specific “as treated” analysis in the appendix of Katz et al. (3). The crossover rate of 35%, however, reveals that a proportion of patients do not benefit from conservative treatment. The clinical scores for these patients improved only following arthroscopic surgery.

We believe it is more important to again point out that the information in the appendix shows that arthroscopic meniscus surgery is also effective for higher grade stages of osteoarthritis (Kellgren–Lawrance Grade 3).

Overall, we agree with our colleague Dr. Müller-Rath that the available data only superficially show the equivalence of both therapies (operative versus non-operative).

Critical analysis of the scientific data shows that arthroscopic partial meniscectomy still has a role in the treatment of meniscal lesions.

We also thank Dr Sauerland for agreeing with our statements. The studies published to date show that arthroscopic partial meniscectomy can be beneficial. However, the evidence regarding the subset of patients who benefited from this intervention is not as clear as indicated in the comment. Locking symptoms were un-

fortunately not predictive for the success of operative treatment.

The only indication that patients with locking symptoms benefit from a surgical procedure is actually found in the work of Herrlin et al. (4), as clinical experience shows that flap tears provoke significant locking symptoms.

It seems entirely plausible to us to conclude that arthroscopy is beneficial based on the crossover of study patients (of up to 35%) from conservative treatment to arthroscopically resective surgery. Indeed, the clinical scores for these patients improved after surgery.

We agree with the Sauerland group, however, that it is important to keep in mind the strong placebo effects of surgery. This applies in fact also to a sham operation. Therefore, future studies should if possible not incorporate sham operations into the comparison group.

When faced with a lack of success, the threshold for undergoing a second surgery is much higher than that for switching from a non-operative therapy to surgery.

In summary, a critical analysis of prospective randomized studies supports the conclusion that arthroscopic partial meniscectomy still plays an important role. This is true for knee joints with and without signs of arthrosis.

DOI: 10.3238/arztebl.2016.0361

REFERENCES

1. Petersen W, Achtnich A, Lattermann C, Kopf S: The treatment of non-traumatic meniscus lesions: a systematic review comparing arthroscopic partial meniscectomy with non-surgical treatment. *Dtsch Arztebl Int* 2015; 112: 705–13.
2. Sihvonen R, Paavola M, Malmivaara A, et al.: Arthroscopic partial meniscectomy versus sham surgery for a degenerative meniscal tear. *N Engl J Med* 2013; 369: 2515–24.
3. Katz JN, Brophy RH, Chaisson CE, et al.: Surgery versus physical therapy for a meniscal tear and osteoarthritis. *N Engl J Med* 2013; 368: 1675–84. Erratum in: *N Engl J Med* 2013; 15: 683.
4. Herrlin SV, Wange PO, Lapidus G, Hållander M, Werner S, Weidenhielm L: Is arthroscopic surgery beneficial in treating non-traumatic, degenerative medial meniscal tears? A five year follow-up. *Knee Surg Sports Traumatol Arthrosc* 2013; 21: 358–64.

Prof. Dr. med. Wolf Petersen

Klinik für Orthopädie und Unfallchirurgie am

Martin Luther Krankenhaus, Berlin

Wolf.Petersen@pgdiakonie.de

Conflict of interest statement

Prof. Petersen has received royalties for patents with the companies Karl Storz and Otto Bock, consultancy fees from Karl Storz, Otto Bock, and aap Implantate AG, and lecture fees from aap Implantate AG.