using computerized software on a 1:1 basis, stratified for gender. Nine out of 60 (15%) – five from the intervention arm and four from the control arm – were lost to follow-up. The groups were well-balanced at baseline for demographic and clinical variables.

The primary outcome, assessed by independent raters, was psychological distress, measured by GHQ-12 with scores being the total sum across 12 items (possible range 0-36). Other outcomes included: functioning, measured using the 12-item interviewer-administered screener version of the WHODAS 2.0; and post-traumatic stress symptoms, measured using the PTSD Checklist for DSM-5 (PCL-5)⁸, which is a 20-item checklist corresponding to the twenty DSM-5 PTSD symptoms in the last week, with items rated on a 0-4 scale (possible range 0-80).

The intervention had high uptake, with 22/30 (73%) completing all sessions. The intervention arm showed improvement in functioning (mean WHODAS 2.0 scores reduced from 17.7 ± 9.2 to 6.6 ± 6.1 vs. 17.0 ± 10.5 to 11.3 ± 10.4 in controls) and in post-traumatic stress symptoms (mean PCL-5 scores reduced from 34.2 ± 20.1 to 9.8 ± 9.1 vs. 32.3 ± 17.1 to 19.5 ± 18.5 in controls). Due to skewed distribution and variance heterogeneity of the outcome variable, log-linear regression was carried out. After adjustment of baseline scores, the results showed a reduction of 90% in geometric mean within the intervention group (95% CI: 90.4%-91.7%, $p=0.04$) in WHODAS 2.0 scores and a reduction of 92% (95% CI: 91.2%-92.3%, $p=0.02$) in post-traumatic stress symptoms. There was no significant change in GHQ-12 scores. On qualitative evaluation of a sub-sample of participants and primary care staff, we found that the intervention was perceived as useful, and was successfully integrated into primary care centres.

As this was a pilot study with a small sample size, recruited through primary care physician referral, and no power calculations were carried out, the findings and their generalizability warrant a cautious interpretation. However, a successful conduction in challenging settings, with adequate enrolment rate, a low drop-out, and balanced randomization provides evi-

dence that RCTs are feasible in such settings. The intervention delivery through non-specialists with no prior mental health care experience and the encouraging results demonstrate the feasibility of the task shifting approach, and are consistent with previous reports^{9,10}. The results of this pilot study should encourage further adaptation and large-scale fully-powered RCTs of this new, transdiagnostic psychological intervention⁴.

Atif Rahman^{1,2}, Naila Riaz³, Katie S. Dawson⁴, Syed Usman Hamdani², Anna Chiumento¹, Marit Sijbrandij⁵, Fareed Minhas⁶, Richard A. Bryant⁴, Khalid Saeed⁷, Mark van Ommeren⁸, Saeed Farooq³

¹University of Liverpool, Liverpool, UK; ²Human Development Research Foundation, Islamabad, Pakistan; ³Lady Reading Hospital, Peshawar, Pakistan; ⁴University of New South Wales, Sydney, Australia; ⁵VU University Amsterdam, Amsterdam, The Netherlands; ⁶Institute of Psychiatry, Rawalpindi, Pakistan; ⁷Mental Health and Substance Abuse Unit, Regional Office for the Eastern Mediterranean Region, World Health Organization, Cairo, Egypt; ⁸Department of Mental Health and Substance Abuse, World Health Organization, Geneva, Switzerland

The authors alone are responsible for the views expressed in this letter, which do not necessarily represent the views, decisions or policies of the institutions with which they are affiliated. The authors are grateful to M. Dherani and D. Wang for statistical advice. M. van Ommeren and S. Farooq are joint last authors. The study was funded by the Office of Foreign Disaster Assistance.

1. Steel Z, Chey T, Silove D et al. JAMA 2009;302:537-49.
2. Van Ommeren M, Saxena S, Saraceno B. BMJ 2005;330:1160-1.
3. World Health Organization. mhGAP Mental Health Gap Action Programme. Geneva: World Health Organization, 2008.
4. Dawson K, Bryant R, Harper M et al. World Psychiatry 2015;14:354-8.
5. Murray LK, Dorsey S, Bolton P et al. Int J Ment Health Syst 2011;5:30.
6. Goldberg DW. A user's guide to the General Health Questionnaire. Windsor: NFER-Nelson, 1988.
7. World Health Organization. Measuring health and disability: manual for WHO Disability Assessment Schedule WHODAS 2.0. Geneva: World Health Organization, 2010.
8. Weathers FW, Litz BT, Keane TM et al. The PTSD Checklist for DSM-5 (PCL-5). www.ptsd.va.gov.
9. van Ginneken N, Tharyan P, Lewin S et al. Cochrane Database Syst Rev 2013;11:CD009149.
10. Rahman A, Fisher J, Bower P et al. Bull World Health Organ 2013;91:593-601.

DOI:10.1002/wps.20312

Treating post-traumatic stress disorder by resource activation in Cambodia

There is a need for effective, low-threshold psychotherapeutic treatments in post-conflict settings¹. However, systematic outcome research on site is still extremely rare. To address this problem we integrated rigorous research procedures into a humanitarian program, the so called Mekong Project, and conducted a randomized controlled trial for the treatment of post-traumatic stress disorder (PTSD) in Cambodia. In short, the Mekong Project aims at establishing independent psychotherapeutic services in several Southeast Asian countries via the systematic training of local health professionals and offering free of charge psychological help to traumatized civilians.

Cambodia is one of the least developed countries in Asia, facing many challenges (e.g., poor standards of health and

education, rural exodus, and political instability). Mental health morbidity in Cambodia is high. It has been found that 53.4% of the Cambodian population suffer from a mental disorder, with anxiety and PTSD being the most frequent (40.0% and 28.4% respectively)². Thus, although some stability has returned to the country during the past decades, there are urgent mental health care needs, including the need for individualized psychiatric services.

Our aim was to test the efficacy of a non-confrontational psychotherapeutic treatment for PTSD. The therapy includes two main treatment principles described in treatment manuals: resource-oriented trauma therapy and resource installation with eye movement desensitization and reprocessing

(EMDR) (short: ROTATE). ROTATE aims at strengthening resilience and coping capacities by activating positive personal resources, and largely draws on psychodynamic principles of the therapeutic relationship. It includes a variety of imaginative resource-activating methods^{3,4} as well as resource development and installation, an EMDR technique aiming at systematically developing and anchoring resources using alternating bilateral stimulation⁵. ROTATE has several advantages: a) it can be safely applied even to complex trauma conditions, with no major side effects being observed so far; b) instead of solely focusing on PTSD symptoms, it also considers the mental comorbidities typically found in these clients, notably depression and anxiety; c) it is especially suitable for clients from non-Western countries, as traditional healing resources like mindfulness strategies can be integrated in an overall framework of resource activation; d) its basic elements can easily be taught, even to paraprofessionals.

Our trial was carried out in cooperation with the Royal University of Phnom Penh and was located in Phnom Penh City and the nearby Kandal Province. Help-seeking outpatients screening positively for PTSD (PTSD Check List ≥ 44 ⁶) were eligible for inclusion. We allowed for comorbid mental health disorders except for psychosis, organic brain disorder, cognitive impairment, dementia, acute suicidality, and acute need for treatment.

Overall, 800 patients were screened for eligibility, of whom 86 (mean age 27 years, 61% female) fulfilled the selection criteria and were randomly assigned to either 5 weekly sessions of ROTATE (N=53) or a 5-week waiting list control group (N=33). Symptoms were measured before and after the intervention (or waiting period). Assessments were performed via personal interview by an investigator blind to treatment allocation. All patients in the control group were offered treatment after the end of the waiting period. The primary outcome was PTSD symptom change on the Indochinese version of the Harvard Trauma Questionnaire (HTQ)⁷. The PTSD scale of the HTQ includes 16 items reflecting the DSM-IV criteria for PTSD. Secondary outcomes included depression, anxiety and social functioning. All applied instruments have been validated for the Cambodian population⁷. The therapy was provided by six Cambodian psychologists who had completed a 3-year course in trauma therapy as part of the Mekong Project.

Based on previous findings of psychological therapies for PTSD⁸, we expected ROTATE to be superior to waiting list with a between group effect size of at least $d=0.65$ on the primary outcome. To detect this difference with a power of 0.80 at $\alpha=0.05$, 2-sided test, 2×40 patients were required. Unfortunately, the concept of randomization, especially being randomized to a waiting list, was very difficult for some clients. As a consequence, randomization failed in 38 patients, leading to an unbalanced allocation ratio (1.6:1), with an overrepresentation of patients randomized to treatment. The trial stopped when the necessary sample size to achieve a power of 0.80 was reached. Data were analyzed by general linear regression models, controlling for baseline symptom severity. The drop-out

rate during the intervention was very low (N=2, one in each group), thus only completer data were analyzed (N=84).

Most frequent types of trauma were traffic accidents (24%), domestic violence (23%) and sexual abuse (16%). Patients receiving ROTATE showed significant reductions in PTSD symptoms compared to the waiting list (baseline adjusted means post-treatment: 1.39, 95% CI: 1.23-1.54 for ROTATE, and 2.86, 95% CI: 2.66-3.06 for waiting list, $p<0.00001$). The between-group effect size was large ($d=2.59$). The within-group effect size was also large for ROTATE ($d=4.43$), while it was moderate in the control group ($d=0.52$). No harms were reported.

We conclude that a treatment focusing on stabilization rather than confrontation, by establishing a secure patient-therapist relationship, applying stabilization techniques, and putting an emphasis on a patient's own resources, significantly reduced symptoms of PTSD in comparison to a waiting list.

The strengths of our study are the following: a) it was conducted on site by local psychologists, which meant that communication between therapists and patients was natural and no interpreters were needed; b) therapists and patients had similar cultural backgrounds, so that culture specific interpretations of symptoms could be taken into account, a factor that has been identified as vital in the therapeutic work with Cambodian patients⁹; c) local psychologists were trained in ROTATE, which is expected to facilitate patient access to a psychological treatment in a country struggling with insufficient mental health care.

Conducting a randomized controlled trial in a developing country is challenging. Nevertheless, we were able to show that the implementation of such a trial was possible and that this specific form of trauma therapy was well accepted by therapists and patients. Our results are preliminary but promising. Further research is required to corroborate the findings.

Christiane Steinert¹, Peter J. Bumke², Rosa L. Hollekamp¹, Astrid Larisch¹, Falk Leichenring¹, Helga Mattheß³, Sek Sisokhom⁴, Ute Sodemann², Markus Stingl¹, Ret Thearom⁴, Hana Vojtova⁵, Wolfgang Wöller⁶, Johannes Kruse^{1,7}

¹Clinic for Psychosomatic Medicine and Psychotherapy, University of Giessen, Giessen, Germany; ²Trauma Aid Germany, Berlin, Germany; ³Psychotraumatology Institute Europe, Duisburg, Germany; ⁴Department of Psychology, Royal University of Phnom Penh, Phnom Penh, Cambodia; ⁵Slovak Institute for Psychotraumatology and EMDR, Trenčin, Slovakia, and Center for Neuropsychiatric Research of Traumatic Stress, Charles University, Prague, Czech Republic; ⁶Hospital for Psychosomatic Medicine and Psychotherapy, Rhein-Klinik, Bad Honnef, Germany; ⁷Clinic for Psychosomatic Medicine and Psychotherapy, University of Marburg, Marburg, Germany

This study was carried out as part of an ODA Project (Official Development Assistance, Project 2010.1572.6, Program for Trauma Treatment in Thailand, Burma, Cambodia and Indonesia, Mekong Projekt) and was financially supported by the German Federal Ministry for Economic Cooperation and Development and the German branch of EMDR-Europe. The authors thank all involved patients for their participation in the study. The therapists responsible for treating the patients were O. Plaktin, S. Samchet, Y. Sotheary, P. Bunna, T. Nary, and E. Nil.

1. Tol WA, Barbui C, Galappatti A et al. *Lancet* 2011;378:1581-91.
2. de Jong JT, Komproe IH, Van Ommeren M. *Lancet* 2003;361:2128-30.
3. Reddemann L. Psychodynamic imaginative trauma therapy, PITT – Manual. Stuttgart: Klett-Cotta, 2011.

4. Wöller W, Leichsenring F, Leweke F et al. *Bull Menninger Clin* 2012;76:69-93.
5. Korn DL, Leeds AM. *J Clin Psychol* 2002;58:1465-87.
6. Blanchard EB, Jones-Alexander J, Buckley TC et al. *Behav Res Ther* 1996; 34:669-73.
7. Mollica RF, Caspi-Yavin Y, Bollini P et al. *J Nerv Ment Dis* 1992;180:111-6.
8. Bisson J, Andrew M. *Cochrane Database Syst Rev* 2007;CD003388.
9. Hinton DE, Otto MW. *Cogn Behav Pract* 2006;13:249-60. DOI:10.1002/wps.20303

High burden of subthreshold DSM-5 post-traumatic stress disorder in U.S. military veterans

A substantial proportion of individuals worldwide develop post-traumatic stress disorder (PTSD) following exposure to traumatic events¹⁻³. Although the epidemiology of PTSD has been widely studied¹⁻³, fewer studies have examined subthreshold PTSD, defined as experiencing clinically significant symptoms of PTSD but not meeting full diagnostic criteria for the disorder. With the field of psychiatry increasingly moving towards a dimensional perspective of mental disorders, it is important to understand the burden of subthreshold manifestations of these disorders.

The lifetime prevalence of subthreshold PTSD has ranged from 3.6 to 25.6%^{2,4-6}. While not a formal diagnosis, subthreshold PTSD is associated with elevated rates of comorbid psychiatric disorders, suicidality, and physical health problems compared to trauma-exposed individuals without subthreshold or threshold PTSD^{2,4-6}. To date, however, only two studies have examined the epidemiology of subthreshold PTSD as defined using the DSM-5. The first analyzed data from the World Health Organization World Mental Health Surveys and found that the prevalence of subthreshold PTSD ranged from 0.7 to 4.6%, depending on the definition used. Further, individuals with subthreshold PTSD were 2.5-5 times more likely to have a comorbid mood or anxiety disorder compared to trauma-exposed controls⁷. This study was limited by the operationalization of PTSD, which was derived from a DSM-IV module and did not include the new DSM-5 symptoms. The second study of a national sample of Vietnam veterans found that the prevalence of current subthreshold PTSD ranged from 1.9 to 5.7%, and that the comorbidity between DSM-5 subthreshold PTSD and comorbid disorders ranged from 0.7 to 30.9%⁸. While these studies provide important insight into the prevalence and correlates of subthreshold DSM-5 PTSD, additional population-based data are needed to better understand the burden of this condition.

We analyzed data from the National Health and Resilience in Veterans Study (NHRVS), a contemporary, nationally representative cohort of U.S. military veterans, to examine the prevalence and clinical correlates of DSM-5 subthreshold PTSD. The NHRVS, conducted in 2013, surveyed 1,484 veterans aged 20+. The sample was ascertained from KnowledgePanel, a nationally representative survey research panel representing approximately 98% of U.S. households. Post-stratification weights were applied to permit generalizability of results to the U.S. veteran population. Study constructs were assessed with the following tools: Trauma History Screen, PTSD Checklist for DSM-5 (PCL-5)⁹, Mini International

Neuropsychiatric Interview and Patient Health Questionnaire-4 for lifetime and current psychopathology, respectively, Fagerström Test for Nicotine Dependence, and Short Form-8 (SF-8) Health Survey for mental and physical functioning¹⁰.

Lifetime PCL-5 responses were used to create a three-group variable: a) no/low PTSD symptoms (defined as endorsement of ≤ 1 PTSD criteria B-E at a severity of "moderate" or higher); b) subthreshold DSM-5 PTSD (defined as endorsement of 2 or 3 B-E criteria, or all 4 B-E criteria but not 1 month symptom duration and/or functional impairment); and c) probable lifetime DSM-5 PTSD (defined as meeting criteria A-G for PTSD). A comparable three-level variable was created for past-month PTSD symptoms, with a score ≥ 38 on the PCL-5 distinguishing between subthreshold and probable PTSD in the absence of past-month symptom duration and functional impairment assessment in the NHRVS. Weighted prevalence of lifetime and past-month subthreshold DSM-5 PTSD was computed in the full sample ($N=1,478$; 6 subjects had missing data). Other analyses were conducted in only trauma-exposed veterans ($N=1,268$). Logistic regression and multivariable analyses of covariance were conducted to examine associations of probable and subthreshold PTSD with comorbid psychiatric disorders and SF-8 scores. Analyses were adjusted for socio-demographic variables, combat veteran status, number of lifetime traumas, and any lifetime mental disorder.

The lifetime and past-month prevalence of subthreshold PTSD was 22.1% and 13.5%, respectively, and higher than the prevalence of lifetime (8.0%) and past-month (4.5%) probable PTSD. The prevalence of lifetime subthreshold PTSD was higher in women than in men (30.3% vs. 21.2%, $X^2=10.3$, $p=0.006$) and, although the prevalence of lifetime probable PTSD decreased across age groups (20.8% in 18-34 year olds to 1.9% in 75+ year olds), the prevalence of subthreshold PTSD remained relatively stable across all but the 75+ age group (21.1% to 26.6%).

Lifetime subthreshold PTSD was associated with a greater likelihood of all lifetime (i.e., major depressive, social anxiety, alcohol and drug use disorders) and current (i.e., major depressive and generalized anxiety disorders, suicidal ideation) psychiatric outcomes, except nicotine dependence, relative to veterans reporting no/low symptoms (adjusted odds ratio, AOR range from 1.7 for lifetime alcohol use disorder to 4.9 for current generalized anxiety disorder). Veterans with probable PTSD had a greater likelihood of all outcomes relative to veterans with no/low symptoms, and these associations were numerically larger in magnitude relative to the subthreshold PTSD group (AOR range