Published in final edited form as:

J Biomech. 2016 May 3; 49(7): 1214–1220. doi:10.1016/j.jbiomech.2016.02.029.

Do Mechanical Strain and TNF- α Interact To Amplify Pro-Inflammatory Cytokine Production in Human Annulus Fibrosus Cells?

Morakot Likhitpanichkul¹, Olivia M Torre¹, Jadry Gruen¹, Benjamin A Walter^{1,2}, Andrew C Hecht¹, and James C Iatridis^{1,*}

¹Leni and Peter W. May Department of Orthopedics, Box 1188, Icahn School of Medicine at Mount Sinai, New York, NY 10029

²Department of Biomedical Engineering, The City College of New York, New York NY

Abstract

During intervertebral disc (IVD) injury and degeneration, annulus fibrosus (AF) cells experience large mechanical strains in a pro-inflammatory milieu. We hypothesized that TNF- α , an initiator of IVD inflammation, modifies AF cell mechanobiology via cytoskeletal changes, and interacts with mechanical strain to enhance pro-inflammatory cytokine production. Human AF cells (N=5, Thompson grades 2–4) were stretched uniaxially on collagen-I coated chambers to 0%, 5% (physiological) or 15% (pathologic) strains at 0.5 Hz for 24 hours under hypoxic conditions with or without TNF- α (10 ng/mL). AF cells were treated with anti-TNF- α and anti-IL-6. ELISA assessed IL-1 β , IL-6, and IL-8 production and immunocytochemistry measured F-actin, vinculin and α -tubulin in AF cells. TNF- α significantly increased AF cell proinflammatory cytokine production compared to basal conditions (IL-1 β : 2.0 $\pm$ 1.4 to 84.0 $\pm$ 77.3, IL-6: 10.6 $\pm$ 9.9 to 280.9 $\pm$ 214.1, IL-8: 23.9 $\pm$ 26.0 to 5125.1 $\pm$ 4170.8 pg/ml for basal and TNF- α treatment, respectively) as expected, but mechanical strain did not. Pathologic strain in combination with TNF- α increased IL-1 β , and IL-8 but not IL-6 production of AF cells. TNF- α treatment altered F-actin and α -tubulin in AF cells, suggestive of altered cytoskeletal stiffness. Anti-TNF- α (infliximab) significantly inhibited pro-inflammatory cytokine production while anti-IL-6 (atlizumab) did not. In conclusion, TNF- α altered AF cell mechanobiology with cytoskeletal remodeling that potentially sensitized AF cells to mechanical strain and increased TNF- α -induced pro-inflammatory cytokine production. Results suggest an interaction between TNF- α and mechanical strain and future mechanistic studies are required to validate these observations.

*Corresponding Author: James C. Iatridis, Ph.D., Icahn School of Medicine at Mount Sinai, Department of Orthopaedics, Box 1188, One Gustave Levy Place, New York, NY 10029, james.iatridis@mssm.edu, Phone: 212-241-1517.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Conflict of Interest Statement

The authors declare that they have no competing interests.

Keywords

Annulus Fibrosus; Intervertebral Disc Degeneration; TNF- α ; Mechanical Stretch; Mechanotransduction; Infliximab

Introduction

Intervertebral disc (IVD) disorders contribute to back pain which is a leading cause of global disability (Cheung et al., 2012; Jacobs, 2011; Vos et al., 2012). Painful IVD degeneration involves complex interactions between mechanical loading and inflammation yet few studies investigate these interactions. For example, excessive spinal loads that induce large dynamic bending moments and asymmetric loads are predictive of spinal functional impairment and pain (Marras et al., 2010). Annulus fibrosus (AF) defects and delamination induce strain 'concentrations', cell death, and pro-inflammatory cytokine production (Adams et al., 2015; Korecki et al., 2008; Michalek et al., 2010; Ulrich et al., 2007; Walter et al., 2011). In contrast, dynamic loading is considered healthy since it can enhance IVD matrix production; although this healthy response to load is altered with degeneration perhaps because dynamic compression facilitates transport of pro-inflammatory cytokines into healthy IVDs where they can persist and induce biomechanical changes (Sowa et al., 2012; Stokes and Iatridis, 2004; Wuertz et al., 2009; Walter et al., 2015a).

TNF- α and IL-1 β are key pro-inflammatory cytokines that are increased in IVD degeneration and known to be produced by cells of the IVD and immune system (Allen et al., 2011; Le Maitre et al., 2007; Olmarker and Rydevik, 2001; Risbud and Shapiro, 2014; Seguin et al., 2005). TNF- α can stimulate the production of pro-inflammatory cytokines, matrix degrading enzymes, and pain mediators (Millward-Sadler et al., 2009; Purmessur et al., 2013; Seguin et al., 2005; Weiler et al., 2005). TNF- α is also implicated in painful spine conditions with infliximab and other TNF- α inhibitors showing improved pain and behavioral responses in clinical trials and animal studies (Allen et al., 2011; Likhitpanichkul et al., 2015; Risbud and Shapiro, 2014; Shamji et al., 2010; Nakamae et al., 2011). However, clinical trial results using infliximab are mixed possibly due to inadequate dosing or delivery (Risbud and Shapiro, 2014) highlighting a need for improved knowledge of how TNF- α affects IVD cells.

Physiological strains stimulated anti-catabolic effects on AF cells in vitro while pathological strains resulted in detrimental responses (Gilbert et al., 2010; Sowa et al., 2011; Sowa et al., 2012; Gawri et al., 2014). Pathologically high mechanical strain on AF cells exposed to IL-1 β induced greater catabolic gene expression compared to mechanical strain alone (Sowa et al., 2011). The roles that healthy and pathological strains play on AF cell pro-inflammatory response and cytoskeleton structure are unknown and improved knowledge of IVD cell mechanobiology under pro-inflammatory conditions may provide new insights on pathophysiology.

The objectives of this study were: 1) to investigate the interaction between TNF- α and tensile strain levels on human AF cells in vitro, with the hypothesis that TNF- α treatment interacts with pathological strains to amplify down-stream pro-inflammatory cytokines

IL-1 β , IL-6, and IL-8 via cytoskeletal changes; and 2) to test the effectiveness of infliximab, an anti-TNF- α drug, and atilizumab, an anti-IL-6 drug, on human AF cells with the hypothesis that infliximab is more effective than atilizumab in inhibiting the interacting effects of TNF- α and high strain. Cells were subjected to 5% and 15% strain to represent physiological and pathological strains, as previously defined for AF cells (Gilbert et al., 2011; Rannou et al., 2000; Sowa et al., 2011). Pro-inflammatory cytokines and cytoskeletal changes are our primary output measurements. In addition to TNF- α and IL-1 β , we measured IL-8 levels and IL-6. IL-8 is associated with the recruitment of immune cells including macrophages, which can further intensify the matrix breakdown and result in granulation tissue formation (Burke et al., 2002; Kim et al., 2008). IL-6 is up regulated in IVD injury, degeneration and sciatica (Andrade et al., 2013; Ulrich et al., 2007; Wuertz and Haglund, 2013), and has been considered a clinical target since it can enhance the effects of TNF- α and IL-1 β (Studer et al., 2011).

Methods

Primary human AF cells were isolated from 5 human IVDs from surgery and autopsy (mean $\pm$ SD age: 50 $\pm$ 16 years) with degeneration grades 2–4 (Thompson et al., 1990). For cell isolation outer AF tissue was macroscopically dissected, minced and digested for 1 hour in 2 g/L pronase (Fisher-Scientific, Waltham, MA) and 4 hours in collagenase type I (Fisher-Scientific, Waltham, MA) in high glucose Dulbecco's Modified Eagles Media (DMEM). The cell suspension was filtered (100 μ m) and cell pellets were washed with PBS. To accelerate proliferation, cells were expanded under normoxic conditions (37°C, 5% CO₂, 20% O₂) in high glucose DMEM (Fisher-Scientific, Waltham, MA), 10% FBS (Fisher-Scientific, Waltham, MA), 50 μ g/mL ascorbic acid (Fisher-Scientific, Waltham, MA), 1% penicillin/streptomycin (Fisher-Scientific, Waltham, MA), 2 μ L/mL primocin (Invivogen, San Diego, CA), at 37°C, 5% CO₂ with media changes each 3–4 days. Experimental media conditions better simulated IVD physiologic conditions by using hypoxia (37°C, 5% CO₂, 5% O₂) as well as serum-free media, high glucose DMEM, 1% ITS (Fisher-Scientific, Waltham, MA), 50 μ g/mL Ascorbic acid, 1% Pen/Strep, and 2 μ L/mL primocin. For experiments, AF cells (passage 1–2; 100,000 cells/mL) were seeded onto silicone-membrane chambers (STREX, B-Bridge, Cupertino, CA) coated with rat-tail collagen I (BD Biosciences, San Jose, CA) for a one day incubation prior to stretch to enable cell confluency and equilibration under experimental conditions.

For TNF- α treatment, 10 ng/mL TNF- α (Fisher-Scientific, Waltham, MA) was added to serum-free media immediately prior to incubation. This TNF- α concentration was shown to increase proinflammatory cytokine and matrix degrading enzyme production in IVD cells (Millward-Sadler et al., 2009; Walter et al., 2015b). For mechanical treatment, Strex chambers were exposed to physiological (5%) or pathological (15%) cyclic tensile uniaxial strain at a 0.5 Hz frequency for 24 hours, and controls were unstrained.

Enzyme linked immunosorbant assay (ELISA) specific for human TNF- α , IL-1 β , IL-6, and IL-8 (MSD N45025B-1, Rockville, MD) assessed concentrations of pro-inflammatory cytokines within culture medium 24 hours after treatment.

Immunocytochemistry was performed on cells 24 hours after initiation of treatments to determine acute effects on F-actin, vinculin, and α -tubulin (N=3). AF cells were fixed using 16% paraformaldehyde, and stained for F-actin, α -tubulin, and vinculin (F-actin:Phalloidin-TRITC: Sigma Cat#P1951, 1:100; α -tubulin: α -tubulin-Alexa Flour 488: eBioScience Cat#53-4502-80, 1:50; vinculin:mouse anti-vinculin: Sigma Cat#V4505, 1:100; Alexa Flour 488 goat anti-mouse: Life Technologies Cat#A1001, 1:200). Viability was determined with live/dead staining (2 μ L/mL calcein, 1 μ L/mL ethidium homodimer, 1 μ L/5mL Hoechst). Apoptosis was measured by caspase-3 activity (rabbit cleaved caspase-3:Cell signaling Cat# mAB9664, 1:100; Alexa Flour 594 goat anti-rabbit:Invitrogen Cat#A-11037, 1:1000). Imaging was performed using confocal (Leica SP5 DM, Wetzlar, Germany) and epifluorescence imaging (AxioImager Z1, Zeiss, Jena, Germany).

Anti-inflammatory treated cells were exposed to 10 ng/mL TNF- α and 1000 ng/mL infliximab (anti-TNF- α) or 1000 ng/mL anti-IL-6 added to fresh serum-free media immediately prior to incubation or application of tensile strain. Infliximab and atilizumab doses were selected based on pilot 3 day dose response studies showing largest effect with no observed cell death.

Cytokine concentrations were normalized to paired controls and compared between treatment groups. Student's paired t-tests were performed to compare cytokine concentrations in response to unstrained basal and TNF- α treatment (N=5). One-way ANOVA with Tukey's post-hoc tests for multiple comparisons were performed to compare cytokine concentrations between 0%, 5%, and 15% strains in the presence of TNF- α (N=5). $p < 0.05$ was considered significant.

Results

TNF- α treatment significantly increased IL-1 β , IL-6 and IL-8 concentrations (IL-1 β : 84.15 ± 77.3 , IL-6: 280.9 ± 214.1 , IL-8: 5125.1 ± 4170.8 pg/ml) compared to basal media, which had low concentrations (IL-1 β : 1.9 ± 1.4 , IL-6: 10.6 ± 9.9 , IL-8: 23.9 ± 26.0 pg/ml) (Figure 1). Mechanical strain at 5% and 15% showed no significant effect on pro-inflammatory cytokine production compared to unstrained groups. Combined TNF- α treatment with 15% strain showed a significant increase in IL-1 β and IL-8 concentrations at 15% compared to 5% strain (Figure 2).

TNF- α treatment on AF cells increased F-actin intensity compared to basal, however no changes in vinculin staining were apparent (Figure 3). The α -tubulin staining demonstrated a more diffusely connected microtubule network for the TNF- α treated cells compared to basal. In basal medium, no apparent differences in F-actin or vinculin staining of AF cells at 5% and 15% strain were observed compared to unstrained (Figure 4). Cells subjected to TNF- α and 15% strain exhibited cytoskeletal changes similar to TNF- α alone (Figure 5). Cells subjected TNF- α and 15% strain had increased cell roundedness, however, no staining for dead cells or cleaved caspase-3 was observed for any groups.

Infliximab non-significantly decreased IL-1 β and IL-8 levels compared to TNF- α -stimulated AF cells undergoing 15% strain, and significantly decreased IL-1 β and IL-8 levels compared to anti-IL-6 treatment (Figure 6).

Discussion

The AF cell niche changes following injury and degeneration to create large deformations and defects that expose AF cells to high levels of pro-inflammatory cytokines and pathological strains. The most important findings were that human AF cells exposed to TNF- α responded to pathological strains with increased IL-1 β and IL-8 production and a modified AF cytoskeletal network with enhanced stress fiber formation from actin polymerization and disrupted microtubule network. Results suggested that TNF- α induced cytoskeletal changes in human AF cells suggestive of enhanced mechanosensitivity, although this hypothesis requires further testing. Anti-TNF- α treatment inhibited pro-inflammatory cytokine production compared to anti-IL-6, suggesting anti-inflammatory treatments with appropriate targets and dose can inhibit these effects.

Little data exists on AF cytoskeletal changes to pro-inflammatory cytokines. TNF- α increased chondrocyte cell stiffness due to increases in F-actin staining (Chen et al., 2015) that was similar to our observations and together suggest TNF- α induced AF cell stiffening could be responsible for heightened AF cell mechanosensitivity. TNF- α and lipopolysaccharide induced permanent alterations in nucleus pulposus cytoskeletal mechanics (Maidhof et al., 2014), suggesting TNF- α -induced cytoskeletal changes may be generalizable to other pro-inflammatory conditions and long-lasting; however it is impossible to compare these NP cell-volume and hydraulic permeability values to cytoskeletal measurements in this study. Similar TNF- α induced cytoskeletal changes including re-organization of F-actin (Koukouritaki et al., 1999; Petrache et al., 2001) and microtubules (Petrache et al., 2003) was observed in other cell types. Disruption to microtubule networks, as observed in this study, could reduce uniformity of the load distribution within the cytoskeleton leading to higher stress concentration at the connection with the nucleus (Maniotis et al., 1997), providing a counter-hypothesis to cell-stiffening as a cause for heightened mechanosensitivity. Vinculin, an intracellular attachment protein linking integrins on the cell membrane to the F-actin cytoskeleton, showed no notable changes in response to TNF- α , suggesting limited changes in vinculin-dependent cell matrix connectivity although it is possible that other cell adhesion proteins may be affected by TNF- α . In the context of the literature, the alterations in cytoskeletal structures observed here support the concept that heightened mechanosensitivity of AF cells following TNF- α treatment resulted from AF cell stiffening. Future studies are required to validate these observations with quantitative data and to mechanistically test which cytoskeletal structures are responsible for these observed effects.

AF cell strain is known to interact with pro-inflammatory cytokines, and the current results help explain the finding that AF cells obtained from a rabbit IVD degeneration model demonstrated enhanced pro-inflammatory responses to mechanical and inflammatory stimuli (Sowa et al., 2012). Specifically, AF cells in degenerated rabbit IVDs may have been chronically exposed to TNF- α resulting in heightened mechanosensitivity. Low magnitudes

but not high magnitudes of strain had a somewhat protective effect of a pro-inflammatory challenge with IL-1 β (Sowa et al., 2011). Excessive pro-inflammatory cytokine exposure is particularly harmful to the IVD due to its low cellularity and limited repair potential (Kandel et al., 2008; Zhu et al., 2012). For example, IVDs in organ culture subjected to TNF- α resulted in cell senescence, irrecoverable matrix changes with proteoglycan degradation, a catabolic shift in gene expression, and biomechanical alterations (Purmessur et al., 2013; Walter et al., 2015a; Walter et al., 2014). The current results are consistent with the literature that pro-inflammatory cytokine production by IVD cells are likely to be an important contributor to the inflamed environment associated with IVD degeneration with the potential to permanently alter IVD cell mechanobiology.

Anti-TNF- α was more effective than anti-IL-6 as it inhibited direct and indirect effects of TNF- α on the AF cell pro-inflammatory response. Infliximab inhibits pro-inflammatory cytokine production by binding to and sequestering extracellular TNF- α . Atlizumab targets the IL-6 receptor and was selected to see if autocrine IL-6 signaling contributed to pro-inflammatory cytokine production since TNF- α and IL-6 were previously shown to act synergistically (Studer et al., 2011). Atlizumab did not have an effect in our cell culture studies suggesting that few interactions between TNF- α and IL-6 were present in our system. An alternate anti-IL-6 drug capable of binding and sequestering IL-6 or greater doses may have larger effects.

Future mechanistic studies with additional pharmacologic blocking and a larger set of inflammatory outcome measures would enhance understanding of the complex interactions between inflammation and mechanobiology. Results show changes in actin fibers and microtubule network rearrangement after 24 hours justifying this relevant time point for measurements, although additional timepoints of cytoskeletal analyses with more anti-inflammatory treatments would allow broader interpretations. A greater sample size is also warranted to address the large human variance and validate the observed findings.

Acknowledgments

Research reported in this publication was supported by the National Institute of Arthritis and Musculoskeletal and Skin Diseases of the National Institutes of Health under Award Numbers R01 AR057397 and R01 AR064157. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

Abbreviations

AF	annulus fibrosus
IVD	intervertebral disc
IL	interleukin
TNF-α	tumor necrosis factor- α
DMEM	Dulbecco's Modified Eagles Media
ELISA	Enzyme linked immunosorbant assay

References

- Adams MA, Lama P, Zehra U, Dolan P. Why do some intervertebral discs degenerate, when others (in the same spine) do not? *Clin Anat.* 2015; 28:195–204. [PubMed: 24753325]
- Allen KD, Shamji MF, Mata BA, Gabr MA, Sinclair SM, Schmitt DO, Richardson WJ, Setton LA. Kinematic and dynamic gait compensations in a rat model of lumbar radiculopathy and the effects of tumor necrosis factor-alpha antagonist. *Arthritis research & therapy.* 2011; 13:R137. [PubMed: 21871102]
- Andrade P, Hoogland G, Garcia MA, Steinbusch HW, Daemen MA, Visser-Vandewalle V. Elevated IL-1beta and IL-6 levels in lumbar herniated discs in patients with sciatic pain. *European spine journal : official publication of the European Spine Society, the European Spinal Deformity Society, and the European Section of the Cervical Spine Research Society.* 2013; 22:714–720.
- Burke JG, Watson RW, McCormack D, Dowling FE, Walsh MG, Fitzpatrick JM. Spontaneous production of monocyte chemoattractant protein-1 and interleukin-8 by the human lumbar intervertebral disc. *Spine.* 2002; 27:1402–1407. [PubMed: 12131736]
- Chen C, Xie J, Rajappa R, Deng L, Fredberg J, Yang L. Interleukin-1beta and tumor necrosis factor-alpha increase stiffness and impair contractile function of articular chondrocytes. *Acta biochimica et biophysica Sinica.* 2015; 47:121–129. [PubMed: 25520178]
- Cheung KM, Samartzis D, Karppinen J, Luk KD. Are "patterns" of lumbar disc degeneration associated with low back pain?: new insights based on skipped level disc pathology. *Spine.* 2012; 37:E430–E438. [PubMed: 22466575]
- Gawri R, Rosenzweig DH, Krock E, Ouellet JA, Stone LS, Quinn TM, Haglund L. High mechanical strain of primary intervertebral disc cells promotes secretion of inflammatory factors associated with disc degeneration and pain. *Arthritis research & therapy.* 2014; 16:R21. [PubMed: 24457003]
- Gilbert HT, Hoyland JA, Freemont AJ, Millward-Sadler SJ. The involvement of interleukin-1 and interleukin-4 in the response of human annulus fibrosus cells to cyclic tensile strain: an altered mechanotransduction pathway with degeneration. *Arthritis research & therapy.* 2011; 13:R8. [PubMed: 21276216]
- Gilbert HT, Hoyland JA, Millward-Sadler SJ. The response of human anulus fibrosus cells to cyclic tensile strain is frequency-dependent and altered with disc degeneration. *Arthritis and rheumatism.* 2010; 62:3385–3394. [PubMed: 20617521]
- Jacobs, JJ. *Spine: Low Back and Neck Pain.* In: Jacobs, JJ.; Andersson, GBJ.; Bell, JE.; Weinstein, SL.; Dormans, JP.; Furman, MB.; Lane, N.; Puzas, JE.; Williams St. Clair, E.; Yelin, EH., editors. *The Burden of Musculoskeletal Diseases in the United States: Prevalence, Societal and Economic Cost.* 2nd. Rosemont, IL: Bone and Joint Initiative USA; 2011.
- Kandel R, Roberts S, Urban JP. Tissue engineering and the intervertebral disc: the challenges. *European spine journal : official publication of the European Spine Society, the European Spinal Deformity Society, and the European Section of the Cervical Spine Research Society.* 2008; 17(Suppl 4):480–491.
- Kim JH, Studer RK, Sowa GA, Vo NV, Kang JD. Activated macrophage-like THP-1 cells modulate anulus fibrosus cell production of inflammatory mediators in response to cytokines. *Spine.* 2008; 33:2253–2259. [PubMed: 18784630]
- Korecki CL, Costi JJ, Iatridis JC. Needle puncture injury affects intervertebral disc mechanics and biology in an organ culture model. *Spine.* 2008; 33:235–241. [PubMed: 18303454]
- Koukouritaki SB, Vardaki EA, Papakonstanti EA, Lianos E, Stournaras C, Emmanouel DS. TNF-alpha induces actin cytoskeleton reorganization in glomerular epithelial cells involving tyrosine phosphorylation of paxillin and focal adhesion kinase. *Molecular medicine.* 1999; 5:382–392. [PubMed: 10415163]
- Le Maitre CL, Hoyland JA, Freemont AJ. Catabolic cytokine expression in degenerate and herniated human intervertebral discs: IL-1beta and TNFalpha expression profile. *Arthritis research & therapy.* 2007; 9:R77. [PubMed: 17688691]
- Likhitpanichkul M, Kim Y, Torre OM, See E, Kazezian Z, Pandit A, Hecht AC, Iatridis JC. Fibrinogenipin annulus fibrosus sealant as a delivery system for anti- TNFalpha drug. *The spine journal : official journal of the North American Spine Society.* 2015; 15:2045–2054. [PubMed: 25912501]

- Maidhof R, Jacobsen T, Papatheodorou A, Chahine NO. Inflammation induces irreversible biophysical changes in isolated nucleus pulposus cells. *PLoS one*. 2014; 9:e99621. [PubMed: 24936787]
- Maniotis AJ, Chen CS, Ingber DE. Demonstration of mechanical connections between integrins, cytoskeletal filaments, and nucleoplasm that stabilize nuclear structure. *Proceedings of the National Academy of Sciences of the United States of America*. 1997; 94:849–854. [PubMed: 9023345]
- Marras WS, Lavender SA, Ferguson SA, Splittstoesser RE, Yang G. Quantitative dynamic measures of physical exposure predict low back functional impairment. *Spine*. 2010; 35:914–923. [PubMed: 20354470]
- Michalek AJ, Funabashi KL, Iatridis JC. Needle puncture injury of the rat intervertebral disc affects torsional and compressive biomechanics differently. *European spine journal : official publication of the European Spine Society, the European Spinal Deformity Society, and the European Section of the Cervical Spine Research Society*. 2010; 19:2110–2116.
- Millward-Sadler SJ, Costello PW, Freemont AJ, Hoyland JA. Regulation of catabolic gene expression in normal and degenerate human intervertebral disc cells: implications for the pathogenesis of intervertebral disc degeneration. *Arthritis research & therapy*. 2009; 11:R65. [PubMed: 19435506]
- Nakamae T, Ochi M, Olmarker K. Pharmacological inhibition of tumor necrosis factor may reduce pain behavior changes induced by experimental disc puncture in the rat: an experimental study in rats. *Spine*. 2011; 36:E232–E236. [PubMed: 21037531]
- Olmarker K, Rydevik B. Selective inhibition of tumor necrosis factor- α prevents nucleus pulposus-induced thrombus formation, intraneural edema, and reduction of nerve conduction velocity: possible implications for future pharmacologic treatment strategies of sciatica. *Spine*. 2001; 26:863–869. [PubMed: 11317106]
- Petrache I, Birukova A, Ramirez SI, Garcia JG, Verin AD. The role of the microtubules in tumor necrosis factor- α -induced endothelial cell permeability. *American journal of respiratory cell and molecular biology*. 2003; 28:574–581. [PubMed: 12707013]
- Petrache I, Verin AD, Crow MT, Birukova A, Liu F, Garcia JG. Differential effect of MLC kinase in TNF- α -induced endothelial cell apoptosis and barrier dysfunction. *American journal of physiology. Lung cellular and molecular physiology*. 2001; 280:L1168–L1178. [PubMed: 11350795]
- Phillips KL, Jordan-Mahy N, Nicklin MJ, Le Maitre CL. Interleukin-1 receptor antagonist deficient mice provide insights into pathogenesis of human intervertebral disc degeneration. *Ann Rheum Dis*. 2013; 72:1860–1867. [PubMed: 23396662]
- Purmessur D, Walter BA, Roughley PJ, Laudier DM, Hecht AC, Iatridis J. A role for TNF α in intervertebral disc degeneration: a non-recoverable catabolic shift. *Biochemical and biophysical research communications*. 2013; 433:151–156. [PubMed: 23438440]
- Rannou F, Poiraudeau S, Foltz V, Boiteux M, Corvol M, Revel M. Monolayer annulus fibrosus cell cultures in a mechanically active environment: local culture condition adaptations and cell phenotype study. *J Lab Clin Med*. 2000; 136:412–421. [PubMed: 11079469]
- Risbud MV, Shapiro IM. Role of cytokines in intervertebral disc degeneration: pain and disc content. *Nature reviews. Rheumatology*. 2014; 10:44–56. [PubMed: 24166242]
- Seguin CA, Pilliar RM, Roughley PJ, Kandel RA. Tumor necrosis factor- α modulates matrix production and catabolism in nucleus pulposus tissue. *Spine*. 2005; 30:1940–1948. [PubMed: 16135983]
- Shamji MF, Setton LA, Jarvis W, So S, Chen J, Jing L, Bullock R, Isaacs RE, Brown C, Richardson WJ. Proinflammatory cytokine expression profile in degenerated and herniated human intervertebral disc tissues. *Arthritis and rheumatism*. 2010; 62:1974–1982. [PubMed: 20222111]
- Sowa G, Coelho P, Vo N, Bedison R, Chiao A, Davies C, Studer R, Kang J. Determination of annulus fibrosus cell response to tensile strain as a function of duration, magnitude, and frequency. *Journal of orthopaedic research : official publication of the Orthopaedic Research Society*. 2011; 29:1275–1283. [PubMed: 21671263]
- Sowa GA, Coelho JP, Vo NV, Pacek C, Westrick E, Kang JD. Cells from degenerative intervertebral discs demonstrate unfavorable responses to mechanical and inflammatory stimuli: a pilot study.

- American journal of physical medicine & rehabilitation / Association of Academic Physiatriests. 2012; 91:846–855. [PubMed: 22760106]
- Stokes IA, Iatridis JC. Mechanical conditions that accelerate intervertebral disc degeneration: overload versus immobilization. *Spine*. 2004; 29:2724–2732. [PubMed: 15564921]
- Studer RK, Vo N, Sowa G, Ondack C, Kang J. Human nucleus pulposus cells react to IL-6: independent actions and amplification of response to IL-1 and TNF-alpha. *Spine*. 2011; 36:593–599. [PubMed: 21178846]
- Thompson JP, Pearce RH, Schechter MT, Adams ME, Tsang IK, Bishop PB. Preliminary evaluation of a scheme for grading the gross morphology of the human intervertebral disc. *Spine*. 1990; 15:411–415. [PubMed: 2363069]
- Ulrich JA, Liebenberg EC, Thuillier DU, Lotz JC. ISSLS prize winner: repeated disc injury causes persistent inflammation. *Spine*. 2007; 32:2812–2819. [PubMed: 18246002]
- Vos T, Flaxman AD, Naghavi M, Lozano R, Michaud C, Ezzati M, Shibuya K, et al. Years lived with disability (YLDs) for 1160 sequelae of 289 diseases and injuries 1990–2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet*. 2012; 380:2163–2196. [PubMed: 23245607]
- Walter BA, Korecki CL, Purmessur D, Roughley PJ, Michalek AJ, Iatridis JC. Complex loading affects intervertebral disc mechanics and biology. *Osteoarthritis and cartilage / OARS, Osteoarthritis Research Society*. 2011; 19:1011–1018.
- Walter BA, Likhitpanichkul M, Illien-Junger S, Roughley PJ, Hecht AC, Iatridis JC. TNFalpha transport induced by dynamic loading alters biomechanics of intact intervertebral discs. *PloS one*. 2015a; 10:e0118358. [PubMed: 25734788]
- Walter BA, Likhitpanichkul M, Nasser P, Hecht AC, Iatridis J. Transient TNF α Exposure Permanently Alters Intervertebral Disc Biomechanics in a Dynamic Compression Bioreactor System. *Transactions of the Orthopaedic Research Society Poster # 0698*. 2014
- Walter BA, Purmessur D, Likhitpanichkul M, Weinberg A, Cho SK, Qureshi SA, Hecht AC, Iatridis JC. Inflammatory Kinetics and Efficacy of Anti-inflammatory Treatments on Human Nucleus Pulposus Cells. *Spine*. 2015b; 40:955–963. [PubMed: 25893355]
- Weiler C, Nerlich AG, Bachmeier BE, Boos N. Expression and distribution of tumor necrosis factor alpha in human lumbar intervertebral discs: a study in surgical specimen and autopsy controls. *Spine*. 2005; 30:44–53. discussion 54. [PubMed: 15626980]
- Wuertz K, Godburn K, MacLean JJ, Barbir A, Donnelly JS, Roughley PJ, Alini M, Iatridis JC. In vivo remodeling of intervertebral discs in response to short- and longterm dynamic compression. *Journal of orthopaedic research : official publication of the Orthopaedic Research Society*. 2009; 27:1235–1242. [PubMed: 19274755]
- Wuertz K, Haglund L. Inflammatory mediators in intervertebral disk degeneration and discogenic pain. *Global spine journal*. 2013; 3:175–184. [PubMed: 24436868]
- Zhu Q, Jackson AR, Gu WY. Cell viability in intervertebral disc under various nutritional and dynamic loading conditions: 3d finite element analysis. *Journal of biomechanics*. 2012; 45:2769–2777. [PubMed: 23040882]

Figure 1. TNF-α increased pro-inflammatory cytokine production while stretch did not
 (A) TNF-α (10 ng/mL) treatment alone significantly increased pro-inflammatory cytokine concentrations produced by human AF cells (N=5) in the absence of stretch. (B) Mechanical strain at physiological (5%) or pathological (15%) levels in the absence of TNF-α treatment exhibited no significant effect on pro-inflammatory cytokine concentration as compared to basal media. (C) Absolute values of pro-inflammatory cytokine concentrations measured in basal media. (mean±SD; ***p<0.005)

	Pro-inflammatory cytokine concentration (pg/mL)		
Unstrained TNF- α media	IL-1 β	IL-6	IL-8
	84.0 $\pm$ 77.3	280.9 $\pm$ 214.1	5125.1 $\pm$ 4170.8

Figure 2. TNF- α and pathological mechanical strain interacted to substantially increase pro-inflammatory cytokine production

(A) Production of pro-inflammatory cytokines by TNF- α treated human AF cells (N=5) in the presence of 15% strain significantly increased for IL-1 β and IL-8 compared to 5% strain but not 0% strain. (B) This was particularly notable given the high cytokine concentrations present for the unstrained TNF- α group. (mean $\pm$ SD; $p < 0.05$)

Figure 3. TNF- α increased F-actin staining and microtubule disruption

Immunocytochemistry of F-actin and vinculin on human AF cells treated with basal media and TNF- α . F-actin staining intensity increased with TNF- α and there was no difference in vinculin staining. (B) An extensive microtubule network was evident using α -tubulin staining on AF cells for the basal media group (yellow arrows). The microtubule network was disrupted with TNF- α treatment, as evidenced by more diffuse α -tubulin staining which was particularly evident at higher optical zoom. (Magnification = 60X, Scale = 25 μ m).

Figure 4. TNF- α but not mechanical stretch induced F-actin changes in human AF cells
 Immunocytochemistry for F-actin and vinculin in human AF cells seeded at 100,000 cells/mL immediately after 24 hours of stretch with and without TNF- α . F-actin and vinculin staining exhibited no apparent differences in cytoskeletal organization across all strain magnitudes. TNF- α treatment increased F-actin staining intensity but there were no discernable differences between strains in cytoskeletal structure with TNF- α treatment. (Magnification = 20X, Scale = 100 μ m).

Figure 5. Microtubules were disrupted in AF cells following TNF- α treatments and stretch
 An extensive microtubule network was clearly demonstrated with α -tubulin staining under all mechanical strain conditions in the basal media group (yellow arrows). Microtubules were disrupted in AF cells subjected to TNF- α treatment with more diffuse α -tubulin staining.

B

	Pro-inflammatory cytokine concentration (pg/mL)		
TNF-α+15% strain	IL-1β	IL-6	IL-8
	103.6 ± 64.1	273.6 ± 173.9	5907.9 ± 3046.7

Figure 6. Pro-inflammatory cytokine production was reduced by anti-TNF-α but not anti-IL-6
 Infliximab decreased the concentrations of pro-inflammatory cytokines produced by AF cells (N=5) exposed to TNF-α (10 ng/mL) and 15% stretch. There was a significant effect on IL-8 concentration for cells treated with anti-TNF-α, but not for anti-IL-6. Anti-IL-6 had a significantly smaller reduction of pro-inflammatory cytokine levels. There was a significant difference between the effects of Anti-TNF-α and Anti-IL-6 on the concentration of pro-inflammatory cytokines. (mean±SD; *p=0.01, #p=0.08)