Published in final edited form as:

J Empir Res Hum Res Ethics. 2016 April ; 11(2): 106–114. doi:10.1177/1556264616648589.

Parental estimation of their child's increased type 1 diabetes risk during the first two years of participation in an international observational study: Results from the TEDDY study

Ulrica Swartling^a, Kristian Lynch^b, Laura Smith^{b,c}, Suzanne Bennett Johnson^d, and the TEDDY Study Group^{*}

^aDepartment of Clinical Sciences, Lund University, CRC, Skåne University Hospital, Malmö, Sweden

^bHealth Informatics Institute, University of South Florida, Tampa FL, USA

^cDepartment of Pediatrics, USF Diabetes Center, University of South Florida, Tampa, FL, USA

^dDepartment of Behavioral Sciences and Social Medicine, Florida State University College of Medicine, Tallahassee FL, USA

Abstract

This study assessed mothers' and fathers' perception of their child's risk of getting type 1 diabetes (T1D) during the first two years of their participation in The Environmental Determinants of Diabetes in the Young (TEDDY) study. TEDDY parents were informed of their child's increased genetic risk for T1D at study inception. Parent perception of the child's risk was assessed at 3, 6, 15 and 27 months of age. In families with no history of T1D, underestimation of the child's T1D risk was common in mothers (> 38%) and more so in fathers (> 50%). The analyses indicated that parental education, country of residence, family history of T1D, household crowding, ethnic minority status, and beliefs that the child's T1D risk can be reduced were factors associated with parent risk perception accuracy. Even when given extensive information about their child's T1D risk, parents often fail to accurately grasp the information provided. This is particularly true for fathers, families from low socioeconomic backgrounds, and those with no family history of T1D. It is important to develop improved tools for risk communication tailored to individual family needs.

Keywords

risk perception; screening; children; type 1 diabetes; informed consent

Type 1 diabetes (T1D) is one of the most common chronic diseases in childhood, and its incidence is rapidly increasing world-wide (D.I.A.M.O.N.D. Project Group, 2006; Patterson, Dahlquist, Gyurus, et al, 2009; Search For Diabetes In Youth Study Group, 2006). T1D is a

Corresponding author: Ulrica Swartling, PhD, Lund University, Department of Clinical Sciences, Division of Diabetes and Celiac disease, CRC, Skåne University Hospital (SUS), Jakob Waldenströms gata 35, 205 02 Malmö, Sweden. ulrica.swartling@med.lu.se, phone 46-733344714.

*Members of the TEDDY Study Group are listed in Appendix

serious chronic disease that occurs following the autoimmune destruction of insulin producing cells in the pancreas. Currently, T1D cannot be prevented or cured, but must be treated with daily administration of exogenous insulin throughout the life of the patient. If not well managed, T1D can lead to serious long-term complications such as eye, kidney, and nerve disease (American Diabetes Association, 2015a,b). This is in contrast to type 2 diabetes, a much more common condition accounting for 90–95% of all diabetes cases, in which lifestyle factors, such as diet and physical activity, are known to contribute to the development of the disease. T1D is usually diagnosed in under or normal weight children while T2D is typically diagnosed in individuals who are overweight. Individuals with type 2 diabetes are often treated with lifestyle changes and oral medications, although some may also be prescribed insulin as their condition progresses (American Diabetes Association, 2015a,b).

Past research has elucidated the HLA genotypes that predispose individuals to T1D. However, the concordance rate for monozygotic twin is only 30–50%, suggesting that environmental factors also play a role in the pathogenesis of T1D (Pociot, et al., 2010). Therefore, it is hypothesized that a combination of an individual's genetic predisposition and environmental factors (e.g., viruses, microbiomal factors, stress, etc) likely contribute to the development of T1D (Eringsmark & Lernmark, 2013). However, in order to identify the environmental factors that contribute to the development of T1D large-scale screening studies follow infants at genetic risk for T1D for many years and collect relevant data through biologic samples and questionnaires (The TEDDY Study group, 2007).

However, there are a number of ethical and psychological issues surrounding newborn genetic screening for T1D (Ross, 2003; Roth, 2001; Stolt et al, 2005, Swartling & Helgesson, 2008). Testing children for T1D risk could increase distress in parents, which in turn could negatively effect the child. The published literature suggests that while parents (mostly mothers) often initially report distress when informed of their child's increased risk, this distress dissipates over time (Johnson et al, 1990; Johnson & Tercyak, 1995; Hummel, Ziegler, & Roth, 2004; Simonen et al, 2006; Johnson et al, 2004, 2007, 2011; Kerruish et al, 2007; Carmichael et al, 2003). This literature also suggests that many parents engage in one or more behaviors in an effort to prevent the disease, most common being increased monitoring behaviors and dietary changes (Hendrieckx et al, 2002; Baughcum et al, 2005; Heshka et al, 2008; Smith et al, 2014).

Communicating risk information is often difficult as the child's increased risk does not guarantee the child will necessarily get the disease. Simply asking families whether they understand the information provided is insufficient; individuals often state they understand information, when in fact their understanding is inaccurate (Stolt et al, 2005; Swartling & Helgesson, 2005). Studies suggest that individuals who are very familiar with T1D, because they live with someone who has it, understand the risk information better than individuals with no immediate family members living with the disease (Carmichael et al, 2003; Johnson & Tercyak, 1995).

Accurate understanding of risk information is important for informed participant decision-making in research studies. Previous studies have shown that T1D risk perception accuracy

is associated with increased anxiety or worry, efforts to prevent the disease, and study retention (Hendrieckx et al, 2002; Baughcum et al, 2005; Hood et al, 2006; Lernmark et al, 2012; Smith et al, 2014) while underestimation of risk has been associated with study drop out (Johnson et al, 2011). However, our current understanding of parental risk perception accuracy is limited due to the fact that most studies have focused on relatively small samples of mothers familiar with T1D over a relatively short period of time.

The Environmental Determinants of Diabetes in The Young (TEDDY) study provides a unique opportunity to study factors associated with both mothers' and fathers' risk perception accuracy over time in a large multi-national study of parents from the general population with no immediate family history of T1D as well as parents whose at-risk child has a first degree relative with T1D. All TEDDY children are at increased genetic risk for T1D and all TEDDY parents were informed of this increased risk at the time they joined TEDDY. We report here mothers' and fathers' risk perception accuracy during their first two years of TEDDY participation and factors associated with risk perception accuracy.

Methods

The TEDDY Study

TEDDY is a prospective multinational cohort study supported by the National Institutes of Health investigating the environmental determinants of T1D in 8676 genetically at-risk children identified at birth. Most TEDDY children (89%) come from the general population with no immediate family history of T1D; the remaining TEDDY children are families where the children have a first-degree relative with T1D. Three centers are located in the United States (Colorado, Florida/Georgia, Washington) and three centers are located in Europe (Finland, Germany, Sweden). TEDDY children are seen every three months for the first four years and biannually thereafter. Data collected at each study visit include biological samples (e.g., blood, nasal swabs), diet records, and a wide range of demographic and psychological measures (TEDDY Study Group, 2007). The relevant Institutional Review Board at each TEDDY site approved the TEDDY study and protocol.

Study population

Parent questionnaire data available as of April 30, 2013 were obtained from 3 (mothers only), 6, 15, and 27-month study visits. On this date all TEDDY subjects were 36 months or older. Excluded from analysis were 2090 children whose last TEDDY visit was prior to 27 months of age ($n = 2090$), children who were determined to be HLA ineligible ($n = 39$), and children who became antibody positive at or before the 27 month study visit ($n = 447$). Selected for analysis were the questionnaire responses of 5953 families (5769 mothers and 5247 fathers) who completed at least one questionnaire at the 6 or 27 month study visits (excluding 118 families). Most were families from the general population (89%; $n=5293$), with the remaining 660 families (11%) having a known family history of T1D.

Risk communication protocol in TEDDY

After genetically screening the infant at birth, eligible families were contacted by phone (between 6–12 weeks after the test) and informed about their child's increased T1D genetic

risk and eligibility for the TEDDY study. During the call, the parent was given a numerical estimate of the child's risk in a standardized manner; this information was subsequently reiterated in a letter sent to the parent's home. The numerical risk given varied by country, based on country-specific epidemiological data, and whether the child belonged to the general population or had a close relative with T1D. For example, a mother from the general population in the U.S. and Germany was informed that out of 100 children with her child's genetic risk, approximately 3 children would develop T1D. In Sweden and Finland, where there is higher incidence of T1D, general population mothers were informed that out of 100 children with her child's genetic risk, approximately 7 children would develop T1D. A mother of a U.S. or German child with a known family history of T1D was informed that out of 100 children with her child's genetic risk, 14 would develop T1D. The numerical estimate about the TEDDY child's risk for T1D was followed by comparison information about risk for T1D in children without their child's genetic risk: 1 out of 300 would develop T1D.

There were some national variances in risk communication: Sweden and Finland (but no other country) did not make any distinction between whether the child came from the general population or had a first degree relative when informing the parents but instead gave the same numerical risk value (7/100). However, in Finland the risk information was complemented with a brochure containing more detailed risk estimates.

Upon request, parents of children with a family history of T1D were given more detailed risk estimates based on whether the family member with T1D was a mother, father or sibling. All families were told that the purpose of TEDDY was to identify the environmental trigger(s) of T1D in genetically at-risk children and if environmental trigger(s) were identified, this information could be used in the future to develop methods of preventing the disease. If families asked if there was anything they could do to prevent T1D in their child, they were told that no means of prevention is currently available and that they should encourage the development of healthy habits in the child.

At each TEDDY visit, blood was drawn for T1D autoantibodies. At all sites, results were sent home by mail except for Sweden where test results were provided at the next study visit. Even when the child tested negative for T1D autoantibodies, the parents were told that the child's risk for T1D had not changed and the child remained at increased risk for T1D (reiterating the risk estimates).

Measures

Risk perception accuracy—Risk perception accuracy was assessed by the following item at the 3- (mothers only), 6-, 15-, and 27-month study visits: *Compared to other children, do you think of your child's risk for developing diabetes is (mark only one answer) - much lower, somewhat lower, about the same, somewhat higher, much higher.* Parents answering “much lower”, “somewhat lower” or “about the same” were classified as inaccurate while parents answering “somewhat higher” or “much higher” were classified as accurate.

Sociodemographic and maternal life style measures—Child sociodemographic variables included the child's gender (male, female), only child status (yes/no); ethnic

minority status (yes/no) (U.S. definition: TEDDY child's mother's first language is not English or she was not born in the U.S. or the child is a member of an ethnic minority group; European definition: TEDDY child's mother's first language or country of birth is other than the TEDDY country in which the child resides); FDR status (yes/no).

Parent sociodemographic variables included mother's age at birth (years); parent's education (basic primary education; some trade school or college; graduated college); marital status (married or living together; single parent); country of residence (U.S., Finland, Germany and Sweden); household crowding (number of persons living in the household divided by the number of rooms in the house). As the crowding distribution was skewed, it was rescored to normalize the distribution (1= 0.00–0.49; 2=0.50–0.59; 3= 0.60–0.75; 4= 0.76–1.00; 5= > 1.00).

Maternal lifestyle behaviors included whether the mother smoked during pregnancy (yes/no) or after having given birth (measured at 9 months) (yes/no), whether the mother worked during pregnancy (yes/no) or after birth (measured at 9 months) (yes/no), and whether the mother drank alcohol during pregnancy (yes/no).

Parental belief that T1D risk can be reduced—At the 6-, 15- and 27- month study visits, parents were asked if they believed they could do something to reduce their child's risk for developing T1D. Parents were asked to agree or disagree to three statements on a 5-point Likert scale (1= strongly agree to 5 = strongly disagree): (1) “I can do something to reduce my child's risk of developing diabetes”, (2) “Medical professionals can do something to reduce my child's risk for developing diabetes”, and (3) “It is up to chance or fate whether my child develops diabetes”. Responses to statements 1 and 2 were reversed scored and then summed with the response to statement 3 so that higher scores indicated greater belief that risk of T1D could be reduced. This 3 item scale score exhibited high internal consistency (Cronbach's alpha coefficient of 0.821 for mothers and 0.793 for fathers).

Statistical analyses

The percent of mothers and fathers with accurate risk perception at 3, 6, 15, and 27 months were calculated. McNemar's test was used to test for differences between mothers and fathers and also within parent across time. Chi-square was used to test for differences between families from the general population and families with a history of T1D. Logistic regression was used to identify factors independently associated with having an accurate versus inaccurate risk perception. Separate regression models were fitted for mothers and fathers at 6 months and 27 months using sociodemographic variables, maternal lifestyle behaviors, and parent belief that the child's T1D risk can be reduced. Factors with p-values less than 0.05 were considered statistically significant and retained. Statistical analyses were conducted using SAS (version 9.4).

Results

Table 1 provides the risk perception accuracy data for mothers and fathers by T1D family history status and across time by study visit. Most parents who underestimated their child's risk believed that their child's T1D risk was “about the same” as another child; few parents

selected responses indicating that they believed their child's risk to be lower than a comparison child. At 6 months, the majority of general population mothers (61.3%) had accurate risk perceptions while the majority of general population fathers (53.2%) had inaccurate risk perceptions ($\chi^2 = 241.9$, $p < .0001$). There was no significant change in risk perception accuracy for mothers or fathers from the general population across time. Mothers where the child had a family history of T1D were far more likely to have accurate risk perceptions (89% at 3 and 86% at 27 months) than mothers from the general population (61% at 3 and 60% at 27 months; $\chi^2 = 188.1$, $p < .0001$) and their risk perception accuracy did not change across time. Mothers where the child had a family history of T1D were also more likely to hold accurate risk perceptions compared to fathers with the same background ($\chi^2 = 37.3$, $p < .0001$). In addition, fathers where the child had a family history of T1D were more likely to have accurate risk perceptions than fathers from the general population ($\chi^2 = 166.4$, $p < .0001$). The majority of both mothers and fathers where the child had a family history of T1D held accurate risk perceptions.

Separate multiple logistic regressions were fitted for mothers and fathers to identify the independent factors associated with accurate risk perception at 6 and 27 months. Table 2 provides the significant results for mothers. The results were essentially the same at 6 and 27 months. As expected, mothers where the child had a family history of T1D were more likely to hold accurate risk perceptions than general population mothers (87% versus 60%). College educated mothers were more likely to hold accurate risk perceptions compared to those with a primary school education (71% versus 44%). Mothers from Finland (72%) and Germany (75–77%) were also significantly more likely to hold accurate risk perceptions compared to mothers from the U.S. (61–63%) and Sweden (55–56%). Child ethnic minority status and household crowding were associated with poorer risk perception accuracy. Approximately 50% of mothers of ethnic minority children were accurate compared to 65% of mothers of nonminority children. Approximately 60% of mothers from the most crowded household were accurate compared to 67% of those from the least crowded household. Only child status had a weak but significant effect and was associated with poorer risk perception accuracy. Strong beliefs that the child's T1D risk could be reduced was actually associated with poorer risk perception accuracy. For example, at 3 months, only 58% of mothers with scores in the highest quartile on this measure were accurate compared to 73% of those with scores in the lowest quartile. We found no significant associations between child gender, maternal age, marital status, or maternal life style factors and risk perception accuracy.

Table 3 provides the results of the logistic regression for fathers at 6 and 27 months. In most respects, the results mimic the findings for mothers. The most important factors associated with risk perception accuracy for fathers were T1D family history status, country, and father's education ($p < .0001$). Greater household crowding was also associated with poorer risk perception accuracy ($p < .01$ at 6 months and $p < .0001$ at 27 months) and like mothers, fathers who held strong beliefs that their child's T1D risk could be reduced were more likely to hold inaccurate risk perceptions ($p < .0001$ at 27 months). The child's ethnic minority status and only child status were not significant predictors for fathers, in contrast to their significant effects for mothers. These findings were consistent across time.

Discussion

To our knowledge this is the largest study to date of risk perception accuracy in parents of infants genetically at-risk for a serious disease (T1D). The study is unique in that it assessed risk perception accuracy: (1) of both mothers and fathers; (2) of parents from the general population as well as those with T1D immediate family members; (3) of parents living in four different countries; (4) at multiple time points over a two year period.

In TEDDY, parents are fully informed of their child's increased genetic risk for T1D at study inception. Further, children are regularly tested for T1D autoantibodies and parents are informed of these test results. In cases of negative antibody test results, parents are told that the child's risk for T1D has not changed and the child remains at increased risk for T1D. Despite these extensive and repeated efforts to inform parents of their child's increased risk, a large proportion of both mothers and fathers, particularly from the general population, underestimated their child's risk. Only 60% of mothers from the general population and less than half of general population fathers had accurate perceptions of their child's risk. In contrast, parents where the child had a family history of T1D had significantly more accurate risk perceptions. More than 80% of these mothers and more than 70% of the fathers had accurate risk perceptions. The high risk perception accuracy rates found in mothers with an T1D immediate family member is consistent with prior reports (Carmichael et al, 2003; Hood et al, 2006). These results are expected since parents from the general population do not have the experience and knowledge about T1D as well as its heritability so familiar to families living with an immediate family member with the disease.

Fathers, regardless of T1D family history, were consistently less accurate than mothers. TEDDY fathers did accompany the child to the study visits less often than mothers (e.g., 29% of TEDDY fathers came to the 6-month TEDDY visit compared to 98% of TEDDY mothers). However, TEDDY father attendance at TEDDY clinic visits was unrelated to risk perception accuracy; fathers who attended TEDDY clinic visits did not have more accurate risk perceptions than fathers who failed to attend. There is ample evidence that mothers are the primary caretakers of most children and make most of the health care decisions for their families (Matoff-Stepp, et al, 2014). Consequently, mothers may be more attentive to health related information relevant to their children.

In TEDDY, we found risk perception accuracy to be highly stable across time, with no decline in risk perception over time previously reported by Carmichael and colleagues (2003) and Hood and colleagues (2006). When the TEDDY protocol was established, the investigators were well aware that risk perception accuracy could decline across time and in response, established a protocol of repeated communication with parents about their child's risk, including sending letters home reiterating information provided on phone calls or at study visits. In this context, it is interesting to note that Sweden was the only TEDDY country that did not regularly send home risk information (giving the results orally at the next visit) and Swedish parents demonstrated the lowest percentages of parents with accurate risk perceptions among TEDDY countries.

In this study, we also found that ethnic minority status, whether one lived in more crowded households, and had less parental education were all factors associated with higher rates of inaccurate risk perception, confirming results published in earlier studies (Carmichael et al, 2003; Hood et al, 2006). These results once again confirm that risk information is difficult to successfully communicate. The results also suggest that risk communication protocols and effective strategies need to be developed for those who are less educated and from lower sociodemographic backgrounds.

The analyses showed that parental belief that a child's T1D risk could be reduced was associated with having a less accurate risk perception. While being accurate in risk perception accuracy and having a belief that risk can be changed are two different things – it may be that people who believe they can reduce risk may underestimate the child's risk more. Perhaps parents who believe the risk can be reduced, view the risk information as less stable and more modifiable, taking the information less seriously and giving it less credence. Previous studies suggest that belief that one can control T1D onset may be linked to increased efforts to prevent the disease, even though no means of preventing T1D currently exists (Smith et al, 2014; Hendrieckx et al, 2002).

This study is limited by the inclusion of families who agreed to join the TEDDY study, with its demanding and intensive protocol. Consequently our findings may not represent risk perception accuracy among families who declined study participation. Nevertheless, our findings suggest that risk information is challenging to present in ways that parents understand. This is particularly the case when parents are not well educated, come from lower sociodemographic backgrounds, and have no immediate family experience with T1D.

This study's results have important implications beyond the care of children with a genetic risk for T1D. Risk communication is a common component of many patient-provider, as well as investigator-subject, communications and this study's findings highlight the difficulty of successfully communicating risk. Development of improved strategies for risk communication tailored to individual and family needs is warranted.

Best practices

A large proportion of parents from the general population underestimate their child's increased genetic risk for type 1 diabetes over an extended period of time (2 years), and fathers underestimate risk more often than mothers. Findings suggest the importance of viewing risk communication as a process, involving both study staff and the participating families. Not having an accurate risk perception may influence informed decision making, so researchers should take notice of situations where parents have formed and continue to have an incorrect understanding of risk. This is particularly important when families have no prior experience with the disease, are members of minority groups, or have less education, or a lower socioeconomic status.

These findings suggest the importance of developing tailored risk communication protocols and effective strategies designed for those who are less educated and from lower sociodemographic backgrounds.

Research Agenda

Different methods of communicating risk (oral, written, and visual) are important tools when informing parents of their child's risk status. These different methods should be used to tailor risk communication strategies for specific populations and test their effectiveness.

Educational Implications

These results are valuable for the staff involved in the TEDDY study and risk communication in general. Tailored information can be designed to help parents develop a correct understanding of their child's risk status and assist those families who hold inaccurate risk perceptions.

Acknowledgments

This study was funded by U01 DK63829, U01 DK63861, U01 DK63821, U01 DK63865, U01 DK63863, U01 DK63836, U01 DK63790, UC4 DK63829, UC4 DK63861, UC4 DK63821, UC4 DK63865, UC4 DK63863, UC4 DK63836, UC4 DK95300, and UC4 DK100238, and Contract No. HHSN267200700014C from the National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK), National Institute of Allergy and Infectious Diseases (NIAID), National Institute of Child Health and Human Development (NICHD), National Institute of Environmental Health Sciences (NIEHS), Juvenile Diabetes Research Foundation (JDRF), and Centers for Disease Control and Prevention (CDC). This work supported in part by the NIH/NCATS Clinical and Translational Science Awards to the University of Florida (UL1 TR000064) and the University of Colorado (UL1 TR001082).

Biographies

Ulrica Swartling is an assistant professor, Department of Clinical Sciences, Diabetes Division, at Lunds University, Sweden. She has conducted research on ethical aspects of involving children in screening for T1D, such as factors related to shared decision-making, the involvement of children, and assent. Dr. Swartling is a member of the Child Engagement subcommittee of the TEDDY study. She was involved in the design and data analysis and took the lead in writing the manuscript.

Kristian Lynch is an Assistant Professor at the Health Informatics Institute at the University of South Florida in Tampa, Florida. His research interests are on epidemiological aspects of type 1 diabetes etiology, including statistical modelling within longitudinal epidemiological research studies. As statistician, he is involved in several of the committees of the TEDDY study. He participated in the design of the current study, including data analysis.

Laura B. Smith is an Assistant Professor in the Department of Pediatrics, the USF Diabetes Center, and the Health Informatics Institute at the University of South Florida in Tampa, Florida. She has conducted research focusing on the psychosocial factors that impact pediatric diabetes management, such as individual child factors and family functioning. She also has research interests in the area of patient-provider interactions, particularly in the treatment of type 1 diabetes. Dr. Smith co-chairs the Child Engagement subcommittee of the current study. She was involved in the data analyses and writing of the current manuscript.

Suzanne Bennett Johnson is a Distinguished Research Professor in the Department of Behavioral Sciences and Social Medicine, Florida State College of Medicine, Tallahassee Florida. Her research interests are on psychosocial aspects of childhood diabetes, including

the impact of genetic testing for type 1 diabetes on children and their families. She chairs the Psychosocial Studies Committee for the TEDDY project. She participated in the conceptualization and design of the current study and was involved in writing the current manuscript.

References

- Baughcum A, Johnson SB, Carmichael S, Lewin A, She J-X, Schatz D. Maternal efforts to prevent type 1 diabetes in at-risk children. *Diabetes Care*. 2005; 28:916–921. [PubMed: 15793195]
- Carmichael SK, Johnson SB, Baughcum A, North K, Hopkins D, Dukes MG, et al. Prospective assessment in newborns of diabetes autoimmunity (PANDA): maternal understanding of infant diabetes risk. *Genetics in Medicine*. 2003; 5:77–83. [PubMed: 12644776]
- D.I.A.M.O.N.D. Project Group. Incidence and trends of childhood type 1 diabetes worldwide 1990–1999. *Diabetes Medicine*. 2006; 23:857–866.
- Helgesson G, Swartling U. Views on data use, confidentiality and consent in a predictive screening involving children. *Journal of Medical Ethics*. 2008; 34:206–209. [PubMed: 18316465]
- Hendrieckx C, De Smer F, Kristoffersen I, Bradley C. Risk assessment for developing type 1 diabetes: intentions of behavioural changes prior to risk notification. *Diabetes Metabolism Research Review*. 2002; 18:36–42.
- Heshka JT, Palleschi C, Howley H, Wilson B, Wells P. A systematic review of perceived risks, psychological and behavioral impacts of genetic testing. *Genetics in Medicine*. 2008; 10:19–32. [PubMed: 18197053]
- Hood K, Johnson SB, Baughcum A, She J, Schatz D. Maternal understanding of infant diabetes risk: differential effects of maternal anxiety and depression. *Genetics in Medicine*. 2006; 8:665–670. [PubMed: 17079884]
- Hummel M, Ziegler A, Roth R. Psychological impact of childhood islet autoantibody testing in families participating in the BABYDIAB study. *Diabetes Medicine*. 2004; 21:324–328.
- Johnson SB. Psychological impact of screening and prediction in type 1 diabetes. *Current Diabetes Report*. 2011; 11:454–459.
- Johnson SB, Lee H-S, Baxter J, Lernmark B, Roth R, Simmell T. The Environmental Determinants of Diabetes in the Young (TEDDY) Study: predictors of early study withdrawal among participants with no family history of type 1 diabetes. *Pediatric Diabetes*. 2011; 12:165–171. [PubMed: 21029290]
- Johnson SB, Baughcum A, Rafkin-Mervis L, Schatz D. For The DPT-1 Study Group. Participant and parent experiences in the oral insulin study of the Diabetes Prevention Trial for Type 1 Diabetes. *Pediatric Diabetes*. 2009; 30:2193–2198.
- Johnson SB, Baughcum A, Hood K, Rafkin-Mervis L, Schatz D. For the DPT-1 Study Group. Participant and parent experiences in the parenteral insulin arm of the Diabetes Prevention Trial for Type 1 Diabetes. *Diabetes Care*. 2007; 30:2193–2198. [PubMed: 17563348]
- Johnson SB, Lernmark B, Baxter J, Roth R, Simell T, Mcleod W. The TEDDY Group. Parent anxiety in response to newborn genetic screening results in the Environmental Determinants of Diabetes in the Young (TEDDY) Study. *Diabetes*. 2007; 56(Suppl 1):A498.
- Johnson SB, Baughcum A, Carmichael S, She J-X, Schatz D. Maternal anxiety associated with newborn genetic screening for type 1 diabetes. *Diabetes Care*. 2004; 27:392–397. [PubMed: 14747219]
- Johnson SB, Tercyak K. Psychological impact of islet cell antibody screening for IDDM on children, adults, and their family members. *Diabetes Care*. 1995; 18:1370–1372. [PubMed: 8721939]
- Johnson SB, Riley W, Hansen C, Nurick M. Psychological impact of islet cell-antibody screening: preliminary results. *Diabetes Care*. 1990; 13:93–97. [PubMed: 2190772]
- Kerruish N, Campbell-Stokes P, Gray A, Merriman T, Robertson S, Taylor B. Maternal psychological reaction to newborn genetic screening for type 1 diabetes. *Pediatrics*. 2007; 120:e324–e335. [PubMed: 17609310]

- Lernmark B, Elding-Larsson H, Hansson G, Lindberg B, Lynch K, Sjoblad S. Parent responses to participation in genetic screening for diabetes risk. *Pediatric Diabetes*. 2004; 5:174–181. [PubMed: 15601359]
- Lernmark B, Lynch K, Ballard L, Baxter J, Roth R, Simell T, Johnson SB. Reasons for staying as a participant in the Environmental Determinants of Diabetes in the Young (TEDDY) longitudinal study. *Journal of Clinical Trials*. 2012; 2(2):100–114.
- Matoff-Stepp S, Applebaum B, Pooler J, Kavanagh E. Women as health care decision-makers: implications for health care coverage in the United States. *Journal of Health Care Poor Underserved*. 2014; 25:1507–1515.
- Patterson C, Dahlquist G, Gyurus E, Soltesz G. For The EURODIAB Study Group. Incidence trends for childhood type 1 diabetes in Europe during 1989–2003 and predicted new cases 2005–20: a multicentre prospective registration study. *Lancet*. 2009; 373:2027–2033. [PubMed: 19481249]
- Pociot F, Akolkar B, Concannon P, Erlich HA, Julier C, Morahan G, Nerup J. Genetics of type 1 diabetes: what's next? *Diabetes*. 2010; 59:1561–1571. [PubMed: 20587799]
- Liese AD, D'agostino RB Jr, Hamman RF, Kilgo PD, Lawrence JM, Liu LL, et al. Search for Diabetes In Youth Study Group. The burden of diabetes mellitus among US youth: prevalence estimates from the SEARCH for Diabetes in Youth Study. *Pediatrics*. 2006; 118:1510–1518. [PubMed: 17015542]
- Ross LF. Minimizing risk: the ethics of predictive diabetes mellitus screening research in newborns. *Archives of Pediatric Adolescence Medicine*. 2003; 157:89–95.
- Roth R, Lynch K, Lernmark B, Baxter J, Simell T, Smith L, Swartling U, Ziegler AG, Johnson SB. The TEDDY Study Group. Maternal anxiety about a child's diabetes risk in the TEDDY study: the potential role of life stress, postpartum depression, and risk perception. *Pediatric Diabetes*. 2015; 16(4):287–298. [PubMed: 25082392]
- Roth LR. Psychological and ethical aspects of prevention trials. *Journal of Pediatric Endocrinology Metabolism*. 2001; 14(Suppl. 1):669–674. [PubMed: 11393561]
- Simonen P, Korhonen T, Simell T, Keskinen P, Karkkainen M, Knip M, et al. Parental reactions to information about increased genetic risk of type 1 diabetes mellitus in infants. *Archives of pediatric adolescence medicine*. 2006; 160:1131–1136.
- Smith LB, Lynch KF, Baxter J, Lernmark B, Roth R, Simell T, Johnson SB. The TEDDY Study Group. Factors associated with maternal-reported actions to prevent type 1 diabetes in the first year of the TEDDY study. *Diabetes Care*. 2014; 37:325–331. [PubMed: 24041684]
- Stolt UG, Helgesson G, Liss P-E, Svensson T, Ludvigsson J. Information and informed consent in a longitudinal screening involving children: a questionnaire survey. *European Journal of Human Genetics*. 2005; 13:376–383. [PubMed: 15657607]
- Swartling U, Helgesson G. Self-assessed understanding as a tool for evaluating consent: reflections on a longitudinal study. *Journal of Medical Ethics*. 2008; 34:557–562. [PubMed: 18591294]
- TEDDY Study Group. The Environmental Determinants of Diabetes in the Young (TEDDY) study: study design. *Pediatric Diabetes*. 2007; 8:286–298. [PubMed: 17850472]

Appendix: The Teddy Study Group

Colorado Clinical Center

Marian Rewers, M.D., Ph.D., PI^{1,4,5,6,10,11}, Kimberly Bautista¹², Judith Baxter^{9,10,12,15}, Ruth Bedoy², Daniel Felipe-Morales, Brigitte I. Frohnert, M.D.¹⁴, Patricia Gesualdo^{2,6,12,14,15}, Michelle Hoffman^{12,13,14}, Rachel Karban¹², Edwin Liu, M.D.¹³, Jill Norris, Ph.D.^{2,3,12}, Adela Samper-Imaz, Andrea Steck, M.D.^{3,14}, Kathleen Waugh^{6,7,12,15}, Hali Wright¹². University of Colorado, Anschutz Medical Campus, Barbara Davis Center for Childhood Diabetes.

Georgia/Florida Clinical Center

Jin-Xiong She, Ph.D., PI^{1,3,4,11}, Desmond Schatz, M.D.*^{4,5,7,8}, Diane Hopkins¹², Leigh Steed^{12,13,14,15}, Jamie Thomas*^{6,12}, Janey Adams*¹², Katherine Silvis², Michael Haller, M.D.*¹⁴, Melissa Gardiner, Richard McIndoe, Ph.D., Ashok Sharma, Joshua Williams, Gabriela Foghis, Stephen W. Anderson, M.D.[^], Richard Robinson. Center for Biotechnology and Genomic Medicine, Georgia Regents University. *University of Florida, ^Pediatric Endocrine Associates, Atlanta.

Germany Clinical Center

Anette G. Ziegler, M.D., PI^{1,3,4,11}, Andreas Beyerlein Ph.D.², Ezio Bonifacio Ph.D.*⁵, Michael Hummel, M.D.¹³, Sandra Hummel, Ph.D.², Kristina Foterek^{¥2}, Mathilde Kersting, Ph.D.^{¥2}, Annette Knopff⁷, Sibylle Koletzko, M.D.^{¶13}, Claudia Peplow¹², Roswith Roth, Ph.D.⁹, Joanna Stock^{9,12}, Elisabeth Strauss¹², Katharina Warncke, M.D.¹⁴, Christiane Winkler, Ph.D.^{2,12,15}. Forschergruppe Diabetes e.V. and Institute of Diabetes Research, Helmholtz Zentrum München, and Klinikum rechts der Isar, Technische Universität München. *Center for Regenerative Therapies, TU Dresden, ¶Dr. von Hauner Children's Hospital, Department of Gastroenterology, Ludwig Maximilians University Munich, ¥Research Institute for Child Nutrition, Dortmund.

Finland Clinical Center

Jorma Toppari, M.D., Ph.D., PI^{¥^1,4,11,14}, Olli G. Simell, M.D., Ph.D.^{¥^1,4,11,13}, Annika Adamsson, Ph.D.^{^12}, Heikki Hyöty, M.D., Ph.D.*^{±6}, Jorma Ilonen, M.D., Ph.D.^{¥¶13}, Sanna Jokipuu[^], Tiina Kallio[^], Miia Kähönen^{μ±}, Mikael Knip, M.D., Ph.D.*^{±5}, Annika Koivu^{¥^}, Mirva Koreasalo*^{±§2}, Kalle Kurppa, M.D., Ph.D.*^{±13}, Maria Lönnrot, M.D., Ph.D.*^{±6}, Elina Mäntymäki^{¥^}, Katja Multasuo^{μ±}, Juha Mykkänen, Ph.D.^{^¥3}, Tiina Niininen^{±*12}, Mia Nyblom*[±], Petra Rajala[^], Jenna Rautanen^{±§}, Anne Riikonen*[±], Minna Romo^{¥^}, Satu Simell, M.D., Ph.D.^{^±13}, Tuula Simell, Ph.D., Ville Simell^{^¥13}, Maija Sjöberg^{¥^12,14}, Aino Stenius^{μ±12}, Maria Särnä[^], Sini Vainionpää[^], Eeva Varjonen^{¥^12}, Riitta Veijola, M.D., Ph.D.^{μ±14}, Suvi M. Virtanen, M.D., Ph.D.*^{±§2}, Mari Vähä-Mäkilä[^], Mari Åkerlund*^{±§}. ¥University of Turku, *University of Tampere, ^μUniversity of Oulu, ^Turku University Hospital, Hospital District of Southwest Finland, [±]Tampere University Hospital, ^²Oulu University Hospital, [§]National Institute for Health and Welfare, Finland, ¶University of Kuopio.

Sweden Clinical Center

Åke Lernmark, Ph.D., PI^{1,3,4,5,6,8,10,11,15}, Daniel Agardh, M.D., Ph.D.¹³, Carin Andrén Aronsson^{2,13}, Maria Ask, Jenny Bremer, Ulla-Marie Carlsson, Corrado Cilio, Ph.D., M.D.⁵, Emelie Ericson-Hallström, Lina Fransson, Thomas Gard, Joanna Gerardsson, Rasmus Bennet, Monica Hansen, Gertie Hansson¹², Cecilia Harmby, Susanne Hyberg, Fredrik Johansen, Berglind Jonasdottir M.D., Helena Elding Larsson M.D., Ph.D.^{6,14}, Sigrid Lenrick Forss, Markus Lundgren¹⁴, Maria Månsson-Martinez, Maria Markan, Jessica Melin¹², Zeliha Mestan, Kobra Rahmati, Anita Ramelius, Anna Rosenquist, Falastin Salami,

Sara Sibthorpe, Birgitta Sjöberg, Ulrica Swarthling, Ph.D.^{9,12}, Evelyn Tekum Amboh, Erika Trulsson, Carina Törn, Ph.D.^{3,15}, Anne Wallin, Åsa Wimar^{12,14}, Sofie Åberg. Lund University.

Washington Clinical Center

William A. Hagopian, M.D., Ph.D., PI^{1,3,4, 5, 6,7,11,13, 14}, Michael Killian^{6,7,12,13}, Claire Cowen Crouch^{12,14,15}, Jennifer Skidmore², Stephen Ayres, Kayleen Dunson, Rachel Hervey, Corbin Johnson, Rachel Lyons, Arlene Meyer, Denise Mulenga, Elizabeth Scott, Joshua Stabbert, Alexander Tarr, Morgan Uland, John Willis. Pacific Northwest Diabetes Research Institute.

Pennsylvania Satellite Center

Dorothy Becker, M.D., Margaret Franciscus, MaryEllen Dalmagro-Elias Smith², Ashi Daftary, M.D., Mary Beth Klein, Chrystal Yates. Children's Hospital of Pittsburgh of UPMC.

Data Coordinating Center

Jeffrey P. Krischer, Ph.D., PI^{1,4,5,10,11}, Michael Abbondandolo, Sarah Austin-Gonzalez, Sandra Baethke, Rasheedah Brown^{12,15}, Brant Burkhardt, Ph.D.^{5,6}, Martha Butterworth², Joanna Clasen, David Cuthbertson, Christopher Eberhard, Steven Fiske⁹, Dena Garcia, Jennifer Garmeson, Veena Gowda, Kathleen Heyman, Francisco Perez Laras, Hye-Seung Lee, Ph.D.^{1,2,13,15}, Shu Liu, Xiang Liu, Ph.D.^{2,3,9,14}, Kristian Lynch, Ph.D.^{5,6,9,15}, Jamie Malloy, Cristina McCarthy^{12,15}, Wendy McLeod, Steven Meulemans, Chris Shaffer, Laura Smith, Ph.D.^{9,12}, Susan Smith^{12,15}, Noah Sulman, Ph.D., Roy Tamura, Ph.D.^{1,2,13}, Ulla Uusitalo, Ph.D.^{2,15}, Kendra Vehik, Ph.D.^{4,5,6,14,15}, Ponni Vijayakandipan, Keith Wood, Jimin Yang, Ph.D., R.D.^{2,15}. *Past staff: Lori Ballard, David Hadley, Ph.D.* University of South Florida.

Project scientist

Beena Akolkar, Ph.D.^{1,3,4,5, 6,7,10,11}. National Institutes of Diabetes and Digestive and Kidney Diseases.

Other contributors

Kasia Bourcier, Ph.D.⁵, National Institutes of Allergy and Infectious Diseases. Thomas Briesse, Ph.D.^{6,15}, Columbia University. Suzanne Bennett Johnson, Ph.D.^{9,12}, Florida State University. Eric Triplett, Ph.D.⁶, University of Florida.

Committees

¹Ancillary Studies, ²Diet, ³Genetics, ⁴Human Subjects/Publicity/Publications, ⁵Immune Markers, ⁶Infectious Agents, ⁷Laboratory Implementation, ⁸Maternal

Studies, ⁹Psychosocial, ¹⁰Quality Assurance, ¹¹Steering, ¹²Study Coordinators, ¹³Celiac Disease, ¹⁴Clinical Implementation, ¹⁵Quality Assurance Subcommittee on Data Quality.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Mothers and fathers risk perception accuracy at 3 (moms only) 6, 15 and 27 months by GP/FDR status

Table 1

Parent	FDR/GP	Age Study Visit Month	No	Inaccurate estimation of risk			Accurate estimation of risk		
				Much lower	Some- what lower	About the same	Inaccurate estimation	Some- what higher	Much higher
			N	%	%	%	% (total)	%	% (total)
Mothers	GP	3	5248	3.0	5.5	31.2	38.7	57.3	3.0
		6	5130	3.0	5.5	31.2	39.7	57.3	3.0
		15	4896	3.1	4.8	32.7	40.5	56.9	2.6
		27	4852	2.5	4.6	32.8	40.0	57.1	3.0
Fathers	FDR	3	655	0.0	0.9	10.4	11.3	67.1	20.0
		6	639	0.5	0.6	11.7	12.8	67.1	20.0
		15	601	0.8	0.5	14.3	15.6	70.4	14.0
		27	549	0.5	0.5	12.8	13.8	71.8	14.4
Fathers	GP	6	4659	4.4	7.0	41.9	53.2	44.2	2.6
		15	4414	3.8	6.2	42.8	52.8	45.0	2.3
		27	4134	3.3	6.2	42.3	51.8	46.2	2.0
		6	588	1.0	2.2	21.8	25.0	58.8	16.2
	FDR	15	553	1.3	2.4	23.9	27.5	59.7	12.8
		27	489	0.6	2.2	23.1	26.0	63.2	10.8
									74.0

FDR = TEDDY child has a first degree relative with T1D

GP = TEDDY child has no first degree relative with T1D

Logistic regression results for mothers' risk perception accuracy at 6 and 27 month study visits

Table 2

Factor	6 months (n = 5577)				27 months (n = 5034)			
	n (%accurate)	OR	95%CI	p- value	n (%accurate)	OR	95%CI	p- value
First degree relative with T1D								
no	4968 (60.1)	1.00	ref		4509 (59.9)	1.00	ref	
yes	609 (86.7)	4.17	3.25 – 5.37	<0.001	525 (86.5)	4.14	3.17 – 5.41	<0.001
Country of residence								
US	2114 (62.9)	1.09	0.94 – 1.26		1957 (61.3)	0.91	0.83 – 1.14	
Finland	1258 (72.2)	1.75	1.48 – 2.07		1093 (72.1)	1.64	1.38 – 1.9	
Germany	358 (74.6)	1.58	1.19 – 2.09		272 (76.5)	1.69	1.21 – 2.31	
Sweden	1847 (54.7)	1.00	ref	<0.001	1712 (55.9)	1.00	ref	<0.001
Child ethnic minority								
no	4820 (64.9)	1.00	ref		4363 (64.6)	1.00	ref	
yes	757 (51.0)	0.70	0.59 – 0.83	<0.001	671 (49.9)	0.70	0.58 – 0.84	<0.001
Mother education								
Primary education	1033 (44.5)	1.00	ref		919 (44.1)	1.00	ref	
Some college/trade school	1338 (59.3)	1.47	1.24 – 1.76		1159 (57.1)	1.43	1.17 – 1.71	
Finished college	3206 (70.5)	2.52	2.15 – 2.94	<0.001	2956 (70.6)	2.71	2.30 – 3.20	<0.001
Only child								
no	3218 (63.5)	1.00	ref		2876 (63.1)	1.00	ref	
yes	2359 (62.4)	0.87	0.84 – 0.94	0.02	2158 (62.0)	0.87	0.76 – 0.98	0.03
Household crowding								
		0.89	0.84 – 0.94	<0.001		0.94	0.91 – 0.97	<0.001
Belief that T1D risk can be reduced								
		0.93	0.91 – 0.96	<0.001		0.92	0.87 – 0.98	0.01

Logistic regression results for fathers' risk perception accuracy at 6 and 27 month study visits

Table 3

Study Visit		6 months (n = 5016)				27 months (n = 4454)			
Factors		n (% accurate)	OR	95% CI	p-value	n (% accurate)	OR	95% CI	p-value
First degree relative with T1D									
	no	4460 (47.0)	1.00	ref		3984 (48.3)	1.00	ref	
	yes	556 (75.9)	3.28	2.65 – 4.05	<0.001	470 (73.8)	2.90	2.31 – 3.64	<0.001
Country residence									
	US	1827 (46.4)	0.65	0.56 – 0.76		1610 (46.5)	0.59	0.50 – .69	
	Finland	1130 (53.5)	0.94	0.80 – 1.12		1003 (56.6)	1.01	0.85 – 1.21	
	Germany	342 (68.1)	1.37	1.05 – 1.80		255 (65.9)	1.20	0.89 – 1.64	
	Sweden	1717 (48.4)	1.00	ref	<0.001	1586 (49.6)	1.00	ref	<0.001
Child ethnic minority									
	no	4390 (51.3)	1.00	ref		3923 (52.3)	1.00	ref	
	yes	626 (42.5)	0.91	0.76 – 1.10	0.33	531 (41.6)	0.90	0.73 – 1.10	0.23
Father education									
	Primary education	1232 (35.4)	1.00	ref		1061 (35.1)	1.00	ref	
	Some college/trade	1272 (48.6)	1.65	1.38 – 1.96		1121 (47.4)	1.61	1.34 – 1.95	
	Finished college	2512 (58.2)	2.63	2.25 – 3.07	<0.001	2272 (60.2)	2.94	2.49 – 3.47	<0.001
	Household crowding		0.93	0.88 – 0.98	<0.01		0.90	0.85 – 0.96	0.001
Belief that T1D risk can be reduced			0.97	0.95 – 1.00	0.05		0.95	0.92 – 0.97	<0.001