4. Asaria RH, Kon CH, Bunce C, Charteris DG, Wong D, Khaw PT, et al. Adjuvant 5-fluorouracil and heparin prevents proliferative vitreoretinopathy : Results from a randomized, double-blind, controlled clinical trial. *Ophthalmology* 2001; 108:1179-83.
5. Scheer S, Morel C, Poisson F, Bonnel S, Touzeau O, Sahel JA, et al. Prevention of proliferative vitreoretinopathy using 5-FU & heparin: clinical efficacy: *J Fr Ophthalmol* 2005; 28:701-6.
6. Garcia RA, Sanchez JG, Arevalo JF. Combined 5-fluorouracil, low-molecular-weight heparin, and silicone oil in the management of complicated retinal detachment with proliferative vitreoretinopathy grade C. *Ophthalmic Surg Lasers Imaging* 2007; 38: 276-82.
7. Wilkinson CP. Prevention of retinal detachment. In: Ryan SJ, Ed-in-chief, *Retina*. 4th Ed, Philadelphia, Elsevier 2006; 2107 – 19.
8. Charteris DG, Aylward GW, Wong D, Groenewald C, Asaria RH, Bunce C. PVR Study Group. A randomized controlled trial of combined 5-fluorouracil and low-molecular-weight heparin in management of established proliferative vitreoretinopathy. *Ophthalmology* 2004; 111:2240-5.
9. Wickham L, Bunce C, Wong D, McGurn D, Charteris DG. Randomized controlled trial of combined 5-Fluorouracil and low-molecular-weight heparin in the management of unselected rhegmatogenous retinal detachments undergoing primary vitrectomy. *Ophthalmology* 2007; 114:698-704.

Quiz

Picture Quiz

Lt Col AR Rajan*, AA Bhavet†

MJAFI 2010; 66 : 150

A nine year old girl presented with six days history of fever. On examination she had right cervical lymphadenopathy and an erythematous macular rash over her trunk and face. She was febrile (39.4°C) with proportionate tachycardia. Systemic examination including cardiovascular system was normal. Over the next 48 hours the fever persisted and she also developed bilateral nonexudative bulbar conjunctivitis (Fig.1). Examination of oropharyngeal mucosa revealed cheilitis, oral congestion and a strawberry tongue (Fig. 2). By the seventh day of fever she had also developed

erythematous swelling over her palms and soles. This was followed by desquamation seen at the fingertips of her hands and feet (Fig. 3). Haematological investigations revealed a leucocyte count of 14,600/mm³ with predominant neutrophils (72%). Her haemoglobin was 10.8g/dl. The erythrocyte sedimentation rate (ESR) was raised (46mm in 1st hour by Wintrobe's method) and the C-reactive protein was elevated (12mg/dl). Her platelet count was 4,25,000/mm³. All biochemical parameters and her electrocardiogram were normal..

What is your diagnosis?

Fig. 1 : Nonexudative bilateral bulbar conjunctivitis.

Fig. 2 : Mucosal changes including fissured lips, strawberry tongue and oropharyngeal congestion.

Fig. 3 : Desquamation of periungual skin .

Answer to Picture Quiz : pg. 166

*Classified Specialist (Paediatrics), Command Hospital (CC), Lucknow -226002. †Senior Research Scientist, CSMMU, Lucknow.