Clinicopathological Spectrum of Renal Biopsies in Children

Sqn Ldr AK Garg*, Col M Kanitkar*, Wg Cdr V Venkateshwar#

Abstract

Background: Renal biopsy has revolutionized the study of glomerular diseases. A retrospective analysis of 104 consecutive renal biopsies performed in children at a tertiary care referral centre over five years is presented.

Methods : All the biopsies were performed non-ultrasound guided by a single consultant nephrologist. Trucut needles were used in the initial few years and a Magnum biopsy gun (Bard) over subsequent three years. There were 66 boys and 38 girls.

Result: A male predominance occurred in the older and younger patients. The male: female ratio was 2.2:1, 1:1, and 2.7:1 for the age groups below five years, 5-10 years and above 10 years respectively. All patients tolerated the biopsy well and success rate was 94%. There were minimal complications in the form of post biopsy haematuria (33.3%). Haematuria was mild in most of the cases and settled down within 24 hours. None required transfusion. However, 60% patients had mild discomfort in the form of local pain. There was no mortality, infection or renal loss. The most common indication for a kidney biopsy was nephrotic syndrome. Out of 104 biopsies, 85 were in children with nephrotic syndrome. The commonest primary renal pathology was mesangial proliferative glomerulonephritis (38%), minimal change disease (19%), focal segmental glomerulosclerosis (15%) and membranoproliferative glomerulonephritis (7%).

Conclusion: Renal biopsy is a safe procedure in experienced hands and the commonest indication for a biopsy in children remains nephrotic syndrome.

MJAFI 2010; 66 : 216-219

Key Words : Nephrotic syndrome; Biopsy gun; Haematuria; Mesangial proliferative glomerulonephritis; Minimal change disease

Introduction

Since its first introduction in 1951, renal biopsy has revolutionized the study of glomerular diseases [1]. With the help of light and electron microscopy and immunofluorescent examination of renal tissues, the renal pathologist has been able to classify glomerulonephritides, to describe their clinico-pathological correlation, natural history and to study their pathogenesis [2]. The clinicians utilize the information for the diagnosis and monitoring of treatment of their patients. The pattern of glomerular diseases may differ in different population groups. We undertook a retrospective review of renal biopsies in children with an aim of analyzing the clinico-pathological spectrum, the complications and the success rate.

A spring-loaded, automated, cutting-needle biopsy "gun" was developed in the early 1980's [3]. It was quickly adopted for renal biopsies because of its ease of use, decreased risk of renal laceration and lessened pain reported by patients [4]. The use of the biopsy gun, in combination with advanced imaging techniques, primarily ultrasound (USG), has led to an increase in safety and yield [5-9].

Material and Methods

This was a retrospective study of children who underwent percutaneous renal biopsy at a tertiary care paediatric nephrology referral unit over five years. All the biopsies were performed by single consultant nephrologist and were non-USG guided. Disposable Trucut needles were used in initial few years, however over the past three years a Magnum™ automatic biopsy gun (Bard®) with disposable needles was used. The case records of these patients were reviewed to collect demographic data, clinical features, indications of renal biopsy and pathological diagnosis. Complications during the procedure were analysed.

All renal biopsies were routinely processed in a standard technique for light microscopy, electron microscopy and immunofluorescence study. Renal tissue was considered adequate for diagnosis if it contained more than four glomeruli. They were reported by the pathologists of our institution as per World Health Organisation (WHO) classification [10].

Clinical features were recorded according to a set of predetermined criteria as follows: gross haematuria was defined as urine grossly red with red blood cells (RBC) and microscopic haematuria as the presence of >5 RBC per high power field. Proteinuria was defined as morning urine albustix 2+ or more, or protein excretion rate of more than 4 mg/m²/hour. Nephrotic syndrome was defined as massive proteinuria

*Graded Specialist (Paediatrics), 7 AF Hospital, Kanpur. *Professor & Head, #Reader (Dept of Paediatrics), AFMC, Pune-40.

Received : 07.05.09; Accepted : 05.05.10 E-mail : mkanitkar15@gmail.com

of >40 mg/m²/hour, spot protein-creatinine ratio >2 , presence of edema and serum albumin concentration of <2.5 g/dl. Nephritic-nephrotic syndrome was defined as nephrotic syndrome with the presence of atypical features such as hypertension, persistent microscopic or gross haematuria, impaired renal function, low complement C₃. Suspected nephritis was considered in the presence of the above features but a normal serum albumin level of >2.5 g/dl. Renal insufficiency was defined as glomerular filtration rate of <90 ml/minute/1.73 m² and further classified using the chronic kidney disease (CKD) staging. Steroid resistance was defined as persisting proteinuria despite four weeks of prednisolone at dose of 2 mg/kg/day. Steroid dependence was defined as relapse of nephrotic syndrome on alternate day prednisolone or within two weeks of stopping prednisolone therapy. Henoch-Schönlein purpura was diagnosed when two or more of the following were present: vasculitic purpura, abdominal pain or gastrointestinal bleeding, joint pain and urinary abnormalities. Systemic lupus erythematosus (SLE) was diagnosed by the criteria of American College of Rheumatology (ACR).

The investigations carried out on each patient prior to a renal biopsy included the haemoglobin (Hb), prothrombin time (PT), activated partial thromboplastin time (APTT), platelet count and renal function test. Biopsy was done in the paediatric ward under local anaesthesia and sedation with midazolam and ketamine.

Results

From January 2003 to December 2008, a total of 104 children

Table 1

Age and gender distribution of children undergoing renal biopsy (n=104)

Age (years)	Sex		Total (%)
	Female	Male	
1-5	14	31	45 (43.5)
6-10	18	19	37 (35.5)
>10	6	16	22 (21)

Table 2

Correlation of histopathology with indications for biopsy (n=104)

Pathology	MesPGN	MCNS	MPGN	FSGS	DPGN	ESRD	Lupus	Inadequate	Others*	Total
Indication										
Haematuria	03							02	02	07
ARF						01			02	03
Hypertension	02							01		03
Lupus							04			04
CKD									01	01
Proteinuria	-		-	-		01	-			01
Nephrotic	35	19	07	15	03	-	-	03	03	85
Total	40	19	07	15	03	02	04	06	08	104
(%)	(38)	(18.2)	(6.7)	(14.4)	(2.8)	(1.9)	(3.8)	(5.7)	(7.6)	(100)

*Others: IgA nephropathy-01; Crescentic GN-02; Oxalate nephropathy-01; Chronic pyelonephritis-01; Interstitial nephritis-03.

ESRD: end stage renal disease; MesPGN: mesangial proliferative glomerulonephritis; MCNS: minimal change nephritic syndrome; MPGN: membranoproliferative glomerulonephritis; FSGS: focal segmental glomerulosclerosis; DPGN: diffuse proliferative glomerulonephritis; ARF: acute renal failure; CKD: chronic kidney disease

underwent renal biopsy at this centre. Their age at biopsy ranged from 12 months to 15 years, with 45 patients below five years of age, 37 patients between 5-10 years and 22 patients above 10 years. There were 66 boys and 38 girls. A strong male predominance occurred in the older and younger patients. The male: female ratio was 2.2: 1, 1:1, and 2.7:1 for the age groups below five years, 5-10 years, and above 10 years respectively (Table 1).

The most common indication for renal biopsy was nephrotic syndrome with either atypical presentation or steroid resistance. Out of 104 biopsies, 85 were in children with nephrotic syndrome. Among other indications seven had persistent haematuria and four had lupus.

As shown in Table 2, the commonest primary renal pathology were mesangial proliferative glomerulonephritis (MesPGN-38%) and minimal change disease (MCD-18.2%) while the tissue was inadequate in 5.7% cases.

When the indication for a biopsy was other than nephrotic syndrome, the biopsy revealed MesPGN in five (26%) cases, Lupus in four (21%) cases and crescentic glomerulonephritis and ESRD in two (10.5%) cases each.

Clinical indication and pattern of renal pathology in nephrotic children

A total of 85 children with nephrotic syndrome were biopsied. Of these, 51 patients were biopsied at onset because of atypical presentation, namely age of onset <1 year (eight cases), presence of nephritic-nephrotic features (11 cases) and initial steroid resistance (32 cases). The remaining 34 patients were biopsied at a later stage of their illness. The majority had either MesPGN (41%) or MCNS (22%) as depicted in Table 3.

Efficacy and adequacy of renal biopsy

The sample was adequate in majority of the 104 biopsies undertaken with inadequate tissue in only six samples. In 49% samples the yield was between 5-10 glomeruli and in 45% glomeruli were >10 .

Table 3**Correlation of indications for biopsy to histopathology in nephrotic syndrome (n=85)**

Pathology	MesPGN	MCNS	MPGN	FSGS	DPGN	Interstitial	Inadequate	Total	%
Indication									
Cyclosporine use	04	01				03		08	9.4
Nephritic onset	03	02	03	01	02			11	13
ISR	14	06		09	01		02	32	37.6
LSR	06	04		04				14	16.4
Early/Late onset	02	04	01	01				08	9.4
SD	04	02					01	07	8.2
ARF	02		03					05	5.9
Total	35	19	07	15	03	03	03	85	100.0

MesPGN: mesangial proliferative glomerulonephritis; MCNS: minimal change nephritic syndrome; MPGN: membranoproliferative glomerulonephritis; FSGS: focal segmental glomerulosclerosis; DPGN: diffuse proliferative glomerulonephritis; ISR: initial steroid resistant; LSR: later steroid resistant ; SD: steroid dependant ; ARF: acute renal failure.

Table 4**Complications following a renal biopsy**

Adverse effect	Present		Age (years)			Sex	
	Yes	No	1-5	5-10	>10	Female	Male
Haematuria	33.3%	66.7%	50%	31%	10%	53%	23%
Discomfort	60%	40%	75%	63%	30%	80%	50%
Haematuria + discomfort	31%	38%					

Complications

Post-biopsy haematuria was noted in 33.3% of which 53% were girls and 23% were boys (Table 4). Haematuria was mild in most of the cases and settled down within 24 hours. However, 6.6% of children required urinary bladder irrigation for a retained clot. None required transfusion. Out of 104 cases 32 (30%) had deranged renal functions before biopsy but the incidence of haematuria was same as in the children with normal renal functions. Some 60% patients had mild discomfort in the form of local pain which was more common in girls (80%) than boys (50%). While 38% patient neither had haematuria nor discomfort, 31% patient had both. There was no mortality, post-biopsy sepsis or renal loss (Table 4).

Discussion

This was a retrospective study carried out from the records and clinical data available with the department of paediatrics at our centre from 2003 to 2008. Total 104 cases records were studied. Of these, 85 cases (81.5%) were children with nephrotic syndrome having either an atypical presentation or a protocol biopsy two years after therapy with calcineurin inhibitors while 19 cases (18.5%) underwent a biopsy for other reasons. This correlates with a similar study by Lubna et al [11] who reported a retrospective study of 108 cases wherein nephrotic syndrome remained the major indication (83.3%) for biopsy.

This data does not represent a true incidence of glomerular diseases in children. Firstly, not all children with renal diseases underwent renal biopsy, such as those

with typical post-streptococcal nephritis, typical nephrotic syndrome or isolated microscopic haematuria. For nephrotic syndrome, no biopsies were performed for children aged one to ten years, without atypical features such as gross haematuria, hypertension, impaired renal function and low complement levels or those who responded to empirical steroid therapy. Similarly no children with typical acute nephritic syndrome with evidence of recent streptococcal infection were biopsied. The same was true for children with single episode of isolated gross haematuria or transient microscopic haematuria.

From our selective approach to biopsy children with nephrotic syndrome, we had a smaller percentage (22%) of minimal change disease compared to 76% of unselected cases in the report of the International study of kidney disease in childhood (ISKDC) [12]. On the other hand, we had a higher percentage (41%) of diffuse mesangial proliferative glomerulonephritis compared to 2.5% reported in ISKDC or 6.8% reported by Habib et al [13].

Our study correlates well with various other studies where selective approach was used for biopsy. Minimal change has been reported in 23-25%, MesPGN 17.6-24%, FSGS 14.8% and MPGN 10.2% [11,14,15]. Different studies have reported variable success rate of biopsy in children ranging from 76-89% [16]. Our success rate was 94% inspite of the procedure having been undertaken without USG guidance. A success rate

of 95-100% has been reported when the procedure is USG guided [11,17].

All patients tolerated the biopsy well. There were minimal complications in the form of post biopsy haematuria in 33.3%. In other similar studies, hematuria following a biopsy is present in about 35% of patients, but gross hematuria is seen in less than 0.5% of patients. A perirenal hematoma is found in as many as 65% of patients, depending upon the diligence of the search, because most are silent. Transfusion is required in less than 1% of biopsies, renal loss in less than 0.1% of cases and loss of life is extremely rare [18-21].

In conclusion, renal biopsy is safe in children under experienced hands even when performed without an ultrasound guided technique. The most common indication for a renal biopsy remains nephrotic syndrome. Haematuria is the commonest complication occurring in 33.3% cases. Common pathological lesions seen in renal biopsy of children are MesPGN, MCNS, FSGS and MPGN in decreasing order of incidence.

Conflicts of Interest

None identified

Intellectual Contribution of Authors

Study Concept : Sqn Ldr AK Garg, Col M Kanitkar

Drafting & Manuscript Revision : Wg Cdr V Venkateshwar, Col M Kanitkar

Statistical Analysis : Sqn Ldr AK Garg, Wg Cdr V Venkateshwar

Study Supervision : Col M Kanitkar

References

- Iversen P, Brun C. Aspiration biopsy of the kidney. *Am J Med* 1951; 11: 324-30.
- Habib R. A story of glomerulopathies: a pathologist's experience. *Pediatr Nephrol* 1993; 7: 336-46.
- Lindgren PG. Percutaneous needle biopsy: a new technique. *Acta Radiol* 1982; 23: 653-6.
- Burstein DM, Korbet SM, Schwartz MM. The use of the automatic core biopsy system percutaneous renal biopsies: a comparative study. *Am J Kidney Dis* 1993; 22: 545-52.
- Nicholson ML, Wheatley TJ, Doughman TM, White SA, Morgan JDT, Furness PN. A prospective randomized trial of three different sizes of core-cutting needle for renal transplant biopsy. *Kidney Int* 2000; 58: 390-5.
- Khajehdehi P, Junaid SMA, Salinas-Madrigal L, Schmitz PG, Bastani B. Percutaneous renal biopsy in the 1990s: safety, value, and implications for early hospital discharge. *Am J Kidney Dis* 1999; 34: 9-7.
- Manno C, Strippoli GF, Arnesano L, Bonifati C, Campobasso N, Gesualdo L, Schena FP. Predictors of bleeding complications in percutaneous ultrasound-guided renal biopsy. *Kidney Int* 2004; 66: 1570-7.
- Preda A, Van Dijk LC, Van Oostaijen JA, Pattynama PMT. Complication rate and diagnostic yield of 515 consecutive ultrasound-guided biopsies of renal allografts and native kidneys using a 14-gauge biopsy gun. *Eur Radiol* 2003; 13: 527-30.
- Tang S, Li JHC, Lui SL, Chang TM, Cheng IKP, Lai KN. Free-hand, ultrasound-guided percutaneous renal biopsy: experience from a single operator. *Eur Radiol* 2002; 41: 65-9.
- Churg J, Bernstein J, Glasscock RJ. Renal disease: classification and atlas of glomerular diseases. 2nd edition, Igaku-Shoin, New York, 1995.
- Lubna AL, John Amoushi, KS Ramprasad. Percutaneous renal biopsy and its findings in children and adolescent in Saudi Arabia: Saudi J Kidney Dis Transplant 1997; 8: 289-93.
- International Study of Kidney disease in children. Nephrotic syndrome in children: Prediction of histopathology from clinical and laboratory characteristics at time of diagnosis. *Kidney Int* 1978; 13: 159.
- R Habib, C Kleinknecht. The primary nephrotic syndrome of childhood: Classification and clinicopathologic study of 406 cases. In: SC Sommers, Editor. Pathology annual. Boston, Appleton-Century-Crofts 1971; 165.
- Abdurrahman MB. Percutaneous renal biopsy in a developing country: Experience with 300 cases. *Ann Trop Paediatr* 1984; 4: 25.
- Al-Rashed SA, AL Mugerien MM, AlSallouin AA. Childhood renal disease in Saudi Arabia: A clinicopathological study of 167 cases. *Int Uro Nephrol* 1996; 28: 607-13.
- Edelmann CM Jr, Greifer I. A modified technique for percutaneous needle biopsy of the kidney. *J Pediatr* 1976; 70: 81-6.
- Bollam Rengaswamy, Mahalingam Vijay Kumar, Nageshwaran Prahalad. Experience of renal biopsy in children with nephrotic syndrome. *Paed Nephrol* 2006; 21: 286-8.
- Preda A, Van Dijk LC, Van Oostaijen JA, Pattynama PMT. Complication rate and diagnostic yield of 515 consecutive ultrasound-guided biopsies of renal allografts and native kidneys using a 14-gauge Biopsy gun. *Eur Radiol* 2003; 13: 527-30.
- Hergesell O, Felten H, Andrassy K, Kühn K, Ritz E. Safety of ultrasound-guided percutaneous renal biopsy - retrospective analysis of 1090 consecutive cases. *Nephrol Dial Transplant* 1998; 13: 975-7.
- Whittier WL, Korbet SM. Timing of complications in percutaneous renal biopsy. *J Am Soc Nephrol* 2004; 15: 142-7.
- Lin WC, Yang Y, Wen YK, Chang CC. Outpatient versus inpatient renal biopsy: a retrospective study. *Clin Nephrol* 2006; 66: 17-24.