Large Bowel Obstruction by Anomalous Congenital Band

Sqn Ldr A Kumar*, Gp Capt TS Ramakrishnan⁺, Wg Cdr S Sahu[#]

MJAFI 2009; 65 : 378-379

Key Words : Congenital band; Large bowel obstruction

Introduction

The incidence of obstruction of large bowel is 3-4 times less frequent than the small bowel. The main causes of large bowel obstruction are malignancy, volvulus, diverticulae, inflammatory bowel disease and pseudo-obstruction [1]. Anomalous congenital bands (ACB) causing large bowel obstruction is a rare occurrence and to best of our knowledge, only three such cases has been reported in literature [2-4]. There is no report of congenital band causing both large bowel obstruction and jejunal ischemia. We report an unusual case of intestinal obstruction in a 76 year old lady who had an ACB obstructing the transverse colon causing a closed loop obstruction, caecal perforation and ischemia of proximal jejunum along with formation of pneumatosis cystoides intestinalis (PCI).

Case Report

A 76 year old lady was admitted with a history of crampy mid abdominal pain on and off for the last one month. She had constipation, vomiting and abdominal distention of two days duration prior to presentation. She was passing flatus occasionally and there was no history of fever. She had no significant past illnesses and had not undergone any abdominal surgery. On examination, she was afebrile with signs of mild dehydration. She had central abdominal distention with tenderness in right iliac fossa and bowel sounds were tympanic. There was no palpable lump or rebound tenderness. The hernial orifices were normal. Per rectal examination was non contributory.

Complete blood counts and biochemical parameters revealed hemoglobin of 7.1 g/dl with normal leucocyte count and serum electrolyte levels. Abdominal radiograph (erect and supine) revealed a few dilated small bowel loops with no significant air fluid level. However the ascending and transverse colon were dilated and loaded with faecal matter. A contrast enhanced computed tomography scan (CECT) of abdomen revealed progressive dilatation of the colon up to the distal third of transverse colon with faeces (Fig. 1). The descending colon was collapsed. An abdominal radiograph

done following the CT scan showed dilated ascending and transverse colon, which was highlighted by the retained contrast given for CECT abdomen, with an abrupt cut of the distal transverse colon (Fig. 2).

The patient was taken up for emergency laparotomy wherein she was found to have an ACB forming a loop around the distal transverse colon (Fig. 3) and causing closed loop obstruction with caecal perforation. The colon proximal to obstruction was loaded with inspissated faeces and was dusky. This band was arising from the right paravertebral region and looping around the transverse colon, ending at the root of mesentery. This band was in addition, causing compression of blood supply to the proximal jejunum leading to gangrene and PCI. There was no associated intestinal malrotation and the band contained blood vessels. The ACB was divided between ligatures and extended right hemicolectomy, resection of gangrenous jejunum with jejuno-jejunostomy was done. The patient was stable till the second post operative day when she had an acute coronary event and died.

Discussion

Congenital bands cause 3% of all intestinal obstruction and almost always lead to small bowel obstruction [5]. In adults, obstruction due to bands is even rarer. ACBs are bands that have no identifiable embryological or acquired basis. A literature search for ACBs causing large bowel obstruction yielded only three cases – that of a band between right lobe of liver and ascending colon [2], a case of congenital mesocolic band causing sigmoid colon obstruction [3] and another case of a parietocolic band obstructing the descending colon. [4].

There are four types of congenital peritoneal band. These bands are associated with intestinal malrotation, usually causing duodenal compression and are fibrous in nature [6]. ACBs, as in our case, are not associated with malrotation and contain blood vessels. These bands may represent a congenital mesenteric anomaly [2].

A congenital band causing large bowel obstruction

*Graded Specialist (Surgery), ⁺Senior Advisor (Surgery and Orthopaedics), 7 Air Force Hospital, Kanpur. [#]Graded Specialist (Radiology), Institute of Aerospace Medicine, Vimanpura, Bangalore.

Fig. 1 : CECT abdomen revealing dilated proximal part of transverse colon with fecal content and scanty contrast (arrow heads).

Fig. 2 : Radiograph of abdomen showing the dilated proximal loop of transverse as well as ascending colon (arrow head).

and becoming symptomatic so late in life is rare. Possibly, the obstruction was not severe to cause significant compromise to the lumen but the inspissated faeces may have led to the band becoming symptomatic and ultimately causing closed loop obstruction and caecal perforation. In addition this case is peculiar because of development of ischemia of proximal jejunum due to compression of blood vessels by the band and the development of PCI in the gangrenous segment of jejunum. The most common cause of PCI is bowel necrosis due to bowel ischemia, infarction, necrotizing enterocolitis, neutropenic colitis, volvulus and sepsis [7].

Despite the availability and wide use of modern imaging techniques, preoperative diagnosis of a band causing the obstruction is very difficult to establish. Plain films are nonspecific while a barium series may provide clues to narrow the differential diagnosis. In the present case, radiograph following CECT abdomen gave more information which was akin to a barium study. A CT

Fig. 3 : Intra-operative photograph showing the congenital anomalous band (arrow heads) causing complete obstruction of the transverse colon.

scan of abdomen is also unlikely to detect a band. A diagnostic laparoscopy is a valuable aid to diagnosis and definitive management [8].

Surgical treatment is the cornerstone of management. With the advent of minimally invasive surgery, laparoscopy may be safe and feasible in the diagnosis and treatment of a congenital band [8].

In conclusion, the possibility of an anomalous congenital band must be considered in the differential diagnosis of elderly patients with symptoms and signs of bowel obstruction in those with no history of abdominal surgery, trauma or clinical hernia.

Conflicts of Interest

None identified

References

1. Singh M, Monson JRT. Large bowel obstruction. In: Recent Advances in Surgery. 25th ed. Johnson CD, Taylor I. The Royal Society of Medicine Press Limited 2002; 117-34.
2. Akgur FM, Tanyel FC, Buyukpamukcu N, Hicsonmez A. Anomalous congenital bands causing intestinal obstruction in children. *J Pediatr Surg* 1992; 27:471-3.
3. Lin DS, Wang NL, Huang FY, Shih SL. Sigmoid adhesion caused by a congenital mesocolic band. *J Gastroenterol* 1999; 34 : 626-8.
4. Aphinives P, Pukkem A, Bhudisawasdi V. Descending colon obstruction by a parietocolic band. *J Med Assoc Thai* 2002; 85: 1042-45.
5. Perry JF, Smith A, Yonehiro EG. Intestinal obstruction caused by adhesions; a review of 388 cases. *Ann Surg* 1955; 142: 810.
6. Wayne ER, Burrington JD. Extrinsic duodenal obstruction. *Surg Gynecol Obstet* 1973; 136:87-91.
7. Pear B L. Pneumatosis intestinalis: A review. *Radiology* 1998; 207: 13-9.
8. Wu JM, Lin HF, Chen KH, Tseng LM, Huang SH. Laparoscopic diagnosis and treatment of acute small bowel obstruction resulting from a congenital band. *Surg Laparosc Endosc Percutan Tech* 2005; 15:294-6.