4. Bhattacharya P, Chandra PK, Datta S, et al. Significant increase in HBV, HCV, HIV and syphilis infections among blood donors in west Bengal, East India 2004-2005: Exploratory screening reveals high frequency of occult HBV infection. *World J*

Gastroenterol 2007; 13:3730-3.

Surg Cdr CN Chaudhari*

*Reader, (Dept of Microbiology), AFMC, Pune- 411040.

REPLY

I would like to thank the reader for thought provoking and extremely relevant points that have been raised in response to our study.

I completely agree that the percentage of voluntary donors in our study at 14.3% is very low. However, due to sustained efforts at recruiting voluntary donors in association with various voluntary blood donation organizations, the percentage of voluntary donors in the blood bank of the hospital where the study was carried out has progressively increased to 25.99% in 2006, 35.9% in 2007 and 42.27% in 2008, the years subsequent to the study period.

The recommendation of replacement donors to meet the requirement of blood in the setting of Armed Forces based on the results of this study was made only because of the shortage of voluntary donors, since both groups come from presumably a similar background with a similar risk of transfusion transmitted disease in Armed Forces. We agree that data from one study cannot be extrapolated to the entire Armed Forces Medical Services and

that, in the larger national perspective, voluntary donations are to be encouraged.

The hypothesis of better socioeconomic status of blood donors was only made in the context of replacement donors who are the relatives of serving personnel. Since they are widely scattered, often coming from great distances to donate blood for their relatives, it will be extremely difficult to recruit and retain them as voluntary donors for the Armed Forces, though they can be motivated to donate blood for society in general.

In the future, as the reader has pointed out, the use of newer tests and strategies will undoubtedly lead to further improvements in the safety of blood transfusion.

Col A Chattoraj*

*Classified Specialist (Pathology), Transfusion Centre (EC), Alipore Road, Kolkata- 700027

Journal Scan

Desai G, Babu GN, Rajkumar RP, Chadra PS. More questions than answers! Clinical dilemmas in psychopharmacology in pregnancy and lactation. *Indian Journal of Psychiatry* 2009; 51: 26-33.

The practice of psychiatry has undergone a sea change in the last one decade with newer understanding of underlying biochemical mechanisms of the disorders and introduction of a plethora of psychopharmacological agents. Psychiatric disorders during pregnancy and postpartal period are common. In view of small family norms and all pregnancies being precious, a thorough understanding of teratogenicity of psychotropics and their possible adverse effects on the new born during lactation is a must by all the clinicians. This will help them to take an informed decision on treatment and also in help safeguarding the physician from avoidable litigations.

The authors have covered this complex issue in a simple and

lucid way. They have done an extensive literature search and explained the matter with examples from clinical practice. They have reviewed most of the commonly prescribed neuroleptics, antidepressants and benzodiazepines from the risk benefit perspective. The required follow-up and investigations like ultra sonography and amniocentesis wherever required have also been suggested in the article. The authors have done an excellent job and this paper is a must read not only for psychiatrists but also for the gynecologists and family physicians involved in the care of pregnant and lactating women.

Contributed by

Col PS Bhat*, Col AK Mitra⁺, Col A Anand[#]

*Reader (Department of Psychiatry), AFMC, Pune. [#]Classified Specialist (Psychiatry), ⁺Senior Advisor (Psychiatry), Command Hospital, Lucknow, UP-226002