A Case of Intramedullary Epidermoid Cyst

Lt Col MN Swamy*

MJAFI 2008; 64 : 72-73

Key Words : Intramedullary cyst; Epidermoids

Introduction

Intramedullary epidermoids constitute about 0.7 % of all spinal tumours [1]. Most spinal epidermoids are intradural extramedullary in location and are usually associated with spinal dysraphism. Intramedullary epidermoids are often seen in the dorsal region and less commonly in the lumbosacral region [2]. They may rarely assume giant proportion [3].

Case Report

A 17 year old youth presented with a five year history of progressive spastic paraparesis. The weakness was distal to

proximal. The weakness initially involved the right lower limb and then the left. There was no history suggestive of axial muscle weakness. The bowel and bladder functions were not impaired. At the time of admission the patient was ambulant without any assistance. On examination the power in the lower limbs was 4+/5. The lower limb deep tendon jerks were brisk and the plantars were extensor. The patient had loss of posterior column sensations below the level D6. Magnetic resonance imaging of the dorsal spine showed an oblong shaped well defined intramedullary lesion at the level of D4. The lesion was iso-intense on T1 and partly hyper intense on T2 weighted MR images. There was a halo surrounding the lesion without enhancement (Fig. 1).

Fig. 1 : T1 and T2 weighted images of dorsal epidermoid

Fig. 2 : Photo micrographs of epidermoid

*Classified Specialist (Surgery and Neurosurgery) Command Hospital (SC) Pune-40

Received : 05.10.2006; Accepted : 03.05.2007

Email : drnarayanswamy@gmail.com

The tumour was approached by a three level laminectomy extending from D4 to D6. A distinct bulge of the dura was noted at D4. On opening the dura, the cord was seen to be distended at this level but the tumour was not visible on the surface of the cord. A mid line myelotomy was performed which showed a pearly white tumour. The lesion was well encapsulated and total excision could be achieved.

The tumour was sent for histopathological evaluation which confirmed the diagnosis of an epidermoid (Fig. 2).

The patient had an uneventful post operative course. He recovered well and at the end of three months was capable of running and cycling. Clinical examination showed power of 5/5 in both the lower limbs although residual spasticity persisted.

Discussion

Epidermoid cysts of the spinal cord are rare lesions. They account for 0.7% of all intradural spinal cord lesions with a reported incidence ranging from 0.2 to 1% [1, 2]. These tumours are commonly intradural and extra medullary in location. Our patient had the rare intramedullary variant. A giant variant of spinal epidermoids has been described in literature [3].

Epidermoid cysts are postulated to have their origin from misplaced nests of epidermal cells. These cell nests occur along the lines of neural tube closure and a patent communication with exterior may persist which may act as portal of entry of infection. Intramedullary abscesses associated with epidermoid cysts have been reported [4]. Another acquired variant of epidermoid cyst associated with repeated lumbar punctures has also been described [5].

The classical presentation is progressive spastic paraparesis in a young individual. Early involvement of bladder and urinary tract infections as the initial presentation has been described [6]. Imaging

characteristics of intramedullary epidermoids may be misleading [1]. In this patient the imaging was evaluated preoperatively at a neuro radiological forum, where the consensus of opinion favoured a diagnosis of intra medullary angiomatous malformation. The intra operative impression of an intramedullary epidermoid was later confirmed by histopathological examination.

Epidermoid tumours are benign developmental lesions and the primary management modality is surgical. Cases of recurrent intramedullary epidermoids have been controlled by radio therapy as an alternative to surgery [7]. These lesions respond favourably to total surgical excision with a good outcome.

Conflicts of Interest

None identified

References

1. Chandra PS, Manjari T, Devi BI, Chandramouli BA, et al. Intramedullary Spinal Epidermoid Cyst. *Neurology India* 2000; 48:75-7.
2. Sudersan L, Paul T R, Raju B V, Ramakrishnamurthy T, et al. Cyst of the Central Nervous System: A Clinical Pathologic study of 145 cases. *Neurology India* 2001;49: 237-42.
3. Cataltepe O, Berker M, Aklan N. A Giant Intramedullary Spinal Epidermoid Cyst of the Cervico Thoracic Region. *Pediatric Neurosurgery* 2000; 40: 120-23.
4. Mongia S, Indira Devi B, Shaji KR, Hegde T. Spinal Subdural Epidermoids - A Separate Entity. *Neurology India* 2000; 50: 529-31.
5. Shigeeen M, Norileta K, Murari N, Takeshi K, et al. Acquired Lumbar Epidermoid Cyst in adult. *Neuralgia Medico-Chirurgica* 2005; 45: 271-9.
6. Ferrara P, Costa S, Rigante D, Mule A, et al. Intramedullary epidermoid cyst presenting with abnormal urologic manifestation. *Spinal Cord* 2003 ; 41: 645-8.
7. Bretz A, Vandam B, Dirk S, et al. Intra Spinal Epidermoid Cyst successfully treated with Radiotherapy. *Neurosurgery* 2003;53:1429-32.

Answers to MCQs

1. (d)	2. (b)	3. (c)	4. (b)
5. (b)	6. (d)	7. (a)	8. (d)
9. (a)	10. (a)		