Unusual Complication of Ventriculoperitoneal Shunts: Anal Extrusion

Dear Editor,

We read with great interest the case report, 'Unusual complication of ventriculoperitoneal shunts: Anal extrusion' published in MJAFI 2007; 63(1): 82-4.

Anal extrusion of the abdominal end of a shunt catheter is a very rare complication but has significant scope for serious and life threatening sequelae, as very well brought out in the paper. The most common clinical presentation is meningitis, which occurs in about 43% of these cases. Fewer than 25% of such cases may present with peritonitis [1]. Various mechanisms have been suggested with regard to the pathogenesis of the perforation viz. foreign body reaction, pressure necrosis of intestinal wall by the tube, and silicon tube allergy. The catheter most commonly associated with perforations, is the Ralmondi spring coiled catheter. The introduction of softer, more flexible silastic tubing has reduced, but not totally eliminated the incidence of bowel perforation [2].

However, a few points in the paper need to be clarified. Standard teaching recommends a different approach to such cases. A small incision is given over the shunt catheter over the abdomen (previous abdominal insertion site could also be chosen), the catheter is located, amputated and exteriorised. The anal protruding distal segment is then pulled out from the anal opening, pre empting any chance of peritoneal or subcutaneous track contamination with luminal microorganisms. Sequential cerebrospinal fluid (CSF) analysis is done daily from the exteriorised catheter and when infection has been conclusively ruled out, revision into a fresh abdominal site can be done [3]. However, some surgeons believe such an anally extruded shunt to be a potentially contaminated system in spite of normal CSF reports and prefer to remove the whole shunt assembly and place a fresh shunt on the contra lateral site.

This thinking is actually reflected in the algorithm described by the authors (Fig. 3) and we fail to understand why the same

algorithm was not applied in Case 2. The authors have described removal of lower end of an anal extruded shunt catheter through a neck incision. The procedure of pulling out the entire distal catheter through a neck incision potentially causes contamination of not only the peritoneal cavity but also the subcutaneous track up to the neck incision. As CSF analysis showed no evidence of infection, replacement with a fresh distal catheter was done at the same sitting. Leaving behind and using the original ventricular catheter and tunnelling a fresh abdominal catheter through such a potentially contaminated track at the same sitting in spite of normal CSF reports is fraught with danger. This is clearly reflected in the complications and morbidity that followed in Case 2.

It is amazing how a recurrent extrusion could take place within 48 hours (Case 2). As described elegantly in the pathophysiology by the authors, the process of bowel penetration is slow and takes place over a period of time. This repeat extrusion within 48 hours could probably be explained by inadvertent intra luminal placement of abdominal catheter during the repeat procedure [3].

References

1. Sharma A, Pandey AK, Diya B, et al. Management of ventriculo-peritoneal shunt protruding through anus. *Ind J Surg* 2006; 68: 173.
2. Hornig GW, Shillito J Jr. Intestinal perforation by peritoneal shunt tubing: report of two cases. *Surg Neurol* 1990; 33: 288-90.
3. R Michael Scott. Shunt Complications. In: RH Wilkings, SS Rengachary, editors. *Neurosurgery*. 2nd edition. New York: McGraw Hill, 1996: 3655-64.

Wg Cdr SGS Datta*, Col HC Pathak*

*Graded Specialist, *Senior Advisor (Surgery & Neurosurgery), Command Hospital, Air Force, Bangalore.

Reply

Dear Editor,

We thank the authors for their comments on the paper, 'Unusual complication of Ventriculoperitoneal shunt: Anal extrusion'.

We are in full agreement with the observation that irrespective of clinical manifestation of meningitis or whether the CSF is clear and negative on microscopic examination, under no circumstances should a fresh shunt be placed at the time of initial presentation.

Why we failed to apply the same algorithm in Case 2? The algorithm was made after our experience with Case 2. The very purpose of this manuscript was to highlight our failure to adhere to established management principles in Case 2 and bring to the notice of the medical fraternity the host of problems that we faced as a result of it, along with the attendant morbidity that the patient was subjected to. The message is very clear. Wait out a period of observation to let even low grade infection manifest itself, in which case it is not important whether the shunt is pulled out in a retrograde manner from the neck or from the subcostal region of the abdomen

where the shunt tube dips into the peritoneal cavity.

However, we feel that the retrograde pulling for the removal of an anal extrusion of a shunt will not cause contamination of the peritoneal cavity since the fibrosed shunt tube track is supposed to be fused to the serosa of the intestine.

We have no answer as to how the anal extrusion of the shunt occurred within forty eight hours. We have documented what happened in this patient. However, it is just not possible that there was an inadvertent intra-luminal positioning of the shunt catheter since while placing a VP shunt, the peritoneal cavity is opened by an incision in the subcostal region and the peritoneal end of the shunt tubing is placed in the peritoneal cavity in the suprahepatic compartment.

Col R Handa (Retd)*, Col R Kale*, Col MM Harjai*

*Ex-Classified Specialist (Paediatric Surgery), Command Hospital (EC), Kolkata. *Senior Advisor (Paediatric Surgery), AH (R&R), Delhi Cantt.