or ADP alone (Type-III). The aggregation studies required for diagnosis of sticky platelet syndrome was not done in this study as facilities of an aggregometer was not available then. The same has now been installed in the centre and is being included in future studies on thrombophilia cases.

- (d) The association of oral contraceptives with thrombophilia is a well known fact. In this study, out of 42 patients, 19 patients were female in the age group of 02 years to 73 years. Of these, five patients were on oral contraceptives for durations varying from 6 months to 2 yrs, of whom only one patient had APCR.
- (e) Malignancy is an important cause of thrombophilia which has been emphasised in this article and elsewhere also in studies from our centre [3].

Systemic Sclerosis with Early Pregnancy

Dear Editor,

This is with reference to the case report titled 'Anaesthetic Management of a case of Early Pregnancy with Systemic Sclerosis' (MJAFI 2005; 61: 287-8). The authors are complimented for the successful management of medical termination of early pregnancy in an old case of Systemic Sclerosis. However, the readers will benefit if few points could be clarified further.

1. It is mentioned in the article that the measurement of arterial pressure can be difficult and misleading especially with a reduced forearm blood flow. However, in the case report, the readers do not get to know how the arterial pressure was monitored, what the site was and whether it correlated clinically. The readers would benefit by knowing various possible methods of measurement of blood pressure and also, oxygen saturation, without which the basic monitoring in such a high risk case would be incomplete.
2. This patient appears to be a case of difficult airway with features of restricted temporomandibular joint mobility, Mallampatti III and 3cms inter-incisor distance. It is mandatory that when such a case is administered general anaesthesia (GA), a difficult airway cart be ready with various airway adjuncts like various sized endotracheal tubes, laryngeal mask airway,

References

1. Ahmed RPH, Gupta PK, Kannan M et al. Factor V Leiden – the commonest molecular defect in arterial and venous thrombophilia in India. *Thrombosis Research* 2003; 110: 19-21.
2. Gupta PK, Ahmed RPH, Kannan M et al. Protein C system defects in Indian children with thrombosis. *Ann Haematol* 2005, 84 : 85-8.
3. Gupta PK, Charan VD, Kumar H. Cancer related thrombophilia : Clinical importance and management strategies. *JAPI* 2005, 53:877-81

Brig M Jaiprakash

Dy Comdt, CH(SC), Pune-40

combitube or fiberoptic bronchoscope, notwithstanding the expected duration of surgery.

3. Regional techniques are acceptable for peripheral surgeries in patients with Raynaud's phenomenon [1]. In this case, a paracervical, caudal or a low subarachnoid blockade could have been considered using limited doses of local anaesthetic. The patient can be counselled properly, explaining the risks of GA. Distortion of regional anatomy may cause difficulty in spinal procedures but a paramedian approach in sitting position, usually gives successful outcome.
4. It would have been a greater concern for the anaesthesiologist had the patient reported for a Caesarean section or any other major surgery. The discussion of the same would have been quite interesting.

References

1. Stoelting RK. Peripheral vascular disease. In: Stoelting RK, Dierdorf SF, editors. *Anaesthesia and co-existing diseases*. 3rd ed. Churchill Livingstone. 1998; 123-30.

Maj R Goyal

Graded Specialist (Anaesthesia), 164 Military Hospital, C/o 99 APO.

REPLY

Dear Editor,

We wish to thank the respondent for the interest shown in the article.

1. Measurement of arterial pressure is a challenge, specially if the duration of surgery is long and it involves blood loss. Vasoconstriction tends to make measurement of blood pressure by auscultation unreliable, necessitating the use of an ultrasonic blood pressure sensor [1]. We used a standard adult Non Invasive Blood Pressure cuff on the right arm to measure BP during the procedure. It was reliable in our case and correlated well clinically. The thighs or legs can be used as alternate sites for blood pressure measurement keeping in mind the size of the cuff and adequacy of circulation. As discussed invasive BP monitoring should be avoided specially in the presence of Raynaud's phenomenon. Preventing the occurrence of Raynaud's phenomenon has been discussed in the article. Keeping the patient warm is of utmost importance in this

situation. Despite measures taken by us, measurement of oxygen saturation was not possible. Depending on the type of surgery and likely duration, invasive monitoring by performing arterial blood gas analysis may be considered. The location and site of arterial puncture require careful selection.

2. Our patient had a difficult airway and we anticipated it. Mask ventilation however seemed possible. A difficult airway cart as suggested was kept ready and available. It did not have combitubes or a fiberoptic laryngoscope, as they were not available in our hospital at the time of performing this procedure. We were prepared to perform an emergency tracheostomy if the situation so demanded.
3. Regional techniques are more acceptable and were offered to the patient. There was no distortion of regional anatomy in our patient (for which a spinal by paramedian approach could have been performed). Despite comprehensive counselling our patient agreed only for a general anaesthesia.