Case of Objective Tinnitus

Lt Col (Mrs) RS Bhadauria*, Col YS Sarma⁺

MJAFI 2005; 61 : 391-392

Key Words: Palatal myoclonus, Tinnitus

Introduction

Tinnitus is defined as phantom auditory sensation without an acoustic generator. In certain instances there may well be an acoustic generator in the form of underlying pathology. Objective tinnitus is audible to an observer whereas subjective tinnitus is not. It can be treated and the experiences prove to be rewarding. Subjective tinnitus on the other hand can not be demonstrated and is difficult to treat.

Case Report

A 53 year old chageman from a factory in Jabalpur reported with complaints of a ticking sound in the left ear over last 4 years. The sound was intermittent and could be reportedly heard by his children who described it as the sound of a ticking clock! He gave history of nasal allergy and discharge from left ear 6 years back. There was no history of deafness or vertigo. Whenever he reported for the ticking sound in the ear he was managed as a case of Eustachian catarrh.

His general examination was noncontributory. Both the tympanic membranes were intact but the left showed evidence of old healed perforation. There was no visible movement of the tympanic membrane when observed under magnification.

Tuning fork tests were normal with weber lateralized to the left. Hearing was 20ft in both ears. A ticking sound could be heard coming from the left ear on going close to the ear. Posterior rhinoscopy revealed bilateral vertical movement of the soft palate synchronous with the ticking sound. Systemic examination, Pure tone audiometry and impedance studies did not reveal any significant abnormality.

CT Scan of posterior fossa and brain stem revealed a 7.5x4mm well defined slightly hypodense area in posterior part of left basal ganglia and calcification of the pineal gland and choroids plexus of both lateral ventricles. MRI did not reveal any significant abnormality indicating artifact.

A diagnosis of ESSENTIAL PALATAL MYOCLONUS was made as no secondary cause could be detected, and the patient was put on tablet sodium valporate 600mg daily. This reduced the tinnitus a great deal and gave him much relief. He reduced the dose after 2-3 months to 400 mg daily and still

felt comfortable with only occasional tinnitus. After a period of 6 months he stopped the medication altogether and reported tinnitus only when he developed an upper respiratory tract infection.

Discussion

Myoclonus refers to brief, involuntary twitching of a muscle or a group of muscles. It describes a symptom and is not a diagnosis. Familiar examples are hiccups and sleep starts that are experienced when a person is drifting off to sleep. Misdiagnosis of the sound as an auditory hallucination can result in prolonged treatment with antipsychotic medication [1].

Palatal myoclonus (PM) is characterized by rhythmic involuntary movements of the soft palate usually at frequencies of 1 to 2 per second. This causes the tensor and levator palati muscles to pull on the Eustachian tube. The resulting sound is transmitted directly to the middle ear. It is difficult to say if this is associated with middle ear myoclonus i.e. tensor tympani muscle contractions and stapedius muscle myoclonus [2]. Irregular twitching in admittance recordings during manual tympanometry may be more commonly seen with middle ear myoclonus. In our patient impedance studies were normal. Automatic impedance audiometer was used and therefore the irregular tracings expected in such a case could not be seen. This condition is also called palatal tremor or palatal nystagmus and is a type of segmental myoclonus [3]. It may be audible to the examiner by placing a stethoscope over the patient's ear [4]. The condition can occur at any age but is common in the fourth decade. It has been described in infants of 18 months and the oldest reported case is 91 years old. It persists during sleep and has been implicated in sleep apnoea.

Essential PM may be more variable than secondary PM. It can often be stopped by the patient, modified by neck position or eliminated on mouth opening. The eyes are rarely involved and it often resolves

*Classified Specialist (ENT), *Senior Advisor (Medicine and Oncology), Military Hospital, Jabalpur.

spontaneously [5]. Basically, all lesions which interrupt the hypothetical circuit (lateral superior cerebellar peduncle, brachium conjunctivum and dentate nucleus) will result in PM. This circuit is called the triangle of Guillain and Mollaret (Fig 1). Most common finding is hypertrophic degeneration of the inferior olivary nucleus involving the Guillain Mollaret triangle [6]. Most theories propose an involvement of the pathways connecting the cerebellum to the inferior olivary nucleus. Palatal myoclonus secondary to cerebral disease is believed to be a result of spontaneous discharge of an enlarged inferior olive due to ipsilateral brainstem disease often involving the eyes [7].

About 70% of PM is secondary to cerebral disease. There is increased glucose metabolism in the medulla which suggests that the inferior olivary nucleus, or the surrounding areas are hyper metabolic and can be pacemaker for the Palatal myoclonus [8].

Treatment consists of antiepileptics such as clonazepam, primidone and sodium valproate. Our patient discontinued medication after 6 months and reported occasional tinnitus during an upper respiratory infection or congestion, which implies that these may be the triggering factors in the generation of palatal myoclonus. Scan of literature does not mention the period for which these drugs should be taken. Recently, botulinum toxin injections have yielded some success when each half is injected alternately to prevent nasal regurgitation of fluids [9].

The recommended imaging is thin sections through the medulla, preferably a T1-MRI of the posterior fossa with gadolinium. Olivary hypertrophy can occur without palatal myoclonus and vice versa but they are frequently associated. Manually operated impedance audiometer would help to show the classical irregular twitching pattern during admittance [10].

Palatal myoclonus is a clinical diagnosis and differential diagnosis is Whipple's disease associated with an oculomasticatory myorhythmia. In these cases eyes have a pendular vergence oscillation accompanying paralysis of vertical gaze and there are contractions of the masticatory muscles. Intestinal biopsy establishes the diagnosis in Whipples. Other causes of objective tinnitus are tumors in the middle ear cleft with or without intracranial involvement.

Our patient had gone from pillar to post for his problem and remained undiagnosed for 4 years, as probably the PM may not have been manifest at the time he reported or was missed. Though movement of the soft palate was bilateral, tinnitus was experienced only in the left ear, probably because Weber was lateralized to the left ear on account of old healed CSOM. Having got an

Fig. 1 : Guillain-Mollaret triangle : Area bounded by red nucleus, inferior olive and dentate nucleus of the cerebellum

explanation for the troublesome complaint and seen the Palatal Myoclonus on the video the patient was much relieved and discontinued the medication once the frequency reduced. This stresses the fact that explanation of a troublesome condition is half the treatment and many patients may do well without any medication.

References

1. Marym N, Gholam R, Mir-Sepasi. Essential Palatal Myoclonus Psychiatric Symptoms. Roozbeh Hospital, Tehran University of Medical Sciences, Tehran, Iran. Shiraz E-Medical journal 2000 Oct; Vol 3; No.4.
2. Oliveria CA, Negreiros Junior J, Cavalcante IC, Bahmad Junior F, Venosa AR. Palatal and middle ear myoclonus: a cause for objective tinnitus. *Int Tinnitus J* 2003; 9(1): 37-41.
3. Jankovic J, Pardo R. Segmental myoclonus: Clinical and pharmacological study. *Arch Neurol* 1986; 43: 1025-31.
4. Vieregge P, Klein C, Gherking E, Kortke D, Kompf D. The diagnosis of "essential palatal tremor". *Neurology* 1997; 49: 248-9.
5. Coles RRA. Tinnitus and its management. In Dafydd Stephens, Alan GK, John G, editor's. *Scott-Brown's Otolaryngology: Adult Audiology*. 6th ed. London: Butterworths. 1997; 2/18/4.
6. Yokota T, Hirashima F, Ito Y, Tanabe H, Furukawa T, Tsukagoshi H. Idiopathic palatal myoclonus. *Acta Neurol Scand* 1990; 81: 239-42.
7. Matsuo F, Ajax ET. Palatal Myoclonus and denervation supersensitivity of the central nervous system. *Ann Neurol* 1979; 5: 72-8.
8. Dubinsky RM, Hallet M, Di Chiro G, Fulham M, Schwankhaus J. Increased glucose metabolism in the medulla of patients with palatal myoclonus. *Neurology* 1991; 41: 557-62.
9. Bogucki A. Botulinum toxin injections under electromyography guidance. *Neurol Neurochir Pol* 1998; 32: 85-93.
10. ME Lutman. Diagnosticaudiometry. In Dafydd Stephens, Editor. *Scott-Brown's Otolaryngology: Adult Audiology*. 6th ed. London: Butterworths. 1997; 2/12/12-14.