EARLY CONGENITAL SYPHILIS: FORGOTTEN BUT NOT YET GONE

Dear Editor,

Congenital syphilis often involves multiple organ systems of body and may be confused with many other diseases of the newborn. With lot of stress on prevention of HIV infection, little concern is being raised about the increasing number of babies born with congenital syphilis. We have encountered two such cases within a span of 5 months emphasizing that syphilis is forgotten but not yet gone.

First baby was a 26 day old female neonate, born to fifth gravida, unbooked mother with history of three abortions, admitted with complaints of excessive crying and paucity of movements of both lower limbs and left upper limb of about 20 days duration. There was no history of abnormal movements, apneic spells, or trauma. Baby maintained a posture of flexed lower limbs at hip and knee. Left elbow and both knee joints were tender. She had pallor and firm hepatosplenomegaly of 3 cm and 2 cm respectively. There were no other positive clinical findings. Her Hb was 9.7 gm/dl with normochromic normocytic anaemia. TORCH titre and HIV test were negative. There was no evidence of tuberculosis in mother and baby. VDRL was reactor for both infant and mother in dilutions of 1:128 and 1:32 respectively and blood TPHA was positive for both. Kiddygram revealed metaphyseal fuzziness of distal end of both femur, metaphyseal fractures at lower end of humerus. Cerebrospinal fluid analysis was normal and VDRL and TPHA were negative. Baby and parents were treated adequately. Follow up VDRL showed a downfall titre and radiological resolution was seen after 10 months.

Second case comprised a female neonate born to booked and immunized primigravida mother at 35 weeks gestation was found to have firm hepatomegaly of 5 cm and splenomegaly of 3 cm at birth. No other clinical signs were detected. Antenatal period was uneventful. There was no history of tuberculosis in mother or close relative. VDRL in the first trimester of pregnancy was non-reactor. Her hemogram, platelet count, and liver enzymes were within normal limit. Blood culture was sterile. Torch titre and HIV test were negative. VDRL was reactor in dilutions of 1:128 and 1:32 for infant and mother respectively. TPHA was positive in both mother and child. Cerebrospinal fluid study was within normal limit and VDRL was also negative. Kiddygram showed cat bite appearance in lower end of both femur and evidence of periostitis in left humerus. Newborn was managed with Inj sodium penicillin for 14 days. Follow up VDRL titre declined and repeat X-ray after 12 months

showed resolution.

The commonest effect of maternal syphilis on the fetus is probably abortion and perinatal deaths may be to the tune of around 40% if left untreated [1]. Only 22% babies become symptomatic and manifestations appear mostly within the first few weeks of life [1]. Splenomegaly and pseudoparalysis, which form part of diagnostic pentad, were present in these two cases [2]. Radiological evidence of skeletal involvement has been reported in 80% cases [3]. The disappearance of these radiological features takes nearly six to twelve months; hence such babies should be followed up for six to twelve months. A high index of suspicion is emphasized since congenital syphilis may mimic any neonatal infection. WHO has reported high rates of syphilis among pregnant women from several countries in eastern Asia [4]. It is now believed that congenital syphilis can occur at any time in gestation, with the risk of fetal infection increasing as the stage of pregnancy advances. Therefore, STS being non-reactor in first trimester does not rule out syphilis and high-risk partners should undergo STS in the last trimester and before delivery. Follow up of the infant should be incorporated in routine paediatric care.

References

1. Kolivras A, De Maubeuge J, Song M, Hansen V, Toppet V, Van Herreweghe I. A case of early congenital syphilis. *Dermatol* 2002;204(4):338-40.
2. Karthikeyan K, Thappa DM. Early congenital syphilis in new millennium. *Paediatr Dermatol* 2002;19(3):275-6.
3. Radolff JD, Sanchez PJ, Schulz KF, Murphy KF. Congenital syphilis. In: Holmes King K, Sparling PF, Mardh Per-andres, Lemon SM, Stamm WE, Piot Peter, Wasserheit JN, editors. Sexually transmitted diseases. 3rded. Mc Graw Hill (USA, New York) 1999;1165-89.
4. WHO. Global prevalence and incidence of selected curable sexually transmitted infections: overview and estimates. WHO/HIV-AIDS/2001.02 and WHO/CDS/CSR/EDC/2001.10. Geneva:WHO, 2001.

Col PL Prasad*, Surg Lt Cdr AK Yadav*, Col MPS Sawhney#

*Senior Advisor (Paediatrics), Command Hospital (Central Command), Lucknow, *Graded Specialist (Paediatrics), INHS Dhanvantri, Naval Hospital, Port Blair, #Senior Adviser (Derm & Ven), Base Hospital, Barrackpore.

PATIENT SAFETY & HEALTH CARE

Dear Editor,

Reference is made to two letters to editor pertaining to patient safety and health care [1, 2]. Following is offered as comments on the subject.

Every doctor during his life time of practising medicine has to focus on a single theme. Shall we redefine this theme as "better attitude and care towards patients"? If accepted this theme cannot be achieved on fixed and rigid ideas.

Rapidly changing medical science has direct bearing on the chosen theme and will keep it fluid requiring the "provider" to be always in "learning mode". The present system of learning and updating by medical officers in Armed Forces is dominated by tradition and institution, which is not always justified. Unlike a bucket, the

human mind does not fill up in a passive manner. Analysing further the present system at grass root level of health care is more "retrospective" rather than "prospective" based learning. Judicious use of evidence based practice of medicine is almost non-existent except in some tertiary care hospitals.

In the quest to remain focused on the chosen theme, alternate learning strategies in the form of group based and problem based learning techniques need to be adopted at all levels of Armed Forces Medical Services. Group based learning fosters critical thinking, improves and promotes clinical judgement. Participants learn from each other and their interaction adds to the richness and depth of learning experience. This learning strategy inculcates a sense of cooperation and has the potential to change the present culture of "name it, blame it, shame it". Problem based learning is closely

linked to evidence based practice.

As active members of clinician group, we have attempted above alternate learning strategies in two hospitals i.e Command Hospital (Air Force) and No 5 Air Force Hospital. The initial fear, hesitation and sluggishness have given way to enthusiasm and active participation. We have reaped rich dividends as applicable to day to day clinical practice. Incorporating clinical audit meeting into these learning strategies has brought about more cohesiveness among peer groups, better referral system and above all true & effective team management. Scientific usage of antibiotics and thereby reducing nosocomial infection and precious funds has been the other benefits.

These alternate learning strategies as applied to health care could easily be adopted at every level to bring about the desired level of

patient safety.

“We think too much about effective methods of teaching and not enough about effective methods of learning.” - John Carolous SJ.

References

1. Taneja N. Building safer healthcare systems: A case for Error-in-Medicine Curriculum in medical training. MJAFI, 2003, 59; 273.
2. Taneja N. Patient safety: Can the Armed Forces show the way? MJAFI, 2003, 59; 274.

Air Cmde TS Raghu Raman

Air Officer Commanding, No 5 Air Force Hospital, C/o 99 APO

REPLY

The author agrees that the ‘problem based learning strategy’ as proposed inculcates a sense of cooperation and is closely linked to evidence based practice. This will definitely lead to a higher standard of patient care, as far as clinical decision making is concerned. Such an approach in the form of regular clinical meetings is already in vogue in the Armed Forces Medical Services. Whether such a strategy has the potential to change the culture of ‘name it, blame

it, shame it’ is doubtful. Much more attitude changes are required at the highest levels if a change in organizational culture is to be brought about in healthcare.

Wg Cdr N Taneja

Classified Specialist (Aviation Medicine), Institute of Aerospace Medicine, Bangalore.

ANSWERS TO MCQS

1-a	2-c	3-b	4-d	5-d
6-d	7-a	8-a	9-b	10-c
11-b	12-d	13-c	14-d	15-c
16-a	17-b	18-a	19-d	20-a
21-d	22-c	23-b	24-c	25-a