SELF PROCUREMENT OF DRUGS

Dear Editor,

I wish to highlight through this letter the problem of self procurement of most drugs including dangerous and restricted drugs in our country.

79 year old ex-serviceman, an old case of generalized seizures was admitted in a state of coma, on 08 May 2000. Coma was preceded by 3-4 generalized seizures. He had flaccid paralysis of all the 4 limbs without any localizing or lateralizing signs. Blood sugar level was 30mg%. History given by relatives revealed that the patient had not taken his regular anti epileptic medication dilantin (diphenyl hydantoin) for one week. On 07 May 2000 evening he procured some medication from local market in his town and consumed 3 tablets of the same. His hypoglycaemia failed to respond to aggressive management with 100 ml of 50% dextrose followed by IV drip of 10% dextrose and he expired 17 hours after admis-

sion.

There is a strong likelihood that the patient consumed 3 tablets of daonil (glybenclamide) in place of dilantin by mistake which led to prolonged hypoglycaemia and fatal outcome, a known complication of daonil in elderly. This suspicion is corroborated though not confirmed conclusively by relatives, as no strip of incriminated tablets could be produced by them.

It is emphasized therefore, that the fatal outcome in this case could have been avoided if the chemist had insisted on production of written prescription by the patient instead of making available the drug on verbal request for purchase

Lt Col SHIV KUMAR

Classified Specialist (Medicine), Military Hospital, Gwalior.

DAILY COST OF MEDICINES ISSUED TO PATIENTS IN A MEDICAL INSTITUTION

Dear Editor,

At the highest level in the Armed Forces Medical Services, data would exist in terms of the total expenditure involved for the treatment provided to serving personnel and their families or the per capita planned expenditure on a serving personnel. However, such data cannot be translated into calculating the average cost of medicines issued daily to patients in a medical care institution. The author could not find such a reported study and it was thus planned to calculate the average daily cost of medicines issued at our Station Sick Quarters (SSQ).

This SSQ is a 15 bedded non dieted SSQ with 4 Medical Officers and an average daily sick report of 100 patients. Patients on long term treatment viz. Hypertension/Diabetes etc collect their medicines on two days of the week i.e. Tuesdays and Fridays. The cost was calculated based on the average price of a particular product as mentioned in Drug Today [1]. Data was collected for a full working week to obviate any bias of sick report on a particular day of the week. Cost was calculated only for medicines received from AFMSDs and issued from the Dispensary and thus does not include medicines issued to ward patients, IV fluids, bandages etc which are not issued from the Dispensary on a day to day basis.

The average daily sick report for the period analysed was 103, OR's families constituting the bulk of the sick report. The average daily cost of medicines for this period was Rs 4420/- day and there is no reason why not to presume that this would be an average figure for the SSQ. If this amount is analysed, Cap Amoxycillin as an

individual drug cost Rs.538.00 (Cap Amoxycillin average cost Rs 3/- for 250 mg capsule) and Rs 347.00 for Syp Amoxycillin (average cost Rs 16/- for 30 ml bottle). As a group antibiotics at Rs 1154/- for capsules/tablets and Rs 459/- for syrups headed the expenditure list.

The above analysis has given few important inputs. Average cost of medicines issued to each patient was about Rs 40/- per patient, excluding the cost of laboratory investigations/dressings etc. Antibiotics constituting 30% of the total cost of medicines issued, points to the frequency of use of antibiotics at primary health care level. The use of a particular antibiotic in preference to other, may be equally effective, yet cheaper antibiotics is also another aspect which could be looked into. It also gives us a knowledge of the expenditure involved in the SSQ. A monthly medicine consumption of Rs 1.25 lakhs is no small amount in a medium sized SSQ like this. Such an analysis provides us the basic data to carry out a specific audit within a medium sized primary health institution in terms of sick report pattern, pattern of consumption of medicines and prescribing habits of medical officers.

Reference

Drug Today. Lorina Publication (India) Inc. Delhi Jan-Mar 2000.

Wg Cdr NARINDER TANEJA^{*}, Sqn Ldr SK SINHA[†]

^{*}Classified Specialist (Aviation Medicine), 15 Squadron Air Force, C/o 56 APO, [†]Medical Officer, 2254 Sqn Air Force, C/o 56 APO.

PROBLEM OF NON DEFLATABLE FOLEY'S CATHETER BALLOON

Dear Editor,

Of late with the use of Indian made Foley's catheters non deflation of Foley's catheter bulb has become a common problem. Standard text books describe various methods like (i) cut the valve of the balloon (ii) injecting liquid paraffin down the main drainage channel (iii) passing urethral catheter stylet down the balloon channel (iv) puncture of balloon with 19g spinal needle

passed suprapubically, transvaginally or transrectally.

However, in female patients, I have used a simpler and safer technique in over 20 cases in the last few years. The method is as follows:

1. Clean vulva with Savlon & saline
2. Place the patient in lithotomy position with thighs spread out.
3. With the left hand pull the Foleys catheter and keep it pulled

with gentle traction.

4. With the gloved (right) hand, thin disposable syringe needle is passed along side the Foley's catheter per urethrally. Invariably the needle punctures the balloon and the fluid from balloon is seen coming out through the needle. The Foley's catheter comes out on

its own with traction.

Lt Col PV RAMA MOHAN

Classified Specialist (Surgery), Military Hospital, Secunderabad

NECROTIZING FASCITIS

Dear Editor,

This refers to the letter to editor 'Necrotizing Fascitis' published in Medical Journal Armed Forces India [1]. We would like to share our experience in successfully managing a case of Fournier's gangrene in our hospital.

A 67 year old ex-serviceman, neglected by his grown up children was brought to this hospital with history of weakness of right side of his body-10 days and swelling of penoscrotal region with foul smelling discharge - 3 days. He was unable to walk and passed stools and urine in bed. Examination revealed a frail old man who had foul smell emanating from him. He had tachycardia with normal blood pressure. Temperature on admission was 99.4°F. Pallor was present. He had right sided hemiparesis. Local examination revealed an edematous scrotum with edema extending on to the penis. There were multiple superficial ulcers. Urine and faeces contaminated the scrotum and perineal region. Skin over the 2/3 of the scrotum was gangrenous and had bullous vesicles. Crepitus was present over the scrotum and penis. He was managed as a case of Fournier's gangrene.

Investigations revealed haemoglobin of 8.6 gm% with mild leucocytosis. Blood sugar fasting and post prandial were 232 mg% and 330 mg% respectively. Blood cultures and pus cultures were sterile. Screening test for HIV was negative.

He was managed with IV fluids, broad-spectrum antibiotics (3rd generation cephalosporins, amino glycosides and metronidazole). Repeated extensive surgical debridement was undertaken which left the testes uncovered. Indwelling Foley's catheter was used for

urinary diversion. Injection plain insulin (8 units) after each meal was used to control blood sugar. With these measures, the patient gradually improved. The infection subsided and penoscrotal wound became healthy. Secondary suturing was undertaken after about four weeks. He was discharged from hospital after 6 weeks and has remained well thereafter.

Fournier's gangrene is a rapidly progressive genital infection that is urological emergency. It was originally described in five healthy men by the French venereologist AJ Fournier in late 1880's. It is an uncommon disease with mortality as high as 60% in some series [2]. Extensive surgical debridement, broad-spectrum antibiotics, and control of diabetes mellitus with nutritional support help in a favourable outcome. The elderly diabetic who is socially neglected, nutritionally depleted and has an immunocompromised status is most vulnerable to this life threatening disease. Test for HIV should be done in all cases, as Fournier's gangrene may be a presenting feature of an undiagnosed HIV infection [3].

References

1. Lt Col Manmohan Harjai. Necrotizing Fascitis. MJAFI 2000;56:273-4.
2. Resnick MI, Benson MA. editors. Manual of clinical problems in urology. 1989;262-4.
3. Hotter JT. Fournier's gangrene as the presenting sign of an undiagnosed human immunodeficiency virus infection. J Urol 1996;155:291-2.

Maj SK MAURYA*, Lt Col FB BHOT*

*Graded Specialist(Surgery), *Classified Specialist (Anaesthesiology), Military Hospital Yol, Himachal Pradesh - 176 052.

Reply

Dear Editor,

At the outset I congratulate the surgical team of MH Yol for managing such a dreaded disease at periphery with excellent result. As I have already mentioned in my published letter necrotizing fascitis is caused by mixed aerobic-anaerobic bacteria and begins with a breach in the integrity of a mucous membrane barrier, such as the mucosa of the gastrointestinal or genitourinary tract. The predisposing factors include peripheral vascular disease, diabetes mellitus, surgery, and penetrating injury to the abdomen. In the above-mentioned patient, diabetes mellitus and nutritional deprivation were the predisposing factors. It is a rapidly spreading destructive disease of the fascia, usually attributed to group A *Streptococcus pyogenes* infection, but can also be caused by anaerobic bacteria, including *Peptostreptococcus* and *Bacteroides* species or may occur as part of gas gangrene caused by *Clostridium perfringens*. This life-threatening condition is diverse in presentation and severity and offers a great challenge to the treating clinician. Early diagnosis may be difficult when pain or unexplained fever is the only presenting

manifestation while patients in the later stages of the disease usually undergo septic shock and multiorgan failure. Early and aggressive surgical exploration is essential to save the life of the patient. The patient mentioned in the text was very well managed but if faecal or urinary contamination hinders in the healing process and recovery of the patient, then diversion of faecal and urinary streams may be necessary to prevent contamination. The importance of suspecting an undiagnosed human immunodeficiency virus infection in these patients is already highlighted in my published letter [1].

References

- Hotter JT. Fournier's gangrene as the presenting sign of an undiagnosed human immunodeficiency virus infection. J Urol 1996;155:291-2.

Lt Col Man Mohan Harjai

Reader, Paediatric Surgeon, Department of Surgery, Armed Forces Medical College, Pune 411 040.
e.mail : harjai101@hotmail.com