with gentle traction.

4. With the gloved (right) hand, thin disposable syringe needle is passed along side the Foley's catheter per urethrally. Invariably the needle punctures the balloon and the fluid from balloon is seen coming out through the needle. The Foley's catheter comes out on

its own with traction.

Lt Col PV RAMA MOHAN

Classified Specialist (Surgery), Military Hospital, Secunderabad

NECROTIZING FASCITIS

Dear Editor,

This refers to the letter to editor 'Necrotizing Fascitis' published in Medical Journal Armed Forces India [1]. We would like to share our experience in successfully managing a case of Fournier's gangrene in our hospital.

A 67 year old ex-serviceman, neglected by his grown up children was brought to this hospital with history of weakness of right side of his body-10 days and swelling of penoscrotal region with foul smelling discharge - 3 days. He was unable to walk and passed stools and urine in bed. Examination revealed a frail old man who had foul smell emanating from him. He had tachycardia with normal blood pressure. Temperature on admission was 99.4°F. Pallor was present. He had right sided hemiparesis. Local examination revealed an edematous scrotum with edema extending on to the penis. There were multiple superficial ulcers. Urine and faeces contaminated the scrotum and perineal region. Skin over the 2/3 of the scrotum was gangrenous and had bullous vesicles. Crepitus was present over the scrotum and penis. He was managed as a case of Fournier's gangrene.

Investigations revealed haemoglobin of 8.6 gm% with mild leucocytosis. Blood sugar fasting and post prandial were 232 mg% and 330 mg% respectively. Blood cultures and pus cultures were sterile. Screening test for HIV was negative.

He was managed with IV fluids, broad-spectrum antibiotics (3rd generation cephalosporins, amino glycosides and metronidazole). Repeated extensive surgical debridement was undertaken which left the testes uncovered. Indwelling Foley's catheter was used for

urinary diversion. Injection plain insulin (8 units) after each meal was used to control blood sugar. With these measures, the patient gradually improved. The infection subsided and penoscrotal wound became healthy. Secondary suturing was undertaken after about four weeks. He was discharged from hospital after 6 weeks and has remained well thereafter.

Fournier's gangrene is a rapidly progressive genital infection that is urological emergency. It was originally described in five healthy men by the French venereologist AJ Fournier in late 1880's. It is an uncommon disease with mortality as high as 60% in some series [2]. Extensive surgical debridement, broad-spectrum antibiotics, and control of diabetes mellitus with nutritional support help in a favourable outcome. The elderly diabetic who is socially neglected, nutritionally depleted and has an immunocompromised status is most vulnerable to this life threatening disease. Test for HIV should be done in all cases, as Fournier's gangrene may be a presenting feature of an undiagnosed HIV infection [3].

References

1. Lt Col Manmohan Harjai. Necrotizing Fascitis. MJAFI 2000;56:273-4.
2. Resnick MI, Benson MA. editors. Manual of clinical problems in urology. 1989;262-4.
3. Hotter JT. Fournier's gangrene as the presenting sign of an undiagnosed human immunodeficiency virus infection. J Urol 1996;155:291-2.

Maj SK MAURYA*, Lt Col FB BHOT*

*Graded Specialist(Surgery), *Classified Specialist (Anaesthesiology), Military Hospital Yol, Himachal Pradesh - 176 052.

Reply

Dear Editor,

At the outset I congratulate the surgical team of MH Yol for managing such a dreaded disease at periphery with excellent result. As I have already mentioned in my published letter necrotizing fascitis is caused by mixed aerobic-anaerobic bacteria and begins with a breach in the integrity of a mucous membrane barrier, such as the mucosa of the gastrointestinal or genitourinary tract. The predisposing factors include peripheral vascular disease, diabetes mellitus, surgery, and penetrating injury to the abdomen. In the above-mentioned patient, diabetes mellitus and nutritional deprivation were the predisposing factors. It is a rapidly spreading destructive disease of the fascia, usually attributed to group A *Streptococcus pyogenes* infection, but can also be caused by anaerobic bacteria, including *Peptostreptococcus* and *Bacteroides* species or may occur as part of gas gangrene caused by *Clostridium perfringens*. This life-threatening condition is diverse in presentation and severity and offers a great challenge to the treating clinician. Early diagnosis may be difficult when pain or unexplained fever is the only presenting

manifestation while patients in the later stages of the disease usually undergo septic shock and multiorgan failure. Early and aggressive surgical exploration is essential to save the life of the patient. The patient mentioned in the text was very well managed but if faecal or urinary contamination hinders in the healing process and recovery of the patient, then diversion of faecal and urinary streams may be necessary to prevent contamination. The importance of suspecting an undiagnosed human immunodeficiency virus infection in these patients is already highlighted in my published letter [1].

References

- Hotter JT. Fournier's gangrene as the presenting sign of an undiagnosed human immunodeficiency virus infection. J Urol 1996;155:291-2.

Lt Col Man Mohan Harjai

Reader, Paediatric Surgeon, Department of Surgery, Armed Forces Medical College, Pune 411 040.
e.mail : harjai101@hotmail.com