Letters to the Editor

CONGENITAL TORTICOLLIS

Dear Editor,

This is with reference to the case report Congenital Torticollis 'MJAFI 2000; 56: 55-56.

Congenital torticollis persisting into adolescence and adulthood usually reveals fibrosis of not only the sternocleidomastoid muscle, but also the layers of the deep fascia including the carotid sheath [1]. It can be associated with ipsilateral facial hemihypoplasia, plagiocephaly, head turned away from the side of the mass and occasionally ipsilateral trapezius atrophy or cervical vertebral abnormality which should be ruled out by an X-ray of the cervical spine [2]. Good results can only be obtained if, along with the sternocleidomastoid muscle, the layers of the deep fascia and carotid sheath can be carefully divided followed by intensive physiotherapy and a

plaster cast over a long period of time.

I had the experience of assisting in the surgery of a young probable rejected in recruitment because of congenital torticollis. The lower end of sternocleidomastoid was divided. The results were far from satisfactory.

References

1. Congenital Torticollis. Farquharsons' Textbook of Operative Surgery, 8th ed, pg 291.
2. Cervical Anomalies, Sabiston Textbook of Surgery, 19th ed Vol-2, pg 1263.

Maj KJ SINGH

Graded Specialist (Surgery), 153 General Hospital, C/O 56 APO

Reply

Dear Editor,

I thank Major KJ Singh for reading the article and expressing his views on the management of congenital torticollis persisting into adolescence. Congenital torticollis is a condition due to developmental aplasia of the muscles and the other soft tissue structures on the affected side of the neck may be involved with the passage of time. [1] I agree with the reader that it may be associated with ipsilateral facial hemihypoplasia, plagiocephaly, head turned away from the side of the mass and occasional ipsilateral trapezius atrophy or cervical vertebral abnormality [2] which were ruled out by x-ray of the cervical spine. Cervical spine subluxations associated with childhood torticollis are usually rotatory, occur at the C1-C2 level, and are best known as atlanto-axial rotatory subluxations. [3] If the subluxation persists, it may be described as an atlanto-axial rotatory fixation [4].

In this particular patient, on exploration there was no shortening of scalenus anterior or contracture of the soft tissue e.g. carotid sheath or deep fascia, hence only the heads of sternocleidomastoid were divided. Results of surgery were satisfactory in this patient and there was no recurrence of the deformity on follow up. This was

possible due to intensive physiotherapy and wearing of torticollis harness. It is important to safeguard the internal jugular vein, spinal accessory nerve and phrenic nerve. Open surgery is preferred as subcutaneous tenotomy is a blind technique and should be avoided in children.

References

1. Saunders RA, Roberts EL. Abnormal head posture in patients with fourth cranial nerve palsy Am Orthop J 1995;45:24-33.
2. R. Kendrick State, Jefery C, Posnick, Derek C, Armstrong. Cervical spine subluxation associated with congenital muscular torticollis and craniofacial asymmetry. Plastic and Reconstructive Surgery 1993;91:1187-95.
3. Bredenkamp JK, Maceri DR. Inflammatory torticollis in children. Arch. Otolaryngol. Head Neck Surgery 1990;116:310-4.
4. Fielding JW, Hawkins RJ. Atlanto-axial rotatory fixation. J Bone Joint Surg (Am) 1977;59-A:37-44.

Lt Col S CHAWLA

Classified Specialist (Surgery), Military Hospital, Bhopal-462031.

INCIDENCE AND ETIOLOGY OF RESPIRATORY DISTRESS IN NEWBORN

Dear Editor,

This is in reference to the article titled 'Incidence and Etiology of Respiratory distress in newborn'[1].

Following comments are offered for review by authors.

1. The clinical diagnosis of respiratory distress in a neonate is usually assessed by the Downe's score which is as under :-

Parameter	0	1	2
Cyanosis	None	In room air	In 40% FIO ₂
Retractions	None	Mild	Severe
Grunting	None	Audible with stethoscope	Audible without stethoscope
Air entry	Clear	Decreased/delayed	Barely audible
Respiratory rate	< 60	60-80	> 80 or apnea

Score : > 4 = Clinical respiratory distress; monitor arterial blood gases

8 = Impending respiratory failure

This scoring criterion is usually adopted for making initial diagnosis and also for management protocols.

2. There are several reports on the incidence of respiratory distress syndrome in Indian literature. In fact in a similar study from the same center [2], the authors have observed similar pattern of etiological factors (respiratory distress was hyaline membrane disease (18%), followed by wet lung syndrome (14%), meconium aspiration (12%), asphyxia (12%) and septicemia (8%). In 8 babies, a lung biopsy (postmortem) was done to confirm the diagnosis. In a study on neonatal autopsy, the authors correlated clinical and autopsy findings. There were 23 neonatal autopsy studies out of 43 neonatal deaths during the period from Jan 1991 to Sept 93. Common antemortem diagnosis included meconium aspiration syndrome, respiratory distress syndrome and aspiration syndrome.