Published in final edited form as:

J Pharmacol Exp Ther. 2008 April ; 325(1): 37–46. doi:10.1124/jpet.107.128215.

Design of Mutant β_2 Subunits as Decoy Molecules to Reduce the Expression of Functional Ca^{2+} Channels in Cardiac Cells

Sabine Télémaque, Swapnil Sonkusare, Terrie Grain, Sung W. Rhee, Joseph R. Stimers, Nancy J. Rusch, and James D. Marsh

Departments of Internal Medicine (S.T., T.G., J.D.M.) and Pharmacology and Toxicology (S.S., S.W.R., J.R.S., N.J.R.), College of Medicine, University of Arkansas for Medical Sciences, Little Rock, Arkansas

Abstract

Calcium influx through long-lasting (“L-type”) Ca^{2+} channels (Ca_V) drives excitation-contraction in the normal heart. Dysregulation of this process contributes to Ca^{2+} overload, and interventions that reduce expression of the pore-forming α_1 subunit may alleviate cytosolic Ca^{2+} excess. As a molecular approach to disrupt the assembly of $\text{Ca}_V1.2$ (α_{1C}) channels at the cell membrane, we targeted the Ca^{2+} channel β_2 subunit, an intracellular chaperone that interacts with α_{1C} via its β interaction domain (BID) to promote $\text{Ca}_V1.2$ channel expression. Recombinant adenovirus expressing either the full β_2 subunit (Full- β_2) or truncated β_2 subunit constructs lacking either the C terminus, N terminus, or both (N-BID, C-BID, and BID, respectively) fused to green fluorescent protein were developed as potential decoys and overexpressed in HL-1 cells. Fluorescence microscopy revealed that the localization of Full- β_2 at the surface membrane was associated with increased Ca^{2+} current mainly attributed to $\text{Ca}_V1.2$ channels. In contrast, truncated N-BID and C-BID constructs showed punctate intracellular expression, and BID showed a diffuse cytosolic distribution. Total expression of the α_{1C} protein of $\text{Ca}_V1.2$ channels was similar between groups, but HL-1 cells overexpressing C-BID and BID exhibited reduced Ca^{2+} current. C-BID and BID also attenuated Ca^{2+} current associated with another L-type Ca^{2+} channel, $\text{Ca}_V1.3$, but they did not reduce transient Ca^{2+} currents attributed to Ca_V3 channels. These results suggest that β_2 subunit mutants lacking the N terminus may preferentially disrupt the proper localization of L-type Ca^{2+} channels in the cell membrane. Cardiac-specific delivery of these decoy molecules in vivo may represent a gene-based treatment for pathologies involving Ca^{2+} overload.

The predominant voltage-gated Ca^{2+} channels (Ca_V) in cardiac cells are the dihydropyridine-sensitive $\text{Ca}_V1.2$ (α_{1C}) channels that mediate long-lasting (“L-type”) Ca^{2+} current as a critical step in excitation-contraction coupling (Bers, 2002). The $\text{Ca}_V1.2$

Address correspondence to: Dr. Sabine Télémaque, Division of Cardiovascular Medicine, Department of Internal Medicine, College of Medicine, University of Arkansas for Medical Sciences, 4301 W. Markham, #832, Little Rock, AR 72205. stelemaque@uams.edu. S.T. and S.S. contributed equally to this work.

A preliminary version of some of these results has been presented in abstract form at the following conference: Sonkusare S, Stimers JR, Grain T, Marsh JD, and Télémaque S (2006) Expression of β_2 subunit mutants alters calcium currents in HL-1 cells (*FASEB J* 20:A1114); *Experimental Biology Meeting 2006*; 2006 Apr 1–5; San Francisco, CA. American Society for Pharmacology and Experimental Therapeutics, Bethesda, MD.

The online version of this article (available at <http://jpet.aspetjournals.org>) contains supplemental material.

channels are multiprotein complexes composed of a large, pore-forming α_{1C} subunit and accessory β and $\alpha_2\text{-}\delta$ subunits. The expression level of functional $\text{Ca}_v1.2$ channels is regulated by the β subunits (β_1 , β_2 , β_3 , and β_4), which are cytoplasmic proteins encoded by four different genes, including multiple splice variants (Foell et al., 2004). The β_2 subunit is predominantly expressed in rat ventricular myocytes, although other gene families have been reported. It is noteworthy that β_2 acts as a molecular chaperone that escorts α_{1C} to the sarcolemma to localize functional $\text{Ca}_v1.2$ channels at the surface membrane (Pragnell et al., 1994; De Waard et al., 1996; Opatowsky et al., 2003). Indeed, we recently reported that overexpression of β_2 in isolated feline ventricular myocytes significantly increases Ca^{2+} influx and cardiac contractility (Chen et al., 2005). Although mutations in the β interaction domain (BID) of β_2 were found to abolish α_{1C} - β interaction (De Waard et al., 1994, 1996), two additional protein-interaction domains (SH3 and GK) were recently identified that also may interact with α_{1C} subunit (Chen et al., 2004; Opatowsky et al., 2004; Van Petegem et al., 2004). Thus, it has been suggested that BID may not bind directly to α_{1C} , but instead it is fundamentally essential for the structural integrity and bridging of the SH3 and GK domains.

Although physiologically essential to normal cardiac function, $\text{Ca}_v1.2$ channels also have been implicated in pathological processes involving Ca^{2+} overload, including cardiac ischemia and myocardial stunning. In these conditions, an elevated cytosolic Ca^{2+} ($[\text{Ca}^{2+}]_{\text{cyt}}$) in the myocardial cells is associated with a loss of myofilament sensitivity to Ca^{2+} (Bolli and Marban, 1999). Moreover, Ca^{2+} overload also plays an important role in cardiac apoptosis and necrosis (Aon et al., 2003). Even under conditions in which $[\text{Ca}^{2+}]_{\text{cyt}}$ is within a normal range, it may be desirable to modestly depress $[\text{Ca}^{2+}]_{\text{cyt}}$ as a therapeutic intervention to slow atrioventricular conduction in atrial fibrillation or to modestly depress contractility in hypertrophic cardiomyopathy. Currently, calcium channel blockers represent the drug therapies of choice for these conditions, but several features of their properties, including their short duration of action and vascular dilator effects, limit their usefulness.

In this respect, a gene therapy approach designed to reduce the availability of β_2 subunits required for the expression of $\text{Ca}_v1.2$ channels may provide for the highly targeted and long-term relief of calcium overload. Accordingly, we engineered recombinant adenoviral constructs expressing truncated β_2 to provide proof of concept that mutated β_2 could potentially act as a dominant-negative construct to reduce $\text{Ca}_v1.2$ channels at the surface membrane. In the present study, we expressed the recombinant adenovirus constructs in HL-1 cells, a cell line derived from AT-1 mouse atrial tumor lineage that displays many normal markers of cardiac biology. HL-1 cells spontaneously depolarize and express ion channels required for generating action potentials (White et al., 2004), and they retain the signaling pathways to respond appropriately to inotropic and chronotropic agonists (Claycomb et al., 1998). The activity of the truncated β_2 constructs was compared with the effect of full-length β_2 (Full- β_2), which we have shown enhances the expression of functional $\text{Ca}_v1.2$ channels in feline cardiac myocytes (Chen et al., 2005). The goal was to retain the ability of the β_2 decoys to bind to α_{1C} but to eliminate the amino and/or carboxyl termini of β_2 that may be required for the expression of functional α_{1C} subunits in the plasma membrane, thereby “trapping” $\text{Ca}_v1.2$ channels inside the cell. We also evaluated the effect of the β_2 subunit mutants on Ca^{2+} current attributed to two other types of Ca^{2+}

channels important in pacemaker activity or contractile function. The $\text{Ca}_v1.3$ channels (α_{1D}) mediate L-type current at mid-voltages, and they are approximately 10-fold less sensitive to 1,4-dihydro-2,6-dimethyl-4-(2-nitrophenyl)-3,5-pyridinedicarboxylic acid dimethyl ester (nifedipine)-induced block than $\text{Ca}_v1.2$ channels at negative membrane potentials (Koschak et al., 2001). HL-1 cells also express Ca_v3 channels that activate at much lower voltages than L-type channels, and they mediate transient (“T-type”) current associated with normal pacemaker activity in cardiac cells (Xia et al., 2004; Zhang et al., 2005).

Materials and Methods

Culture of HL-1 Cells

HL-1 cells (kindly provided by Dr. W. C. Claycomb, Louisiana State University, Baton Rouge, LA) were grown and maintained in Claycomb medium (SAFC Biosciences, Lenexa, KS) supplemented with penicillin-streptomycin (1×) and 2 mM L-glutamine (Invitrogen, Carlsbad, CA). After trypsinization, dissociated cells were either plated in standard 60-mm tissue culture dishes ($0.8\text{--}1.0 \times 10^6$ cells/ml) for mRNA and/or protein expression assays or in glass culture dishes (MatTek, Ashland, MA) for electrophysiology ($60\text{--}80 \times 10^3$ cells/ml). Cells were then infected for 24 to 48 h with one of four different GFP-fused β_2 subunit constructs, including Full- β_2 , or truncated β_2 lacking either the N terminus (C-BID), the C terminus (N-BID), or both termini (BID only) or with the control GFP construct. The optimal dose of crude viral stock (multiplicity of infection, 50–100) needed to optimize protein expression without visible cell death was determined by fluorescent imaging and immunoblot techniques. Experimental conditions were developed that produced >95% infection of HL-1 cells for each of the constructs studied, as described previously (Fan et al., 2003; Telemaque-Potts et al., 2003; Chen et al., 2005). All studies on HL-1 cells were performed between 24 and 48 h after transfection.

Detection of Ca_L Channel Subunit Gene Expression in HL-1 Cells

To identify the endogenous β subunits expressed in HL-1 cells, total RNA was isolated from untreated HL-1 cells using a TRIzol reagent (Invitrogen). First-strand cDNA was synthesized using SuperScript III reverse transcriptase (Invitrogen). Primers for PCR reactions were designed to detect $\text{Ca}_v1.2$ (α_{1C}) and different β subunit isoforms (β_1 , β_{2a} , β_{2b} , β_3 , and β_4). Sense and antisense primers were as follows: α_{1C} : 5'-CCGCCCCTACCAAGATCAAC-3', 5'-TCGTGTCATTGACAATGCGG-3', covering cDNA bases 2536 to 2791 (256 bp; GenBank accession number NM_012517.1 [rat]); β_1 : 5'-TCCAGAAGAGCGGCATGTCCC-3', 5'-AGGACGTACTCCCGTCTGAC-3', covering cDNA bases 5 to 140 (136 bp; GenBank accession number NM_031173.2 [mouse]); β_{2a} : 5'-CGAGTACGGGTGTCCTATG, 5'-CGTCCTATCCACCAGTCAT-3', covering cDNA bases 31 to 320 (290 bp; GenBank accession number NM_053851.1 [rat]); β_{2b} : 5'-AGGCAGTTGGTGTCTTCTC-3', 5'-CGTCCTATCCACCAGTCAT-3', covering cDNA bases 10 to 323 (314 bp; GenBank accession number AF423193 [rat]); β_3 : 5'-CATCCCTGGACTTCAGAAC-3', 5'-TGGTAGGCATCTGCATAGTC-3', covering cDNA bases 1125 to 1301 (177 bp; GenBank accession number NM_007581.2 [mouse]); and β_4 : 5'-CTCACCATATCCCACAGCAA-3', 5'-TCCGGGTAATCTTCTTCCACCA-3', covering

cDNA bases 1266 to 1478 (213 bp; GenBank accession number NM_001037099.1 [mouse]).

Reactions contained 2 μ l of cDNA, 10 pmol of each primer, 0.3 mM dNTPs, 1.0 mM MgSO_4 , and 0.5 units of *Pfx* DNA polymerase (Invitrogen) mixed with 1 $\times$ enhancer buffer in a total volume of 25 μ l. Amplification steps included initial denaturation at 94°C for 3 min, followed by 30 cycles (denaturation for 30 s at 94°C; annealing 30 s at 53°C; extension 30 s at 68°C). A final step for product extension was performed at 68°C for 5 min. The PCR products (5 μ l) were separated using a 1.2% agarose gel stained with ethidium bromide for visualization. All oligonucleotide sequences were directed toward the mouse/rat sequences of the Ca^{2+} channel subunits.

Generation of β_2 Subunit Constructs

A plasmid encoding the rat heart/brain β_{2a} subunit cDNA, a generous gift from Dr. E. Perez-Reyes (Perez-Reyes et al., 1992), was used as a template for generating three PCR-based mutations of the β_{2a} subunit. In brief, cDNA encoding the N-BID fragment (bp 376–1177), BID fragment (bp 867–1177), and C-BID fragment (bp 867–2427) were amplified, digested, and directionally cloned into pEGFP-C1 vector (Clontech, Mountain View, CA), an expression vector driven by a cytomegalovirus (CMV) promoter. All β_2 subunit constructs were cloned in frame with an enhanced GFP reporter gene located at the amino-terminal end of the construct. A schematic representation of the various β_2 subunit adenoviral constructs is illustrated in Fig. 1.

Preparation of Recombinant Adenovirus Construct

Recombinant adenovirus constructs were prepared based on the methods described by He et al. (1998) and made commercially available (AdEASY) by Stratagene (La Jolla, CA). The CMV-GFP- β subunit fragments were gel-purified and subcloned into the pShuttle vector. Then, it was linearized and cotransformed into *Escherichia coli* BJ5183 cells with pAdEasy-1, the adenoviral backbone. A CMV-GFP lacking any β subunit construct also was subcloned into a pShuttle vector to construct a control virus (Fan et al., 2003). Recombinants were then linearized and transfected into human embryonic kidney (HEK) 293 cells (American Type Culture Collection, Manassas, VA) using FuGENE 6 reagent (Roche Molecular Systems, Alameda, CA). Virus-containing medium from HEK 293-infected cells was collected 3 to 6 days after infection, and proper expression of GFP-fused β subunits was confirmed by PCR and immunodetection. High-titer viral stock was generated by several rounds of amplification in HEK 293 cells and by purification by cesium chloride gradient ultracentrifugation (Niranjan et al., 1996; Telemaque-Potts et al., 2003). Final viral stock titer was determined by plaque assay ($1\text{--}3 \times 10^8$ plaque-forming unit/ μ l).

Immunodetection

Forty-eight hours after infection, cells were washed twice with ice-cold phosphate-buffered saline (PBS), and then they were lysed in Triton lysis buffer containing 50 mM HEPES, 150 mM NaCl, 1 mM EGTA, 1.5 mM MgCl_2 , 1% Triton X-100, 1% glycerol, 1 $\times$ Sigma phosphatase inhibitor cocktail, and 1 $\times$ Sigma protease inhibitor cocktail (Sigma-Aldrich, St. Louis, MO). Cell lysates were incubated on ice for 15 to 20 min and centrifuged at 4°C for

15 min at 14,000*g*. The supernatants were then collected. Protein concentration in supernatants was measured by the Bradford method (Bio-Rad, Hercules, CA). Equal protein samples (40 µg of total protein) were loaded on 3 to 8% NuPAGE Tris acetate gels (Invitrogen), and they were run and transferred to a polyvinylidene fluoride membrane. After blocking the membrane (Tris-buffered saline solution containing 10% nonfat dry milk and 0.1% Tween 20, immunoblotting was performed using mouse α_{1C} monoclonal antibody (1:200; NeuroMabs, Davis, CA), mouse β -actin monoclonal antibody (1:5000; Sigma-Aldrich), and mouse GFP monoclonal antibody (1:2000; Invitrogen). Corresponding secondary horseradish peroxidase-conjugated antibodies were used (1:10,000–1:20,000 dilutions; Jackson ImmunoResearch Laboratories Inc., West Grove, PA). Additional Western blots were probed with polyclonal antibodies using methods described previously (Pesic et al., 2004). These antibodies were obtained commercially (Alomone Labs, Jerusalem, Israel) and directed against Cav1.2 (α_{1C}), Cav1.3 (α_{1D}), Cav3.1 (α_{1G}), and Cav3.2 (α_{1H}). Protein bands were visualized by enhanced chemiluminescence, and the blots were exposed to Hyperfilm (GE Healthcare, Chicago, IL). Band densities were quantified by densitometry (Gel-Doc System; Bio-Rad), analyzed with Bio-Rad image analysis software, and they were normalized to β -actin expression.

Fluorescent Localization of Recombinant Adenovirus Proteins

Infected cells grown on glass-bottomed tissue culture dishes for 24 h after infection were washed, fixed in 4% paraformaldehyde, and kept in the dark at 4°C until fluorescent imaging. The images were acquired using a spinning disk confocal CARV II unit (BD Biosciences, San Jose, CA) on a Zeiss 200M microscope (Carl Zeiss, Thornwood, NY) fitted with a 100 $\times$ /1.4 numerical aperture objective and a Retiga EXi camera (QImaging, Surrey, BC, Canada).

Whole-Cell Patch Clamp

Functional expression of Ca²⁺ channels was evaluated using the whole-cell configuration of the patch-clamp technique (Hamill et al., 1981). After 24 h of infection with the different β_2 constructs, culture dishes were mounted on the stage of an inverted microscope (Nikon, Tokyo, Japan), and HL-1 cells expressing GFP were identified by epifluorescence microscopy and selected for electrophysiological recordings. Cells were initially bathed in a solution containing 145 mM NaCl, 5.4 mM KCl, 1 mM CaCl₂, 1 mM MgCl₂, 10 mM HEPES, and 5.5 mM dextrose. After formation of a conventional gigaohm seal, the patched membrane was ruptured to obtain the whole-cell configuration. Cells were then bathed in a solution that consisted of 20 mM BaCl₂, 115 mM *N*-methyl-D-glucamine, 10 mM HEPES, 1 mM MgCl₂, and 5.5 mM dextrose (pH 7.4; 300 mOsm). The pipette solution consisted of 120 mM cesium aspartate, 30 mM cesium chloride, 0.5 mM GTP, 5 mM MgATP, 5 mM creatine phosphate, and 5 mM sodium pyruvate (pH 7.2; 300 mOsm). Data acquisition and pulse generation were performed using pClamp software (Molecular Devices, Sunnyvale, CA) and an Axopatch 200A amplifier interfaced with a Digidata-1320 acquisition system (Molecular Devices). The data were sampled at 2 to 10 kHz, and they were filtered at 1 to 5 kHz. Capacitance and series resistance compensations were used, and leak current was linearly subtracted using a P/4 protocol, with a subtracting holding potential of –100 mV. Currents were normalized to cell size by dividing by cell capacitance (in pA/pF). Data

analysis was performed using Clampfit (Molecular Devices) and Origin (OriginLab Corp., Northampton, MA), as described previously (Stimers et al., 2003). Activation curves were fit by the Boltzmann equation: $I/I_{\max} = A_2 + (A_1 - A_2)/(1 + \exp((V - V_{1/2})/s))$, where V is the command voltage, $V_{1/2}$ is the potential of half-maximal activation, s is the slope factor, A_1 is the maximum value of the ratio $I/I_{\max}$ (fixed at 1), and A_2 is the minimum value of the ratio $I/I_{\max}$. Steady-state inactivation curves also were fit according to the Boltzmann function $I/I_{\max} = A_2 + (A_1 - A_2)/(1 + \exp((V - V_{1/2})/s))$, to obtain half-maximal inactivation values.

Chemicals

Unless otherwise stated, all standard chemical reagents were purchased from Sigma-Aldrich.

Statistical Analysis

Data are reported as mean $\pm$ S.E.M. Statistical comparisons were performed using one-way analysis of variance followed by post hoc Student-Newman-Keuls test. Values of P less than 0.05 were considered significant.

Results

Characterization of Ca²⁺ Channel Subunits in HL-1 Cells

Before introducing β subunit decoys, we determined the native profile of β subunits in HL-1 cells. Transcripts for the Ca_v1.2 (α_{1C}) subunit and for multiple β subunits, including β_1 , β_2 , β_3 , and β_4 , were detected (Fig. 2A). Subsequently, we confirmed the presence of nifedipine-sensitive, high-threshold Ca²⁺ channel current (I_{Ca}) attributed to Ca_v1.2 channels in these cells. Exposure of HL-1 cells to 10 μ M nifedipine ($n = 4$) resulted in a marked inhibition of I_{Ca} elicited by stepwise 10-mV depolarizing pulses between -40 to $+50$ mV (Fig. 2B). In these recordings, the I_{Ca} resistant to block by nifedipine was attributed to the presence of Ca²⁺ channel types other than Ca_v1.2 (Xia et al., 2003; Hansen et al., 2004; Zhang et al., 2005).

Adenovirus Infection of HL-1 Cells

Initial experiments assessed the potential for cytotoxicity of the adenoviral treatment. The multiplicity of infection was adjusted to achieve $>95\%$ cell transduction by the adenovirus (determined by GFP fluorescence) at 48 h after viral treatment, with minimal ($<10\%$) cytopathic effects visible as cell detachment, rounding, or blebbing. There was no evident difference in cytopathic effect between HL-1 cells treated with a control adenovirus (without construct) and the wild-type or mutant constructs. To confirm that adenoviral infection with GFP-fused constructs containing β_2 subunits resulted in expression of the appropriate protein, immunoblots were performed on cell lysates collected from HL-1 cells 48 h after adenoviral infection. The use of an antibody directed against GFP confirmed that all GFP-fused β_2 constructs were successfully expressed in the infected HL-1 cells (Fig. 3, lanes 2–6), whereas no immunoreactivity was detected in untreated HL-1 cells (Fig. 3, lane 1). Full- β_2 was detected as an ~ 106 -kDa protein (lane 2), whereas truncation of the amino terminus resulted in detection of C-BID at ~ 88 kDa (lane 3). Likewise, N-BID with a truncated carboxyl terminus was detected at ~ 66 kDa (lane 4), and the isolated BID was detected at ~ 42 kDa (lane 5). Control infection with GFP resulted in the expected ~ 30 -kDa protein (lane

6). Immunoblots were also performed to assess the expression level of Ca_v1.2 (α_{1C}) in untreated HL-1 cells compared with HL-1 cells infected with GFP alone or GFP-fused β_2 constructs (Fig. 3B). The averaged data from six independent blots indicated that the protein expression level of α_{1C} was similar between untreated cells (lane 1), cells infected with GFP-fused Full- β_2 , C-BID, N-BID, or BID constructs (lanes 2 to 5), or cells infected with GFP alone (lane 6), suggesting that overexpression of the β_2 constructs did not alter the total number of Ca_v1.2 channels in the HL-1 cells. Likewise, overexpression of the β_2 constructs did not change the total number of Ca_v1.3 channels in the HL-1 cells (data not shown).

Cellular Localization of Various β_2 Subunit Adenoviral Constructs

Cells were examined by fluorescence microscopy 24 h after treatment with adenoviral vectors. Preliminary experiments demonstrated that for adenoviral infection with either the Full- β_2 construct or various mutated constructs, gene expression determined by GFP fluorescence was already present 24 h after viral infection. Maximal *in vivo* fluorescence was observed by 48 h, which was confirmed by immunoblots. Although the duration of gene expression was not formally investigated in this study, strong fluorescence was maintained for at least 5 days under similar experimental conditions. We postulated that overexpression of Full- β_2 subunit would promote the normal trafficking of Ca_v1.2 channels to the plasma membrane. However, we anticipated that truncated β_2 subunits would retain the ability to bind to the α_{1C} subunit but display abnormal protein trafficking, and thereby not localize Ca_v1.2 channels at the surface membrane. Compared with untreated HL-1 cells (Fig. 4A), confocal images of HL-1 cells infected with GFP revealed a diffuse pattern of fluorescence in the cytosol (Fig. 4B), as reported previously in isolated cardiomyocytes (Telemaque-Potts et al., 2003; Chen et al., 2005). In contrast, overexpression of Full- β_2 produced sarcolemmal localization (Fig. 4C, arrows), whereas C-BID and N-BID fusion proteins seemed primarily localized to intracellular vesicles (Fig. 4, D and E, respectively). Overexpression of BID produced a diffuse reticular pattern suggestive of cytosolic localization (Fig. 4F). Thus, the truncated β_2 subunits showed punctate or diffuse intracellular localization that was distinctly different from the sarcolemmal localization of full-length β_2 . We also attempted to localize endogenous α_{1C} subunit by immunostaining, using two different commercially available antibodies against Ca_v1.2 channels, but we were unable to detect a specific signal at the cell membrane (Supplemental Figs. A and B).

Effect of Overexpression of β_2 Adenoviral Constructs on I_{Ca}

Subsequently, we compared the impact of overexpressing the β_2 adenoviral constructs (24 h after infection) on whole-cell I_{Ca} elicited by 10-mV depolarizing steps from a holding potential of -40 mV to potentials between -40 and +50 mV (Fig. 5A). Initial studies suggested that this pulse protocol primarily activated high-threshold, nifedipine-sensitive Ca_v1.2 channels (Fig. 2B). Compared with cells infected with GFP alone (Fig. 5A), overexpression of Full- β_2 markedly increased the membrane density of I_{Ca} (Fig. 5B) as reflected in the enhanced amplitude of the current-voltage (I-V) curve (Fig. 5F). Current densities were ~2.5-fold higher at the peak of the I-V curve in Full- β_2 -overexpressing cells compared with the cells overexpressing GFP alone (-17.9 ± 3.0 and -7.2 ± 0.6 pA/pF, respectively; $n = 10-12$). To test the hypothesis that truncated β_2 subunits serve as dominant-negative mutants, cells were infected with adenoviral constructs expressing N-BID, C-BID,

or BID. Recordings of whole-cell I_{Ca} in these cells are depicted in Fig. 5 (C–E, respectively), and average I–V curves are plotted in Fig. 5F ($n = 7$ –10). Cells infected with N-BID (Fig. 5C) showed I_{Ca} densities similar to GFP-infected cells (Fig. 5A), as verified by similar I–V curves. However, I_{Ca} was markedly reduced in C-BID- and BID-overexpressing cells (Fig. 5, D and E, respectively) compared with GFP alone (Fig. 5A). Overexpression of truncated C-BID and BID significantly decreased I_{Ca} density at the peak of the I–V curve to -2.7 ± 0.33 and -2.5 ± 0.7 pA/pF, respectively, compared with -7.2 ± 0.6 pA/pF in cells infected with GFP alone. Although Ca^{2+} channel β subunits can influence Ca^{2+} current kinetics, we did not observe consistent changes in the average inactivation rate constant (599 ± 50 ms) determined for the peak Ca^{2+} current elicited at +20 mV between HL-1 cells overexpressing the GFP, C-BID, and BID constructs.

Effect of Overexpression of C-BID and BID on Voltage-Dependent Activation and Inactivation of I_{Ca}

Although we hypothesized that C-BID and BID reduced I_{Ca} in HL-1 cells by retaining $Ca_v1.2$ channels intracellularly, we investigated the possibility that voltage-dependent gating also was altered. The voltage dependence of activation was evaluated by holding cells at -55 mV and imposing 10-mV depolarizing steps between -50 and $+20$ mV. Peak current at each voltage was plotted as a ratio of maximal current elicited at $+20$ mV, and the data were fit by a Boltzmann equation to obtain $V_{1/2}$. The resulting activation curves in Fig. 6A revealed no significant difference ($n = 4$ –5) in $V_{1/2}$ values between cells overexpressing GFP (-14 ± 1 mV), C-BID (-12 ± 1 mV), or BID (-13 ± 1 mV). Inactivation was assessed by subjecting cells to 1400-ms prepulse voltages (10-mV steps) between -50 and 30 mV from a holding potential of -55 mV. Cells were briefly returned to -55 mV for 20 ms, and then they were pulsed to $+20$ mV to assess the effect of inactivating prepulses on channel availability. Peak current at each prepulse voltage was plotted as the ratio of maximal current elicited from a holding potential of -55 mV. Figure 6B shows data fit with the Boltzmann function, and it reveals that $V_{1/2}$ values were not significantly different ($n = 4$ –5) for HL-1 cells overexpressing GFP (-7 ± 2 mV), C-BID (-3 ± 2 mV), and BID (-5 ± 1 mV).

Effect of Overexpression of C-BID and BID on Ca^{2+} Channels Other than $Ca_v1.2$

The finding that C-BID and BID suppressed functional $Ca_v1.2$ channels raised the important question of whether this dominant-negative effect was $Ca_v1.2$ -specific, or whether it extended to other types of Ca^{2+} channels. In particular, L-type Ca^{2+} currents contributed by $Ca_v1.3$ channels (α_{1D}) have been reported in HL-1 cells (Xia et al., 2004). The $Ca_v1.3$ channels can be distinguished from high-threshold $Ca_v1.2$ channels by their activation at mid-voltages and their lower sensitivity to nifedipine at negative membrane potentials (Koschak et al., 2001; Xu and Lipscombe, 2001; Xia et al., 2004; Zhang et al., 2005; Striessnig et al., 2006). Low-threshold T-type Ca^{2+} currents ascribed to Ca_v3 (α_{1G} and α_{1H}) channels also have been described in HL-1 atrial cells (Xia et al., 2003; Hansen et al., 2004; Yang et al., 2005). To collectively activate L-type and T-type channels to obtain “total current”, we imposed progressive 10-mV depolarizing steps over a wide voltage range from -80 to $+50$ mV. Total current was suppressed in HL-1 cells overexpressing C-BID and BID compared with those expressing only the GFP construct (Fig. 7A). Subsequently, the addition of $10 \mu M$ nifedipine revealed two components of total I_{Ca} . The nifedipine-resistant,

residual I_{Ca} showed peak activation at +10 mV, revealing lower threshold Ca^{2+} current (Fig. 7B). Digital subtraction of this I_{Ca} component from total current revealed the nifedipine-sensitive I_{Ca} attributed to $Ca_v1.2$ channels that maximally activated at +30 mV (Fig. 7C). It is noteworthy that the I–V curve for the low-threshold, nifedipine-resistant I_{Ca} was significantly suppressed by C-BID and BID (Fig. 7B), suggesting that the dominant-negative effect of these β_2 subunit mutants extends to Ca^{2+} channels other than $Ca_v1.2$.

Based on these findings, we questioned whether C-BID and BID act nonspecifically to suppress both T-type and L-type Ca^{2+} currents. It is noteworthy that the surface localization of L-type channels, including $Ca_v1.2$ and $Ca_v1.3$, is thought to rely at least partly on β subunits (Catterall et al., 2005). In contrast, T-type channels are missing the known motifs required for β subunit interaction (Perez-Reyes, 1998, 2003; Catterall et al., 2005). To confirm the expression of both $Ca_v1.3$ and Ca_v3 channels in the HL-1 cells, and thus a possible contribution to nifedipine-resistant I_{Ca} , we probed Western blots with α_1 subtype-specific antibodies using mouse brain as a positive control in each of the left lanes (Fig. 8A). In addition to detecting $Ca_v1.2$, α_1 subunits corresponding to $Ca_v1.3$, $Ca_v3.1$ and $Ca_v3.2$ were evident. Subsequently, we applied a depolarizing pulse between –80 and –10 mV to cells overexpressing GFP, C-BID, or BID to maximally activate low-threshold, T-type (Ca_v3) channels but avoid activation of mid- ($Ca_v1.3$) and high ($Ca_v1.2$)-threshold L-type channels (Fig. 8B, top traces). To further isolate T-type current, (1*S*,2*S*)-2-[[[3-(2-benzimidazolyl)propyl] methylamino]ethyl]-6-fluoro-1,2,3,4-tetrahydro-1-isopropyl-2-naphthyl methoxyacetate dihydrochloride hydrate (mibefradil) (7 μ M), a preferential blocker of T-type channels (Martin et al., 2000; Xia et al., 2003) was applied and peak mibefradil-sensitive current elicited at –10 mV was calculated by digital subtraction (Fig. 8B, bottom traces). This transient current was fully intact after BID and C-BID overexpression, suggesting that the decoy effect of the β_2 subunit mutants does not extend to T-type channels (Fig. 8). Rather, the dominant-negative effect of C-BID and BID on nifedipine-resistant current apparently reflects an action on $Ca_v1.3$ channels or other Ca^{2+} channel types.

Discussion

The results of this study provide initial evidence that gene transfer of truncated β_2 subunits into HL-1 cells prevents the proper localization of $Ca_v1.2$ channels in the surface membrane. We took advantage of the chaperone function of the β_2 subunit to alter functional Ca_L channel expression using overexpressed truncated β_2 subunit mutants in HL-1 cells. These mutants were designed to bind the pore-forming α_{1C} subunits but prevent their proper insertion at the surface membrane form functional Ca_L channels. We have demonstrated that two engineered β_2 subunit mutants, C-BID and BID, can serve a dominant-negative function in HL-1 cells, disrupt sarcolemmal localization of $Ca_v1.2$ channels, and specifically decrease I_{Ca} . Thus, the findings establish the principle that engineered β_2 subunits represent a viable molecular approach for reducing the number of functional $Ca_v1.2$ channels in cardiomyocytes. In our experiments, the adenoviral delivery system seems to be highly effective in delivering the β_2 subunit constructs to the HL-1 cells, and the cognate proteins were readily detected. We previously used adenoviral delivery to introduce the full-length β_2 subunit into isolated feline ventricular myocytes to confirm the important role of this regulatory subunit in promoting $Ca_v1.2$ channel expression (Chen et al., 2005). In this

earlier study performed in adult feline cardiomyocytes, we established that adenoviral-mediated gene delivery can indeed alter $\text{Ca}_v1.2$ channel expression and function within 24 to 48 h, a time course similar to that for HL-1 cells.

In the present study, we also examined the dominant-negative effects of C-BID and BID on additional Ca^{2+} channels coexpressed in HL-1 cells with $\text{Ca}_v1.2$. In our cells, mibefradil-sensitive, T-type current was not attenuated by either β_2 subunit mutant, consistent with reports that Ca_v3 channels do not have binding motifs to permit β subunit interaction (Perez-Reyes, 1998, 2003; Catterall et al., 2005). However, other components of nifedipine-resistant current were suppressed by gene delivery of C-BID and BID. One target probably was the $\text{Ca}_v1.3$ channel that provides L-type current at mid-voltages in HL-1 cells, and it is 10-fold less sensitive to nifedipine than $\text{Ca}_v1.2$ channels. Although precise identification of $\text{Ca}_v1.3$ current is difficult due to the lack of a specific antagonist, members of the Ca_v1 gene family, including $\text{Ca}_v1.3$, share an α -interactive domain that enables α_1 - β interaction (Catterall et al., 2005). Thus, a shared feature of the Ca_v1 gene family may be susceptibility to dominant-negative β_2 mutants. In this regard, one potential therapeutic benefit of our mutants may include the treatment of supraventricular tachyarrhythmias, because atrial cells seem to densely express $\text{Ca}_v1.3$ channels, which contribute to normal rhythmicity (Zhang et al., 2005). In contrast, $\text{Ca}_v1.3$ channels are only sparsely expressed in ventricular tissues, and $\text{Ca}_v1.2$ channels seem to be the primary pathway for voltage-gated Ca^{2+} influx. Thus, disruption of expression of $\text{Ca}_v1.2$ channels by our β_2 subunit mutants may be beneficial in pathologies involving Ca^{2+} overload as a strategy to depress $[\text{Ca}^{2+}]_{\text{cyt}}$ and restore normal contractility.

Our findings also provide initial insight into the interactions between the α_{1C} subunit and truncated β_2 subunit decoys that may be required to interrupt the trafficking of functional Ca_L channels to the surface membrane. In contrast to the observation that overexpression of the full-length β_2 subunit at the surface membrane was associated with increased I_{Ca} , the highly distinctive vesicular patterns for N-BID and C-BID localization and the reticular pattern for BID suggest disruption of proper localization of Ca_L channel subunits. Indeed, each of the truncated β_2 subunits was designed to retain the central BID sequence necessary for α_{1C} - β interaction, but at least one of the two additional protein-interaction domains (SH3 and GK) recently reported to be required for channel assembly was eliminated (Chen et al., 2004; Opatowsky et al., 2004; Van Petegem et al., 2004). Thus, the SH3 domain was absent in C-BID, a significant portion of the GK domain was missing in N-BID, and the SH3 domain and a significant portion of the GK domain were missing in BID. In each case, the localization of β_2 subunits was abnormal, supporting previous reports that the β subunits rely on at least three domains in the molecule to confer structural integrity.

Although all three truncated β_2 subunits resulted in altered cellular localization, only C-BID and BID acted as dominant-negative subunits by markedly depressing I_{Ca} in the HL-1 cells. It seems logical to assume that the overexpression of the two successful β_2 mutants, C-BID and BID, overwhelmed interactions between the endogenous α_{1C} and β subunits to sequester α_{1C} subunit. However, the abnormal vesicular localization of N-BID did not confer a decoy function, suggesting that it did not successfully compete for the α_{1C} subunit although it shared the BID binding site with the C-BID and BID negative constructs. This lack of

dominant-negative effect of N-BID is consistent with previous observations that a short N-terminal isoform of β_1 subunit can modulate the gating of cardiac $\text{Ca}_v1.2$ channels, without promoting membrane trafficking of the channel complex (Cohen et al., 2005). Interestingly, a recent report demonstrated a link between the $\text{Ca}_v1.2$ activity and the cell endocytic machinery. In that study, a reduction in the number of channels present at the plasma membrane was observed when the SH3 domain of the β subunit (also located in the N terminus) was expressed in oocytes (Gonzalez-Gutierrez et al., 2007). The removal of the channels from the membrane by this β_2 -SH3 construct is not consistent with our observation that N-BID had no effect on Ca^{2+} currents. This suggests that additional portions of the N terminus of the β subunit may be involved in the regulation of the turnover of the L-type Ca^{2+} channels. Alternatively, it is possible that the spatial cellular localization of N-BID in these vesicle-like compartments may prevent binding interaction with the endogenous α_{1C} protein.

Our study has several limitations. First, the HL-1 cells represent an atrial cell line that expresses robust Ca^{2+} currents and other markers of cardiac function, suggesting that it is a useful model in which to study Ca^{2+} channel regulation (Claycomb et al., 1998; White et al., 2004; Xia et al., 2004). However, the effectiveness of our dominant-negative β_2 constructs clearly may differ between HL-1 cells and native cardiac cells that may express different levels of endogenous β subunits or even process the mutant β subunits differently. Indeed, the success of our β_2 subunit mutants to reduce Ca^{2+} current in cardiac myocytes will depend on the rate of L-type channel turnover in the surface membrane *in vivo*. The precise turnover rate is unknown, but the fact that changes in Ca^{2+} current in response to norepinephrine or thyroid hormone are evident within 24 to 48 h suggests that the membrane turnover of L-type Ca^{2+} channels is a highly dynamic process (Kim et al., 1987; Maki et al., 1996). Second, we did not attempt to pinpoint the cellular sites involved in the aberrant localization of the C-BID, N-BID, and BID mutants. Further detailed colocalization studies will be needed to identify the cellular compartments or organelles that account for these expression patterns.

Although our initial attempt to detect by immunostaining the endogenous α_{1C} subunit in the HL-1 cells was not successful (Supplemental Figs. A and B), we believe that colocalization of the endogenous α_{1C} subunit with the GFP-tagged β_2 subunit constructs would provide important insight into the mechanism by which these β_2 subunits affect Ca^{2+} channel function. Finally, structure-function studies are needed to optimize the design of dominant-negative β_2 constructs to obtain high-affinity molecules that serve as effective decoys for the L-type Ca^{2+} channels as a prelude to expressing it specifically in heart.

It is noteworthy that new molecular approaches designed to down-regulate L-type Ca^{2+} channel subunits in cardiac myocytes represent alternatives to the traditional use of organic Ca^{2+} channel blocking drugs to attenuate $[\text{Ca}^{2+}]_{\text{cyt}}$. The concept of directly titrating the number of Ca^{2+} channel proteins rather than altering the open-state probability of an existing Ca^{2+} channel population may have advantages, particularly if coupled to a cardiac-specific delivery system to provide durability of effect. For example, work by others shows that *in vitro* gene knockdown of the β_2 subunit using RNA interference technology significantly decreased calcium transients in neonatal rat cardiomyocytes, prevented an increase in

relative cell size, and abrogated phenylephrine-induced protein synthesis in a cellular model of hypertrophy (Cingolani et al., 2007). Furthermore, in an aortic-banded rat model of left ventricular hypertrophy, the same authors showed that knockdown of the β_2 subunit was capable of attenuating cardiac hypertrophy without compromising systolic performance, suggesting that the targeted suppression of pathologies involving Ca^{2+} overload may be possible (Cingolani et al., 2007). The adenoviral vectors used in our study are primarily useful in vitro for proof-of-concept, because they are not regarded as ideal clinical vectors due to their immunogenic nature. However, other vectors including adeno-associated viruses containing cardiac-specific promoters may permit targeted knockdown of L-type channels in the heart.

In summary, the data presented in the current study do support the concept that adenovirus-mediated gene transfer of dominant-negative β_2 subunit mutants can effectively and specifically modulate L-type calcium channel expression and function in the HL-1 cell model system of cardiomyocytes.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank Drs. Zhen Wang and Larisa Buzdugan for excellent technical support, and GibAnn Berryhill for assistance in preparing the manuscript.

This work was supported in part by the University of Arkansas for Medical Sciences (UAMS) Department of Internal Medicine Departmental Research Endowments (to S.T. and J.D.M.), UAMS Department of Pharmacology and Toxicology Research Funds (to J.R.S.), the Department of Veterans Affairs in the form of a VA Merit Award (to J.D.M.), and United States Public Health Service Grants R01 HL-68406 and R01 HL-050818 (to N.J.R.) from the National Institutes of Health (NIH). The University of California Davis/National Institute of Neurological Disorders and Stroke (NINDS)/National Institute of Mental Health (NIMH) NeuroMab Facility (mouse α_{1C} subunit antibody) is supported by NIH Grant U24 NS-050606. The mouse α_{1C} subunit monoclonal antibody (clone L57/46) used in the present study was developed by and/or obtained from the University of California Davis/NINDS/NIMH NeuroMab Facility and maintained by the Department of Pharmacology, School of Medicine, University of California, Davis, CA.

ABBREVIATIONS

Ca_v	Ca^{2+} channel
BID	β interaction domain
SH3	Src homology 3
GK	guanylate kinase
GFP	green fluorescence protein
Full-β_2	full β_2 subunit
Ca_L	L-type Ca^{2+} channel
PCR	polymerase chain reaction

bp	base pair(s)
CMV	cytomegalovirus
HEK	human embryonic kidney
I_{Ca}	Ca ²⁺ channel current
I–V	current-voltage
V_{1/2}	half-maximal activation

References

- Aon MA, Cortassa S, Marban E, O'Rourke B. Synchronized whole cell oscillations in mitochondrial metabolism triggered by a local release of reactive oxygen species in cardiac myocytes. *J Biol Chem*. 2003; 278:44735–44744. [PubMed: 12930841]
- Bers DM. Cardiac excitation-contraction coupling. *Nature*. 2002; 415:198–205. [PubMed: 11805843]
- Bolli R, Marban E. Molecular and cellular mechanisms of myocardial stunning. *Physiol Rev*. 1999; 79:609–634. [PubMed: 10221990]
- Catterall WA, Perez-Reyes E, Snutch TP, Striessnig J. International Union of Pharmacology. XLVIII. Nomenclature and structure-function relationships of voltage-gated calcium channels. *Pharmacol Rev*. 2005; 57:411–425. [PubMed: 16382099]
- Chen X, Zhang X, Kubo H, Harris DM, Mills GD, Moyer J, Berretta R, Potts ST, Marsh JD, Houser SR. Ca²⁺ Influx-induced sarcoplasmic reticulum Ca²⁺ overload causes mitochondrial-dependent apoptosis in ventricular myocytes. *Circ Res*. 2005; 97:1009–1017. [PubMed: 16210547]
- Chen Y-H, Li MH, Zhang Y, He LL, Yamada Y, Fitzmaurice A, Shen Y, Zhang H, Tong L, Yang J. Structural basis of the subunit interaction of voltage-gated Ca²⁺ channels. *Nature*. 2004; 429:675–680. [PubMed: 15170217]
- Cingolani E, Ramirez Correa GA, Kizana E, Murata M, Cho HC, Marban E. Gene therapy to inhibit the calcium channel beta subunit: physiological consequences and pathophysiological effects in models of cardiac hypertrophy. *Circ Res*. 2007; 101:166–175. [PubMed: 17556655]
- Claycomb WC, Lanson NA, Stallworth BS, Egeland DB, Delcarpio JB, Bahinski A, Izzo N. HL-1 cells: a cardiac muscle cell line that contracts and retains phenotypic characteristics of the adult cardiomyocyte. *Proc Natl Acad Sci U S A*. 1998; 95:2979–2984. [PubMed: 9501201]
- Cohen RM, Foell JD, Balijepalli RC, Shah V, Hell JW, Kamp TJ. Unique modulation of L-type Ca²⁺ channels by short auxiliary beta1d subunit present in cardiac muscle. *Am J Physiol Heart Circ Physiol*. 2005; 288:H2363–H2374. [PubMed: 15615847]
- De Waard M, Pragnell M, Campbell KP. Ca²⁺ channel regulation by a conserved beta subunit domain. *Neuron*. 1994; 13:495–503. [PubMed: 8060623]
- De Waard M, Scott VE, Pragnell M, Campbell KP. Identification of critical amino acids involved in alpha1-beta interaction in voltage-dependent Ca²⁺ channels. *FEBS Lett*. 1996; 380:272–276. [PubMed: 8601439]
- Fan QI, Vanderpool KM, O'Connor J, Marsh JD. Decoy calcium channel beta subunits modulate contractile function in myocytes. *Mol Cell Biochem*. 2003; 242:3–10. [PubMed: 12619859]
- Foell J, Balijepalli R, Delisle B, Yunker A, Robia S, Walker JW, McEnery M, January C, Kamp TJ. Molecular heterogeneity of calcium channel beta-subunits in canine and human heart: evidence for differential subcellular localization. *Physiol Genomics*. 2004; 17:183–200. [PubMed: 14762176]
- Gonzalez-Gutierrez G, Miranda-Laferte E, Neely A, Hidalgo P. The Src homology 3 domain of the beta-subunit of voltage-gated calcium channels promotes endocytosis via dynamin interaction. *J Biol Chem*. 2007; 282:2156–2162. [PubMed: 17110381]
- Hamill O, Marty A, Neher E, Sakmann B, Sigworth F. Improved patch-clamp techniques for high resolution current recording from cells and cell free membrane patches. *Pflugers Arch*. 1981; 391:85–100. [PubMed: 6270629]

- Hansen JP, Chen RS, Larsen JK, Chu PJ, Janes DM, Weis KE, Best PM. Calcium channel $[\gamma]$ subunits are unique modulators of low voltage-activated (Cav3.1) calcium current. *J Mol Cell Cardiol.* 2004; 37:1147–1158. [PubMed: 15572045]
- He TC, Zhou S, da Costa LT, Yu J, Kinzler KW, Vogelstein B. A simplified system for generating recombinant adenoviruses. *Proc Natl Acad Sci U S A.* 1998; 95:2509–2514. [PubMed: 9482916]
- Kim D, Marsh JD, Smith TW. Effects of thyroid hormone on slow Ca channel function in cultured chick ventricular cells. *J Clin Invest.* 1987; 80:88–94. [PubMed: 2439547]
- Koschak A, Reimer D, Huber I, Grabner M, Glossmann H, Engel J, Striessnig J. α_1D (Cav1.3) Subunits can form L-type Ca^{2+} channels activating at negative voltages. *J Biol Chem.* 2001; 276:22100–22106. [PubMed: 11285265]
- Maki TM, Gruver EJ, Davidoff AJ, Izzo N, Toupin D, Colucci W, Marks AR, Marsh JD. Regulation of calcium channel expression in neonatal myocytes by catecholamines. *J Clin Invest.* 1996; 97:656–663. [PubMed: 8609220]
- Martin RL, Lee JH, Cribbs LL, Perez-Reyes E, Hanck DA. Mibefradil block of cloned T-type calcium channels. *J Pharmacol Exp Ther.* 2000; 295:302–308. [PubMed: 10991994]
- Niranjan V, Telemaque S, deWit D, Gerard RD, Yanagisawa M. Systemic hypertension induced by hepatic overexpression of human preproendothelin-1 in rats. *J Clin Invest.* 1996; 98:2364–2372. [PubMed: 8941655]
- Opatowsky Y, Chen CC, Campbell KP, Hirsch JA. Structural analysis of the voltage-dependent calcium channel beta subunit functional core and its complex with the α_1 interaction domain. *Neuron.* 2004; 42:387–399. [PubMed: 15134636]
- Opatowsky Y, Chomsky-Hecht O, Kang MG, Campbell KP, Hirsch JA. The voltage-dependent calcium channel β subunit contains two stable interacting domains. *J Biol Chem.* 2003; 278:52323–52332. [PubMed: 14559910]
- Perez-Reyes E, Castellano A, Kim HS, Bertrand P, Bagstrom E, Lacerda AE, Wei X, Birnbaumer L. Cloning and expression of a cardiac/brain beta subunit of the L-type calcium channel. *J Biol Chem.* 1992; 267:1792–1797. [PubMed: 1370480]
- Perez-Reyes E. Molecular characterization of a novel family of low voltage-activated, T-type, calcium channels. *J Bioenerg Biomembr.* 1998; 30:313–318.
- Perez-Reyes E. Molecular physiology of low-voltage-activated t-type calcium channels. *Physiol Rev.* 2003; 83:117–161. [PubMed: 12506128]
- Pesic A, Madden JA, Pesic M, Rusch NJ. High blood pressure upregulates arterial L-type Ca^{2+} channels: is membrane depolarization the signal? *Circ Res.* 2004; 94:e97–e104. [PubMed: 15131006]
- Pragnell M, DeWaard M, Mori Y, Tanabe T, Snutch TP, Campbell KP. Calcium channel beta-subunit binds to a conserved motif in the I–II cytoplasmic linker of the α_1 -subunit. *Nature.* 1994; 368:67–70. [PubMed: 7509046]
- Stimers JR, Ha J, Rock TM. Electrical remodeling caused by hypertrophy in HL-1 cells (Abstract). *Circulation.* 2003; 108:IV56.
- Striessnig J, Koschak A, Sinnegger-Brauns MJ, Hetzenauer A, Nguyen NK, Busquet P, Pelster G, Singewald N. Role of voltage-gated L-type Ca^{2+} channel isoforms for brain function. *Biochem Soc Trans.* 2006; 34:903–909. [PubMed: 17052224]
- Telemaque-Potts S, John SK, Grain T, Potts JT, Marsh JD. Expression of beta2 subunit mutants alters localization of L-type calcium channels in rat adult cardiomyocytes (Abstract). *FASEB J.* 2003; 17:A537.
- Van Petegem F, Clark KA, Chatelain FC, Minor DL. Structure of a complex between a voltage-gated calcium channel beta-subunit and an alpha-subunit domain. *Nature.* 2004; 429:671–675. [PubMed: 15141227]
- White SM, Claycomb W, Constantin PE. Cardiac physiology at the cellular level: use of cultured HL-1 cardiomyocytes for studies of cardiac muscle cell structure and function. *Am J Physiol Heart Circ Physiol.* 2004; 286:H823–H829. [PubMed: 14766671]
- Xia M, Salata JJ, Figueroa DJ, Lawlor A-M, Liang HA, Liu Y, Connolly TM. Functional expression of L- and T-type Ca^{2+} channels in murine HL-1 cells. *J Mol Cell Cardiol.* 2004; 36:111–119. [PubMed: 14734053]

- Xia M, Imredy JP, Santarelli VP, Liang HA, Condra CL, Bennett P, Koblan KS, Connolly TM. Generation and characterization of a cell line with inducible expression of Cav3.2 (T-type) channels. *Assay Drug Dev Technol*. 2003; 1:637–645. [PubMed: 15090236]
- Xu W, Lipscombe D. Neuronal Ca(V)1.3 α 1 L-type channels activate at relatively hyperpolarized membrane potentials and are incompletely inhibited by dihydropyridines. *J Neurosci*. 2001; 21:5944–5951. [PubMed: 11487617]
- Yang Z, Shen W, Rottman JN, Wikswo JP, Murray KT. Rapid stimulation causes electrical remodeling in cultured atrial myocytes. *J Mol Cell Cardiol*. 2005; 38:299–308. [PubMed: 15698836]
- Zhang Z, He Y, Tuteja D, Xu D, Timofeyev V, Zhang Q, Glatter KA, Xu Y, Shin HS, Low R, et al. Functional roles of Cav1.3(α 1D) calcium channels in atria: insights gained from gene-targeted null mutant mice. *Circulation*. 2005; 112:1936–1944. [PubMed: 16172271]

Fig. 1.
Schematic representation of rat cDNA constructs expressing Full- β_2 and truncated β_2 subunits N-BID, BID, and C-BID. EGFP, enhanced GFP.

Fig. 2.
A, agarose gel of PCR products corresponding to the α_{1C} subunit of $\text{Ca}_v1.2$ and native β_1 , β_{2a} , β_{2b} , β_3 , and β_4 subunits in HL-1 cells. GAPDH, glyceraldehyde-3-phosphate dehydrogenase. B, representative whole-cell I_{Ca} recorded from an HL-1 cells before (left) and after (right) exposure to 10 μ M nifedipine for 2 min. Similar results were obtained in three other nifedipine-treated cells.

Fig. 3.

Immunoblot of total cell lysates from HL-1 cells (40 μ g protein/lane) infected with five different adenoviruses expressing GFP alone or GFP-fused β_2 subunit constructs. Lane 1, untreated cells; lane 2, cells infected with Full- β_2 (~106 kDa); lane 3, C-BID (~88 kDa); lane 4, N-BID (~66 kDa); lane 5, BID (~42 kDa); and lane 6, GFP (~30 kDa). A, immunodetection using GFP antibody. B, gel blots depicting expression of α_{1C} (240- and ~200-kDa bands) and β -actin (~42 kDa, internal loading control) using specific antibodies. Data are shown as mean $\pm$ S.E.M. of six independent experiments.

Fig. 4. Fluorescence microscopic images of HL-1 cells show a different localization pattern for each β_2 subunit construct at 24 h after infection. Untreated cells (A) and cells infected with GFP (B), Full- β_2 (C), C-BID (D), N-BID (E), and BID (F). Arrows indicate the cell membrane localization of fluorescence. Scale bar, 10 μ m.

Fig. 5.

Effect of adenovirus-mediated overexpression of Full- β_2 subunit and different truncated β_2 subunit constructs on I_{Ca} primarily attributed to $Ca_v1.2$ channels in HL-1 cells. A, representative raw current traces (-30 to +20 mV) of whole-cell I_{Ca} measured in cells expressing GFP (A), Full- β_2 (B), N-BID (C), C-BID (D), or BID (E). Average $I-V$ data were normalized to cell capacitance. Cells were held at -40 mV, and they were given a series of 200-ms pulses from -40 to +50 mV, using progressive 10-mV depolarizing steps. F, $I-V$ relationship shows I_{Ca} as a function of membrane potential at which it was elicited in cells

expressing GFP (■), Full- β_2 (▽), N-BID (○), C-BID (□), or BID (●). Average peak I_{Ca} was plotted versus voltage. Data are shown as mean $\pm$ S.E.M. §, $P < 0.05$ between Full- β_2 -infected cells and cells infected with GFP alone ($n = 10$ – 12). *, $P < 0.05$ between BID-infected cells and cells infected with GFP alone ($n = 10$ each). #, $P < 0.05$ between C-BID-infected cells and cells infected with GFP alone ($n = 7$ – 10).

Fig. 6.

Effect of adenovirus-mediated overexpression of different truncated β_2 subunit constructs on voltage-dependent activation and inactivation of Ca^{2+} channels primarily attributed to $\text{Ca}_v1.2$. A, activation curve showing normalized currents elicited at various command potentials (-50 to $+20$ mV) from a holding potential of -55 mV. $I/I_{\max}$ value was calculated at each voltage, and it was plotted against the voltage. The solid lines represent fits with the Boltzmann function. There was no significant difference in $V_{1/2}$ between cells overexpressing GFP compared with C-BID and BID ($n = 4-5$). B, voltage-dependent inactivation of Ca^{2+} channel currents using a double-pulse protocol. Prepulse voltage steps (-50 to 30 mV) from a holding potential of -55 mV were imposed, and channel availability evaluated by a subsequent voltage pulse from -55 to $+20$ mV. $I/I_{\max}$ value was calculated at each voltage and plotted against the voltage. Data fit with the Boltzmann function revealed similar half-inactivation voltages for GFP-, C-BID-, and BID-overexpressing cells. Data values are expressed as mean $\pm$ S.E.M. ($n = 4-5$).

Fig. 7.

Calcium current separation using nifedipine to determine the effect of C-BID and BID overexpression on different Ca^{2+} current components in HL-1 cells. Currents were elicited using 200-ms depolarizing steps (-60 to 50 mV) from a holding potential of -80 mV, in 10 -mV increments. The currents were normalized to cell capacitance. The resulting I-V relationships are shown for cells overexpressing GFP, BID, or C-BID before (A, total current) and after (B, nifedipine-resistant current) exposure of cells to 10 μM nifedipine. C, I-V relationship for nifedipine-sensitive Ca^{2+} current was obtained by subtracting the nifedipine-resistant current from total current. All I-V relationships (A–C) were significantly suppressed by overexpression of BID and C-BID compared with the control construct GFP ($n = 4$ – 5 each point).

Fig. 8.

A, Western blot confirming that HL-1 cells express L-type (Ca_v1.2 and Ca_v1.3) and T-type (Ca_v3.1 and Ca_v3.2) Ca²⁺ channels. The molecular masses for the proteins of interest were Ca_v1.2, 190 to 240 kDa; Ca_v1.3, 250 kDa; Ca_v3.1, 240 kDa; and Ca_v3.2, 230 kDa. Br, brain. B, low-threshold current was elicited by progressive 10-mV pulses from -80 to -10 mV in control solution in HL-1 cells overexpressing GFP, C-BID, and BID (top traces). The transient current was reduced by 7 μ M mibefradil, a preferential blocker of T-type channels (bottom traces). C, bar diagram shows the peak mibefradil-sensitive Ca²⁺ current density at -10 mV, which was calculated as the difference between the peak Ca²⁺ current density at -10 mV before and after the treatment with mibefradil. There was no significant difference in mibefradil-sensitive Ca²⁺ current among the three groups of cells ($n = 4-5$).