Published in final edited form as:

Contemp Clin Trials. 2015 July ; 43: 172–178. doi:10.1016/j.cct.2015.06.003.

FAMily CEntered (FACE) advance care planning: Study design and methods for a patient-centered communication and decision-making intervention for patients with HIV/AIDS and their surrogate decision-makers

Allison L. Kimmel^a, Jichuan Wang^{a,b}, Rachel Scott^c, Linda Briggs^d, and Maureen E. Lyon^{a,b}

Allison L. Kimmel: akimmel@childrensnational.org; Jichuan Wang: jiwang@cnmc.org; Rachel Scott:

Rachel.k.scott@medstar.org; Linda Briggs: lbriggs@gundersonhealth.org; Maureen E. Lyon: mlyon@cnmc.org

^aCenter for Translational Science, Children's National Health System, 111 Michigan Avenue NW, Washington, DC, USA

^bThe George Washington University School of Medicine and Health Sciences, 2300 I Street NW, Washington, DC, USA

^cMedStar Washington Hospital Center, 110 Irving St NW, Washington, DC, USA

^dGunderson Lutheran Medical Foundation, Inc., 1900 South Avenue, La Crosse, WI, USA

Abstract

Although the Human Immunodeficiency Virus/Acquired Immunodeficiency Syndrome (HIV/AIDS) has become a chronic illness, disease-specific advance care planning has not yet been evaluated for the palliative care needs of adults with HIV/AIDS. This prospective, longitudinal, randomized, two-arm controlled clinical trial aims to test the efficacy of FAMily CEntered Advance Care Planning among adults living with AIDS and/or HIV with co-morbidities on congruence in treatment preferences, healthcare utilization, and quality of life. The FAMily CEntered intervention arm is two face-to-face sessions with a trained, certified facilitator: Session 1) Disease-Specific Advance Care Planning Respecting Choices Interview; Session 2) Completion of advance directive. The Healthy Living control arm is: Session 1) Developmental/Relationship History; Session 2) Nutrition. Follow-up data will be collected at 3, 6, 12, and 18-month post-intervention. A total of 288 patient/surrogate dyads will be enrolled from five hospital-based, out-patient clinics in Washington, District of Columbia. Participants will be HIV positive and 21 years of age; surrogates will be 18 years of age. Exclusion criteria are homicidality, suicidality, psychosis, and impaired cognitive functioning. We hypothesize that this intervention will enhance patient-centered communication with a surrogate decision-maker about end of life treatment preferences over time, enhance patient quality of life and decrease health care utilization. We further hypothesize that this intervention will decrease health disparities for Blacks in completion

Corresponding author: Allison L Kimmel, Center for Translational Science, Children's National Health System, 111 Michigan Avenue NW, Washington, DC, 20010, USA. akimmel@childrensnational.org; Phone: +1 202 476 3565.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

of advance directives. If proposed aims are achieved, the benefits of palliative care, particularly increased treatment preferences about end-of-life care and enhanced quality of life, will be extended to people living with AIDS.

Keywords

HIV/AIDS; Advance care planning; Decision making; Family intervention; Quality of life; Palliative care

Introduction

Despite the efficacy and availability of antiretroviral therapy and the associated longer life expectancy, there were 1,285 deaths among persons with HIV in the District of Columbia (DC) between 2008 and 2012, with 41% of the deaths HIV-related [1]. Blacks in DC make up under half of DC residents but account for 75% of all residents living with HIV [1]. Likewise, among individuals living with HIV in 2012, 85.5% of deaths were Black residents [1]. Blacks nationwide are half as likely as Whites to have completed advance directives [2]. Barriers may include the historical experience of discrimination against Blacks by health care institutions, or misinterpretation of Do Not Resuscitate (DNR) orders as euthanasia or an attempt to deny beneficial care [3–6].

The negative consequences of no or poor Advance Care Planning (ACP) include unmet care or delivery of unnecessary or unwanted care [7–9], dismissing the importance of non-relative caregivers such as gay partners [10–16], loss of respect for autonomy and loss of decision-making capacity of the person living with HIV/AIDS (PLWHA) [17]. End of life discussions are often deferred to the physician or patients are handed an advance directive document by a clerk at the time of hospital admission. Often, patients have little comprehension of what advanced directives are. A surrogate decision maker might be identified, but with no communication between the surrogate and the patient about values and goals of care. Advance Care Planning can significantly reduce these negative consequences by optimizing quality of life (QOL) [18–26], facilitating autonomy, and increasing access to information and choice [27]. Disease Specific-Advanced Care Planning (DS-ACP) has demonstrated increased congruence in treatment preferences between patients and caregivers and increased likelihood that patients' preferences will be honored [28–31]. However, the benefits of ACP for adult PLWHA have not yet been demonstrated through DS-ACP [32–35].

Objectives of the FACE ACP Study

Our goal is to test the implementation of ACP among adult people living with AIDS or HIV and co-morbidities (e.g. Hepatitis C) in Washington, DC. We developed and tested a successful HIV-specific ACP program for HIV positive teens and their families, the Family CEntered (FACE) ACP [28–30] in consultation with Co-Investigator Linda Briggs, who was critical in developing adult models of ACP for other serious medical conditions [31, 36–37]. FACE bridges important gaps in communication research with PLWHA. Our intervention

promotes an active ACP decision-making process focused on the patient/surrogate dyad to address end of life treatment choices.

Aims and hypotheses

Aim 1—To determine the efficacy of FACE ACP on congruence in treatment preferences between PLWHA and their surrogates over time. The aim of congruence is to ensure that adult patients have a voice in their medical treatment even in the event when the patient cannot speak for him or herself, and that the surrogate decision maker knows what the patient wants and has made a commitment to honor the patient's wishes. *We hypothesize that the development of congruence may not be homogeneous and FACE may influence the pattern of congruence development. We further hypothesize that different patterns of congruence development may have different effects on health care utilization. Lastly, we hypothesize that compared to control, FACE participants will better maintain congruence over time.*

Aim 2—To determine the efficacy of FACE on key components of QOL for PLWHA. *We hypothesize FACE participants will increase or better maintain psychosocial QOL compared to controls.*

Aim 3—To minimize health disparities in ACP between Blacks and non-Blacks and identify factors associated with disparities. *We hypothesize Blacks in the FACE intervention will complete advance directives at a rate comparable to non-Blacks, and at significantly greater rates compared to controls.*

Study design

Overview

This study is a prospective, longitudinal, two-arm randomized controlled clinical trial (RCCT). This trial will test the efficacy of the FACE intervention on study outcomes over 18 months post-intervention. PLWHA/surrogate dyads (N=288) will be enrolled and randomized to the FACE ACP intervention or Healthy Living Control (HLC) condition at a ratio of 2:1 (N=192 FACE ACP dyads and N=96 HLC dyads) with the goal of obtaining data from approximately 202 dyads at 18 months post-intervention (Figure 1. FACE Consort Diagram).

Inclusion of women and minorities

Gender distribution is approximately 50% male, 49% female and 1% transgender. Racial distribution is approximately 83% Black. Ethnic distribution among all racial groups is approximately 8% Hispanic or Latino.

Participant dyads

During enrollment, PLWHA will choose a surrogate decision-maker, using the Disease Specific ACP guidelines [38]. Patient/surrogate dyads commit to being in the study regardless of which arm they are randomly assigned.

Sites

Through recruitment at five hospital-based, out-patient clinics in Washington, DC, we will meet our enrollment goals and have adequate statistical power to increase generalizability of findings. Block randomization by study site to condition will control for idiosyncrasies of site-specific effects. Each study site's interdisciplinary teams are experienced in research recruitment, specifically having successfully recruited and retained PLWHA, including Black PWLHA [39].

Eligibility, recruitment, consent, randomization

The study has been approved by the five participating sites' Institutional Review Boards (IRB) and will be reviewed twice a year by the FACE Safety Monitoring Committee. Study research assistants (RA) in consultation with Health Care Providers (HCP) approach potentially eligible participants. HCPs are not present at the time of approach to prevent any perception of coercion. PLWHA and surrogate decision-makers complete written informed consent. Participant eligibility is verified prior to completing baseline study measures. Inclusion and exclusion criteria are presented in Table 1.

Once consent is obtained, RAs administer the screening assessments. The consent and screening process takes approximately 45 minutes. The baseline visit also lasts approximately 45 minutes. Based on the randomized group assigned, FACE ACP and HLC participants will receive two 60–90 minute sessions scheduled one week apart. Sessions 1 and 2 will be followed by process questionnaires administered by the RA-Assessor. Post-intervention follow-up visits at months 3, 6, 12 and 18 will be conducted to assess outcomes.

Dyads are randomized to either FACE ACP or HLC groups using a randomly permuted block design and a 2:1 ratio, controlling for site and perinatal verse other modes of transmission. Randomization is performed via RedCAP at the coordinating site once the dyad has completed baseline procedures.

Assessment and outcome measures

Outcome assessments will be conducted at screening, baseline, following sessions 1 and 2, and post intervention follow-up visits, with the exception of the Statement of Treatment Preferences, which is administered at the time of Session 1. See Table 2 for schedule of study assessment and measures.

Interventions

Research team

Each site has a Co-Investigator (Co-I) and three research assistants (RAs), the RA-Interventionist, RA-Control, and RA-Assessor. Site Co-Is oversee site activities and provide weekly, face-to-face supervision of the RAs. Research Assistants are trained to competency criteria and assist with recruitment, screening, enrollment and baseline screening measure collection. Booster training sessions for RAs are held as needed. Only the RA-Assessor is permitted to administer post-randomization assessments.

FACE ACP intervention

In Session 1, the RA-Interventionist, through the Respecting Choices interview ® [38], facilitates conversation about goals, values and experiences which inform decision-making about EOL treatment preferences. In Session 2, the RA-Interventionist facilitates completion of an advance directive. See Table 3. Sessions are conducted in a private office. Session 1 is audio or video-taped to monitor fidelity to the protocol and possible contamination between the study arms. Review is provided by the Principle Investigator (PI) or selected Co-I (Briggs). Trained/certified facilitators include mental health professionals and graduate students in public health, psychology, counseling and nursing.

Healthy Living Control (HLC)

The HLC also uses a dyadic approach; however, participants do not receive the ACP intervention curriculum. Dyads assigned to the HLC condition also meet twice, one week apart, also for approximately 60 minutes each session. See Table 4.

Data analysis

Statistical analyses will use outcome measures scores (e.g. congruence in treatment preferences, QOL, healthcare utilization) and time varying-covariates variables (Highly Active Antiretroviral Therapy [HAART] adherence, religion/spirituality). Descriptive statistics will characterize study participants overall and within intervention groups. Multivariate statistical analyses will assess the efficacy of FACE. Intent-to-treat (ITT) analysis recognizes noncompliance and protocol deviations and includes all randomized participants in final analysis, allowing for imitation of clinical practice [40]. The GMM [41–45], a popular mixture model designed for longitudinal data analysis, will be used to examine the pattern of outcome development trajectories. We intend to identify possible distinctive unknown a priori latent classes of outcome development trajectories over time (e.g. congruence in treatment preferences between PLWHA and surrogates); classify individuals into their most likely latent trajectory classes; assess how the typology of outcome development differs between the FACE and control groups; and examine how the trajectory pattern influences the distal outcome (e.g. health care utilization. Both time-invariant covariates (e.g. demographic variables) [46–47] and time-varying covariates (e.g. HAART adherence and religion/spirituality) will be controlled in the GMM model.

Analytical plans

Aim 1—Congruence in decision-making for EOL care will be measured based on agreement (i.e. both patient and his/her surrogate choose the same option) on the Statement of Treatment Preferences. Kappa coefficients will be applied to assess chance-adjusted agreement between patient and surrogate responses. Latent Growth Model (LGM) will be applied to explore the trajectory of congruence development over time [48–50]. Then the GMM will be used to test Hypothesis A by examining the heterogeneity of congruence development [51–52]. Figure 2 shows the GMM, in which y_0 – y_3 are repeated measures of congruence at different time points.

Latent growth intercept and latent growth slope are two latent growth factors, representing the initial level of congruence and rate of change in congruence, respectively. The time scores in the model are set to 0 for Session 1 visit, and 1 for the first follow-up visit (month 3) for the purpose of model identification. Instead of assuming a linear or nonlinear polynomial function (e.g. quadratic or cubic), time scores are set free (denoted by * in Figure 2) to be model estimated. As such, the pattern of change or type of congruence development trajectory will be determined by data. The latent class in the model is a categorical latent variable that captures the pattern of congruence development trajectories. Time-invariant covariates (e.g. race) will be used to predict the trajectories of the congruence development over time, as well as the pattern of congruence development trajectories, and time-varying covariates (HAART adherence, religion/spirituality) may be included to predict the level of congruence at different time points. Hypothesis B will be tested by regressing a distal outcome (e.g., health care utilization) on the latent class of congruence development trajectories (see Figure 2). To test Hypothesis C, we will regress the latent growth slope factor and the latent class variable on FACE ACP, controlling for covariates. The impact of the FACE on the rate of change in congruence over time is allowed to vary across different classes of congruence development trajectories. In Figure 2, dotted lines indicate interactions between covariates and the latent class variable. As attrition is inevitable in longitudinal studies, robust model estimator (e.g. MLR) using the full information maximum likelihood (FIML) will be used for model estimation. Importantly, missing at random (MAR), instead of missing completely at random (MCAR), can be assumed [52–57].

Aim 2—Similarly, the GMM will be used to examine the development trajectory of QoL and effects of FACE on its trajectory. The same time-invariant covariates (age, gender, race, education) will be controlled, while time-varying covariates (HAART adherence, religion/spirituality) will be included in the model to predict measures of QOL at different time points.

Aim 3—The rate of completing advance directives among participants at baseline and each follow-up time point will be documented and the rate disparity between Black and non-Black participants at each of these time point will be described and tested by intervention groups using Chi-square statistics and logistical regression.

Sample size and power

In longitudinal studies, the number of repeated measurements plays an important role in statistical power, because there is a tradeoff between the sample size and the number of repeated measurements [58]. Various sample sizes were estimated corresponding to various combinations of effect size and autocorrelation. For a modest autocorrelation of 0.30 and a small effect size of 0.25, the estimated sample size for a longitudinal analysis to achieve a power of 0.80 at 0.05 level is about N=188 at each of the 4 observation time points. Our proposed baseline sample (N=288) will ensure a large enough statistical power for statistical analyses using traditional models. With application of the advanced latent growth analytic models for longitudinal data analysis, the statistical power would be larger than the traditional analytical methods for a given sample size [58–59].

Discussion

FACE offers five advancements over current standard of care: 1) promotes shared decision-making with families; 2) integrates the evidence-based Disease-Specific ACP curriculum [38]; 3) differs from previously published research on advance directive documentation alone [60–61] by involving the patient/surrogate decision-maker in conversations about treatment preferences, then sharing them with the primary HCP; 4) is grounded in Leventhal's theory of self-regulation and illness representations [62–64]; and 5) acknowledges those who prefer to have their doctor or family make these decisions for them.

Recruitment for dyadic EOL studies has challenges such as identifying a surrogate, scheduling patients and surrogates, and dealing with conflicts that may arise [65–66]. To date, 15% of PLWHA interested in participating could not identify potential surrogate they trusted enough to make decisions for them. An additional site was added to meet enrollment goals. Hiring staff committed to flexibility assisted in addressing scheduling problems. Conflicts are managed by involvement of study ethicist or chaplain. Selection bias may occur as individuals may be hesitant to participate in EOL studies, so demographic data and reason for declining are collected from those who chose not to participate. Threats to validity may arise due to RA's personal values about death and dying. Fidelity procedures are in place to process RA's emotional reactions to implementing the intervention. Finally, some adults who wanted to participate could not, because their chosen surrogate decision-maker was their child who had participated in our adolescent study.

Conclusion

This study will provide replicable, evidence-based model for facilitating decisions about EOL care informed by the patient's representation of illness and lived experience with the involvement of their chosen surrogate decision-maker. Surrogates can then be assured they honored their loved one's wishes, and patients can feel supported to the end, trusting that their families were there.

Acknowledgments

This research was made possible by grant number 1R01NR14-052-03 from the National Institute of Nursing Research (NINR) at the National Institutes of Health. Its contents are solely the responsibility of the authors and do not necessarily represent the official views of NINR. This research was also supported in part by CTSI-CN NCRR Grant # UL1RR031988. We thank our families for agreeing to participate. We thank the FACE Palliative Care Consortium, including Linda Briggs, David Parenti, Laura Coster, Debra Benator, Fred Gordon, Rachel Scott, Leon Lai, Jichuan Wang, Connie Trexler, and Lawrence J. D'Angelo, as well as our consultants Bruce Rapkin, Bea Krauss and Linda Koenig, for their dedicated efforts toward the completion of this study. Special thanks also to our clinical coordinators: Brittney Lee from the coordinating center at Children's National Health System; Ginny Levin from George Washington University Medical Faculty Associates; Chelsea Tanous from Georgetown University Hospital; Leah Squires and Melissa Turner from Veteran's Medical Center; Patricia Tanjutco from MedStar Washington Hospital Center.

Abbreviations

HIV	human immunodeficiency virus
AIDS	acquired immunodeficiency syndrome

PC	palliative care
ACP	advance care planning
FACE	FAMILY CEntered
PLWHA	people living with HIV/AIDS

References

1. Annual Epidemiology & Surveillance Report. Washington, DC: 2013. [http://doh.dc.gov/sites/default/files/dc/sites/doh/page_content/attachments/2013 Annual Report Final Edit.pdf](http://doh.dc.gov/sites/default/files/dc/sites/doh/page_content/attachments/2013%20Annual%20Report%20Final%20Edit.pdf).
2. AARP North Carolina End of Life Care Survey: African American Members. North Carolina American Association of Retired Person; 2003. <http://www.cchospice.org/siteres.aspx?resid=75b95d85-55ef-4873-ad09-258a8b5fab84>
3. Mazanec P, Tyler MK. Cultural considerations in end-of-life care. *Am J Nurs*. 2003; 103(3):50–59. [PubMed: 12626940]
4. Krakauer EL, Crenner C, Fox K. Barriers to optimum end-of-life care for minority patients. *J Am Geriatr Soc*. 2002; 50(1):182–190. [PubMed: 12028266]
5. Crawley LM, Marshall PA, Lo B, Koenig BA. Strategies for culturally effective end-of-life care. *Ann Intern Med*. 2002; 136(9):673–679. [PubMed: 11992303]
6. Emanuel EJ, Fairclough DL, Emanuel LL. Attitudes and desires related to euthanasia and physician-assisted suicide among terminally ill patients and their caregivers. *J Am Med Assoc*. 2000; 284(19):2460–2468.
7. Blank, RH.; Merrick, JC., editors. *End-of-Life Decision Making*. Cambridge: MIT Press; 2004.
8. Krug R, Karus D, Selwyn PA, Raveis VH. Late-stage HIV/AIDS patients' and their familial caregivers' agreement on the palliative care outcome scale. *J Pain Symptom Manage*. 2010; 39(1):23–32. [PubMed: 19782531]
9. Sacajiu G, Raveis V, Selwyn PA. Patients and family care givers' experiences around highly active antiretroviral therapy (HAART). *AIDS Care*. 2009; 12(21):1528–1536. [PubMed: 20024732]
10. Wenger NS, Kanouse DE, Collins RL, Gifford AL, Bozzette SA. End-of-life discussions and preferences among persons living with HIV. *J Am Med Assoc*. 2001; 285(22):2880–2890.
11. Fins, JJ.; Schiff, ND. *Brief : The Afterlife of Terri Schiavo*. Garrison, New York: 2005. p. 8 <http://www.medscape.com/viewarticle/511647>
12. Chandan U, Richter L. Strengthening families through early intervention in high HIV prevalence countries. *AIDS Care*. 2009; 21(S1):76–82. [PubMed: 22380981]
13. Cherlin E, Fried T, Prigerson HG, Schulman-Green D, Johnson-Hurzeler R, Bradley EH. Communication between physicians and family caregivers about care at the end of life: When do discussions occur and what is said? *J Palliat Med*. 2005; 8(6):1176–1185. [PubMed: 16351531]
14. Perkins HS. Controlling Death : The False Promise of Advance Directives. *Ann Intern Med*. 2007; 147(1):51–57. W - 8-W - 9. [PubMed: 17606961]
15. Braun UK, Beyth RJ, Ford ME, McCullough LB. Voices of African American, Caucasian, and Hispanic surrogates on the burdens of end-of-life decision making. *J Gen Intern Med*. 2008; 23(3):267–274. [PubMed: 18172738]
16. Hansen L, Archbold P, Stewart B. Role strain and ease in decision-making to withdraw or withhold life support for elderly relatives. *J Nurs Scholarsh*. 2004; 3(36):233–238. <http://www.ncbi.nlm.nih.gov/pubmed/15495492>. [PubMed: 15495492]
17. Woods SP, Moore DJ, Grant I. Cognitive neuropsychology of HIV-associated neurocognitive disorders HIV-associated neurocognitive disorders. *Neuropsychol Rev*. 2009; 19(2):152–168. [PubMed: 19462243]
18. HIV Care: An Agenda for Change. Princeton, New Jersey: 2008. Workgroup of Palliative and End-of-life care in HIV/AIDS; p. 1-40.

19. Skarupski KA, Fitchett G, Leon CFMDe. Daily Spiritual Experiences in a Biracial, Community-based Population of Older Adults Abstract Objectives. *Aging Ment Health*. 7(14):779–789.
20. Tarakeshwar N PhD, Vanderwerker LC, Prigerson HG. Religious Coping is Associated with the quality of life of patients with advanced Cancer. *J Palliat Med*. 2006; 9(3):646–657. [PubMed: 16752970]
21. Pargament KI, Koenig HG, Tarakeshwar N, Hahn J. Religious Struggle as a predictor of mortality among medically ill elderly patients. *Arch Intern Med*. 2007; 161(15):1881–1885. [PubMed: 11493130]
22. Lee CC, Czaja SJ, Schulz R. The moderating influence of demographic characteristics, social support, and religious coping on the effectiveness of a multicomponent psychosocial caregiver intervention in three racial ethnic groups. *J Gerontol B Psychol Sci Soc Sci*. 2010; 65B(2):185–194. [PubMed: 20056684]
23. Ironson G, Stuetzle R, Ironson D, et al. View of God as benevolent and forgiving or punishing and judgmental predicts HIV disease progression. *J Behav Med*. 2011; 34(6):414–425. [PubMed: 21340531]
24. Trevino KM, Pargament KI, Cotton S, et al. Religious coping and physiological, psychological, social, and spiritual outcomes in patients with HIV/AIDS: cross-sectional and longitudinal findings. *AIDS Behav*. 2010; 14(2):379–389. [PubMed: 18064557]
25. Pargament KI, McCarthy S, Shah P, et al. Religion and HIV: A review of the literature and clinical implications. *South Med J*. 2004; 97(12):1201–1209. [PubMed: 15646758]
26. Lyon ME, Garvie PA, Kao E, et al. Spirituality in HIV-infected adolescents and their families: Family CEntered (FACE) Advance Care Planning and medication adherence. *J Adolesc Heal*. 2011; 48(6):633–636.
27. O'Neill, JF.; Selwyn, PA.; Schietinger, H. A Clinical Guide to Supportive and Palliative Care for HIV/AIDS. United States Health Resources and Services Administration; 2003.
28. Lyon ME, Garvie PA, Briggs L, He J, McCarter R, D'Angelo LJ. Development, feasibility, and acceptability of the Family/Adolescent-Centered (FACE) advance care planning intervention for adolescents with HIV. *J Palliat Med*. 2009; 12(4):363–372. [PubMed: 19327074]
29. Lyon ME, Garvie PA, McCarter R, Briggs L, He J, D'Angelo LJ. Who will speak for me? Improving end-of-life decision-making for adolescents with HIV and their families. *Pediatrics*. 2009; 123(2):e199–e206. [PubMed: 19171571]
30. Lyon ME, Garvie PA, Briggs L, et al. Is it safe? Talking to teens with HIV/AIDS about death and dying: a 3-month evaluation of Family Centered Advance Care (FACE) planning - anxiety, depression, quality of life. *HIV/AIDS Res Palliat Care*. 2010; 2:27–37.
31. Briggs, La; Kirchhoff, KT.; Hammes, BJ.; Song, M-K.; Colvin, ER. Patient-centered advance care planning in special patient populations: a pilot study. *J Prof Nurs*. 2004; 20(1):47–58. [PubMed: 15011193]
32. HIV Surveillance Report: Diagnoses of HIV Infection and AIDS in the United States and Dependent Areas. Atlanta, GA: 2012.
33. Trans-NIH Plan for HIV-Related Research. Washington, DC: 2014. http://www.oar.nih.gov/strategicplan/fy2014/pdf/FY2014_reducing_related.pdf
34. Grady PA. Introduction: Papers from the National Institutes of Health State-of-the-Science Conference on Improving End-of-Life Care. *J Palliat Med*. 2005; 8(S1):S1–S3. [PubMed: 16499457]
35. Greenberg B, McCorkle R, Vlahov D, Selwyn PA. Palliative care for HIV disease in the era of highly active antiretroviral therapy. *J Urban Heal*. 2000; 77(2):150–165.
36. Detering KM, Hancock AD, Reade MC, Silvester W. The impact of advance care planning on end of life care in elderly patients: randomised controlled trial. *BMJ*. 2010; 340:c1345. [PubMed: 20332506]
37. Song M, Ward SE, Connolly MC. Randomized Controlled Trial of SPIRIT: An effective approach to preparing African-American dialysis patients and families for end of life. *Res Nurs Health*. 2009; 32(3):260–273. [PubMed: 19205027]
38. Briggs, LA.; Hammes, BJ. Disease-Specific Patient Centered Advance Care Planning (DS-ACP) Manual. La Crosse, WI: Gundersen Lutheran Medical Foundation, Inc.;

39. Lyon ME, Woodward K. Nonstigmatizing ways to engage HIV-positive African-American teens in mental health and support services: a commentary. *J Natl Med Assoc.* 2003; 95(3):196–200. [PubMed: 12749679]
40. Gupta SK. Intention-to-treat concept: A review. *Perspect Clin Res.* 2011; 2(3):109–112. <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3159210/?report=reader>. [PubMed: 21897887]
41. Fuzhong L, Duncan TE, Duncan SC, Acock A. Latent Growth Modeling of Longitudinal Data: A Finite Growth Mixture Modeling Approach. *Struct Equ Model A Multidiscip J.* 2001; 8(4):493–530.
42. Muthén, BO. Latent variable analysis: Growth mixture modeling and related techniques for longitudinal data. In: Kaplan, D., editor. *Handbook of Quantitative Methodology for the Social Sciences*. Newbury Park, CA: 2004. p. 345–368.
43. Wang, J. *Methods and Applications of Mixture Models. Advances and Applications in Statistical Sciences*; 2010.
44. Muthén B, Muthén LK. Integrating person-centered and variable-centered analyses: growth mixture modeling with latent trajectory classes. *Alcohol Clin Exp Res.* 2000; 24(6):882–891. [PubMed: 10888079]
45. Bates ME. Integrating person-centered and variable-centered approaches in the study of developmental courses and transitions in alcohol use: Introduction to the special section. *Alcohol Clin Exp Res.* 2000; 24(6):878–881. [PubMed: 10888078]
46. Roberts CB, Couper DJ, Chang PP, James SA, Rosamond WD, Heiss G. Influence of life-course socioeconomic position on incident heart failure in blacks and whites: The Atherosclerosis Risk in Communities Study. *Am J Epidemiol.* 2010; 172(6):717–727. [PubMed: 20696652]
47. Smith AK, McCarthy EP, Paulk E, et al. Racial and ethnic differences in advance care planning among patients with cancer: impact of terminal illness acknowledgment, religiousness, and treatment preferences. *J Clin Oncol.* 2008; 26(25):4131–4137. [PubMed: 18757326]
48. McArdle, JJ.; Anderson, ER. Latent growth models for research on aging. In: Birren, JE.; Schaie, KW., editors. *Handbook of Psychology of Aging*. San Diego: Academic Press; 1999. p. 21–44.
49. Meredith W, Tisak J. Latent curve analysis. *Psychometrika.* 1990; 55:107–122.
50. Duncan SC, Duncan TE. Modeling incomplete longitudinal substance use data using latent variable growth curve methodology. *Multivariate Behav Res.* 1994; 29:313–338. [PubMed: 26745232]
51. Muthén, LK.; Muthén, BO. *Mplus: Statistical Analysis with Latent Variables User's Guide. Version 6*. Los Angeles: Muthén & Muthén;
52. Hausman JA, Wise DA. Attrition Bias in Experimental and Panel Data: The Gary Income Maintenance Experiment. *Econometrica.* 1979; 47(2):455–474.
53. Verbeek M. On the estimation of a fixed effects model with selectivity bias. *Econ Lett.* 1990; 34(3): 267–270.
54. Verbeek M, Nijman T. Testing for selectivity bias in panel data models. *Int Econ Rev (Philadelphia).* 33(3):681–703.
55. Wang, JF.; Carlson, RG.; Shi, P. 134th Annual Meeting of the American Public Health Association. Boston: 2006. Dealing with non-random attrition in longitudinal studies: Application of sample selection models.
56. Zabel JE. Estimating fixed and random effects models with selectivity. *Econ Lett.* 1992; 40(3): 269–272.
57. Greene, WH. *LIMDEP: Econometric Modeling Guide*. 9th. Plainview: Economic Software, Inc.; 2002.
58. Diggle, PJ.; Heagerty, PJ.; Liang, K-Y.; Zeger, SL. *Analysis of Longitudinal Data*. 2nd. Oxford: Oxford University Press; 2002.
59. Muthen BO, Curran PJ, Brown H, Hussong A, Khoo S. General Longitudinal Modeling of Individual Differences in Experimental Designs : A Latent Variable Framework for Analysis and Power Estimation. 1997; 2(4):371–402.
60. Solomon MZ, Sellers DE, Heller KS, et al. New and lingering controversies in pediatric end-of-life care. *Pediatrics.* 2005; 116(4):872–883. [PubMed: 16199696]

61. SUPPORT Investigators. A controlled trial to improve care for seriously ill hospitalized patients: The study to understand prognoses and preferences for outcomes and risks of treatments (SUPPORT). *J Am Med Assoc.* 1995; 274(20):1591–1598. doi:
62. Lau RR, Bernard TM, Hartman KA. Further explorations of common-sense representations of common illnesses. *Heal Psychol.* 1989; 8(2):195–219.
63. Leventhal, HH.; Nerenz, DR.; Steele, DJ. Illness representations and coping with health threats. In: Baum, A.; Taylor, SE.; Singer, JE., editors. *Handbook of Psychology and Health: Social Psychological Aspects of Health.* Hillsdale, New Jersey: Erlbaum; 1984. p. 219-252.
64. Donovan HS, Ward S. A representational approach to patient education. *J Nurs Scholarsh.* 2001; 33(3):211–216. [PubMed: 11552546]
65. Quinn C, Dunbar SB, Clark PC, Strickland OL. Challenges and strategies of dyad research: cardiovascular examples. *Appl Nurs Res.* 2010; 23(2):e15–e20. [PubMed: 20420989]
66. Friedman DB, Johnson KM, Owens OL, et al. Developing partnerships and recruiting dyads for a prostate cancer informed decision making program: lessons learned from a community-academic-clinical team. *J cancer Educ.* 2012; 27(2):243–249. [PubMed: 22528633]

Figure 1.
FACE Consort Diagram

Figure 2.
Growth Mixture Model with covariates and distal outcome

Table 1

Inclusion/exclusion criteria for participants.

Inclusion Criteria			Exclusion Criteria	
<i>Patient</i>				
1	Age 21 or older at time of enrollment		1	In Intensive Care Unit
2	Aware of his/her HIV diagnosis		2	<21 years of age at time of enrollment
3	Outpatients or inpatients (except Intensive Care Unit) with advanced HIV as determined by:		3	Unaware of his/her HIV diagnosis
	a.	Detectable viral load (>200 copies per ml) on two occasions in the last 1-year period	4	Non-English speaking
	b.	CD4 count <200 ever	5	Active psychosis or homicidal or suicidal ideation, determined at screening from a structured questionnaire
	c.	Current opportunistic infection	6	Score consistent with Dementia on HIV Dementia Scale
	d.	HIV-infected with a co-morbidity that may significantly limit life expectancy, including malignancy, cirrhosis, cardiomyopathy, end stage renal/kidney or liver disease, diabetes, HIV Associated Neurological Disorder (HAND), etc.	7	Developmentally delayed
4	English speaking			
5	All ethnic groups			
6	All genders			
<hr/>				
<i>Surrogate</i>				
1	Selected by patient		1	<18 years at the time of enrollment
2	Age 18 or older		2	Does not know patient's HIV status
3	Willingness to discuss problems related to HIV and End of Life		3	Non-English speaking
4	English speaking		4	Active psychosis or homicidal or suicidal ideation, determined at screening from a structured questionnaire
5	Consent to participate		5	Score consistent with Dementia on HIV Dementia Scale
6	Knows HIV status of patient		6	Developmentally delayed

Table 2

Schedule of study assessment and measures.

Measures	Participant Completion	Time Point	Description
Demographic Data Interview	Patient & Surrogate	0	Demographic data collection
HIV Dementia Scale	Patient & Surrogate	0, 3,4,5,6	Rapid screener to identify HIV dementia
Psychological Interview	Patient & Surrogate	0	Questions to screen for homicidality & psychosis
Patient Health Questionnaire PHQ-9 for Depression	Patient & Surrogate	0, 3,4,5,6	Assess symptoms of depression and severity
Statement of Treatment Preferences	Patient & Surrogate	1,3,4,5,6	Adapted using experts and community advisory board, tool to express values and goals related to future decision-making re: frequently occurring situations specific to HIV/AIDS
Medication Adherence Self Report Inventory	Patient	0,3,4,5,6	Validated self-report measure of medication adherence, using the Visual Analogue Scale for estimated adherence in the past month.
Medical Outcomes Study-HIV (MOS-HIV)	Patient & Surrogate	0,3,4,5,6	Measures HIV specific Health Related Quality of Life (HRQOL) general health perceptions; pain; physical, role, cognitive and social functioning; mental health; energy/fatigue; health distress and quality of life
Palliative Care Outcome Scale (POS)	Patient & Surrogate	0,3,4,5,6	Valid and reliable measure of patient's current health, social and psychological status
Brief Multidimensional Measurement of Religiousness/Spirituality - BMMRS-adapted	Patient & Surrogate	0,3,4,5,6	Assesses the construct of spiritual functioning, specifically subscales for Forgiveness, Values & Beliefs will be used for study purposes.
Brief-Religious Coping Questionnaire - Brief RCOPE	Patient and Surrogate	0,3,4,5,6	Assesses positive and negative religious coping methods. Study will use 14-item version.
Satisfaction Scale	Patient & Surrogate	1,2	Process measure developed in pilot study to assess satisfaction
Longitudinal Satisfaction Questionnaire	Patient & Surrogate	3,4,5,6	Process measure to assess longitudinal satisfaction with future decision making
Quality of Patient-Interviewer Communication	Patient & Surrogate	1,2	Process measure assesses quality of communication between Patient/Surrogate and the interviewer
Medical Chart Abstraction	Chart Abstraction	0,3,4,5,6	Mode of transmission, CD4 count, viral load, hospitalization, emergency room visit or dialysis since last study visit
Lyon Advance Care Planning Survey	Patient & Surrogate	0	This is a 31-item outcome measure instrument, developed and adapted with permission

Measures	Participant Completion	Time Point	Description
			from the AARP (2003) and Edinger and Smucker (1992).
Threat Appraisal Scale (TAS)	Patient	0, 3, 4, 5, 6	This measure will retrospectively estimate the patient's threat of HIV when patient learned of HIV diagnosis.

0=Screening & Baseline; 1=Session 1; 2=Session 2; 3=3-mo. follow up; 4=6-mo. follow up; 5=12-mo. follow up; 6=18-mo. follow up

Table 3

Family CEntered Advance Care Planning intervention.

Session 1 Foundation	Session 1 Goals	Session 1* Process
Disease-Specific Advance Care Planning (DS ACP) Interview® (Briggs and Hammes), formerly Respecting Choices Interview ®. After session, dyad is giving Gunderson health information sheets on 1) CPR and 2) Tube Feedings by RA-Interventionist.	<ol style="list-style-type: none"> 1. To facilitate conversations and shared decision-making between the patient and surrogate about palliative care, providing an opportunity to express fears, values, spiritual and other beliefs and goals with regard to death and dying 2. To prepare the surrogate to be able to fully represent the patient's wishes 	<p>Patient and surrogate dyad together: <u>Stage 1</u> assesses the patient's understanding of current medical condition, prognosis, complications; <u>Stage 2</u> explores patient's philosophy regarding EOL decision-making and their understanding of the facts; <u>Stage 3</u> reviews rationale for future medical decisions the patient would want the surrogate to understand/act on; <u>Stage 4</u> uses the Statement of Treatment Preferences to describe clinical situations common to AIDS and related treatment choices; <u>Stage 5</u> summarizes the discussion/need for future discussions as situations/preferences change. Gaps in information are identified and referrals are made. <i>(This session will be videotaped)</i></p>
Session 2 Foundation	Session 2 Goals	Session 2* Process
<p>The Five Wishes© is a legal document that helps a person express how they want to be treated if they are seriously ill and unable to speak for him/herself. Unique among living will and health agent forms - it looks to all of a person's needs: medical, personal, emotional, spiritual.</p> <p><i>* The Site Study Coordinator will be responsible for filing The Five Wishes © in the medical record, sending a copy to their treating physician, and giving a copy to the participant.</i></p>	<ol style="list-style-type: none"> 1. Which person the patient wants to make health care decisions for him/her; 2. The kind of medical treatment the patient wants; 3. How comfortable the patient wants to be; 4. How the patient wants people to treat him/her. 5. What the patient wants loved ones to know; 6. Any spiritual or religious concerns the patient may have. 	<p>Processes, such as labeling feelings and concerns, as well as finding solutions to any identified problem, are facilitated. Appropriate referrals are made to help resolve disagreements over decision-making (e.g. a hospital ethicist or their doctor) or spiritual issues (e.g. a hospital chaplain or their clergy). <i>*These sessions may include other loved ones.</i></p>

Table 4

Healthy Living Control condition.

Session 1 Foundation	Session 1 Goals	Session 1 Process
<u>Developmental History</u> Adapted Barkley Developmental History form will be administered, with all medical questions removed from the developmental history to prevent any risk of contamination with the experimental FACE condition. After session, dyad is given Gunderson health information sheets on 1) CPR and 2) Tube Feedings by RA-Control.	To take a non-medical developmental history. The RA-Control will conduct the session in a structured interview format to provide a control for time (approximately 45–60 minutes) and attention. For surrogates who do not know the patient's developmental history, a set of relationship questions is asked.	Patient and surrogate dyad participate together: 1. Conversation is facilitated by RA-Control in a private room. 2. The patient will be asked first, then the surrogate, to parallel the structure of the intervention. 3. If surrogate is friend/partner, please refer to questions specific to friend/partner in the script. <i>(This session will be videotaped)</i>
Session 2 Foundation	Session 2 Goals	Session 2 Process
<u>Nutrition Session.</u> Using the American Academy of Pediatrics Bright Futures counseling guides, participants will be asked questions about nutrition.	To provide nutrition information using the American Academy of Pediatrics Bright Futures counseling guides. Content in this review is not related to Advance Care Planning or the FACE intervention.	This structured questionnaire/information will be administered to the patient/surrogate dyad by the trained RA-Control together in a private room. The patient will be asked first, then the surrogate to parallel the structure of the intervention.