CORRESPONDENCE

The Presentation, Diagnosis, and Treatment of Sexually Transmitted Infections

by Prof. Dr. med. Florian M.E. Wagenlehner, Prof. Dr. med. Norbert H. Brockmeyer, Dr. med. Thomas Discher, Prof. Dr. med. Klaus Friese, and Prof. Dr. med. Thomas A. Wichelhaus in issue 1–2/2016

Prof. Dr. med. Dietrich Abeck
München
professorabeck@mytum.de

Conflict of interest statement

Prof. Abeck has received consultant fees, registration fee reimbursement, travel and accommodation expenses, and speaking honoraria from Taurus Pharma.

Morphological Variants Forgotten

I would like to add some comments that I think are important to the successful and important review paper on sexually transmitted infections (1).

Chancroid caused by *Haemophilus ducreyi* should at least be mentioned together with diseases caused by genital, anal, perianal, or oral ulcers. In addition to the differential diagnosis between soft chancre and syphilis, morphological variants of the disease are difficult to distinguish from a herpes infection (“dwarf chancroid”) (2). A recent prospective study from Paris showed a *Haemophilus ducreyi* infection in 3% of cases of genital ulcer (3). The disease is not usually diagnosed. Neither the complex media that is necessary to grow the microaerophile pathogen in culture, nor the necessary nucleic acid amplification methods, are routinely available (4).

Therapy for human papillomavirus diseases is prone to errors. Cyclosporin and 5-fluorouracil are not indicated for the treatment of anogenital warts. Permitted substances for patient self-medication, which are recommended by the large STD specialist societies CDC and IUSTI, are imiquimod, podophyllotoxin, and the purified dry extract of green tea leaves with the active substance epigallocatechin gallate. Only these three substances are mentioned in the current CDC treatment guidelines (5).

Cryotherapy, similar to curettage or treatment with electrocautery, is a surgical procedure and is therefore a destructive method.

DOI: 10.3238/arztebl.2016.0460a

REFERENCES

1. Wagenlehner FME, Brockmeyer NH, Discher T, Friese K, Wichelhaus TA: The presentation, diagnosis and treatment of sexually transmitted infections. Dtsch Arztebl Int 2016; 113: 11–22.
2. Ballard RC, Abeck D, Korting HC, Dangor Y, Braun-Falco O: Morphologic variants of genital ulcer caused by *Haemophilus ducreyi*. Hautarzt 1989; 40: 443–7.
3. Hope-Rapp E, Anyfantakis V, Fouéré S, et al.: Etiology of genital ulcer disease. A prospective study of 278 cases seen in an STD clinic in Paris. Sex Transm Dis 2010; 37: 153–8.
4. O'Farrell N, Lazaro N: UK National Guideline for the management of chancroid in 2014. Int J STD AIDS 2014; 25: 975–83.
5. Workowski KA, Bolan GA: Sexually transmitted diseases treatment guidelines, 2015. MMWR Recomm Rep 2015; 64: 1–137.

Co-factor *Ureaplasma urealyticum*

It is gratifying that at least also *Mycoplasma genitalium* and *Ureaplasma urealyticum* are cited as causative agents of urethritis (1). In my opinion, especially *U. ureal* is underrated as a (co-) agent of urethritis. In 40 years of urological practice, this was by far the most frequent pathogen in smears in recurrent urethritis and cystitis. Treating both infected people (according to an antibiogram) led to, in 90% of cases, an end of recurrences, and happy patients.

U. ureal and *Mycoplasma spp.* play important roles at least as cofactors, when one considers how many patients, and in particular female patients, suffer from recurrent cystitis. Studies in this area are only patchy. From my perspective and practical experience, *U. urealyticum* and *Mycoplasma spp.* play a much greater role than previously recognized.

DOI: 10.3238/arztebl.2016.0460b

REFERENCES

1. Wagenlehner FME, Brockmeyer NH, Discher T, Friese K, Wichelhaus TA: The presentation, diagnosis and treatment of sexually transmitted infections. Dtsch Arztebl Int 2016; 113: 11–22.

Reinhard Laszig
Kiel
Laszig@urologe-kiel.de

Conflict of interest statement

The author declares that no conflict of interest exists.

In Reply:

Unfortunately, space limitations made it impossible to discuss all important sexually transmitted diseases (1). These also include lymphogranuloma venereum, which is caused by *Chlamydia trachomatis* serovar CTL1–3. This disease is endemic to tropical and subtropical countries, and in Europe, it affects especially HIV-infected men who have sex with men (MSM). It is diagnosed by nucleic acid amplification techniques (NAAT), and a positive result should be followed by serovar differentiation. Patients are treated with doxycycline (100 mg orally twice daily for 21 days) or, alternatively, with azithromycin (1.5 g p. o. on days 1, 8, and 15). Azithromycin resistance has also been described. Swab controls (using NAAT) are therefore carried out at six weeks after therapy. Partner treatment should always be carried out (2).

Chancroid (or *ulcus molle*), which is rare in Europe, is caused by *Haemophilus ducreyi*. It is characterized by an ulcer that is usually painful, which provides an important differential diagnosis to the commonly painless *ulcus durum*. Diagnosis can be made only using special culture media or by employing NAAT, which however usually depends on in-house PCRs rather than routinely available equipment. In clinical practice, diagnosing a painful ulcer with regional lymphadenopathy is a diagnosis of exclusion, with a lack of proof of *T. pallidum* and/or herpes. Treatment consists of a single dose of azithromycin (1.5 g p. o.). Concomitant HIV infection frequently leads to treatment failure (2).

Caused by *Klebsiella*, *granuloma inguinale* is a rare, mildly contagious ulcer that results from a chronic granulomatous infectious disease of the genital and perigenital regions. Without treatment, it continuously propagates and may abscess. It is treated with cotrimoxazole (960 mg p. o. twice daily for 21 days) (2).

In regard to treatment of human papilloma virus diseases, it should be noted that only imiquimod, podophyllotoxin and the purified dry extract from green teas leaves with the active substance epigallocatechin gallate are mentioned in the current CDC treatment guideline from 2015 (3). In the AWMF guideline, 5-fluorouracil is also recommended for treating extensive multifocal, anal intraepithelial neoplasia, if this cannot be resolved primarily by ablative treatment (4).

The comment from Reinhard Laszig, that *Ureaplasma urealyticum* and *Mycoplasma genitalium* are also important causative pathogens in urethritis, is correct.

This is accounted for in the new guidelines (3, 5). Further, therapy proposals exist for these, as mentioned in the publication (1). DOI: 10.3238/arztebl.2016.0460c

REFERENCES

1. Wagenlehner FME, Brockmeyer NH, Discher T, Friesse K, Wichelhaus TA: The presentation, diagnosis and treatment of sexually transmitted infections. Dtsch Arztebl Int 2016; 113: 11–22.
2. Bremer V, Brockmeyer N, Coenenberg J, et al.: STI/STD Beratung, Diagnostik und Therapie. 2014, AWMF. www.awmf.org/uploads/tx_szleitlinien/059-006L_S1_STI_STD-Beratung_2015-07.pdf (last accessed on 14 March 2016).
3. Workowski KA, Bolan G: Sexually transmitted diseases treatment guidelines, 2015. MMWR Recomm Rep 2015; 64: 1–137.
4. Brockmeyer N, Degen O, Eldering G, et al.: Anale Dysplasien und Analkarzinome bei HIV-Infizierten: Prävention, Diagnostik, Therapie. 2014, AWMF. www.awmf.org/leitlinien/detail/ll/055-007.html (last accessed on 15 March 2016).
5. Bradshaw CS, Jensen JS, Tabrizi SN, et al.: Azithromycin failure in *Mycoplasma genitalium* urethritis. Emerg Infect Dis 2006; 12: 1149–52.

Prof. Dr. med. Florian M. E. Wagenlehner
Justus-Liebig-Universität Gießen
Wagenlehner@AOL.com

Conflict of interest statement

Prof. Wagenlehner has received consulting fees from Astellas, Bionorica Cubist, Galenus, Leo-Pharma, Merlion, OM-Pharma, Pierre Fabre, Perell Research, Rosen Pharma, and Zambon. He has received honoraria for speaking at continuing medical education events from Astellas, Bionorica Cubist, Galenus, Leo-Pharma, Merlion, OM-Pharma, Pierre Fabre, Rosen Pharma, and Zambon. He has also received payment for carrying out clinical trials on behalf of Astellas, Bionorica, Calixa, Cerexa, Cubist, The German Research Foundation (Deutsche Forschungsgemeinschaft), the European Association of Urology, Galenus, The Hessen State Ministry of Higher Education, Research, and the Arts, Merlion, OM-Pharma, Rosen Pharma, and Zambon.