Published in final edited form as:

J Am Chem Soc. 2016 June 22; 138(24): 7753–7762. doi:10.1021/jacs.6b04040.

Factors That Control the Reactivity of Cobalt(III)–Nitrosyl Complexes in Nitric Oxide Transfer and Dioxygenation Reactions: A Combined Experimental and Theoretical Investigation

Pankaj Kumar^{#†}, Yong-Min Lee^{#†}, Lianrui Hu[‡], Jianwei Chen[‡], Young Jun Park[†], Jiannian Yao[‡], Hui Chen^{*‡}, Kenneth D. Karlin^{*¶}, and Wonwoo Nam^{*†}

[†]Department of Chemistry and Nano Science, Ewha Womans University, Seoul 120-750, Korea

[‡]Beijing National Laboratory for Molecular Sciences, CAS Key Laboratory of Photochemistry, Institute of Chemistry, Chinese Academy of Sciences, Beijing 100190, China

[¶]Department of Chemistry, The Johns Hopkins University, Baltimore, Maryland 21218, United States

[#] These authors contributed equally to this work.

Abstract

Metal–nitrosyl complexes are key intermediates involved in many biological and physiological processes of nitric oxide (NO) activation by metalloproteins. In this study, we report the reactivities of mononuclear cobalt(III)–nitrosyl complexes bearing *N*-tetramethylated cyclam (TMC) ligands, [(14-TMC)Co^{III}(NO)]²⁺ and [(12-TMC)Co^{III}(NO)]²⁺, in NO-transfer and dioxygenation reactions. The Co(III)–nitrosyl complex bearing 14-TMC ligand, [(14-TMC)Co^{III}(NO)]²⁺, transfers the bound nitrosyl ligand to [(12-TMC)Co^{II}]²⁺ via a dissociative pathway, {[[(14-TMC)Co^{III}(NO)]²⁺ → {(14-TMC)-Co···NO}²⁺}, thus affording [(12-TMC)Co^{III}(NO)]²⁺ and [(14-TMC)Co^{II}]²⁺ as products. The dissociation of NO from the [(14-TMC)Co^{III}(NO)]²⁺ complex prior to NO-transfer is supported experimentally and theoretically. In contrast, the reverse reaction, which is the NO-transfer from [(12-TMC)Co^{III}(NO)]²⁺ to [(14-TMC)Co^{II}]²⁺, does not occur. In addition to the NO-transfer reaction, dioxygenation of [(14-TMC)Co^{III}(NO)]²⁺ by O₂ produces [(14-TMC)Co^{II}(NO₃)]⁺, which possesses an O,O-chelated nitrato ligand and where, based on an experiment using ¹⁸O-labeled O₂, two of the three O-atoms in the [(14-TMC)Co^{II}(NO₃)]⁺ product derive from O₂. The dioxygenation reaction is proposed to occur via a dissociative pathway, as proposed in the NO-transfer reaction, and via the formation of a Co(II)–peroxynitrite intermediate, based on the observation of phenol ring nitration. In contrast,

*Corresponding Authors: chen@iccas.ac.cn, karlin@jhu.edu, wwnam@ewha.ac.kr.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/jacs.6b04040.

Procedure for synthesis of **1**, spectroscopic data for **1** and **6**, reactivity data for **1**, crystallographic data for **6**, computational details, and Cartesian coordinates of species in reactions (PDF)

Crystallographic data for [(14-TMC)Co^{II}(NO₃)](BPh₄), complex **6** (CIF)

The authors declare no competing financial interest.

$[(12\text{-TMC})\text{Co}^{\text{III}}(\text{NO})]^{2+}$ does not react with O_2 . Thus, the present results demonstrate unambiguously that the NO-transfer/dioxygenation reactivity of the cobalt(III)–nitrosyl complexes bearing TMC ligands is significantly influenced by the ring size of the TMC ligands and/or the spin state of the cobalt ion.

Graphical abstract

INTRODUCTION

As a free radical nitrogen–oxide molecule, NO (nitric oxide, nitrogen monoxide) is known to play diverse roles in biological processes.¹ In natural systems, the biosynthesis of NO and its various biological and physiological reactions come about through the interaction with metalloproteins of iron and/or copper.² Thus, reactions of NO through its coordination with transition metal centers are of great interest for chemists and biochemists.³ In addition, there is a long history of the study of transition metal–NO interactions, to understand fundamental aspects of the structure, bonding, and reactivity of metal–NO species.³

There are a few reports on the intermolecular transfer of metal-coordinated ligands such as O_2 , O_2^{2-} , and NO,^{4,5} and it has been shown that NO-transfer between two metal complexes can occur, depending on the NO-to-metal binding constant, ligand geometry, and metal oxidation state. The prior literature on metal complex NO-transfer chemistry suggests at least two proposed pathways, but no unambiguous mechanistic conclusion has ever been reached. In this regard, one of the early examples involves the acid-promoted intermolecular transfer of NO from Co^{II} to Cr^{II} , involving the formation of a labile aqua complex at lower pH.^{5a} In another case, protonation of an axial ligand transforms the geometry of coordinated NO toward a more bent conformation, allowing NO-transfer.^{5b} Caulton and co-workers undertook a detailed investigation of nitrosyl transfer from $\text{Co}(\text{NO})(\text{dmgH})_2$ (dmgH = monoanion of dimethylglyoxime (dmgH_2)) to the various metal complexes of Fe, Co, Ni, and Ru.^{5c} For the first time, they suggested a μ -bridged nitrosyl formation prior to the transfer of NO. A dissociative pathway of NO-transfer was described for a $\text{Co}(\text{NO})(\text{dmgH})_2$ complex in the presence of hemoglobin (Hb), with first dissociation of NO followed by formation of HbNO .⁶ Lippard and co-workers also suggested a dissociative pathway of NO-transfer from manganese to iron bearing a tropocoronand ligand framework.⁷ Recently, intermolecular transfer of a ruthenium-bound nitrosyl to a coordinatively unsaturated iron complex was reported to occur through a heterobimetallic μ -bridged nitrosyl intermediate.⁸

Thus, on the basis of insights coming from all of these reactions, two mechanisms have been proposed (Scheme 1, M and M' are different metal ions): one is a dissociative pathway of NO-transfer (Scheme 1a),⁶ and the other involves an associative pathway (Scheme 1b).⁸

In addition to the mechanism described above, the study of the reactions of both NO and O₂ or their derivatives with transition metal complexes has attracted much attention. Since the coordination chemistry of metal–nitrosyl complexes has a long history, various reactions with metal–nitrosyl complexes have been reported.³ These species can be exceedingly stable, but many of them do in fact react with O₂, and metal nitrite or nitrate complexes are frequently obtained as the products, depending on the system at hand.^{9–11} Clarkson and Basolo described for the first time the chemistry of certain cobalt–nitrosyl complexes, proposing that a peroxyxynitrite (PN, [−]OON=O) intermediate may form and thermally decay to give a cobalt nitrite complex product.⁹ Many other examples supported this work.¹⁰ In the case of dinitrosyl-iron species (DNICs), the metal–nitrosyl complex reacts with O₂ to form a metal nitrate complex as the final product, putatively via the isomerization of a PN intermediate.¹¹ It should be noted that, in aqueous chemistry, (i) peroxyxynitrite is the diffusion-controlled reaction product of NO plus superoxide anion (O₂^{•−}),¹² and (ii) the primary decomposition product of peroxyxynitrous acid is nitric acid derived from a very fast isomerization reaction.¹³

In what is turning out to be chemistry that is complementary, the study of the reactivity of metal–O₂ species with NO has been of great interest,^{14,15} in part due to its relevance in understanding the chemistry involved in the enzyme NO dioxygenase (NOD). NOD activity aids the maintenance of appropriate cellular NO balance by converting excesses to give the biologically benign nitrate ion.¹³ Thus, oxyhemoglobin, oxymyoglobin, and bacterial heme–O₂ adducts (“oxy-ferrous” or formally Fe^{III}–superoxide) are NODs. These reactions are thought to proceed through Fe^{III}–PN intermediates, prior to nitrate formation;^{14,15} nitrogen dioxide (NO₂) may otherwise be released.¹⁶ Thus, metal–PN generation may occur via metal–O₂ plus NO or metal–NO plus O₂ reactions, as shown recently in coordination complexes of chromium,^{17a,b} iron,^{17c} copper,¹⁸ hemes,¹⁹ and cobalt–porphyrins.²⁰

Nam and co-workers have described an extensive series of first-row transition metal–superoxo, –peroxo, and –oxo species bearing *N*-tetramethylated cyclam (TMC) ligands,²¹ in which structure, bonding, and reactivity of the metal–oxygen complexes are greatly influenced by the ring size of the macrocycles. In relation to the NOD reactivity, TMC chromium–superoxo, chromium–peroxo, and iron–peroxo complexes were treated with NO or nitrosonium (NO⁺) ion, and in all cases metal–peroxyxynitrite intermediates were implicated as transient intermediates, leading to the formation of nitrite or nitrate metal complexes as final products.¹⁷

We report herein for the first time that the NO-transfer and dioxygen reactivity of Co^{III}–nitrosyl complexes bearing TMC ligands are tuned by the ring size of the supporting ligands and/or the spin state of Co^{II} ion. More specifically, a new Co^{III}–nitrosyl complex bearing a 14-TMC ligand, [(14-TMC)-Co^{III}(NO)]²⁺ (**1**) (14-TMC = 1,4,8,11-tetramethyl-1,4,8,11-tetraazacyclotetradecane), is generated and characterized spectroscopically and structurally. Its reactivities are investigated in the NO-transfer reaction from **1** to [(12-TMC)-Co^{II}]²⁺ (12-

TMC = 1,4,7,10-tetramethyl-1,4,7,10-tetraazacyclododecane) and in the reaction with O₂, and are compared with those of the previously reported [(12-TMC)Co^{III}(NO)]²⁺ (**2**) complex.²² We propose the mechanism of the NO-transfer and dioxygen reactions as occurring via a dissociative pathway. Density functional theory (DFT) calculations further clarify the detailed mechanism of NO-transfer from **1** to [(12-TMC)-Co^{II}]²⁺, supporting the experimental implication of a NO dissociation pathway.

RESULTS AND DISCUSSION

Generation and Characterization of [(14-TMC)-Co^{III}(NO)]²⁺ (**1**)

The Co^{III}-nitrosyl complex bearing a 14-TMC ligand, [(14-TMC)Co^{III}(NO)]²⁺ (**1**), was synthesized by adding NO(g) to a solution of [(14-TMC)Co^{II}]²⁺. As shown in Figure 1a, addition of NO (1 equiv) to the solution of [(14-TMC)Co^{II}]²⁺ under an Ar atmosphere in CH₃CN at -40 °C resulted in the formation of a new species with absorption bands at 395 nm ($\epsilon = 690 \text{ M}^{-1} \text{ cm}^{-1}$) and 555 nm ($\epsilon = 220 \text{ M}^{-1} \text{ cm}^{-1}$), which are similar to those of [(12-TMC)-Co^{III}(NO)]²⁺ (**2**; 368 and 538 nm) and [(13-TMC)-Co^{III}(NO)]²⁺ (375 and 550 nm).²² A solution IR spectrum of **1** exhibited a characteristic peak at 1715 cm⁻¹ (Figure 1a, inset), which is almost identical to NO stretching bands of **2** (1712 cm⁻¹) and [(13-TMC)Co^{III}(NO)]²⁺ (1716 cm⁻¹).²² An electron paramagnetic resonance (EPR) spectrum of **1** is silent, suggesting the trivalency of the cobalt center in **1** (Figure 1b). In addition to EPR, a ¹H NMR spectrum of **1** supports that the oxidation state of cobalt center in **1** is 3+ with a spin state of *S* = 0 (Figure S1 in Supporting Information (SI)). The results described above demonstrate that the formally Co^{III}(NO⁻) or {Co(NO)}⁸ was formed in the reaction of [(14-TMC)Co^{II}]²⁺ with NO.

In addition to the above spectroscopic characterization, the electronic structure of the Co(III)-nitrosyl complexes was investigated by carrying out DFT calculations to understand NO binding to the cobalt center, revealing that both **1** and **2** have an open-shell singlet ground state (Figure 2). The DFT-optimized geometry of **2** is in a good agreement with the previously reported X-ray crystal structure of **2**-(ClO₄)₂ (Figure 2c).²² The triplet states of **1** and **2** are 3.5 and 0.3 kcal/mol higher than the corresponding singlet states, respectively. Moreover, the nearly linear Co(1)-N(5)-O(1) geometry in the DFT-optimized structure of the triplet state of **2** suggests that the triplet state is not likely to be the ground state. We also calculated the quintet states of **1** and **2**, which are higher in energy than the corresponding ground singlet states of **1** and **2** by 6.9 and 14.0 kcal/mol, respectively. The calculated Co(1)-N(5)-O(1) angle of about 150° in the quintet state of **2** is also inconsistent with the X-ray crystal structure of **2**-(ClO₄)₂. The open-shell character of the singlet states of **1** and **2** is depicted by the two singly occupied orbitals as shown in Figure 2, in which the two unpaired electrons couple antiferromagnetically. Thus, all spectroscopic data (Figure 1) with the DFT-optimized structure (Figure 2) support that **1** is a cobalt(III)-nitrosyl complex binding the NO ligand in an end-on fashion.

NO-Transfer Reactions

The reactivity of the Co^{III}-nitrosyl complexes was investigated in NO-transfer reactions under an Ar atmosphere in CH₃CN at -40 °C. Upon addition of one equiv of [(12-

TMC)Co^{II}]2⁺ to a CH₃CN solution of [(14-TMC)Co^{III}(NO)]2⁺ (**1**, black line in Figure 3a), a new UV-vis spectrum was obtained immediately (blue line in Figure 3a), and following this, the blue line spectrum slowly changed to the red line spectrum with an absorption band at 368 nm (Figure 3a, inset, red circles), which corresponds to the characteristic absorption band of [(12-TMC)Co^{III}(NO)]2⁺ (**2**) (Scheme 2, reaction *a*).²² The spectrum with blue line was confirmed to be the sum of UV-vis spectra of **1** and [(12-TMC)Co^{II}]2⁺ (Figure S2a in SI). It should be noted that, in the absence of [(12-TMC)Co^{II}]2⁺, no spectral change was observed (Figure 3a, inset, black circles), indicating that **1** is highly stable under an Ar atmosphere at -40 °C.²³ The final spectrum (red line in Figure 3a), which was confirmed to be the sum of those spectra of **2** and [(14-TMC)Co^{II}]2⁺ (Figure S2b in SI), indicating that **2** was generated quantitatively (>95%) based on the ϵ (M⁻¹ cm⁻¹) value at 368 nm due to **2**, since the [(14-TMC)Co^{II}]2⁺ product has almost no absorption at 368 nm as shown in Figure 1a (pink line). This NO-transfer from **1** to [(12-TMC)Co^{II}]2⁺ for the formation of **2** was confirmed by addition of [(12-TMC)Co^{II}]2⁺ to a solution of **1** in increments of 0.2 equiv in CH₃CN at -40 °C (Figure 4). The spectroscopic titration at 368 nm (Figure 4, inset) for the formation of **2** indicates that the reaction stoichiometry of **1** and [(12-TMC)Co^{II}]2⁺ is 1:1, and 1 equiv of [(12-TMC)-Co^{II}]2⁺ is required for the full conversion of **1** to **2**. In contrast to the NO-transfer from **1** to [(12-TMC)Co^{II}]2⁺, no NO-transfer from **2** to [(14-TMC)Co^{II}]2⁺ was observed (Figure S3 in SI), indicating that the binding affinity of NO to [(12-TMC)Co^{II}]2⁺ is higher than that to [(14-TMC)Co^{II}]2⁺. In addition, the formation rates of **1** and **2** in the reactions of [(14-TMC)Co^{II}]2⁺ and [(12-TMC)Co^{II}]2⁺ with NO, respectively, were determined to be the second-order rate constants of 3.3(3) and 12(1) M⁻¹ s⁻¹, respectively (Figure S4 in SI), indicating that the formation rate of **2** is ~4 times faster than that of **1**.

The intermolecular NO-transfer from **1** to [(12-TMC)-Co^{II}]2⁺ was further confirmed by examination of the associated solution IR, ESI-MS, and EPR spectroscopies. According to solution IR measurements, a peak at 1715 cm⁻¹ for the NO stretching frequency of **1** (Figure 1a, inset) was changed to 1712 cm⁻¹ due to formation of **2** (Figure 5a), which is identical to the NO stretching frequency of **2**,²² upon addition of [(12-TMC)Co^{II}]2⁺ to a solution of **1**. An ESI-MS of the reaction solution exhibited prominent mass peaks at *m/z* = 416.0, 414.2, and 178.1 (Figure 3b), whose mass and isotopic distribution patterns correspond to [(12-TMC)Co^{III}(NO)(ClO₄)]⁺ (calcd *m/z* = 416.1), [(14-TMC)Co^{II}(ClO₄)]⁺ (calcd *m/z* = 414.1), and [(14-TMC)Co^{II}(CH₃CN)]2⁺ (calcd *m/z* = 178.1), respectively, indicating that **2** and [(14-TMC)Co^{II}]2⁺ were formed in this reaction. An EPR spectrum of the final reaction solution revealed that a high-spin cobalt(II) complex formed, which is identical to that of [(14-TMC)Co^{II}]2⁺ (see Figure 1b). Based on the above spectroscopic characterization of the reaction solution, we further conclude that **2** and [(14-TMC)Co^{II}]2⁺ were produced by reacting **1** with [(12-TMC)Co^{II}]2⁺ (Scheme 2, reaction *a*).

Mechanistic Insight into the NO-Transfer Reaction

We then investigated the concentration effect of [(12-TMC)Co^{II}]2⁺ on the NO-transfer reaction of **1** to [(12-TMC)Co^{II}]2⁺ and found that the NO-transfer reaction was independent of the [(12-TMC)Co^{II}]2⁺ concentration (Figure S5 in SI). As shown in Figure 3a, the reaction of **1** with [(12-TMC)Co^{II}]2⁺ did not afford the formation of a new associated

intermediate, indicating that this reaction does not occur via an associative pathway (vide supra). In addition, there is an equilibrium between **1** and a cage molecule, $\{(14\text{-TMC})\text{Co}\cdots\text{NO}\}^{2+}$, prior to the reaction with $[(12\text{-TMC})\text{Co}^{\text{II}}]^{2+}$, as shown in Scheme 3. Since the first step of the reaction (Scheme 3, pathway *a*) is the rate-determining step, the overall rate constant (k_{obs}) of $8.2(7) \times 10^{-4} \text{ s}^{-1}$ (Figure 3a, inset) is equal to the rate constant of the dissociation to the cage molecule (k_1). The second-order rate constant of $3.3(3) \text{ M}^{-1} \text{ s}^{-1}$ determined in the reaction of $[(14\text{-TMC})\text{Co}^{\text{II}}]^{2+}$ with NO (Figure S4 in SI) indicates that the rate of re-formation to **1** (k_{-1}) might be at least more than 10^3 -fold faster than that of dissociation to the cage molecule (k_1). In addition, the rate of the formation of **2** (k_2 ; Scheme 3, pathway *b*) might be faster than that of the re-formation of **1** (k_{-1}), because the NO binding affinity of $[(12\text{-TMC})\text{Co}^{\text{II}}]^{2+}$ is greater than that of $[(14\text{-TMC})\text{Co}^{\text{II}}]^{2+}$. Therefore, as shown in Scheme 3, we propose that the NO-transfer reaction of **1** with $[(12\text{-TMC})\text{Co}^{\text{II}}]^{2+}$ occurs via a dissociative pathway^{6,7} through a putative cage molecule, $\{(12\text{-TMC})\text{Co}^{\text{II}}\cdots\text{NO}\}^{2+}$, and the order of rates for each step is $k_2 > k_{-1} \gg k_1$.

DFT Calculations for Mechanistic Insight into the NO-Transfer Reaction of **1** with $[(12\text{-TMC})\text{Co}^{\text{II}}]^{2+}$

DFT calculations were conducted to investigate why a dissociative cage molecule mechanism is preferred to an associative mechanism in the NO-transfer from **1** to $[(12\text{-TMC})\text{Co}^{\text{II}}]^{2+}$. First, for the dissociative pathway, the calculated NO dissociation profile starting from the ground singlet state of **1** is depicted in Figure 6. We can see that the barrier of NO dissociation is only 3.2 kcal/mol. This small barrier is associated with a large stretching of Co-NO bond in a NO-dissociative transition state (TS) by about 1.4 Å from **1**, as shown in Figure 7. After NO dissociation from **1**, the NO coordination to $[(12\text{-TMC})\text{Co}^{\text{II}}]^{2+}$ is found to be a barrierless process.

In comparison, the NO-transfer profile from **1** to $[(12\text{-TMC})\text{Co}^{\text{II}}]^{2+}$ for the associative pathway is shown in Figure 8. Here, we employed a model of $\{[(14\text{-TMC})\text{Co}^{\text{III}}\text{-NO-Co}^{\text{II}}(12\text{-TMC})](\text{ClO}_4)_3\}^+$. In the absence of three perchlorate counterions, the $[(14\text{-TMC})\text{Co}^{\text{III}}\text{-NO-Co}^{\text{II}}(12\text{-TMC})]^{4+}$ is not stable, and the $[(12\text{-TMC})\text{Co}^{\text{II}}]^{2+}$ fragment dissociates from the $[(14\text{-TMC})\text{Co}^{\text{III}}(\text{NO})]^{2+}$ fragment, possibly due to the Coulombic repulsion between the two +2 charged Co fragments. As shown in Figure 8, the associative mechanism is initiated from the μ -NO-bridged intermediate **3** by NO-transfer from the $[(14\text{-TMC})\text{Co}^{\text{II}}]^{2+}$ fragment to the $[(12\text{-TMC})\text{-Co}^{\text{II}}]^{2+}$ fragment, resulting in intermediate **4** with an NO coordinated to the $[(12\text{-TMC})\text{Co}^{\text{II}}]^{2+}$ fragment by its O-atom. Then, O- to N-atom coordination isomerization occurs, leading to the final NO-transfer product **5**. The highest-lying TS (**TS**₄₋₅) along the reaction profile is 9.3 kcal/mol higher than **3**, which is significantly higher than the 3.2 kcal/mol barrier in the dissociative pathway, indicating that the latter is actually the more preferred NO-transfer pathway from **1** to $[(12\text{-TMC})\text{-Co}^{\text{II}}]^{2+}$.

Considering the experimental results observed in solution IR and EPR experiments and taking account of the UV-vis kinetic measurements as well as the computational results, the reaction for NO-transfer is likely to occur via a dissociative pathway through a cage molecule prior to NO dissociation from cobalt center. Therefore, it is highly logical to assume that NO-transfer between **1** and $[(12\text{-TMC})\text{Co}^{\text{II}}]^{2+}$ takes place through NO

dissociation pathway as shown in Scheme 3. The affinity of cobalt ion toward NO is the key factor and controls the mode of NO-transfer mechanism, which can be explained on the basis of the TMC ring size and the spin state of Co^{II} ion. To understand the TMC ring size effect, we compare the rate of the formation of **1** and **2** (vide supra), suggesting the formation rate of **2** is ~4 time faster than that of **1** (Figure S4 in SI). This explains the high affinity of **2** toward NO and suggests a higher stability of **2** compared to **1**. Our computational results for **1** and **2** also directly showed their relative stabilities, showing that **2** is more stable than **1** by 3.6 kcal/mol in Gibbs free energy. Therefore, this difference of the stability of **1** and **2** is the thermodynamic driving force for the NO-transfer reaction from **1** bearing a larger TMC ligand to $[(12\text{-TMC})\text{Co}^{\text{II}}]^{2+}$ bearing a smaller TMC ligand.

The spin state of the cobalt ions in $[(12\text{-TMC})\text{Co}^{\text{II}}]^{2+}$ and $[(14\text{-TMC})\text{Co}^{\text{II}}]^{2+}$ was also determined. An X-band EPR spectrum of $[(12\text{-TMC})\text{Co}^{\text{II}}]^{2+}$ shows signals at $g_1 = 2.34$, $g_2 = 2.31$, and $g_3 = 2.05$, which are very typical for low spin ($S = 1/2$) Co^{II} species (Figure 5b, black line), while that of $[(14\text{-TMC})\text{Co}^{\text{II}}]^{2+}$ exhibits signals at $g_1 = 7.4$, $g_2 = 2.17$, and $g_3 = 1.50$, which are the characteristic values of a high-spin ($S = 3/2$) Co^{II} species (Figure 1b, pink line). This ground spin state difference between $[(12\text{-TMC})\text{Co}^{\text{II}}]^{2+}$ and $[(14\text{-TMC})\text{Co}^{\text{II}}]^{2+}$ was also sensitively monitored in our DFT calculations (Figure S6 in SI). The spin state of the cobalt centers influences the reactivity toward the NO coordination, indicating that $[(12\text{-TMC})\text{Co}^{\text{II}}]^{2+}$ (low-spin $S = 1/2$) has very high affinity for NO, compared to the $[(14\text{-TMC})\text{Co}^{\text{II}}]^{2+}$ species (high-spin $S = 3/2$). Consequently, as the ring size of the TMC ligand increases, the affinity of the cobalt center toward NO coordination decreases, and the reactivity of Co^{III} -nitrosyl increases, which hence facilitates the NO-transfer from **1** to $[(12\text{-TMC})\text{Co}^{\text{II}}]^{2+}$.

Dioxygenation Reactivity of Co^{III} -Nitrosyl Complex, **1**

In addition to the NO-transfer reactions, the O_2 reactivity of $[14\text{-TMC})\text{Co}^{\text{III}}(\text{NO})]^{2+}$ (**1**) was investigated. Upon addition of an excess of O_2 to a CH_3CN solution of **1**, the absorption band at 395 nm due to **1** decayed with first-order kinetics (Figure 9), indicating that **1** reacted with O_2 , whereas **2** did not show any reactivity toward O_2 as reported previously.²² The first-order rate constant was determined to be $7.6(7) \times 10^{-4} \text{ s}^{-1}$ (Figure S7 in SI), which is quite similar to that obtained in the NO-transfer reaction from **1** to $[(12\text{-TMC})\text{Co}^{\text{II}}]^{2+}$ (vide supra). We also found that the reaction rate of **1** with O_2 was independent of the concentration of O_2 (Figure S7 in SI), indicating that the mechanism of the reaction of **1** with O_2 is the same as that of the NO-transfer reaction from **1** to $[(12\text{-TMC})\text{Co}^{\text{II}}]^{2+}$ (vide supra). There is an equilibrium between **1** and a cage molecule, $\{(14\text{-TMC})\text{Co} \cdots \text{NO}\}^{2+}$, prior to the reaction with O_2 , as shown in Scheme 4. Since the first step of the reaction (Scheme 4, step *a*) is the rate-determining step, the overall rate constant (k_{obs}) of $7.6(7) \times 10^{-4} \text{ s}^{-1}$ is equal to the rate constant of the dissociation to the cage molecule because the rates of re-formation of **1** and reaction with O_2 are much faster than that of the dissociation to the cage molecule.²⁴

The products obtained in the reaction of **1** with O_2 were analyzed by solution IR, ESI-MS, and EPR spectroscopies. Upon addition of O_2 to a solution of **1**, an IR peak at 1715 cm^{-1} for the NO stretching vibration of **1** disappeared, and concomitantly a new IR peak at 1302

cm⁻¹ due to a nitrate NO₃⁻ stretching vibration²⁵ was formed (Figure 9b). ESI-MS spectrum of the reaction solution exhibited prominent mass peaks at $m/z = 377.1$ and 178.1 (Figure 9a, inset), whose mass and isotopic distribution patterns correspond to [(14-TMC)-Co^{II}(NO₃)]⁺ (**6**, calcd $m/z = 377.1$) and [(14-TMC)-Co^{II}(CH₃CN)]²⁺ (calcd $m/z = 178.1$). When ¹⁶O₂ was replaced by isotopically labeled ¹⁸O₂, the mass peak at $m/z = 377.1$ was shifted to $m/z = 381.1$. (blue and red lines in Figure 9a, inset). The observed shift of four mass units indicates that two of the three O atoms in the NO₃⁻ ligand of **6** are derived from O₂. An EPR spectrum of the complete reaction solution exhibited signals at $g_1 = 7.0$, $g_2 = 2.09$, and $g_3 = 1.52$, indicating that a high-spin cobalt(II) complex was the final product formed (Figure S8 in SI).²⁶

Finally, **6** was structurally characterized using X-ray crystallography. Single crystals of **6** were grown from the solution obtained in the reaction of **1** with O₂. The X-ray crystal structure of **6** revealed that the nitrate anion coordinates to Co center in a bidentate fashion to produce a product with pseudo-octahedral geometry (Figure 10). Co–O distances of 2.234 and 2.225 Å indicate that two O atoms of the nitrate anion coordinate quite symmetrically to the Co center. The crystallographic data and selected bond distances and angles for **6** are listed in Tables S1–S3 in the SI.

According to the above results, we propose a reaction mechanism as shown in Scheme 4. The first step should be the dissociation of NO ligand from **1** to form a cage molecule, {(14-TMC)Co···NO}²⁺, indicating that the Co^{III} center was reduced to Co^{II}. Then, this cage molecule, {(14-TMC)Co···NO}²⁺, reacts with O₂ to form a putative Co^{II} peroxynitrite (PN) intermediate, Co^{II}(OON=O⁻) (Scheme 4, step *b*). Subsequently, isomerization of Co^{II}–PN would generate the Co^{II}–NO₃ complex (**6**) (Scheme 4, step *c*),²⁷ as reported previously in metal-bound PN intermediates.^{14,15,28} In the dioxygenation reaction of **1**, we observed the formation of nitrate anion, which is the product formed by the isomerization of PN.^{14,15} Since metal-PNs are known to be extremely unstable,¹⁹ there are only few reports about the metal-bound PN complexes.^{29,30} However, PN is often proven to be present by its ability to effect phenol ring nitration.³¹ Thus, 2,4-di-*tert*-butylphenol (DTBP) was added to a solution of **1** prior to addition of dioxygen to support our proposed mechanism (Scheme 5). We found that 2,4-di-*tert*-butyl-6-nitrophenol (nitro-DTBP) and dimerized DTBP (2,2'-dihydroxy-3,3', 5,5'-tetra-*tert*-butylphenol) were produced with the yields of 60(5)% and 10(4)%, respectively (see Experimental Section). Thus, the experimental results, such as the formation of nitro-DTBP with a good yield, strongly support the proposed reaction mechanism, which is the formation of a Co-PN intermediate in the reaction of **1** with O₂.

CONCLUSION

We have demonstrated for the first time that the NO-transfer and dioxygen reactivity of Co^{III}–nitrosyl complexes bearing *N*-tetramethylated cyclam ligands can be tuned by varying the ring size of the supporting ligands and/or the spin state of Co^{II} ion. In the NO-transfer reaction and the reaction with O₂, the reactivity of [(14-TMC)Co^{III}(NO)]²⁺ (**1**) was compared with that of a smaller ring-size congener, [(12-TMC)Co^{III}(NO)]²⁺ (**2**). We found that the NO moiety in **1** was transferred to [(12-TMC)Co^{II}]²⁺ to form **2**, whereas no NO-transfer reaction occurred between **2** and [(14-TMC)Co^{II}]²⁺; these results were interpreted

as being due to the higher NO binding affinity of $[(12\text{-TMC})\text{Co}^{\text{II}}]^{2+}$ compared to that of $[(14\text{-TMC})\text{Co}^{\text{II}}]^{2+}$. The NO-transfer reaction was independent of the $[(12\text{-TMC})\text{Co}^{\text{II}}]^{2+}$ concentration, leading us to suggest that the mechanism of the NO-transfer from **1** to $[(12\text{-TMC})\text{Co}^{\text{II}}]^{2+}$ occurs via dissociation of the NO ligand from **1** to form a cage molecule, $\{(14\text{-TMC})\text{Co}\cdots\text{NO}\}^{2+}$, prior to its reaction with $[(12\text{-TMC})\text{Co}^{\text{II}}]^{2+}$ to produce **2**. DFT calculations supported the experimental observation for the NO dissociation pathway. In addition to the NO-transfer reaction, the reactivity of **1** toward O_2 was investigated. A cobalt–nitrate complex, $[(14\text{-TMC})\text{Co}^{\text{II}}(\text{NO}_3)]^+$, which was structurally characterized by X-ray crystallography, was produced as a sole product in the reaction of **1** and O_2 , while **2** was inert toward O_2 .²² The reaction rate of **1** with O_2 was independent of the O_2 concentration, indicating that the reaction mechanism of **1** and O_2 is similar to that of the NO-transfer reaction from **1** to $[(12\text{-TMC})\text{Co}^{\text{II}}]^{2+}$, which occurs via a dissociation of NO ligand from **1** to form a cage molecule, $\{(14\text{-TMC})\text{Co}\cdots\text{NO}\}^{2+}$, prior to the reaction with O_2 . The reaction of the cage molecule, $\{(14\text{-TMC})\text{Co}\cdots\text{NO}\}^{2+}$, with O_2 produces a putative peroxynitrite (PN) intermediate, $\text{Co}^{\text{II}}(\text{OON}=\text{O})^-$, which is converted to the final $\text{Co}^{\text{II}}\text{-NO}_3$ product. As a conclusion, we have demonstrated unambiguously that the reactivities of the cobalt(III)–nitrosyl complexes bearing TMC ligands in the NO-transfer and dioxygenation reactions are significantly influenced by the spin state of the cobalt(II) center, caused by the ring size of the TMC ligands. Such explanations have also been provided and published for other first-row transition metal ions, where the chemistries are compared using metal complexes bearing different ring-size TMC ligands.³²

EXPERIMENTAL SECTION

Materials

All chemicals obtained from Aldrich Chemical Co. were of the best available purity and used without further purification unless otherwise indicated. Solvents were dried according to published procedures and distilled under Ar prior to use.³³ The 14-TMC ligand and 1,4,7,10-tetraazacyclododecane compound were purchased from Aldrich Chemical Co. 12-TMC was prepared by reacting excess amounts of formaldehyde and formic acid with 1,4,7,10-tetraazacyclododecane.³⁴ $[(14\text{-TMC})\text{Co}^{\text{II}}]^{2+}$ and $[(12\text{-TMC})\text{Co}^{\text{II}}]^{2+}$ complexes were synthesized and isolated according to the literature methods.^{21b} Co(III)–nitrosyl complexes (**1** and **2**) were synthesized by reported procedures.²²

Instrumentation

UV–vis spectra were recorded on a Hewlett-Packard 8453 diode array spectrometer equipped with a UNISOKU Scientific Instruments for low-temperature experiments. Coldspray ionization time-of-flight mass (CSI-TOF MS) spectral data were collected on a JMS-T100CS (JEOL) mass spectrometer equipped with the CSI source. Typical measurement conditions are as follows: needle voltage, 2.2 kV; orifice 1 current, 50–500 nA; orifice 1 voltage, 0–20 V; ring lens voltage, 10 V; ion source temperature, 5 °C; spray temperature, –40 °C. Electrospray ionization mass spectra (ESI-MS) were collected on a Thermo Finnigan (San Jose, CA, USA) LCQTM Advantage MAX quadrupole ion trap instrument, by infusing samples directly into the source using a manual method. The spray voltage was set at 4.2 kV and the capillary temperature at 80 °C. EPR spectra were recorded

at 5 K using an X-band Bruker EMX-plus spectrometer equipped with a dual-mode cavity (ER 4116DM). Low temperatures were achieved and controlled using an Oxford Instruments ESR900 liquid He quartz cryostat with an Oxford Instruments ITC503 temperature and gas flow controller. Low-temperature solution IR spectra were recorded on Remspec 626 FT-IR spectrometer. ^1H NMR spectra were measured with a Bruker model digital AVANCE III 400 FT-NMR spectrometer. Crystallographic data collections were carried out on a Bruker SMART AXS diffractometer equipped with a monochromator in the Mo $K\alpha$ ($\lambda = 0.71073 \text{ \AA}$) incident beam.

Purification of Nitric Oxide

Nitric oxide gas ($\text{NO}_{(\text{g})}$) was purchased from Dong-A Specialty Gases in Korea and purified as follows:²² $\text{NO}_{(\text{g})}$ was first passed through two columns filled with NaOH beads and molecular sieves to remove higher nitrogen oxide and moisture impurities and collected in a frozen form in a first trap cooled at 77 K with liquid N_2 . Further purification was done by distillation of frozen $\text{NO}_{(\text{g})}$ (as crystalline N_2O_2) by warming at -80°C (acetone/dry ice mixture, -80°C), collecting it in a second trap cooled at 77 K with liquid N_2 . This second trap was again warmed to -80°C , and the highly purified $\text{NO}_{(\text{g})}$ was collected in another vacuumed Schlenk flask fitted with a rubber septum (free from oxygen, after several cycles of vacuum and Ar purging). The $\text{NO}_{(\text{g})}$ should be at high pressure in the Schlenk flask (>1 atm; the septum bulges outward due to high pressures inside the Schlenk flask). Then, $\text{NO}_{(\text{g})}$ was purged to the distilled and degassed CH_3CN (15 mL) in a Schlenk flask, which was already connected with an oil bubbler. $\text{NO}_{(\text{g})}$ purging with vigorous stirring under 1 atm for 20 min is required to make NO-saturated CH_3CN solution. The concentration of NO-saturated in CH_3CN solution at 20°C was approximated to be 14 mM based on published values for $\text{NO}_{(\text{g})}$ solubility.³⁵ According to the literature, $\text{NO}_{(\text{g})}$ solubility in diethyl ether is 27 mM, while for most other non-ethereal organic solvents the values are in the range of 11–18 mM.³⁵

X-ray Crystallography

The single crystals of **6**-(BPh_4), which were formed by anion exchange with BPh_4^- from **6**-(ClO_4) complex, were obtained by slow diffusion of Et_2O into the CH_3CN solution at -30°C . We tried to grow the suitable crystals of **6**-(ClO_4), but we failed to grow them because of high degree of disorder in perchlorate anion. Thus, to grow suitable crystals, ClO_4 anion was exchanged to BPh_4 anion. A single crystal of **6**-(BPh_4) was picked from solution by a nylon loop (Hampton Research Co.) and mounted on a goniometer head in a $\text{N}_2(\text{g})$ cryostream. Data collections were carried out on a Bruker SMART AXS diffractometer equipped with a monochromator in the Mo $K\alpha$ ($\lambda = 0.71073 \text{ \AA}$) incident beam. The CCD data for **6**-(BPh_4) were integrated and scaled using the Bruker-S SAINT software package, and the structure was solved and refined using SHELXTL V 6.12.25.³⁶ The crystallographic data and selected bond distances and angles for **6**-(BPh_4) are listed in Tables S1–S3 in the SI. CCDC-1449724 for **6**-(BPh_4) contains the supplementary crystallographic data for this paper. These data can be obtained free of charge via www.ccdc.cam.ac.uk/data_request/cif (or from the Cambridge Crystallographic Data Centre, 12 Union Rd., Cambridge CB2 1EZ, UK; fax (+44) 1223-336-033, or deposit@ccdc.cam.ac.uk).

Reactivity Studies

All the reactions were run in a 1 cm UV cuvette, by monitoring the UV-vis spectral changes of the reaction solutions, and the rate constants were determined by fitting the changes in absorbance at 393 and 368 nm for **1** and **2**, respectively. Reactions were run at least in triplicate, and the data reported represent the average of these reactions. Complex **1** was prepared in situ by reaction of $[(14\text{-TMC})\text{Co}^{\text{II}}]^{2+}$ with an equal amount of $\text{NO}_{(\text{g})}$ under an Ar atmosphere in CH_3CN at -40°C . **1** (0.50 mM) was then used in different reactivity studies, such as NO-transfer and NO dioxygenation reactions under above reaction conditions. The NO-transfer reactions were carried out by adding different amounts of $[(12\text{-TMC})\text{Co}^{\text{II}}]^{2+}$ (0.50, 2.5, and 5.0 mM) to a solution of **1** (0.50 mM), which was in situ prepared under an Ar atmosphere in CH_3CN at -40°C . UV-vis spectral changes were monitored at 368 nm due to **2**, and rate constants were determined by fitting the changes in absorbance at 368 nm. In NO-transfer reactions, the yields were measured by comparing molar extinction coefficient (ϵ) of **2** at 368 nm with authentic sample.²² The reactions of **1** with O_2 were carried out by purging excess O_2 (0.10 M) to a CH_3CN solution of **1** (0.50 mM) at -40°C . The reactions were followed by monitoring the change in absorbance at 395 nm.

Reaction of **1** with 2,4-di-*tert*-butylphenol (DTBP) were carried out in CH_3CN at -40°C . **1** (2.0 mM, 5.0 mL) was prepared under an Ar atmosphere in CH_3CN at -40°C in a 30 mL Schlenk flask fitted with a septum. Upon addition of DTBP to a solution of **1**, no reaction occurred. However, upon addition of excess oxygen to a solution containing **1** and DTBP in CH_3CN at -40°C , the absorption band at 395 nm due to **1** disappeared. The reaction mixture was stirred for 3 h at -40°C before warming to room temperature. The products were analyzed by injecting reaction mixture directly into a GC and were identified by comparison with authentic samples. The product yields were determined to be 60(5)% for 2,4-di-*tert*-butyl-6-nitrophenol and 10(4)% for 2,2'-dihydroxy-3,3',5,5'-tetra-*tert*-butylbiphenol by comparison against standard curves prepared with authentic samples and using decane as an internal standard.

EPR Spectroscopy

EPR spectra were recorded for different reactions, such as generation of **1**, NO-transfer reaction from **1** to $[(12\text{-TMC})\text{Co}^{\text{II}}]^{2+}$, and dioxygenation reaction of **1**. All the reactions were performed under an Ar atmosphere in CH_3CN at -40°C , and then EPR spectra were recorded at 5 K. In the case of NO-transfer reaction, EPR spectral changes as a function of time (0–170 min) have been recorded at 5K for the reaction of **1** (2.0 mM) with equivalent amount of $[(12\text{-TMC})\text{Co}^{\text{II}}]^{2+}$ in CH_3CN at -40°C . Time-dependent EPR measurements revealed the clear conversion of **1** to **2**.

Solution IR Spectroscopy

Solution IR spectra for samples were recorded with a sophisticated setup, attached to an IR instrument (Remspec model 626) equipped with a very sensitive probe under an Ar atmosphere in CH_3CN at -40°C . The formation of Co(III)–nitrosyl complexes (**1** and **2**), NO-transfer reactions, and dioxygenation reactivity of **1** were followed by monitoring the

characteristic nitrosyl stretching vibration bands. **1** (3.0 mM, 2.0 mL) was in situ prepared under the above experimental conditions/setup and showed the formation of a peak at 1715 cm^{-1} , which corresponds to nitrosyl stretching in solution medium. Solution IR spectral changes for the reaction of **1** (3.0 mM, 2.0 mL) with equal amount of $[(12\text{-TMC})\text{Co}^{\text{II}}]^{2+}$ show the shifting of characteristic nitrosyl stretching of **1** (1715 cm^{-1}) to 1712 cm^{-1} (corresponding to **2**). In addition, solution IR spectral changes for reaction of **1** (3.0 mM, 2.0 mL) with dioxygen were also recorded in CH_3CN at -40°C , showing that the decay of the peak at 1715 cm^{-1} due to **1**, and concomitantly the formation of a new peak at 1302 cm^{-1} , which corresponds to a NO_3^- stretching vibration,²⁵ was observed.

Synthesis of $[(14\text{-TMC})\text{Co}^{\text{II}}(\text{NO}_3)](\text{BPh}_4)$

Upon addition of an aqueous solution of sodium tetraphenylborate (52 mg, 0.15 mmol) to a CH_3OH solution of $[(14\text{-TMC})\text{Co}^{\text{II}}]^{2+}$ (56 mg, 0.1 mmol) at room temperature, a very light-colored precipitate of $[(14\text{-TMC})\text{Co}^{\text{II}}](\text{BPh}_4)_2$ was produced within 10 min. The precipitate was filtered, washed with cold methanol, and then dried over CaCl_2 under vacuum. $[(14\text{-TMC})\text{Co}^{\text{III}}(\text{NO})]^+$ was also generated by reacting $[(14\text{-TMC})\text{-Co}^{\text{II}}](\text{BPh}_4)_2$ (3.0 mM, 10 mL) with an equal amount of NO, and then $[(14\text{-TMC})\text{Co}^{\text{II}}(\text{NO}_3)](\text{BPh}_4)$ was obtained by reacting with excess dioxygen in CH_3CN at -40°C over 3 h. The reaction mixture containing $[(14\text{-TMC})\text{Co}^{\text{II}}(\text{NO}_3)](\text{BPh}_4)$ was layered with Ar-saturated Et_2O for several days at -30°C to obtain the light pink-colored crystals of **6**-(BPh_4).

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGMENTS

The authors gratefully acknowledge financial support from the NRF of Korea through CRI (NRF-2012R1A3A2048842 to W.N.) and GRL (NRF-2010-00353 to W.N.) along with the U.S. National Institutes of Health (to K.D.K.), the National Natural Science Foundation of China (21290194, 21521062, and 21473215 to H.C.), and the Ministry of Science and Technology of China (2012YQ120060 to J.Y.).

REFERENCES

- (1). (a) Furchgott RF. *Angew. Chem., Int. Ed.* 1999; 38:1870. (b) Ignarro LJ. *Angew. Chem., Int. Ed.* 1999; 38:1882. (c) Murad F. *Angew. Chem., Int. Ed.* 1999; 38:1856. (d) Ignarro, LJ. *Nitric Oxide, Biology and Pathobiology*. Academic Press; San Diego, CA: 2000. (e) Richter-Addo GB, Legzdins P, Burstyn J. *Chem. Rev.* 2002; 102:857. [PubMed: 11942780] (f) Wasser IM, de Vries S, Moënné-Loccoz P, Schröder I, Karlin KD. *Chem. Rev.* 2002; 102:1201. [PubMed: 11942794] (g) Möller JKS, Skibsted LH. *Chem. Rev.* 2002; 102:1167. [PubMed: 11942791] (h) Bon CLM, Garthwaite J. *J. Neurosci.* 2003; 23:1941. [PubMed: 12629199] (i) Pepicelli O, Raiteri M, Fedele E. *Neurochem. Int.* 2004; 45:787. [PubMed: 15312973]
- (2). (a) Martin CT, Morse RH, Kanne RM, Gray HB, Malmström BG, Chan SI. *Biochemistry.* 1981; 20:5147. [PubMed: 6271178] (b) Cooper CE, Torres J, Sharpe MA, Wilson MT. *FEBS Lett.* 1997; 414:281. [PubMed: 9315702] (c) Tocheva EI, Rosell FI, Mauk AG, Murphy MEP. *Science.* 2004; 304:867. [PubMed: 15131305] (d) Lehnert, N.; Berto, TC.; Galinato, MGI.; Goodrich, LE. *The Handbook of Porphyrin Science*. Kadish, KM.; Smith, K.; Guillard, R., editors. World Scientific; Hackensack, NJ: 2011. p. 1-247.
- (3). (a) Enemark JH, Feltham RD. *Coord. Chem. Rev.* 1974; 13:339. (b) Richter-Addo, GB.; Legzdins, P. *Metal Nitrosyls*. Oxford University Press; New York: 1992. (c) McCleverty JA. *Chem. Rev.*

2004; 104:403. [PubMed: 14871130] (d) Ford PC, Lorkovic IM. Chem. Rev. 2002; 102:993. [PubMed: 11942785] (e) Heinecke J, Ford PC. Coord. Chem. Rev. 2010; 254:235. (f) Fry NL, Mascharak PK. Acc. Chem. Res. 2011; 44:289. [PubMed: 21361269] (g) Berto TC, Speelman AL, Zheng S, Lehnert N. Coord. Chem. Rev. 2013; 257:244. (h) Tsai M-L, Tsou C-C, Liaw W-F. Acc. Chem. Res. 2015; 48:1184. [PubMed: 25837426] (i) Pulukkody R, Darensbourg M. Acc. Chem. Res. 2015; 48:2049. [PubMed: 26090911] (j) Hunt AP, Lehnert N. Acc. Chem. Res. 2015; 48:2117. [PubMed: 26114618] (k) Fitzpatrick J, Kim E. Acc. Chem. Res. 2015; 48:2453. [PubMed: 26197209] (l) Hematian S, Garcia-Bosch I, Karlin KD. Acc. Chem. Res. 2015; 48:2462. [PubMed: 26244814]

- (4). (a) Cho J, Sarangi R, Annaraj J, Kim SY, Kubo M, Ogura T, Solomon EI, Nam W. Nat. Chem. 2009; 1:568. [PubMed: 20711413] (b) Sastri CV, Oh K, Lee YJ, Seo MS, Shin W, Nam W. Angew. Chem., Int. Ed. 2006; 45:3992.
- (5). (a) Armor J. Inorg. Chem. 1973; 12:1959. (b) Roberts RL, Carlyle DW, Blackmer GL. Inorg. Chem. 1975; 14:2739. (c) Ungermann CB, Caulton KG. J. Am. Chem. Soc. 1976; 98:3862.
- (6). Doyle MP, Pickering RA, Dykstra RL, Cook BR. J. Am. Chem. Soc. 1982; 104:3392.
- (7). Franz KJ, Lippard SJ. Inorg. Chem. 2000; 39:3722. [PubMed: 11196841]
- (8). Metzker G, Lopes PP, da Silva ACH, da Silva SC, Franco DW. Inorg. Chem. 2014; 53:4475. [PubMed: 24738470]
- (9). (a) Clarkson SG, Basolo F. J. Chem. Soc., Chem. Commun. 1972:670. (b) Clarkson SG, Basolo F. Inorg. Chem. 1973; 12:1528.
- (10). (a) Cheng L, Powell DR, Khan MA, Richter-Addo GB. Chem. Commun. 2000:2301. (b) Videla M, Roncaroli F, Slep LD, Olabe JA. J. Am. Chem. Soc. 2007; 129:278. [PubMed: 17212399] (c) Roncaroli F, Videla M, Slep LD, Olabe JA. Coord. Chem. Rev. 2007; 251:1903. (d) Subedi H, Brasch NE. Inorg. Chem. 2013; 52:11608. [PubMed: 24050193]
- (11). Skodje KM, Williard PG, Kim E. Dalton Trans. 2012; 41:7849. [PubMed: 22538296]
- (12). (a) Blough NV, Zafiriou OC. Inorg. Chem. 1985; 24:3502. (b) Nauser T, Koppenol WH. J. Phys. Chem. A. 2002; 106:4084.
- (13). (a) Pfeiffer S, Gorren ACF, Schmidt K, Werner ER, Hansert B, Bohle DS, Mayer B. J. Biol. Chem. 1997; 272:3465. [PubMed: 9013592] (b) Coddington JW, Hurst JK, Lymar SV. J. Am. Chem. Soc. 1999; 121:2438. (c) Lymar SV, Khairutdinov RF, Hurst JK. Inorg. Chem. 2003; 42:5259. [PubMed: 12924897] (d) Goldstein S, Lind J, Merényi G. Chem. Rev. 2005; 105:2457. [PubMed: 15941219] (e) Koppenol WH, Bounds PL, Nauser T, Kissner R, Rüegger H. Dalton Trans. 2012; 41:13779. [PubMed: 23007920] (f) Molina C, Kissner R, Koppenol WH. Dalton Trans. 2013; 42:9898. [PubMed: 23698514]
- (14). (a) Schopfer MP, Wang J, Karlin KD. Inorg. Chem. 2010; 49:6267. [PubMed: 20666386] (b) Ouellet H, Ouellet Y, Richard C, Labarre M, Wittenberg B, Wittenberg J, Guertin M. Proc. Natl. Acad. Sci. U. S. A. 2002; 99:5902. [PubMed: 11959913]
- (15). (a) Gardner PR, Gardner AM, Martin LA, Salzman AL. Proc. Natl. Acad. Sci. U. S. A. 1998; 95:10378. [PubMed: 9724711] (b) Ford PC, Lorkovic IM. Chem. Rev. 2002; 102:993. [PubMed: 11942785] (c) Gardner PR, Gardner AM, Brashear WT, Suzuki T, Hvitved AN, Setchell KDR, Olson JS. J. Inorg. Biochem. 2006; 100:542. [PubMed: 16439024]
- (16). (a) Pestovsky O, Bakac A. J. Am. Chem. Soc. 2002; 124:1698. [PubMed: 11853446] (b) Herold S, Koppenol WH. Coord. Chem. Rev. 2005; 249:499. (c) Su J, Groves JT. Inorg. Chem. 2010; 49:6317. [PubMed: 20666389]
- (17). (a) Yokoyama A, Han JE, Cho J, Kubo M, Ogura T, Siegler MA, Karlin KD, Nam W. J. Am. Chem. Soc. 2012; 134:15269. [PubMed: 22950528] (b) Yokoyama A, Cho K-B, Karlin KD, Nam W. J. Am. Chem. Soc. 2013; 135:14900. [PubMed: 24066924] (c) Yokoyama A, Han JE, Karlin KD, Nam W. Chem. Commun. 2014; 50:1742.
- (18). (a) Maiti D, Lee D-H, Sarjeant AAN, Pau MYM, Solomon EI, Gaoutchenova K, Sundermeyer J, Karlin KD. J. Am. Chem. Soc. 2008; 130:6700. [PubMed: 18457392] (b) Park GY, Deepalatha S, Pui SC, Lee D-H, Mondal B, Narducci Sarjeant AA, del Rio D, Pau MYM, Solomon EI, Karlin KD. J. Biol. Inorg. Chem. 2009; 14:1301–1311. [PubMed: 19662443]
- (19). Kurtikyan TS, Ford PC. Chem. Commun. 2010; 46:8570.

- (20). (a) Kurtikyan TS, Eksuzyan SR, Hayrapetyan VA, Martirosyan GG, Hovhannisyan GS, Goodwin JA. J. Am. Chem. Soc. 2012; 134:13861. [PubMed: 22881578] (b) Kurtikyan TS, Eksuzyan SR, Goodwin JA, Hovhannisyan GS. Inorg. Chem. 2013; 52:12046. [PubMed: 24090349]
- (21). (a) Nam W. Acc. Chem. Res. 2015; 48:2415. [PubMed: 26203519] (b) Cho J, Jeon S, Wilson SA, Liu LV, Kang EA, Braymer JJ, Lim MH, Hedman B, Hodgson KO, Valentine JS, Solomon EI, Nam W. Nature. 2011; 478:502. [PubMed: 22031443]
- (22). Kumar P, Lee Y-M, Park YJ, Siegler MA, Karlin KD, Nam W. J. Am. Chem. Soc. 2015; 137:4284. [PubMed: 25793706]
- (23). The stability of **1** in the absence of substrate (i.e., [(12-TMC) Co^{II}]²⁺) in CH₃CN indicates that the NO-dissociation and NO-transfer reaction could not be a solvent-assisted process (Figure 3a, inset).
- (24). (a) Solc M. Nature. 1966; 209:706. (b) Holleman, AF.; Wiberg, E. Inorganic Chemistry. Academic Press; San Diego: 2001.
- (25). Nakamoto, K. Infrared and Raman Spectra of Inorganic and Coordination Compounds, Part B, Applications in Coordination, Organometallic, and Bioinorganic Chemistry. 6th ed.. John Wiley & Sons, Inc.; Hoboken, NJ: 2009.
- (26). The final product obtained is a Co(II) complex, not a Co(III) complex, indicating that there is a missing electron in the overall reaction of **1** with O₂ to give Co(II)-nitrate complex **6**, which was confirmed by X-ray crystallography, ESI-MS, and EPR spectroscopy. That is the reason why we did not describe a more detailed mechanism and, based only on the product analyses, we proposed the reaction mechanism of **1** with O₂ (see Scheme 4).
- (27). The peroxyxynitrite isomerization to nitrate is generally thought to occur extremely rapidly, sometimes without observable intermediates. It can occur by an initial homolytic cleavage of the PN O–O bond, forming a high-valent metal–oxo and nitrogen dioxide (NO₂) molecule within a cage, followed by re-formation of an N–O bond to give nitrate.
- (28). (a) Owen TM, Rohde J-U. Inorg. Chem. 2011; 50:5283. [PubMed: 21526756] (b) Ford PC, Fernandez BO, Lim MD. Chem. Rev. 2005; 105:2439. [PubMed: 15941218]
- (29). Wick PK, Kissner R, Koppenol WH. Helv. Chim. Acta. 2000; 83:748.
- (30). Tran NG, Kalyvas H, Skodje KM, Hayashi T, Moënné-Loccoz P, Callan PE, Shearer J, Kirschenbaum LJ, Kim E. J. Am. Chem. Soc. 2011; 133:1184. [PubMed: 21244001]
- (31). (a) Gunaydin H, Houk KN. Chem. Res. Toxicol. 2009; 22:894. [PubMed: 19374346] (b) Ferrer-Sueta G, Radi R. ACS Chem. Biol. 2009; 4:161. [PubMed: 19267456]
- (32). (a) Cho J, Sarangi R, Nam W. Acc. Chem. Res. 2012; 45:1321. [PubMed: 22612523] (b) Garcia-Bosch I, Cowley RE, Díaz DE, Siegler MA, Nam W, Solomon EI, Karlin KD. Chem. -Eur. J. 2016; 22:5133. [PubMed: 26919169]
- (33). Armarego, WLF.; Chai, CLL. Purification of Laboratory Chemicals. 6th ed.. Pergamon Press; Oxford, U.K.: 2009.
- (34). Halfen JA, Young VG Jr. Chem. Commun. 2003:2894.
- (35). Young, CL. Solubility Data Series Vol. 8: Oxides of Nitrogen. International Union of Pure and Applied Chemistry (IUPAC); 1981.
- (36). Sheldick, GM. *SHELXTL/PC*, Version 6.12 for Windows XP. Bruker Axis Inc.; Madison, WI: 2001.

Figure 1.

(a) UV-vis spectral changes for the formation of **1** upon addition of NO (1 equiv) to a solution of [(14-TMC)Co^{II}]²⁺ (pink line; 0.50 mM) under an Ar atmosphere in CH₃CN at -40 °C. **1** was generated within 3 min. Inset: Solution IR spectra of **1** (3.0 mM, black line) and [(14-TMC)Co^{II}]²⁺ (3.0 mM, pink line) recorded under an Ar atmosphere in CH₃CN at -40 °C. (b) EPR spectra of **1** (black line) and [(14-TMC)Co^{II}]²⁺ (pink line) recorded in CH₃CN at 5 K.

Figure 2.

(a) DFT-optimized geometry of $[(14\text{-TMC})\text{Co}^{\text{III}}(\text{NO})]^{2+}$ (**1**) in the singlet ground state. (b) The two singly occupied orbitals accommodating two electrons with opposite spins in the singlet ground state of **1**. (c) DFT-optimized geometry of the singlet ground state of **2** compared with the corresponding X-ray crystal structure. H-atoms are omitted for clarity.

Figure 3.

(a) UV-vis spectral changes obtained in the NO-transfer from **1** (0.50 mM; black line) to [(12-TMC)Co^{II}]²⁺ (0.50 mM) to form **2** (red line) through the intermediacy of the blue line spectrum under an Ar atmosphere in CH₃CN at -40 °C. The blue line spectrum was confirmed to be the sum of those spectra of **1** (0.50 mM) and [(12-TMC)Co^{II}]²⁺ (0.50 mM). Inset: reaction time course monitored at 368 nm due to **2** in the NO-transfer reaction (red circles) and that monitored at 395 nm for natural decay of **1** (black circles). (b) ESI-MS spectrum of the reaction solution obtained in the NO-transfer reaction of **1** with [(12-

TMC)Co^{II}]²⁺. The peaks at $m/z = 414.2$, 416.0 , and 178.1 correspond to [(14-TMC)Co^{II}(ClO₄)]⁺ (calcd $m/z = 414.1$), [(12-TMC)Co^{III}(NO)(ClO₄)]⁺ (calcd $m/z = 416.1$), and [(14-TMC)Co^{II}(CH₃CN)]²⁺ (calcd $m/z = 178.1$), respectively.

Figure 4.

UV-vis spectral changes showing the formation of **2** (red line) observed upon addition of $[(12\text{-TMC})\text{Co}^{\text{II}}]^{2+}$ into a solution of **1** (black line) in increments of 0.20 equiv under Ar atmosphere in CH_3CN at -40°C . Inset: spectral titration monitored by following the absorbance at 368 nm due to the formation of **2**.

Figure 5.

(a) Solution IR spectral changes obtained in the reaction of **1** (3.0 mM) and [(12-TMC)Co^{II}]²⁺ (3.0 mM) under an Ar atmosphere in CH₃CN at -40 °C, showing the conversion of the NO stretching frequency due to **1** (black line) to that due to **2** (red line). (b) EPR spectral changes obtained in the reaction of **1** (2.0 mM) and [(12-TMC)Co^{II}]²⁺ (2.0 mM) under an Ar atmosphere in CH₃CN at -40 °C, only showing that [(12-TMC)Co^{II}]²⁺ (black line) was converted to [(14-TMC)Co^{II}]²⁺ (red line) because **1** and **2** are EPR-silent (*S*

= 0). All EPR spectra were recorded at 5 K with different reaction times (0, 5, 35, 70, 105, and 170 min). The inset shows the initial (black line) and final (red line) spectra expanded.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Figure 6.
DFT-calculated reaction profile of NO dissociation from **1** on singlet ground state.

Figure 7. DFT-optimized structures of reactant (**1**), transition state (**TS_{diss-NO}**), and product (**PR_{diss-NO}**) of NO dissociation process (all on singlet ground state, bond distance/angle labeled in angstroms/degrees, H-atoms are omitted for clarity).

Figure 8. DFT-calculated associative reaction profile of NO-transfer from **1** to $[(12\text{-TMC})\text{Co}^{\text{II}}]^{2+}$ initiated from a μ -NO-bridged intermediate $\{[(14\text{-TMC})\text{Co}^{\text{III}}\text{-NO-Co}^{\text{II}}(12\text{-TMC})](\text{ClO}_4)_3\}^+$ (**3**) on ground doublet state.

Figure 9.

(a) UV-vis spectral changes observed in the reaction of **1** (0.50 mM; black line) with O_2 (0.10 M) in CH_3CN at -40°C . Inset: ESI-MS spectrum of the complete reaction solution (black line). The peaks at $m/z = 377.1$ and 178.1 correspond to $[(14\text{-TMC})\text{-Co}^{\text{II}}(\text{NO}_3)]^+$ (**6**) (calcd $m/z = 377.1$) and $[(14\text{-TMC})\text{-Co}^{\text{II}}(\text{CH}_3\text{CN})]^{2+}$ (calcd $m/z = 178.1$). Blue and red lines show the isotope distribution patterns for **6**- ^{16}O (blue) and **6**- ^{18}O (red), which were obtained from the reaction of **1** with $^{16}\text{O}_2$ and $^{18}\text{O}_2$, respectively. (b) Solution IR spectral changes for

the disappearance of **1** (left panel, black line) and the formation of NO_3^- (right panel, blue line) observed in the reaction of **1** (3.0 mM) with O_2 (0.10 M) in CH_3CN at -40°C .

Figure 10. ORTEP diagram of $[(14\text{-TMC})\text{Co}^{\text{II}}(\text{NO}_3)]^+$ (**6**) with 50% probability thermal ellipsoids (gray, C; blue, N; red, O; violet, Co). H atoms have been removed for clarity.

(a) Dissociative pathway

(b) Associative pathway

Scheme 1.

Scheme 2.

Scheme 3.

Scheme 4.

Scheme 5.