Table 1.

Outpatient Clinic	General Neurology (Jan–March 2001) ²	Cognitive Function (Oct 2001–Dec 2001)	General Neurology (Jan–March 2002)
<i>n</i>	198	58 (new 39; FU 19)	211
Heard of NHS Direct?	–	34/56 (= 61%)	120 (= 57%)
Used NHS Direct?	4 (= 2%)	3/34 (= 9%; 5% of all consultations)	26 (= 22%; 12% of all consultations)
Call related to referral?	1/4	2/3	4/26

National Audit Office of increased use of NHS Direct⁴. Since most callers report themselves satisfied with information received^{4,5}, it seems reasonable to infer that calls help to shape patients' health beliefs and expectations, whether appropriately or not. Awareness of calls to NHS Direct may allow clinicians to focus on issues of particular concern to patients, and hopefully thereby improve patient satisfaction with their consultation.

References

1 Donaldson L. Telephone access to health care: the role of NHS Direct. *J Roy Coll Physicians Lond* 2000;**34**:33–35.
2 Larner AJ. Use of Internet medical websites and of NHS Direct by neurology outpatients prior to consultation. *Int J Clin Pract* 2002;**56**:219–21.
3 Abbott J, Alberti H. NHS Direct. *Br J Gen Pract* 2001;**51**:580–581.
4 The Comptroller and Auditor General. *NHS Direct in England*. London: Stationery Office, 2002.
5 O'Cathain A, Munro JF, Nicholl JP, Knowles E. How helpful is NHS Direct? Postal survey of callers. *Br Med J* 2000;**320**: 1035.

AJ LARNER

Consultant Neurologist
Liverpool

Troponin I as a risk stratification tool in the district general hospital

The value of cardiac troponin measurement as a risk stratification tool in patients with suspected acute coronary syndromes and without electrocardiographic ST-segment elevation has been well established¹. Recent guidelines advise troponin estimation in all patients with suspected acute coronary syndromes^{2,3}. Early invasive treatment (coronary angiography and revascularisation where appropriate) is recommended in troponin positive

patients while troponin negative patients can be discharged early with stress testing (before or after discharge)³. Utilising troponin in guiding treatment may have important implications with regard to bed resources, particularly in the district general hospital. We performed a retrospective audit to determine the effect of using troponin I on length of hospital stay in patients admitted with chest pain.

Our hospital, a district general hospital with on-site diagnostic coronary angiography, incorporated troponin I measurement into chest pain management guidelines two years ago. It was suggested that troponin I be measured on admission and 12 hours later in patients presenting with chest pain but without ST-segment elevation. The cut-off level for a normal troponin I (Abbot AxSYM system) was <0.3 mcg/l, the lower limit of detection. The new guidelines suggested that i) troponin negative patients could be discharged earlier (with stress testing as appropriate) ii) troponin positive patients should be referred to the cardiologists for possible inpatient coronary angiography. Median length of hospital stay and clinical characteristics were determined by case note review in a random selection of 229 patients admitted with chest pain during a 6-month period prior to the introduction of troponin measurement. A similar audit was performed in 210 randomly selected patients admitted during a 6-month period after introduction of troponin measurement. Results from both patient groups were compared using the Chi 2 test. Our results demonstrated a shortening in median length of stay (from 3 to 2 days) in patients with non-diagnostic ECGs and cardiac risk factors/prior history of coronary disease. Extrapolated to all chest pain admissions, shows that this shortening

would gain 1,351 bed days per annum (3.7 beds/day). However, this gain was offset by an increase in median length of stay (from 5 to 8 days) for high-risk patients presenting with ischaemic ECG changes. Extrapolation showed that this increase would yield a loss of 1,513 bed days per annum. Troponin positive patients were more likely to undergo in-patient coronary angiography (62% v 35%, P<0.05) and revascularisation (36% v 11%, P<0.05).

In conclusion, our audit shows that using troponin measurement in guiding treatment has important bed resource implications. Adopting troponin measurement reduces length of hospital stay in low-risk patients. However, using troponin status to help identify high-risk patients who require inpatient coronary angiography revascularisation significantly lengthens hospital stay. In hospitals without on-site diagnostic angiography, hospital stay may be further prolonged. We suggest that prompt transfer of high-risk (troponin positive) patients to tertiary cardiac centres will result in net bed-stay savings for district general hospitals.

Acknowledgement: The audit was funded by a grant from The Mason Medical Research Foundation.

References

1 Klootwijk P, Hamm C. Acute coronary syndromes: diagnosis. *Lancet* 1999; Suppl 2:SII 10–5.
2 Department of Health. *National Service Framework for Coronary Heart Disease*. London: DoH, 2000.
3 Bertrand ME, Simoons ML, Fox KAA, et al. Management of acute coronary syndromes: acute coronary syndromes without persistent ST segment elevation. *European Heart Journal* 2000;**21**: 1406–1432.

JAGDIP S SIDHU

ANIL TANEJA

KEITH A STEER

DENNIS J WRIGHT

Department of Medicine and Department of
Chemical Pathology
Northwick Park Hospital
Harrow

Patterns of childbearing amongst female hospital doctors and GPs

The article *Women in hospital medicine* by E Paice (*Clin Med* Sept/Oct 2001 pp344–5) highlights some important issues faced by