Published in final edited form as:

Mol Neurobiol. 2016 September ; 53(7): 4821–4832. doi:10.1007/s12035-015-9407-8.

Examining the neural and astroglial protective effects of cellular prion protein expression and cell death protease inhibition in mouse cerebrocortical mixed cultures

Kevin K.W. Wang^{1,*}, Zhihui Yang¹, Allen Chiu², Fan Lin¹, and Richard Rubenstein^{2,*}

¹Program for Neurotrauma, Neuroproteomics and Biomarkers Research, Departments of Psychiatry, Neuroscience and Physiological Science, McKnight Brain Institute, L4-100, University of Florida, 1149 South Newell Drive, Gainesville, FL 32611, USA

²Laboratory of Neurodegenerative Diseases and CNS Biomarker Discovery, Departments of Neurology and Physiology/Pharmacology, Box #1213, SUNY Downstate Medical Center, 450 Clarkson Avenue, Brooklyn, NY 11203-2098, USA, office: (718) 270-2019, fax: (718) 270-2459

Abstract

Overexpression of cellular prion protein, PrP^C, has cytoprotective effects against neuronal injuries. Inhibition of cell death-associated proteases such as necrosis-linked calpain and apoptosis-linked caspase are also neuroprotective. Here we systemically studied how PrP^C expression levels and cell death protease inhibition affect cytotoxic challenges to both neuronal and glial cells in mouse cerebrocortical mixed cultures (CCM).

Primary CCM derived from three mouse lines expressing no (PrP^C knockout mice, PrPKO), normal (wildtype, wt) or high (tga20) levels of PrP^C were subjected to necrotic challenge [calcium ionophore A23187], and apoptotic challenge [staurosporine (STS)]. CCM originated from tga20 mice provided the most robust neuron-astroglia protective effects against necrotic and early apoptotic cell death (LDH release) at 6 hr, but subsequently lost its cytoprotective effects. In contrast, PrPKO-derived cultures displayed elevated A23187- and STS-induced cell death at 24 hr. Calpain inhibitor SNJ-1945 protected against A23187 challenge at 6 hr in CCM from all three mouse lines; but protected only against A23187 and STS treatment by 24 hr in the PrPKO line. In parallel, caspase inhibitor Z-D-DCB protected against pro-apoptotic STS challenge at 6 and 24 hr.

Furthermore, we also examined α II-spectrin-breakdown products (primarily from neurons), and GFAP-breakdown products (from astroglia) as cytoskeletal proteolytic biomarkers. Overall, it appeared that both neurons and astroglial cells were less vulnerable to proteolytic attack during A23187 and STS challenges in tga20-derived cultures but more vulnerable in PrPKO-derived cultures. In addition, calpain and caspase inhibitors provide further protection against respective protease attacks on these neuronal and glial cytoskeletal proteins in CCM regardless of mouse-line origin. Lastly, some synergistic cytoprotective effects between PrP^C expression and addition of cell death-linked protease inhibitors were also observed.

*Co-corresponding authors: Kevin K.W. Wang, kwang@ufl.edu and Richard Rubenstein, richard.rubenstein@downstate.edu.

Keywords

cellular prion protein; calpain; caspase; necrosis; apoptosis; cytotoxin; cytoprotection; proteases; biomarkers

Introduction

The cellular prion protein (PrP^C) is a host-coded glycoprotein attached to the surface of the cell membrane by a glycosyl-phosphatidylinositol (GPI) anchor. PrP^C binds copper with high affinity and is highly expressed on the surface of neuronal and glial cells [1, 2]. Although PrP^C presumably plays a role in the neuropathology and transmissibility of prion diseases by undergoing a conformational change into an abnormal protease-resistant isoform (PrP^{Sc}) [3, 4], the normal physiologic role of PrP^C is still uncertain. It has been reported that PrP^C plays a diverse role in normal cellular functions including cell survival [5] neuritogenesis [6], neurite outgrowth [7] and myelin maintenance [8]. Furthermore numerous studies have reported that PrP^C functions in neuroprotection by preventing neuronal apoptosis and promoting neuronal survival [1, 2, 8–12].

Calpains are cytosolic located calcium-dependent proteases, which, under physiologic conditions, undergo controlled activation physiologically and potentially mediate neuronal cell signaling and synaptic function such as long-term potentiation [3, 4, 13, 14]. Under pathophysiological conditions, calpain is hyperactivated due to extreme and sustained calcium influx. In turns hyperactivated calpain can damage cytoskeletal proteins, resulting in neurotoxicity [5, 14, 15]. Calpains have significant involvement in the neuropathological mechanisms associated with different types of insults including traumatic brain injury (TBI), stroke, epilepsy and organ injury [6, 16, 17]. The activation of calpain has been associated with several neurodegenerative disorders. For example, calpain alters the processing of various proteins (tau, amyloid precursor protein, several cytoskeletal proteins) associated with Alzheimer's disease (AD). Cell culture studies have shown that calpain activation does not cause significant PrP^C degradation [7, 18]. It has also been shown that calpain inhibitors are neuroprotective [16, 17, 19–22].

Caspases, are calcium-independent proteases, involved in signal transduction, cell-cycle programming and DNA repair and programmed cell death (apoptosis) [23]. Although there are several types of caspases, caspase 3 is mainly linked to neuronal apoptosis [16]. Under conditions of traumatic brain injury (TBI), spinal cord injury, or cerebral ischemia, caspases activate apoptotic cascades. Similar caspase-dependent apoptotic events have been associated with neurodegenerative diseases including AD, Huntington's disease and amyotrophic lateral sclerosis [24]. Therefore, the inhibition of caspase proteolytic activity is neuroprotective since it suppresses the apoptotic component of neuroinjury or neurodegeneration.

Here we have generated neuronal-glial mixed cultures (CCM) from mouse lines expressing different levels of PrP^C to evaluate the neural- and glial-protective properties of both PrP^C and calpain-caspase inhibitors following necrotic and apoptotic challenges.

Methods

CCM cultures and treatments

Primary CCM were obtained from 1 day old mouse pups derived from wt, PrPKO and tga20 mice. Cells were cultured in Dulbecco's modified Eagle's medium/F12 medium containing 10% horse serum (heat-inactivated) and 6% fetal bovine serum (heat-inactivated) in 96-well poly-L-lysine-coated plates similarly as previously described studies with rats [25–28]. Non-neuronal cell division was halted 3 days post plating by adding to the culture medium 35 µg/ml uridine and 15 µg/ml 5-fluoro-2'-deoxyuridine. These treatments did not alter neuronal cell viability. All cultures were maintained in a 37° C oxygenated incubator (21% O₂, 8% CO₂, and 71% N₂) at all times except during media changes. On day 14–16 post-plating maturation, the cultures were washed two times with defined neurobasal medium containing B27 supplement and then subjected to neurotoxic and gliatoxic challenges [20 µM calcium ionophore A23187 (A23), or 1 µM staurosporine (STS)] in the absence or presence of calpain inhibitor SNJ-1945 (SNJ) and caspase inhibitor Z-D-DCB (ZD) [26, 29].

Cell death assay

Cell death of the CCM cultures (either necrosis, or secondary necrosis following apoptosis) was assessed by measuring activity of the cytosolic enzyme lactate dehydrogenase (LDH) released into the medium as described previously [30, 31] at 6 hr and 24 hr after cytotoxic challenges were added. Two-way ANOVA was used for all LDH data analysis.

Western blots and quantification

Cellular proteins were extracted at 24 hr post challenges using lysis buffer (1% Triton X-100, 20 mM Tris-HCl, pH 7.4, 5 mM EGTA, 1 mM dithiothreitol, 1x protease inhibitor cocktail) as described previously for western blotting to independently assess neuronal [αII-spectrin and spectrin breakdown products (SBDPs)] and glial cell injury (GFAP and its BDPs)[32–34]. Briefly, for SDS-polyacrylamide gel electrophoresis (SDS-PAGE), cell lysate (20 µg protein per lane) samples were loaded onto 10–12 lane Tris-Glycine 4–20% mini-gels (Life Technologies/Invitrogen) [29]. Following electrophoresis, separated proteins were laterally transferred to polyvinylidene fluoride (PVDF) membranes (iBlot semi-dry transfer system) at a constant voltage of 12 V for 9 min at ambient temperature. After electrotransfer, the membranes were blocked for 1 hr at ambient temperature in 5% non-fat milk prepared in Tris-buffered saline containing 0.05% Tween-20 (TBST) followed by incubation at 4° C for 3 hr or overnight with primary antibodies [mouse anti-αII-spectrin (Biomol) or rabbit anti-GFAP (Abcam) at 1:2000 dilution in TBST-5% non-fat milk]. This was followed by three washes with TBST, a 2-hr incubation at ambient temperature with a secondary antibody [goat anti-rabbit or anti-mouse IgG conjugated to alkaline phosphatase (Novagen)] and developed with substrate BCIP-NBT reagent (KPL). Housekeeping proteins, β-actin and/or carbonic anhydrase-II (CA-II), were included as sample loading controls. Quantitative evaluation of target protein levels was performed via computer-assisted densitometric scanning (Epson XL3500 scanner, Long Beach, CA, USA) and image analysis with Image J software (version 1.6) (NIH). Two-way or one-way ANOVA were used for all immunoblotting quantification analysis.

Results

PrP is resistant to calpain and Caspase-3 digestion

To first understand the interplay between calpain, caspase and PrPc – we subjected brain lysate from tga20 overexpressed PrPc mice *in vitro* digestion with calpain-1, -2 and caspase-3 (1:50, protease to total protein ratio) Western blots are probed with anti-PrP (30K) MAb E119. We found that no fragments of PrPc were observed and the intact PrPc levels remain the same after digestion (Suppl. Figure 1). As positive controls for calpain and caspase-3 activities, upper part of the blot was also probed with anti- α II-spectrin antibody – which shows signature SBDP150/145 and SBDP150i/120 patterns upon calpain-1, 2 and caspase-3 digestions, respectively [16–17]. This demonstrates that calpain and caspase did not directly target PrPc.

Effects of PrP^C expression on cell death of challenged CCM cultures

For this study, we employed CCM culture from wildtype (wt) mice, PrP-knockout (PrPKO) mice or from PrPc transgenic mice tga20. CCM lysate shows no PrPc protein (30 kDa), while tga20 (tga) CCM lysate showed about 3-fold the levels of PrP^c in wildtype CCM lysate (Suppl. Figure 2).

Following the 14–16 day *in vitro* maturation period of the CCM derived from the three mouse lines, the cultures were subjected to necrotic challenge (A23187; A23), and apoptotic challenge (staurosporine; STS). The cell death protease, calpain, is activated extensively in the presence of both necrotic challenges, and to a lesser extent with pro-apoptotic STS challenge. Caspases, however, are only activated during apoptosis (i.e. STS challenge). We measured cell death by LDH release at an acute (6 hr) and delayed time point (24 h) (Figs. 1, 2).

Based on the 6 h LDH results, A23187 induced rapid and strongest necrosis (Fig. 1). All three cell lines (wt, PrPKO, tga20) have significantly increased levels of LDH release with A23 by 6 h and 24 h, when compared to their respective controls (indicated by *, two way ANOVA) (Figs. 1A). Also, both tga20 CCM, which overexpresses PrP^c in neurons, and PrPKO cells released more LDH than their wildtype (et0 counterpart at 24 h, However, tga20 cells released more LDH than the PrPKO cells at both 6 h and 24 h (indicated by #) (Fig. 1A, and see table on right).

Apoptosis-inducer STS also began to cause cell death by 6 h and 24 h and triggers significant release of LDH by ‘secondary necrosis’ in CCM from wt, PrPKO and tga20 mouse lines, when compared to respective controls, with the exception of Tga with STS for 6 h (Fig. 1B). In addition, with STS treatment, tga20 CCM released less LDH at 6 h than wildtype CCM, while PrPKO CCM released more LDH than tga20 CCM at 24 h (indicated by #) (Fig. 1B, and see table on right).

Effects of calpain and caspase inhibitors on cell death of challenged CCM from wt, PrPKO and tga20

From the LDH data set, we also sought to analyze the effects of calpain and caspase protease inhibitors on CCM cell death as reflected by LDH release. In ANOVA analysis, we observed that at 6 h and 24 h, calpain inhibitor SNJ-1945 (SNJ) protected against LDH against pro-necrotic A23 challenge only in PrPKO CCM indicated by **) (Figure 2A). As expected, pan caspase inhibitor Z-D-DCB (ZD) offered no significant protections against pro-necrosis A23187 challenge. Against pro-apoptotic STS challenge, SNJ also provided protection to PrPKO culture at both 6 h and 24 h (Figure 2B, indicated by **). In parallel, Z-D-DCB (ZD) robustly protected against STS challenge in PrPKO CCM (Fig. 2, indicated by **).

Effects of PrP^C expression levels on cytoskeletal proteolytic biomarkers, SBDPs and GFAP-BDPs, in challenged CCM

CCM were again subjected to necrotic challenge (A23) and apoptotic challenge (STS). We measured cytoskeletal proteolytic damage with (i) α II-Spectrin breakdown products (SBDPs) caused by calpain (SBDP150/SBDP145 for necrosis) and by caspase-3 (SBDP120 for apoptosis) at the 24 hr time point (Wang et al., 2000) (Fig. 3), and with (ii) GFAP-BDPs induced by calpain (GBDP-38K) and by caspase (GBDP-22K) [34] (Fig. 4).

First, relative abundance of α II-Spectrin (and SBDP), GFAP (and GBDP) was examined in mouse cerebrocortical mixed neuron/astroglia culture (CCM) versus mouse primary astroglial culture. α II-Spectrin and GFAP were found to be highly intense bands in control CCM mixed culture samples, while SBDP150/145 and GBDP38K are equally prominent in A23187 (A23)-treated CCM cells (Suppl. Fig. 3). In contrast, in control primary astroglial culture, α II-spectrin was barely visible (even after long exposure) but the intact GFAP band was very intense. In A23187-treated astroglial culture, SBDP150 was again barely detectable, while GFAP-38K was observed as an intense band, in addition to other GFAP-BDPs between 48–40K). Astroglia isolation method from mouse cortex was similarly to previously described method [51]. Based on the data shown, we concluded that SBDPs we observed were predominantly derived from neurons in the CCM, while GFAP and GBDPs were exclusively derived from the astrocytes in the CCM.

In our next set of experiments, we examined the effects of cytotoxic challenges (A23187 and STS) on α II-spectrin integrity in CCM (Figure 3). At 24 hours following A23187 treatment, there were significant increases of necrosis-linked SBDP145 in all three mouse lines (wt, PrPKO and tga20) when compared to respective controls (Fig. 3B, indicated by *). In parallel, apoptosis inducer STS triggered significant formation of both calpain-mediated SBDP145 and caspase-3 mediated SBDP120 in CCM cells from all three mouse lines (Fig. 3B, C, indicated by *).

We also examined astroglia toxicity under the same paradigms (Figure 4). A23187 treatment induced significant increases of calpain-generated GBDP-38K in all three cell lines (wt, PrPKO and tga20) when compared to respective controls (Fig. 4B, indicated by *). STS also caused astroglial injury by 24 h, while triggering both calpain-produced GBDP-38K (Fig.

4B) and caspase-mediated GBDP-22K in CCM from the wt, PrPKO, tga20 lines (Fig. 4B, C).

Effects of calpain/caspase inhibitors on α II-spectrin and GFAP cytoskeletal proteolysis in A23187 and STS challenged CCM from wt, PrPKO and tga20 mice

Next, we examined the effects of inhibition of cell death proteases against α II-spectrin and GFAP cytoskeletal proteolysis in cytotoxin-challenged CCM from wt, PrPKO and tga20 mice. When CCM from the 3 mouse lines were challenged with A23 in the presence of SNJ1945 (SNJ), the SBDP145 was significantly reduced (Fig. 5A, C, E, indicated by P-value). Caspase inhibitor Z-D-DCB (Z-D) had no significant effects on SBDP145 levels in A23-treated cells from all three mouse lines (Fig. 5B, C, E). Z-D exhibited no effects on calpain-mediated SBDP145 in A23187-treated wt and tga20 CCM lines, but elevated its counterpart in PrPKO cells (Fig. 5A, C, E; indicated by blue #),

With STS treatment, SNJ again attenuated SBDP145 formation in all three CCM lines (Fig. 5B, C, E). SNJ had no effects on caspase-produced SBDP120, in CCM from wt or Tga20-derived CCM, but actually elevated SBDP120 (Fig. 5B, D, F; indicated by blue #). Z-D robustly blocked SBDP120 in STS-treated cells from all three mouse lines (Fig. 5B, D, F, indicated by P-value).

As for astroglial cell cytoskeletal proteolysis, we examined the effects of both SNJ and ZD on GFAP-BDP formation (Figure 6). First, Calpain-produced GBDP-38K was strongly inhibited by calpain inhibitor SNJ in both A23- and STS- treated CCM from all three mouse lines (Fig. 6A, C, E; indicated by P-values). Interestingly, SNJ also blocked caspase-mediated GBDP-22K in both CCM from wt and PrPKO lines (Fig. 6B, D, F, indicated by #).

Conversely, caspase inhibitor ZD had no effects on GBDP-38K in A23-treated wt and PRPKO glial cells, but did reduce GBDP-38K in tga20 CCM (Fig. 6A, C, E, indicated by #). As expected, ZD blocked caspase-mediated GBDP-22K in STS-treated glial cells from all three mouse lines (Fig. 6B, D, F). Overall, we observed that in CCM from all three mouse lines, calpain and caspase inhibitors can exert their effects on blocking their respective α II-spectrin and GFAP BDP formation, with some crossover inhibitions also observed. Again, this is consistent with the overall LDH results described above. Some

Effects of different levels of cellular PrPc on CCM responses to A23187 and STS with or without calpain/caspase inhibition

Using two-way ANOVA, we sought to examine CCM responses to A23187 and STS with or without the addition of SNJ or ZD might differ in wt, PrPKO and tga20 mouse lines (Figure 7). We found that SBDP145 levels in A23-, STS- and STS+ZD treatments were significantly higher in PrPKO CCM when compared to wt counterpart. Conversely, SBDP145 levels were significantly lower in A23-, A23+Z-D and STS-treated tag20 CCM, as compared to wt CCM. Lastly, SBDP145 levels were found higher in A23-, A23+ZD an STS in tga20 CCM, as compared to the PrPKO counterparts (Fig. 7A, also see Figure 5).

Similarly, we also found that SBDP120 levels in STS- and STS+SNJ was significantly higher in PrPKO CCM as compared to wt, while SBDP120 levels were significantly lower in

STS- and STS+SNJ treated tag20 CCM, as compared to CCM. In addition, SBDP120 levels were lower in STS- and STS+SNJ treated tag20 CCM, as compared to its PrPKO counterpart (Fig. 7B, also see Fig. 5). There were minimal SBDP120 levels in A23187-treated cells; thus, CCM cell line comparative analysis was not performed.

For GFAP-BDP analysis, We found that GBDP-38K levels in A23-, A23+ZD- and STS- and treatment were significantly higher in PrPKO CCM when compared to wt counterpart. Conversely, GBDP-38K levels are significantly lower in A23-, A23+Z-D and STS- and STS +Z-treated tag20 CCM, as compared to wt CCM. Lastly, GBDP-38K levels were also found higher in A23-, A23+ZD an STS- and STS+ZD-treated tag20 CCM, as compared to the PrPKO counterparts (Fig. 7C, also see Fig. 5).

Similarly, we also found that GBDP-22K levels in STS- and STS+SNJ was significantly higher in PrPKO CCM as compared to wt, while GBDP-22K levels were significantly lower in STS- and STS+SNJ treated tag20 CCM, as compared to CCM. Lastly, GBDP-22K levels were lower in STS- and STS+SNJ treated tag20 CCM, as compared to the PrPKO counterparts (Fig. 7D, also see Fig. 6). There were minimal GBDP-22K in A23187-treated cells; thus, CCM cell line comparative analysis was not performed.

Discussion

PrP^C is a glycoprotein expressed at highest levels in neurons of the central nervous system (CNS) and, to a lesser degree, in glial cells. It is localized on the cell surface and bound to the membrane via a GPI anchor. There have been numerous studies suggesting that PrP^C has a neuroprotective function. Kim et al. [35] reported that, in neural cell culture studies, PrP^C prevents Bax-mediated apoptosis through a caspase-dependent pathway. Further, due to its affinity for copper, PrP^C can prevent oxidative stress by altering the activity of Cu-Zn superoxide dismutase [36]. PrP^C also mediates signal transduction pathways responsible for neurite outgrowth through activation of kinases [37]. In addition to the neuroprotective properties of PrP^C, other studies have shown that PrP^C can also play a role in neurotoxic events through signaling cascades. For example, neurons with PrP^C are more sensitive to STS, a kinase inhibitor linked to apoptosis [38]. Lastly, to date, there is no study on PrP^C expression in neurons with concomitant protection of neighboring astroglia cells.

In parallel, many studies have demonstrated that inhibition of both cell death-linked calpain and caspase protease families are neuroprotective in brain injury [16, 39]. Both proteases are known to express not only in neurons but also in glial cells. However, It is unclear if calpain and caspase also play a role in the cell death process of astroglia cells (glia) during CNS injury conditions. Calpain appears to be universally activated in necrosis and some apoptotic conditions, while caspases are robustly activated in most if not all apoptotic cell death conditions in varying pathological conditions reported [16, 29, 40–43]. We hypothesize that the cytoprotective effects of calpain and/or caspase inhibition is likely extended to glia cells as well.

Thus, in this study, we used CCM from mouse lines expressing different levels of PrP^C (PrP^C non-expressor PrPKO, wt and PrP^C overexpressing tag20) to systemically examine the

potential neuro- and glia-protective properties of both PrP^C and calpain-caspase inhibitors against either pro-necrotic (A23) or pro-apoptotic (STS) challenges. We found that PrP^C overexpression cells (CCM from tga20 mice) provided the most robust neuronal-glia protective effects against necrotic and early apoptotic cell death, while the CCM from PrPKO mice have more extensive combined early neuronal-glia death post-challenges (measured by LDH release at 6 h) (Fig 1A, B). We further note that by 24 hr post-challenge, PrP^C overexpressing cultures had lost the neuronal-glia protective effects. Yet, PrPKO cells still displayed more extensive cell death compared to wildtype cells at 24 hr (Fig 1A, B).

Taking advantage of the established proteolytic SBDPs as neuronal biomarkers [16, 44, 45] and GFAP-BDPs as glial injury biomarkers [34, 46–48], we also examined in more details if PrP^C can protect both cell types against pro-necrotic and pro-apoptotic challenges. α II-spectrin is known to be highly enriched in axons. To confirm that α II-spectrin is primarily a neuronal protein in our culture system, we examined mouse primary CCM (cortical mixed culture of about 50:50 neurons-astrocytes) in parallel with primary mouse astrocytes (> 95% astrocytes). We showed that by western blot, intact α II-spectrin and SBDPs were readily observed in control and A23187-treated CCM, respectively, while α II-spectrin and SBDPs were minimally detected in mouse primary astrocyte culture counterpart (Suppl. Figure 3). In contrast, in both CCM and astrocyte culture, intact GFAP and GFAP-BDP were prominently detected in control and A23187 treatment, respectively (Suppl. Figure 3). Thus SBDPs observed in our experiments were predominantly derived from the neurons in CCM. In addition, our data shows that both the neuronal injury as reflected by SBDP145 levels in pro-necrotic challenges and SBDP120 levels in pro-apoptotic challenge (16) were sensitive and inversely related to the expression levels of PrP^C (Fig 3).

For astroglial cells, we observed the signature 38 kDa fragment (in addition to minor fragment of 44K) in necrotic challenges, as we have described before [34] (Fig 4B). We reported here an additional fragment of GFAP of 22 kDa that we suspect was generated by caspase attack (Fig 4A). We found that glia cell injuries, as reflected by the levels of GBDP-38K and GBDP-22K, are again sensitive and irreversibly related to the levels of PrP^C expression (Fig 4A, B, C). We like to point out that we used a neuronal murine PrP promoter for the PrP transgene in the tga20 mouse line. This explains that in tga20 mouse brain, PrP is expressed found to be expressed in neurons, not astrocytes (Suppl. Figure 4). Thus any glia-protective effects of PrP^C is likely derived from the neuron-glia co-culture environment including the likely increased release of PrP^C into culture media -especially under cytotoxic conditions, as we have described previously [18].

In addition, calpain and caspase inhibitors can apparently provide further protection against necrotic and apoptotic cell death phenotypes, respectively (Fig 2A, B). These inhibitors also inhibit respective protease attacks on two cellular cytoskeletal proteins (α II-spectrin and GFAP) regardless of the CCM culture source (wt, PrPKO or tga20) (Fig 5, 6). SBDP150 and SBDP145 primarily generated in neurons and GBDP-38K generated in glial cells were robustly blocked by calpain inhibitors, while caspase inhibitor blocked SBDP120 formation (Fig 5). Interestingly, we found some Z-D cross-over inhibition of calpain-generated BDPs, such as SBDP145 reduction in A23+ZD-treated tga20 CCM and GBDP-38K reduction in

A23187+ZD-treated tga20 CCM. SNJ also cross-inhibited caspase-generated GBDP-22K in STS+ANJ-treated cells. These suggest interplay between the two proteases.

Overall, this is the first study to address that not only PrP^C expression in neurons is neuroprotective, but also that its expression (and possible release of PrP^C) also has cytoprotective effects on its co-cultured astroglial cells. In addition, calpain and caspase inhibitors can provide further protection against respective protease attacks on these neuronal and glial cytoskeletal proteins. This study also raises an important point about the potential synergistic cytoprotective effects of cell death protease inhibition and PrP^C, as this concept can potentially be explored therapeutically.

There are several limitations in our current study. For example, we do not know the dosage of PrP^C overexpression that is required to provide significant neuro- and/or glia-protective effects above wt cells where there is already a basal level of PrP^C. We also do not know if such neuron- and glia-protective effects of PrP^C are translatable to animal studies of TBI where both neuron and glia cell death are common. If so, will the synergistic effects of protease inhibition and PrP^C expression be preserved under *in vivo* environment? In fact, we are currently conducting such *in vivo* studies to address some of these key questions. Lastly, we submit that it is extremely important to understand the biochemical and subcellular mechanisms for PrP^C to exert its cytoprotective effects on both neurons and on astroglial cells. Based on such understanding, it might become possible in the future to produce PrP^C-mimicking small molecule drugs that can afford similar neuro- and glia-protective effects as potential treatment for various brain injury conditions.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgement

This work was supported, in part, by the SUNY Downstate Medical Center, the University of Florida Department of Psychiatry and McKnight Brain Institute faculty development funds (KKW) and an award from the U.S. Department of Defense (DoD) (RR). For the DoD funding: 1. The U.S. Army Medical Research Acquisition Activity, Fort Detrick, MD 21702-5014 is the awarding and administering acquisition office, 2. This work was supported by the Office of the Assistant Secretary of Defense for Health Affairs through the U.S. Army Medical Research and Materiel Command under award # W81XWH-11-2-0069. Opinions, interpretations, conclusions and recommendations are those of the authors and are not necessarily endorsed by the DoD, and 3. In conducting research using animals, the investigators adhere to the laws of the United States and regulations of the Department of Agriculture.

References

1. Prusiner SB. The prion diseases. *Brain Pathol.* 1998; 8:499–513. [PubMed: 9669700]
2. Salès N, Rodolfo K, Hässig R, et al. Cellular prion protein localization in rodent and primate brain. *Eur J Neurosci.* 1998; 10:2464–2471. [PubMed: 9749773]
3. Caughey B, Chesebro B. Transmissible spongiform encephalopathies and prion protein interconversions. *Adv Virus Res.* 2001; 56:277–311. [PubMed: 11450303]
4. White AR, Enever P, Tayebi M, et al. Monoclonal antibodies inhibit prion replication and delay the development of prion disease. *Nature.* 2003; 422:80–83. [PubMed: 12621436]
5. Krebs B, Wiebelitz A, Balitzki-Korte B, et al. Cellular prion protein modulates the intracellular calcium response to hydrogen peroxide. *J Neurochem.* 2007; 100:358–367. [PubMed: 17241158]

6. Lopes MH, Hajj GNM, Muras AG, et al. Interaction of cellular prion and stress-inducible protein 1 promotes neuritogenesis and neuroprotection by distinct signaling pathways. *The Journal of Neuroscience*. 2005; 25:11330–11339. [PubMed: 16339028]
7. Kanaani J, Prusiner SB, Diacovo J, et al. Recombinant prion protein induces rapid polarization and development of synapses in embryonic rat hippocampal neurons in vitro. *J Neurochem*. 2005; 95:1373–1386. [PubMed: 16313516]
8. Bremer J, Baumann F, Tiberi C, et al. Axonal prion protein is required for peripheral myelin maintenance. *Nat Neurosci*. 2010; 13:310–318. [PubMed: 20098419]
9. Mitteregger G, Vosko M, Krebs B, et al. The Role of the Octarepeat Region in Neuroprotective Function of the Cellular Prion Protein. *Brain Pathology*. 2007; 17:174–183. [PubMed: 17388948]
10. Hoshino S, Inoue K, Yokoyama T, et al. Prions prevent brain damage after experimental brain injury: a preliminary report. *Acta Neurochir Suppl*. 2003; 86:297–299. [PubMed: 14753456]
11. Williams SK, Fairless R, Weise J, et al. Neuroprotective effects of the cellular prion protein in autoimmune optic neuritis. *The American Journal of Pathology*. 2011; 178:2823–2831. [PubMed: 21641403]
12. Couplier M, Messiaen S, Boucreaux D, Eloit M. Axotomy-induced motoneuron death is delayed in mice overexpressing PrPc. *NSC*. 2006; 141:1827–1834.
13. Lynch G, Baudry M. Brain spectrin, calpain and long-term changes in synaptic efficacy. *Brain Res Bull*. 1987; 18:809–815. [PubMed: 3040193]
14. Liu J, Liu MC, Wang KK, et al. Calpain in the CNS: from synaptic function to neurotoxicity. *Sci Signal*. 2008; 1:re1. [PubMed: 18398107]
15. Liu J, Liu MC, Wang KK, et al. Physiological and pathological actions of calpains in glutamatergic neurons. *Sci Signal*. 2008; 1:scisignal-123tr3v1.
16. Wang KK. Calpain and caspase: can you tell the difference?, by kevin K.W. Wang Vol. 23, pp. 20-26 [In Process Citation]. *Trends in Neurosciences*. 2000; 23:59. [PubMed: 10652545]
17. Wang KKW, Po-Wai Y. Calpain inhibition: an overview of its therapeutic potential. *Trends in pharmacological sciences*. 1994; 15:412–419. [PubMed: 7855906]
18. Wang KK, Zoltewicz JS, Chiu A, et al. Release of Full-Length PrP(C) from Cultured Neurons Following Neurotoxic Challenges. *Front Neurol*. 2012; 3:147–147. [PubMed: 23093947]
19. Wang Y, Briz V, Chishti A, et al. Distinct roles for μ -calpain and m-calpain in synaptic NMDAR-mediated neuroprotection and extrasynaptic NMDAR-mediated neurodegeneration. *Journal of Neuroscience*. 2013; 33:18880–18892. [PubMed: 24285894]
20. Takano E, Maki M. [Calpastatin: molecular mechanism of calpain inhibition]. *Tanpakushitsu Kakusan Koso*. 1997; 42:2181–2188. [PubMed: 9366195]
21. Ray SK, Karmakar S, Nowak MW, Banik NL. Inhibition of calpain and caspase-3 prevented apoptosis and preserved electrophysiological properties of voltage-gated and ligand-gated ion channels in rat primary cortical neurons exposed to glutamate. *NSC*. 2006; 139:577–595.
22. Yuen PW, Wang KKW. Calpain inhibitors: novel neuroprotectants and potential anticataract agents. *Drugs of the Future*. 1998; 23:741–750.
23. Deveraux QL, Reed JC. IAP family proteins—suppressors of apoptosis. 1999
24. Nicotera P, Leist M, Manzo L. Neuronal cell death: a demise with different shapes. *Trends in pharmacological sciences*. 1999; 20:46–51. [PubMed: 10101962]
25. DOD-DVBIC. DVBIC Worldwide-TBI 2000–2013 Report. DOD Report 1–5. 2014.
26. Zhang Z, Lerner SF, Liu MC, et al. Multiple α II-spectrin breakdown products distinguish calpain and caspase dominated necrotic and apoptotic cell death pathways. *Apoptosis*. 2009; 14:1289–1298. [PubMed: 19771521]
27. Goldstein LE, Fisher AM, Tagge CA, et al. Chronic Traumatic Encephalopathy in Blast-Exposed Military Veterans and a Blast Neurotrauma Mouse Model. *Science Translational Medicine*. 2012; 4:134ra60–134ra60.
28. McKee AC, Stern RA, Nowinski CJ, et al. The spectrum of disease in chronic traumatic encephalopathy. *Brain*. 2013; 136:43–64. [PubMed: 23208308]

29. Yang Z, Lin F, Robertson CS, Wang K. Dual vulnerability of TDP-43 to calpain and caspase-3 proteolysis after neurotoxic conditions and traumatic brain injury. *Journal of Cerebral Blood Metabol.* 2014; 34:1444–1452.
30. Koh JY, Choi DW. Quantitative determination of glutamate mediated cortical neuronal injury in cell culture by lactate dehydrogenase efflux assay. *J Neurosci Methods.* 1987; 20:83–90. [PubMed: 2884353]
31. Nath R, Probert A Jr, McGinnis KM, Wang KKW. Evidence for activation of caspase-3-like protease in excitotoxin- and hypoxia/hypoglycemia-injured neurons. *J Neurochem.* 1998; 71:186–195. [PubMed: 9648865]
32. Guingab-Cagmat J, Newsom K, Vakulenko A, et al. In vitro MS-based proteomic analysis and absolute quantification of neuronal-glia injury biomarkers in cell culture system. *Electrophoresis.* 2012; 33:3786–3797. [PubMed: 23161537]
33. Zoltewicz JS, Mondello S, Yang B, et al. Biomarkers track damage after graded injury severity in a rat model of penetrating brain injury. *J Neurotrauma.* 2013; 30:1161–1169. [PubMed: 23409698]
34. Zhang Z, Zoltewicz JS, Mondello S, et al. Human traumatic brain injury induces autoantibody response against glial fibrillary acidic protein and its breakdown products. *PLoS ONE.* 2014; 9:e92698. [PubMed: 24667434]
35. Kim B-H, Lee H-G, Choi J-K, et al. The cellular prion protein (PrPC) prevents apoptotic neuronal cell death and mitochondrial dysfunction induced by serum deprivation. *Molecular Brain Research.* 2004; 124:40–50. [PubMed: 15093684]
36. Brown DR, Schulz-Schaeffer WJ, Schmidt B, Kretzschmar HA. Prion protein-deficient cells show altered response to oxidative stress due to decreased SOD-1 activity. *Exp Neurol.* 1997; 146:104–112. [PubMed: 9225743]
37. Chen S, Mangé A, Dong L, et al. Prion protein as trans-interacting partner for neurons is involved in neurite outgrowth and neuronal survival. *Mol Cell Neurosci.* 2003; 22:227–233. [PubMed: 12676532]
38. Paitel E, Sunyach C, da Costa CA, et al. Primary cultured neurons devoid of cellular prion display lower responsiveness to staurosporine through the control of p53 at both transcriptional and post-transcriptional levels. *J Biol Chem.* 2004; 279:612–618. [PubMed: 14570892]
39. Larner SF, Hayes RL, Wang KKW. Proteolytic mechanisms of cell death in the central nervous system. In: Lajtha, A.; Banik, N.; Ray, SK., editors. *Handbook of Neurochemistry.* US: Springer; 2008. p. 249–279.
40. Shahim P, Tegner Y, Wilson DH, et al. Blood biomarkers for brain injury in concussed professional ice hockey players. *JAMA Neurol.* 2014; 71:684–692. [PubMed: 24627036]
41. Villa PG, Henzel WJ, Sensenbrenner M, et al. Calpain inhibitors, but not caspase inhibitors, prevent actin proteolysis and DNA fragmentation during apoptosis. *J Cell Sci.* 1998; 111:713–722. [PubMed: 9472000]
42. Park SY, Tournell C, Sinjoanu RC, Ferreira A. Caspase-3- and calpain-mediated tau cleavage are differentially prevented by estrogen and testosterone in beta-amyloid-treated hippocampal neurons. *NSC.* 2007; 144:119–127.
43. Liu MC, Akle V, Zheng W, et al. Comparing calpain- and caspase-3-mediated degradation patterns in traumatic brain injury by differential proteome analysis. *Biochem J.* 2006; 394:715–725. [PubMed: 16351572]
44. Siman R, Zhang C, Roberts VL, et al. Novel surrogate markers for acute brain damage: cerebrospinal fluid levels correlate with severity of ischemic neurodegeneration in the rat. *J Cereb Blood Flow Metab.* 2005; 25:1433–1444. [PubMed: 15902199]
45. Zhang Z, Mondello S, Kobeissy F, et al. Protein biomarkers for traumatic and ischemic brain injury: from bench to bedside. *Translational Stroke Research.* 2011; 3:1–32. [PubMed: 21666822]
46. Papa L, Lewis LM, Falk JL, et al. Elevated levels of serum glial fibrillary acidic protein breakdown products in mild and moderate traumatic brain injury are associated with intracranial lesions and neurosurgical intervention. *Ann Emerg Med.* 2012; 59:471–483. [PubMed: 22071014]
47. Okonkwo DO, Yue JK, Puccio AM, et al. GFAP-BDP as an acute diagnostic marker in traumatic brain injury: results from the prospective transforming research and clinical knowledge in traumatic brain injury study. *J Neurotrauma.* 2013; 30:1490–1497. [PubMed: 23489259]

48. Wang KK, Moghieb A, Yang Z, Zhang Z. Systems biomarkers as acute diagnostics and chronic monitoring tools for traumatic brain injury. SPIE Defense, Security, and Sensing. 2013 87230O–87230O–15.
49. Neumar RW, Xu YA, Gada H, Guttman RP, Siman R. Cross-talk between calpain and caspase proteolytic systems during neuronal apoptosis. J Biol Chem. 2003 Apr 18; 278(16):14162–14167. 2003 Epub 2003 Feb 7. PMID: 12576481. [PubMed: 12576481]
50. Yang Z, Lin F, Robertson CS, Wang KKW. Dual vulnerability of TDP-43 to calpain and caspase-3 proteolysis following neurotoxic conditions and Traumatic Brain Injury. J. Cereb. Blood Flow Metabol J Cereb Blood Flow Metab. 2014 Sep; 34(9):1444–1452. 2014.
51. Aloisi F, Agresti C, Levi G. Establishment, characterization, and evolution of cultures enriched in type-2 astrocytes. J Neurosci Res. 1988; 21:188–198. [PubMed: 3216420]

Group	Time point	Difference	t	P value
wt-ctr vs PrP KO ctr	6h	0.2012	1.735	ns
	24h	0.159	1.372	ns
wt-ctr vs tga ctr	6h	-0.03433	0.2962	ns
	24h	0.01533	0.1323	ns
PrP KO ctr vs tga ctr	6h	-0.2355	2.032	ns
	24h	-0.1437	1.239	ns
wt-ctr vs wt-A	6h	0.641	5.53	P<0.001
	24h	0.3985	3.438	P<0.01
PrP KO ctr vs PrP KO-A	6h	0.5718	4.933	P<0.001
	24h	0.9647	8.322	P<0.001
tga ctr vs tag A	6h	0.4708	4.062	P<0.001
	24h	0.816	7.039	P<0.001
wt-A vs PrP KO-A	6h	0.132	1.138	ns
	24h	0.7252	6.256	P<0.001
wt-A vs tag A	6h	-0.2045	1.764	ns
	24h	0.4328	3.734	P<0.001
PrP KO-A vs tag A	6h	-0.3365	2.903	P < 0.05
	24h	-0.2923	2.522	P < 0.05

Figure 1. Effect of PrP^C expression levels on LDH release from CCM after 6 and 24 hr of cytotoxin challenge

LDH release as a monitor for cell death in wt, PrPKO and tga20 (tga) mouse-derived CCM subjected to (A) pro-necrotic challenge (20 μ M A23187) and (B) pro-apoptotic challenge (1 μ M staurosporine; STS) for 6hr and 24 h. Wt, PrPKO and tga20-derived CCM that are unchallenged (Control) are also included., (*) indicates LDH release levels higher than respective control. (#) indicates that PrPKO LDH release are higher or tga20 LDH levels are lower than their wt counterparts with either A23187 or STS challenge. N= 4 separate experiments. Significance is defined as $p < 0.05$ (two- way ANOVA). ns = not significant. Those Group comparisons that were significant are highlighted in bold. Group comparisons between two mouse lines that are significant are indicated in blue labels and #.

Figure 2. Effects of calpain and caspase inhibition on LDH release from CCM after 6 and 24 hr of cytotoxin challenge

LDH release as a monitor for cell death in wt, PrPKO and tga20 mouse-derived CCM subjected to pro-necrotic and pro-apoptotic challenges for 6 and 24 hr. wt (WT), PrPKO and tga20 (tga)-derived CCM are subjected to (A) pro-necrosis 20 μ M A23187 [A23], (B) pro-apoptosis staurosporine [STS] with or without 30 μ M SNJ-1945 (SNJ) or caspase inhibitor Z-D-DC (ZD, 50 μ M). (*) indicates that calpain inhibitor SNJ-1945 (30 μ M SNJ) or caspase inhibitor Z-D-DC (50 μ M ZD) treatment yield statistically significant differences in LDH levels compared to the no inhibitor counterparts (with A23 or STS). N= 4 separate experiments. Significance is defined as $p < 0.05$ (two way ANOVA). ns = not significant. Those Group comparisons that were significant are highlighted in bold letters in the table and with * on graph.

Figure 3. Effect of PrP^C expression on cytotoxin-induced, and neuronal injury-linked, αII-spectrin breakdown products formation in mouse CCM

Necrosis inducer A23187 (20 μM) and apoptosis inducer staurosporine (STS 1 μM) were used. (A) Representative immunoblots for αII-spectrin; quantification of (B) SBDP145 levels and (C) SBDP120 levels. Significance is defined as $p < 0.05$ (two-way ANOVA). ns = not significant. N= 4 separate experiments. Those Group comparisons that were significant are highlighted in bold letters in the table and with * on graph.

Figure 4. Effect of PrP^C expression on cytotoxin-induced and glial injury-linked GFAP-BDP formation in mouse CCM

Necrosis inducer A23187 (20 μ M) and apoptosis inducer staurosporine (STS 1 μ M) were used. (A) Representative immunoblots for GFAP, and quantification of (B) GBDP-38K levels and (C) GBDP-22K levels. N= 4 separate experiments. Significance is defined as $p < 0.05$ (two-way ANOVA). ns = not significant. Those Group comparisons that were significant are highlighted in bold letters in the table and with * on graph.

Figure 5. Effect of calpain and caspase inhibitors on cytotoxin-induced SBDP formation in CCM from wt, PrPKO and tga20 mice

Calpain produced SBDP150, SBDP145 and caspase-produced SBDP120 were monitored. Calpain inhibitor SNJ-1945 (SNJ), caspase inhibitor Z-D-DCB (ZD) were used 1 hr prior to cytotoxin A23187 treatment for 24 h with or without SNJ or ZD (A), or STS treatment for 24 h with or without SNJ or ZD (B). N = 4 separate experiments. Significance is defined as $p < 0.05$ (two-way ANOVA). ns = not significant. Those Group comparisons (A23 or STS groups with or without SNJ or ZD) that were significant are highlighted in bold. Group comparisons between two mouse lines that are significant are indicated in blue labels and #.

Figure 6. Effect of calpain and caspase inhibitors on cytotoxin-induced glial GFAP-BDP formation in CCM from wt, PrPKO and tga20 mice

Calpain produced GBDP-38K and putative caspase-produced GBDP-22K were monitored.

(A, C) A23187 treatment for 24 h with or without SNJ-1945 or Z-D-DCB. (B, D) STS treatment for 24 h with or without SNJ-1945 or Z-D-DCB.). N = 4 separate experiments. Significance is defined as $p < 0.05$ (two-way ANOVA). ns = not significant. Those Group comparisons (A23 or STS groups with or without SNJ or ZD) that were significant are highlighted in bold. Group comparisons between two mouse lines that are significant are indicated in blue labels and #.

A

SBDP145			
wt vs PrPKO	Difference	t	P value
A23187	10.1	5.502	P<0.001
A23+SNJ	-0.232	0.1264	ns
A23+ZD	2.9	1.58	ns
STS	7.274	3.962	P<0.01
STS+SNJ	2.85	1.552	ns
STS+ZD	10.54	5.739	P<0.001
wt vs tga	Difference	t	P value
A23187	-10.6	5.774	P<0.001
A23+SNJ	-1.532	0.8345	ns
A23+ZD	-14.2	7.735	P<0.001
STS	-4.926	2.683	ns
STS+SNJ	0.782	0.426	ns
STS+ZD	-10.13	5.519	P<0.001
PrPKO vs tga	Difference	t	P value
A23187	-20.7	11.28	P<0.001
A23+SNJ	-1.3	0.7081	ns
A23+ZD	-17.1	9.315	P<0.001
STS	-12.2	6.646	P<0.001
STS+SNJ	-2.068	1.127	ns
STS + ZD			

B

SBDP120			
wt vs PrPKO	Difference	t	P value
A23187	-0.03333	0.05265	ns
STS	13.1	20.69	P<0.001
STS+SNJ	12.27	19.38	P<0.001
STS+ZD	0.6667	1.053	ns
wt vs tga	Difference	t	P value
A23187	-0.03333	0.05265	ns
STS	-4.3	6.792	P<0.001
STS+SNJ	-4.567	7.213	P<0.001
STS+ZD	0	0	ns
PrPKO vs tga	Difference	t	P value
A23187	0	0	ns
STS	-17.4	27.48	P<0.001
STS+SNJ	-16.83	26.59	P<0.001
STS+ZD	-0.6667	1.053	ns

C

GBDP-38K			
wt vs PrPKO	Difference	t	P value
A23187	13.03	6.092	P<0.001
A23+SNJ	0	0	ns
A23+ZD	12.93	6.045	P<0.001
STS	6.5	3.038	P < 0.05
STS+SNJ	0	0	ns
STS+ZD	4.267	1.994	ns
wt vs tga	Difference	t	P value
A23187	-7.4	3.459	P<0.01
A23+SNJ	0.1	0.04674	ns
A23+ZD	-8.4	3.926	P<0.01
STS	-9.333	4.362	P<0.001
STS+SNJ	0	0	ns
STS+ZD	-7.533	3.521	P<0.01
PrPKO vs tga	Difference	t	P value
A23187	-20.43	9.55	P<0.001
A23+SNJ	0.1	0.04674	ns
A23+ZD	-21.33	9.971	P<0.001
STS	-15.83	7.4	P<0.001
STS+SNJ	0	0	ns
STS+ZD	-11.8	5.515	P<0.001

D

GBDP-22K			
wt vs PrPKO	Difference	t	P value
A23187	0	0	ns
STS	6.473	17.5	P<0.001
STS+SNJ	2.7	7.301	P<0.001
STS+ZD	0.0266	0.07211	ns
wt vs tga	Difference	t	P value
A23187	0	0	ns
STS	-3.35	9.058	P<0.001
STS+SNJ	-1.133	3.065	P < 0.05
STS+ZD	0	0	ns
PrPKO vs tga	Difference	t	P value
A23187	0	0	ns
STS	-9.823	26.56	P<0.001
STS+SNJ	-3.833	10.37	P<0.001
STS+ZD	-0.0266	0.07211	ns

Figure 7. Effects of different levels of cellular PrPc on CCM responses to A23187 and STS with or without calpain/caspase inhibition

For alphaII-spectrin, calpain-produced SBDP145 and caspase-produced SBDP120 were monitored (**A, B**). For GFAAP, calpain produced GBDP-38K and putative caspase-produced GBDP-22K were monitored. (**C, D**). N= 4 separate experiments. Significance is defined as $p < 0.05$ (two- way ANOVA). ns = not significant. Group comparisons between two mouse lines that are significant are highlighted in bold.