CASE REPORT

Kinesiology taping as an adjunct for pain management in cancer?

Gourav Banerjee,^{1,2} Jonathan Rebanks,³ Michelle Briggs,^{1,2} Mark I Johnson^{1,2}¹Faculty of Health and Social Sciences, Leeds Beckett University, City Campus, Leeds, West Yorkshire, UK²Leeds Pallium Research Group, Leeds, UK³Physiotherapy at Castle Clinic and Nidderdale Medical Practice, Knaresborough, North Yorkshire, UK

Correspondence to

Gourav Banerjee,
g.banerjee@leedsbeckett.ac.uk

Accepted 23 June 2016

SUMMARY

We present the case of a 46-year-old woman who developed severe pain described as 'tearing' and 'searing' in the left side of the mid-trapezius region near the thoracic 8 vertebra (T8). The patient had undergone surgery for T8 fracture which had resulted from metastasis (secondary breast cancer). A community nurse referred the patient for physiotherapy assessment and treatment for her musculoskeletal pain and related symptoms that had affected her mobility and functional activities. The patient was treated with soft tissue therapy with the addition of kinesiology taping on follow-up visits. Kinesiology tape was applied over her left side trapezius region and left shoulder. The patient reported significant reductions in pain severity and felt greater control and stability over her left shoulder region, which resulted in better function and overall quality of life measures. She did not report any adverse effects.

BACKGROUND

People with cancer, especially in advanced stages, have to cope with a number of distressing symptoms including pain, fatigue, lack of energy, oedema, dyspnoea and abdominal discomfort that result in reduced quality of life.^{1–4} Management of these symptoms is pharmacotherapy-led with a trade-off between symptom relief and adverse effects. Non-pharmacological interventions are often used in combination with medication to alleviate symptoms.

Kinesiology taping involves the application of elasticated, thin, porous, water-resistant, cotton-based adhesive tape to the skin. It has become popular in recent times through its use by high profile elite sportspeople. The tape and technique was developed in the 1970s for the management and rehabilitation of sports-related musculoskeletal injuries.⁵ Nowadays, kinesiology taping is used by physical and sports therapists, osteopaths, chiropractors and nurses to manage musculoskeletal pain, cancer-related lymphoedema and stroke-related spasticity.^{6–7} Kinesiology tape is available without prescription in a variety of shapes, sizes, colours and patterns (eg, Kinesio Tex Tape, Rocktape) and can be worn for up to 5 days during normal activities of daily living including showering. Kinesiology taping differs from conventional taping and bandaging techniques that often use rigid zinc oxide tape because it does not restrict movement and appears to be better tolerated.⁸ Conventional taping techniques stabilise injuries by reducing loading and restricting movement of injured body parts. In contrast, kinesiology tape is

applied to the skin under mild tension by stretching the tape and/or stretching the skin (eg, by flexing or extending the joint). This provides a pulling force to the skin which may correct articular malalignments and influence proprioception and muscle function.⁹ It is also claimed that kinesiology taping 'lifts' the epidermis to produce regions of decompression beneath the skin and that this improves microcirculation of the blood and lymph that relieves swelling and pain.⁹ In addition, it is claimed that kinesiology taping stimulates low threshold cutaneous mechanoreceptors, especially during movement that could affect proprioception and also modulate pain via inhibition of nociceptive transmission in the central nervous system ('closing the pain gate').¹⁰ Scientific evidence to support these claims is lacking.

Research evidence on the effectiveness of kinesiology taping to improve pain and function is limited to musculoskeletal conditions in non-cancer populations including patellofemoral pain syndrome and rotator cuff tendinopathy. There are at least 19 systematic reviews and meta-analyses with insufficient evidence to judge outcome because of too few randomised controlled trials (RCTs) and those that exist have small sample sizes. A recent systematic review by Lim and Tay¹¹ evaluated kinesiology taping for musculoskeletal pain and disability lasting for more than 4 weeks and included 17 studies. They found that kinesiology taping was superior to minimal or no intervention for musculoskeletal pain and disability that lasts for more than 4 weeks but not superior to other treatment approaches.

Kinesiology taping is currently used by some practitioners to alleviate lymphoedema and associated symptoms in patients with cancer. There are two systematic reviews^{6–7} that have identified only two studies. Tsai *et al*¹² found no differences in excess limb size or water composition between a single-layered bandage and kinesiology tape in 41 patients with decongestive lymphatic therapy for breast-cancer-related moderate-to-severe lymphoedema. Patients reported that they preferred kinesiology tape because it was easier to use, more comfortable and convenient, and allowed longer wearing time. Białoszewski *et al*¹³ found that kinesiology taping added to standard lymphatic massage produced a faster reduction of oedema compared with standard lymphatic massage on its own in 24 patients without cancer with post-operative oedema.

Recently, we evaluated reports of the use of kinesiology taping in cancer.¹⁴ We found seven studies

CrossMark

To cite: Banerjee G, Rebanks J, Briggs M, *et al*. *BMJ Case Rep* Published online: [please include Day Month Year] doi:10.1136/bcr-2016-216439

with a comparison group (including RCTs) and nine case series/reports without a comparison group in breast cancer for lymphoedema and related outcome measures including pain, grip strength, limb range of motion (ROM) and quality of life measures. There was weak evidence for benefits of kinesiology taping as an adjunct to complete decongestive therapy for managing cancer-related lymphoedema.

We describe a patient with secondary breast cancer who was given kinesiology taping treatment to manage severe pain and improve function of the upper limb.

CASE PRESENTATION

A 46-year-old woman diagnosed with primary breast cancer in August 2011 and secondary breast cancer in 2014 underwent spinal surgery in December 2014 due to fracture of the thoracic 8 vertebra (T8) resulting from metastases. A community nurse referred the patient to physiotherapy for assessment and treatment of musculoskeletal-related symptoms that had affected her mobility and function. The patient was seen by the physiotherapist (JR) at the hospice outpatient department on 18 December 2015. The patient's primary symptom was severe pain with sensations of 'tearing' and 'searing' accompanied by stiffness in the left side mid-trapezius region near the site of the fracture. Symptoms were attributed to the consequential effects of the disease and surgery and poor posture resulting from the patient's hypervigilance to pain. The patient was found positive for signs of soft tissue tightness around the region of pain and her symptoms had adversely affected her activities of daily living and quality of life. She was treated by JR with low-level myofascial tissue release techniques including passive stretching of the soft tissue which alleviated her symptoms. The patient returned to clinic distressed and anxious on 12 January 2016 because the symptoms had reappeared.

TREATMENT

The patient was treated with soft tissue therapy techniques by JR during the first consultation on 18 December 2015. This alleviated her symptoms. The patient returned to clinic on 12 January 2016 distressed and anxious, reporting that symptoms had reappeared. In addition to the soft tissue therapy techniques that he had employed earlier, JR decided to try kinesiology tape to manage pain and provide support to the shoulder. Three strips of blue coloured Tiger K Tape (width=5 cm; length ~20 cm) stretched ~20% more than their original length were applied over the left side trapezius region. During the application of kinesiology tape, the patient assumed a position where the cervicothoracic spine remained flexed with the arm hanging forwards, stretching the shoulder (figure 1). The tape was applied directly from a superior to inferior direction over the painful site which corresponded to the affected dermatome.

The patient was provided with basic information about the principles and practice of kinesiology tape and advised to continue wearing tape until her next appointment which was scheduled a week later which was, 19 January 2016. She was prescribed a low intensity exercise plan for muscular lengthening and advised to continue with normal daily living activities including showering or bathing with the tape in situ and to contact clinic if any problems arose. A telephone conversation between JR and the patient on 14 January 2016 confirmed that kinesiology tape was still in situ and in good condition, with minimal lifting of tape edges from the skin. The patient reported a substantial reduction in the severity and occurrence of painful episodes and that the left shoulder girdle area felt more 'stable'. She also reported that she had more control of

Figure 1 Application of three strips of kinesiology tape with ~20% stretch applied from superior to inferior direction with soft tissues in the region of the shoulder girdle and upper back in lengthened (stretched) position.

the shoulder joint complex while performing movements. No adverse effects were reported and the patient reported satisfaction with treatment. The patient was scheduled to return to clinic for follow-up assessment on 19 January 2016.

OUTCOME AND FOLLOW-UP

The patient attended clinic on 19 January 2016 for follow-up and was reassessed by JR. Kinesiology tape was in situ and in good condition, although there was lifting and curling of tape edges from the skin. She reported that pain severity reduced by ~50% compared with pretreatment and that the severity of tearing and searing sensations over her left mid-trapezius area had reduced by ~85%. The kinesiology tape was removed by JR. Inspection of the skin area under the tape revealed mild redness which was attributable to the tape removal. The patient reported that she had not experienced any adverse events. JR applied new kinesiology tape in an identical manner to that described for the first treatment.

The patient attended clinic on 2 February 2016 with the tape in situ. She reported satisfaction with kinesiology tape treatment and that she continued to experience reductions in pain severity and significant improvements in functional activities. The patient reported that she would like to continue using kinesiology taping, and so JR provided information about purchasing tape over the counter (and internet) and then demonstrated how to apply the kinesiology tape. It was decided that the husband would be the best person to apply the tape.

The patient attended clinic on 23 February 2016 and was again assessed by JR. She seemed relaxed. The patient reported that she had purchased kinesiology tape (Rocktape) from a local store in Leeds which was readily available. Her husband had applied the kinesiology tape without difficulty after about every 5 days with 12–24 hours of intermittent break periods before reapplication. The patient explained that intermittent breaks were necessary for moisturising the skin and allowing it to 'breathe' and to avoid getting the skin 'red' and 'sore'. After a few trials with Rocktape and then on subsequent use with Tiger K Tape, the patient stated that Tiger K Tape adheres to the skin better and longer and so was preferable to her.

The patient reported that kinesiology taping continued to be of benefit as she was able to self-manage her pain, which has improved her functional activities and emotional well-being. No adverse effect was reported.

The patient continued to wear kinesiology tape until last contact made with the patient by GB on 19 April 2016.

DISCUSSION

There is a paucity of reports on the use of kinesiology taping to manage musculoskeletal pain in patients with cancer. A pilot study without a comparison group found that kinesiology taping improved asymmetry between the scapula upper ribs and shoulder in 12 patients with postmastectomy breast cancer.¹⁵ Pyszora *et al*^{16–17} have reported four cases of the use of kinesiology taping added to integrated neuromuscular inhibition and myofascial release techniques, and exercise in elderly patients with advanced lung cancer, multiple myeloma and secondary breast cancer. They found improvements in pain, physical function and quality of life, and that the patients were very satisfied with kinesiology taping as a treatment. Kinesiology taping has also been shown to improve cervical pain and shoulder ROM and strength when used as an adjunct to physiotherapy treatment in a middle-aged woman who underwent partial glossectomy and neck dissection following tongue cancer.¹⁸

There is weak evidence that kinesiology taping improves lymphoedema. Case reports and studies without a comparison group suggest that kinesiology taping is beneficial or is as effective as conventional treatments for reducing lymphoedema (eg, compression sleeves) in cancer.^{19–27} These reports also suggest that kinesiology taping may be beneficial for mitigating associated symptoms in the lymphoedema limb including pain,²⁰ muscle tension,¹⁹ feeling of fullness/tightness/heaviness,^{20–22} and improving joint ROM¹⁹ and tissue texture,²¹ with overall improvements in disability²² and activities of daily living.²⁷ Kinesiology tape is comfortable to wear compared with compression garments and may be an alternative option for patients with poor short-stretch bandage compliance or to those that are contraindicated to decongestive compression therapy.^{20–21–23–24–27}

In summary, the patient in our report found kinesiology taping beneficial for managing pain, and for improving function and emotional well-being with no adverse effects. Kinesiology tape is inexpensive, safe, convenient to use and can be administered by a caregiver. Precautions include open wounds, dermatological diseases, allergy to adhesive tape and those particularly relevant in cancer include frail skin susceptible to irritation and damage. Some manufacturers of kinesiology tape have expressed fear of increasing blood flow and spread of cancer cells when using the tape for individuals with active cancer, although there is no evidence to support this claim and this hazard would equally apply to many lymphoedema treatments currently in use. Adverse effects associated with kinesiology taping are minimal.^{28–29}

We hope that this case will catalyse interest on the clinical utility of kinesiology taping as a treatment option for management of pain, swelling and other symptoms in cancer.

Patient's perspective

'Within 48 hours of taping applied, the symptoms had virtually disappeared and I was able to carry out everyday tasks to the same degree as before the initial pain. The use of the tape has been nothing short of miraculous in symptom management. Without the tape I was physically and psychologically distressed, the relief the tape brought helped me to regain movement and the ability to live as best as I can. Without the tape my back and postural muscles felt as if they were permanently in spasm. The tape taught my back muscles where they should sit and brought a much needed feeling of stability.'

Learning points

- ▶ Kinesiology taping is a low cost, simple, easy-to-apply non-pharmacological treatment that is being increasingly used in the management and rehabilitation of musculoskeletal-related injuries including pain in non-cancer settings.
- ▶ The case presented here demonstrates that kinesiology taping may have a role in the management of musculoskeletal pain and related outcome measures including physical function and quality of life in a selection of patients with cancer.
- ▶ Adverse effects associated with kinesiology taping are minimal and could include occurrence of skin soreness of mild-to-moderate intensity in non-allergic persons if kinesiology tape is allowed to remain in situ for too long (ie, >3–5 days).
- ▶ Carefully selected patients may benefit from kinesiology taping as an adjunct to other therapies within the continuum of cancer care, especially as it can be applied by the patient themselves and/or by their carers.
- ▶ Future research on the use and scope of kinesiology taping in cancer is warranted.

Acknowledgements The authors would like to acknowledge the patient who extended full support and cooperation.

Contributors MIJ, MB and GB conceived of the study; JR executed the study at a hospice. This project was funded by a PhD student bursary from the Jane Tomlinson Appeal; MIJ is the grant holder. JR undertook physiotherapy-related assessment and treatment responsibilities of this case. All authors contributed to the writing of the manuscript and approved the final manuscript. No outside contribution was requested or received during the preparation of this article.

Funding The Jane Tomlinson Appeal.

Competing interests None declared.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Teunissen SC, Wesker W, Kruitwagen C, *et al*. Symptom prevalence in patients with incurable cancer: a systematic review. *J Pain Symptom Manage* 2007;34:94–104.
- 2 Fan G, Filipczak L, Chow E. Symptom clusters in cancer patients: a review of the literature. *Curr Oncol* 2007;14:173.
- 3 Raphael J, Ahmedzai S, Hester J, *et al*. Cancer pain: part 1: pathophysiology; oncological, pharmacological, and psychological treatments: a perspective from the British Pain Society endorsed by the UK Association of Palliative Medicine and the Royal College of General Practitioners. *Pain Med* 2010;11:742–64.
- 4 Kroenke K, Zhong X, Theobald D, *et al*. Somatic symptoms in patients with cancer experiencing pain or depression: prevalence, disability, and health care use. *Arch Intern Med* 2010;170:1686–94.
- 5 Kinesio®UK. Kinesio® UK: A Brief History of Kinesio Tex Taping®. 2016 (cited 29 March 2016). <http://www.kinesiotaping.co.uk/history.jsp>
- 6 Morris D, Jones D, Ryan H, *et al*. The clinical effects of Kinesio® Tex taping: a systematic review. *Physiother Theory Pract* 2013;29:259–70.
- 7 Kalron A, Bar-Sela S. A systematic review of the effectiveness of Kinesio taping—fact or fashion? *Eur J Phys Rehabil Med* 2013;49:699–709.
- 8 Chang WD, Chen FC, Lee CL, *et al*. Effects of Kinesio taping versus McConnell taping for patellofemoral pain syndrome: a systematic review and meta-analysis. *Evid Based Complement Alternat Med* 2015;2015:471208.
- 9 Kase K, Wallis J, Kase T, *et al*. *Clinical therapeutic applications of the Kinesio taping method*. 3rd edn. Kinesio USA, LLC, 2013.
- 10 D'Mello R, Dickenson AH. Spinal cord mechanisms of pain. *Br J Anaesth* 2008;101:8–16.
- 11 Lim EC, Tay MG. Kinesio taping in musculoskeletal pain and disability that lasts for more than 4 weeks: is it time to peel off the tape and throw it out with the sweat? A systematic review with meta-analysis focused on pain and also methods of tape application. *Br J Sports Med* 2015;49:1558–66.

- 12 Tsai HJ, Hung HC, Yang JL, *et al.* Could Kinesio tape replace the bandage in decongestive lymphatic therapy for breast-cancer-related lymphedema? A pilot study. *Support Care Cancer* 2009;17:1353–60.
- 13 Białoszewski D, Woźniak W, Zarek S. Clinical efficacy of kinesiology taping in reducing edema of the lower limbs in patients treated with the ilizarov method—preliminary report. *Ortop Traumatol Rehabil* 2008;11:46–54.
- 14 Banerjee G, Rose A, Briggs M, *et al.* Kinesiology taping as a novel adjunct in oncology and palliative care? *British Psychosocial Oncology Society 2016 Annual Conference*; 03 March 2016; Madingley Hall, University of Cambridge, UK: Wiley Online Library, 2016:22.
- 15 Karczewska E, Szlachta P, Pytka K, *et al.* Kinesio taping method in the asymmetry treatment of the shoulder girdle in women after mastectomy—a pilot study. *Eur J Med Technol* 2016;1:10.
- 16 Pyszora A, Graczyk M, Krajnik M. What is the role of a physiotherapist in palliative care? Cases report. *Adv Palliat Med* 2009;8:31–4.
- 17 Pyszora A, Wójcik A, Krajnik M. Are soft tissue therapies and Kinesio taping useful for symptom management in palliative care? Three case reports. *Adv Palliat Med* 2010;9:87–92.
- 18 Courtney-Koro S. Rehabilitation following partial glossectomy and neck dissection for tongue cancer. *Rehabil Oncol* 2004;22:15–20.
- 19 Lipińska A, Śliwiński Z, Kiebzak W, *et al.* The influence of Kinesiotaping applications on lymphoedema of an upper limb in woman after mastectomy. *Fizjo Pol* 2007;7:258–69.
- 20 Pyszora A, Krajnik M. Is Kinesio taping useful for advanced cancer lymphoedema treatment? A case report. *Adv Palliat Med* 2010;9:141–4.
- 21 Finnerty S, Thomason S, Woods M. Audit of the use of kinesiology tape for breast oedema. *J Lymphoedema* 2010;5:38–44.
- 22 Kaya E, Kaplan C, Dandin Ö. Kinesiotaping for breast cancer related lymphedema. *Meme Sagligi Dergisi/J Breast Health* 2012;8:166–8.
- 23 Chou YH, Li SH, Liao SF, *et al.* Case report: manual lymphatic drainage and kinesio taping in the secondary malignant breast cancer-related lymphedema in an arm with arteriovenous (AV) fistula for hemodialysis. *Am J Hosp Palliat Care* 2013;30:503–6.
- 24 Hubert P, Gelais SS, Harris K, *et al.* The effects of SpiderTech™ kinesiology tape on the management of lymphedema in patients that are status post breast cancer treatment. *Rehabil Oncol* 2013;31:47.
- 25 Taradaj J, Halski T, Zdunczyk M, *et al.* Evaluation of the effectiveness of kinesio taping application in a patient with secondary lymphedema in breast cancer: a case report. *Prz Menopauzalny* 2014;13:73–7.
- 26 Navarro-Brazález B, Sánchez-Sánchez B. El vendaje en el tratamiento fisioterapéutico del linfedema secundario a cancer de mama: una serie de casos. *Fisioterapia* 2014;36:49–53.
- 27 Martins Jde C, de Aguiar SS, Fabro EA, *et al.* Safety and tolerability of Kinesio® Taping in patients with arm lymphedema: medical device clinical study. *Support Care Cancer* 2016;24:1119–24.
- 28 Wilson B, Bialocerkowski A. The effects of Kinesiotape applied to the lateral aspect of the ankle: relevance to ankle sprains—a systematic review. *PLoS ONE* 2015;10:e0124214.
- 29 Beatriz LA, Rafael M-M. Kinesio taping and patellofemoral pain syndrome: a systematic review. *Cent Eur J Sport Sci Med* 2015;9:47–54.

Copyright 2016 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- Submit as many cases as you like
- Enjoy fast sympathetic peer review and rapid publication of accepted articles
- Access all the published articles
- Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow