CASE REPORT

Distal coronary embolisation during transcatheter aortic valve implantation

Akira Tsujimura,¹ Naritatsu Saito,¹ Kenji Minakata,² Takeshi Kimura¹¹Department of Cardiovascular Medicine, Graduate School of Medicine, Kyoto University, Kyoto, Japan²Department of Cardiovascular Surgery, Graduate School of Medicine, Kyoto University, Kyoto, Japan

Correspondence to

Dr Naritatsu Saito,
naritatsu@kuhp.kyoto-u.ac.jp

Accepted 27 June 2016

SUMMARY

A 92-year-old woman was admitted to a hospital with severe aortic valve stenosis for transcatheter aortic valve implantation (TAVI). TAVI was performed under general anaesthesia. After balloon valvuloplasty, the patient became hypotensive and transesophageal echocardiography showed severe aortic regurgitation with severely depressed left ventricular wall motion. A 26 mm Sapien XT valve was deployed. However, the ventricular wall motion was still severely depressed. Coronary angiography showed occlusion of the mid-left anterior descending (LAD) artery. After crossing a 0.014" guidewire, manual vacuum aspiration was conducted and multiple emboli were removed. 2 drug-eluting stents were deployed in the LAD artery. Following this, the left ventricular wall motion improved. The patient's failure symptoms improved remarkably after TAVI. Histopathological examination of the aspirated emboli was compatible with a recent thrombus containing platelets, fibrin, erythrocytes and leucocytes. This is the first report to describe a distal coronary embolisation during TAVI with histopathological confirmation of the embolus.

BACKGROUND

Coronary obstruction is one of the most serious complications of transcatheter aortic valve implantation (TAVI).¹ The commonest mechanism of coronary obstruction during TAVI is the displacement of a native valve cusp over the coronary ostium. Distal coronary embolisation during TAVI is very rare. We experienced a rare case of distal coronary embolisation, and the emboli were successfully managed by coronary interventions. This is the first report to describe a distal coronary embolisation during TAVI with histopathological confirmation of the embolus.

CASE PRESENTATION

A 92-year-old woman was admitted to a hospital for syncope and a 3-month history of progressive dyspnoea on exertion. On admission, the patient was diagnosed with severe aortic stenosis (AS) and progressive heart failure. In spite of optimal medical therapy, she remained dyspnoeic (New York Heart Association functional class IV). She was subsequently transferred to the hospital for further management.

INVESTIGATIONS

A transthoracic echocardiography showed presence of severe AS with reduced left ventricular systolic

function (peak aortic jet velocity=5.5 m/s; aortic valve area=0.23 cm²; ejection fraction=20%). Coronary angiography revealed normal results. Her brain natriuretic peptide (BNP) level was 7135 pg/mL. The Society of Thoracic Surgeons 30-day predicted risk of mortality score calculated for surgical aortic valve replacement was 23%. In view of this, she was deemed an extremely high-risk patient for surgical aortic valve replacement, and a TAVI via transfemoral approach was conducted.

TREATMENT

TAVI was performed under general anaesthesia using transesophageal echocardiography (TEE). Venoarterial extracorporeal membrane oxygenation (ECMO) support was initiated via left femoral artery and right femoral vein cannulation as haemodynamic instability was encountered during the induction phase of anaesthesia. Four thousand units of heparin were administered before starting the ECMO support. Activated clotting time was monitored at 30 min intervals and maintained above 300 s during the procedure. Balloon valvuloplasty was performed using a 20 mm balloon catheter under rapid right ventricular pacing at 200 bpm. The patient became hypotensive following predilation, and TEE showed severe aortic regurgitation (AR) with severely depressed left ventricular wall motion. Aortography also demonstrated severe AR and abrupt coronary occlusion in the mid-left anterior descending (LAD) artery ([figure 1A](#)).

A 26 mm Sapien XT valve (Edwards Lifesciences, California, USA) was deployed under rapid pacing at 200 bpm. TEE showed that the valve was implanted in the correct position, with no perivalvular leakage. However, the left ventricular wall motion was still severely depressed. Selective coronary angiography using a 7 Fr FL4 guiding catheter showed occlusion of the mid-LAD artery ([figure 1B](#)). After crossing a 0.014" guidewire, an intravascular ultrasound (IVUS) examination was performed, which showed that the mid-LAD artery was occluded by low echoic thrombus without calcified components. Plaque composition on arterial wall was not significant in the IVUS examination ([video 1](#)). Manual vacuum aspiration using the Eliminate aspiration catheter (Terumo, Tokyo, Japan) was then conducted several times. Multiple emboli were removed ([figure 2A](#)), and the LAD artery was recanalised; however, there was still severe stenosis caused by the remaining emboli. After balloon dilation of the occluded segment, two drug-eluting

CrossMark

To cite: Tsujimura A, Saito N, Minakata K, et al. *BMJ Case Rep* Published online: [please include Day Month Year] doi:10.1136/bcr-2016-216620

Figure 1 Fluoroscopy images during the procedure. (A) Aortography just after predilation showing severe aortic regurgitation and abrupt coronary occlusion in the mid-left anterior descending (LAD) artery (arrow). Selective coronary angiography after valve implantation showing coronary occlusion in the mid-LAD artery (arrow) (B). Final coronary angiography after successful implantation of two drug-eluting stents (C).

Video 1 Intravascular ultrasound imaging of the occluded left anterior descending artery. The artery is filled with low echoic thrombus. Plaque composition on arterial wall is not significant.

stents were deployed using the culotte stent technique: a 3.0×38 mm Promus PREMIER drug-eluting stent (Boston Scientific, Massachusetts, USA) in the first diagonal branch, and a 3.0×24 mm Promus PREMIER stent in the LAD artery (figure 1C). Following this, the left ventricular wall motion improved and the hemodynamics stabilised. ECMO support was weaned

off, and the procedures were completed. No further embolic events occurred postoperatively.

OUTCOME AND FOLLOW-UP

The patient's failure symptoms improved remarkably after TAVI, and her BNP level decreased to 382 pg/mL. Histopathological examination of the aspirated emboli was compatible with a recent thrombus containing platelets, fibrin, erythrocytes and leucocytes. There were no calcified components in the aspirated thrombi (figure 2B, C).

DISCUSSION

TAVI has recently emerged as an effective therapeutic option for patients with symptomatic severe AS. Coronary artery obstruction after TAVI is rare, but can be a fatal complication.¹ The mechanism of coronary obstruction during TAVI is usually the displacement of a native valve cusp over the coronary ostium.¹ Distal coronary embolisation during TAVI is very uncommon. Freixa *et al*² described a case of distal right coronary artery occlusion post-TAVI that was diagnosed after the patient developed chest pain in the recovery room. However, their study lacked histopathological analysis as the embolus could not be aspirated. It was speculated that the embolus might have been a detached aortic valve leaflet. In our case, histopathological examination was available; the embolus in our patient was a recent thrombus that contained erythrocytes, leucocytes and fibrin, which was considered to be formed during the procedure. To the best of our knowledge, this is the first report to describe a distal coronary embolisation during TAVI with histopathological confirmation of the embolus.

Figure 2 Macroscopic and microscopic findings of the aspirated emboli. (A) Gross findings of the aspirated emboli. (B) and (C) microscopic findings of the aspirated emboli compatible with a recent thrombus. Leucocytes, erythrocytes and rich fibrin formation is observed. No calcified components are observed (C).

Learning points

- ▶ Distal coronary embolisation is a rare but possible complication during transcatheter aortic valve implantation (TAVI).
- ▶ TAVI operators should be aware that distal coronary embolisation during TAVI can occur even when activated clotting time is maintained above 300 s, and no thrombus is detected by transesophageal echocardiography.
- ▶ Rapid percutaneous coronary intervention is the first treatment of choice when coronary embolisation occurs.

Acknowledgements The authors would like to thank all the members of the TAVI heart team in Kyoto University Hospital, including Fumiaki Fukumura, Kyokun Uehara, Kazuhisa Sakamoto, Toshiaki Toyota, Taro Nakatsu, Hirotoshi Watanabe, Tomohiko Taniguchi, Junichi Tazaki, Masao Imai and Hiroki Daijo.

Contributors NS was involved in design of the work, data collection, data analysis and interpretation, drafting of the article, critical revision of the manuscript and final approval of the version to be published. AT was involved in data collection, data analysis and interpretation, drafting and critical revision of the manuscript and final approval of the version to be published. KM and TK were involved in drafting, critical revision of the manuscript and final approval of the version to be published.

Competing interests None declared.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Ribeiro HB, Nombela-Franco L, Urena M, *et al.* Coronary obstruction following transcatheter aortic valve implantation: a systematic review. *JACC Cardiovasc Interv* 2013;6:452–61.
- 2 Freixa X, Bonan R, Asgar AW. Unusual coronary occlusion post transcatheter aortic implantation: the importance of clinical assessment. *Can J Cardiol* 2013;29:1014. e5–6.

Copyright 2016 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow