Published in final edited form as:

J Neurovirol. 2016 April ; 22(2): 149–158. doi:10.1007/s13365-015-0379-2.

Effects of HIV and childhood trauma on brain morphometry and neurocognitive function

Georgina Spies¹, Fatima Ahmed-Leitao¹, Christine Fennema-Notestine^{3,4}, Mariana Cherner³, and Soraya Seedat^{1,2}

¹South African Research Chairs Initiative (SARCHI), PTSD program, Department of Psychiatry, Stellenbosch University, South Africa

²MRC Unit on Anxiety and Stress Disorders, Department of Psychiatry, Stellenbosch University, South Africa

³Department of Psychiatry, University of California San Diego, La Jolla, CA, USA

⁴Department of Radiology, University of California San Diego, La Jolla, CA, USA

Abstract

A wide spectrum of neurocognitive deficits characterise HIV infection in adults. HIV infection is additionally associated with morphological brain abnormalities affecting neural substrates that subserve neurocognitive function. Early life stress (ELS) also has a direct influence on brain morphology. However, the combined impact of ELS and HIV on brain structure and neurocognitive function has not been examined in an all-female sample with advanced HIV disease. The present study examined the effects of HIV and childhood trauma on brain morphometry and neurocognitive function. Structural data were acquired using a 3T Magnetom MRI scanner and a battery of neurocognitive tests was administered to 124 women; HIV positive with ELS ($n = 32$), HIV positive without ELS ($n = 30$), HIV negative with ELS ($n = 31$), HIV negative without ELS ($n = 31$). Results revealed significant group volumetric differences for right anterior cingulate cortex (ACC), bilateral hippocampi, corpus callosum, left and right caudate, and left and right putamen. Mean regional volumes were lowest in HIV positive women with ELS compared to all other groups. Although causality cannot be inferred, findings also suggest that alterations in the left frontal lobe, right ACC, left hippocampus, corpus callosum, left and right amygdala, and left caudate may be associated with poorer neurocognitive performance in the domains of processing speed, attention/working memory, abstraction/executive functions, motor skills, learning, and language/fluency with these effects more pronounced in women living with both HIV and childhood trauma. This study highlights the potential contributory role of childhood trauma to brain alterations and neurocognitive decline in HIV infected individuals.

Corresponding author: Dr Georgina Spies, Department of Psychiatry, Faculty of Medicine and Health Sciences, Stellenbosch University, PO Box 241, Cape Town, 8000, Ph: 27-21-9389776, Fax: 27-21-9389739, ggiocos@sun.ac.za.

This work was previously presented at the 27th ECNP Congress, October 18–21 2014, Berlin, Germany.

We have no conflict of interest to declare.

Keywords

HIV; childhood trauma; MRI; brain volumetrics; Freesurfer; neurocognition

Introduction

In 2014, an estimated 5.51 million South Africans were infected with human immunodeficiency virus (HIV), and for women aged 15–49, an estimated 18.5% of the population was HIV-positive (Statistics South Africa 2014). Among South African women, high rates of intimate partner violence, rape, and childhood abuse have been reported (Andersson et al. 2008; Jewkes et al. 2001; Jewkes et al. 2002; Kalichman and Simbayi 2004). Many women infected with HIV in adulthood also have developmental trajectories characterised by trauma. Such trauma has been associated with poor neurocognitive functioning and other neurologic and psychiatric sequelae (Malan-Muller et al. 2013; Spies et al. 2012a; Spies et al. 2012b; Troeman et al. 2011). The brain and behavioral changes that attend early life trauma may, in this vulnerable population, result in additional complications related to the course of HIV disease. However, the impact that early life stress [ELS] may have on the presentation and neuropathology of HIV has received very little attention.

HIV penetrates the blood-brain-barrier early in the course of infection and can be highly neurovirulent (Valcour et al. 2012), infecting nerve cells and resulting in neuroinflammation and neuronal death (Kumar et al. 2009). HIV infected individuals commonly have neurocognitive deficits characterised as HIV-associated neurocognitive disorders (HAND) (Antinori et al. 2007). In addition, morphological brain abnormalities are common (Holt et al. 2012; Letendre et al. 2009) and these abnormalities may be similar across various genetic clades of the virus (Ortega et al. 2013). Studies have reported volumetric, functional, and metabolite abnormalities in HIV-infected individuals most commonly within frontal-striatal brain regions (Ances et al. 2006; Aylward et al. 1993; Becker et al. 2011; Cohen et al. 2010b; Jernigan et al. 1993; Paul et al. 2002; Towgood et al. 2012), and these abnormalities have been linked to poor cognitive performance (Castelo et al. 2006; Cohen et al. 2010a; Harezlak et al. 2011; Melrose et al. 2008; Moore et al. 2006; Paul et al. 2008b). Risk for significant brain involvement, as manifested by both neuroimaging abnormalities and neurocognitive impairments, is higher with more advanced disease stage and with greater immunocompromise (e.g., lower nadir CD4) (Centers for Disease Control and Prevention 1993; Ellis et al. 2011; Heaton et al. 1995). Even with effective antiretroviral therapy (ART) individuals who are HIV-infected continue to demonstrate ongoing aberrations in white and grey matter, as demonstrated by recent neuroimaging studies (Becker et al. 2011; Becker et al. 2012; Cardenas et al. 2009; Cohen et al. 2010a; Cohen et al. 2010b; Jernigan et al. 2011). Recent evidence further suggests that increases in brain white matter and subcortical gray matter abnormalities are linked to CD4 recovery among infected individuals (Fennema-Notestine et al. 2013). The continued presence of brain abnormalities and neurocognitive impairments, despite improvements in treatment, suggest ongoing low grade infection and/or chronic inflammation.

A growing body of evidence indicates that ELS can have pathological brain effects (Andersen et al. 2008; Choi et al. 2009; Cohen et al. 2006; Mehta et al. 2009; Paul et al. 2008a; Seckfort et al. 2008; Stein et al. 1997; Teicher et al. 2004; Tottenham et al. 2010) in otherwise healthy individuals. In clinical samples (e.g. PTSD and major depression) studies of ELS have similarly identified stress-related structural, functional, and metabolite alterations in individuals who have suffered adverse childhood events, such as abuse (Bremner et al. 2003; Bremner et al. 1997; De Bellis et al. 2000; Vythilingam et al. 2002), as well as events in adulthood (e.g. intimate partner violence, IPV) (Fennema-Notestine et al. 2002; Seedat et al. 2005). These alterations include structural abnormalities in the amygdala, hippocampus, corpus callosum, anterior cingulate cortex, and caudate nuclei in individuals with ELS, and in supratentorial cranial vaults in individuals with IPV. Studies in HIV-infected individuals have reported high rates of childhood emotional, physical, and sexual abuse (Allers and Benjack 1991; Cohen et al. 2000; Walton et al. 2011) and ELS may increase the risk for HAND or compound HIV-related neural abnormalities.

To the best of our knowledge, only a single cross-sectional study has directly examined the combined impact of HIV infection and ELS on brain volumes and associated neurocognitive functioning. Clark et al. studied four groups of individuals with and without HIV-infection and/or trauma histories in the United States, where clade B HIV predominates and the majority of individuals are on antiretroviral treatment (Clark et al. 2012). Their findings supported an interaction between HIV status and ELS severity, such that larger amygdala volumes were evident in HIV-infected individuals with high levels of ELS. These abnormalities were associated with neurocognitive dysfunction (Clark et al. 2012). The aforementioned study examined relatively complex groups that differed with respect to education, current stress, depression, lifetime cocaine use, and HCV status, although these variables did not appear to account for the interaction of neurocognitive and neural findings.

It is against this background that the present study investigated the combined effects of HIV infection and childhood trauma on brain structure and neurocognitive performance in an all-female cohort infected with clade C HIV, less than half of whom are on treatment. Our hypothesis was two-fold; firstly that women with the ‘double hit’ of HIV infection and childhood trauma would have significantly altered brain volumes in regions typically impacted by HIV and ELS (caudate, putamen, hippocampus, amygdala, corpus callosum, and anterior cingulate) compared to HIV-positive women without childhood trauma. Secondly, we hypothesized that alterations in brain volumes in these regions in dually affected women would be associated with poorer neurocognitive performance.

Methods

Participants

A total of 124 women tested for HIV status were included; 62 were HIV-positive, 32 with a history of childhood trauma and 30 without, and 62 were HIV-negative, 31 with a history of childhood trauma and 31 without.

Eligibility criteria included: willingness and ability to provide written informed consent, ability to read and write in either English, Afrikaans (a West Germanic language spoken

natively in South Africa), or isiXhosa (one of the indigenous African languages spoken natively in South Africa) at 5th grade level, between 18 and 65 years of age, medically well enough to undergo neuropsychological testing and MRI scanning. Exclusions were: MRI contraindications such as pregnancy or metal in the body, a current or past history of schizophrenia, bipolar disorder or other psychotic disorders, history of substance abuse or dependence, significant previous head injury, demonstrated cognitive impairment on the HIV Dementia Scale, current seizure disorders of any cause, history of CNS infections or neoplasms, hepatitis C positive status, and current use (within the past month) of any psychotropic medication (including antidepressants).

Procedure

The study was approved by the ethics committee of Stellenbosch University, Cape Town, South Africa. All women included in the present study were tested for HIV status at their local health care facility, with HIV status confirmed by means of ELISA. Sixty two (44.6%) women were HIV-positive. Participants were recruited through community health care facilities (VCT sites and HIV units) in and around the Cape metropole of South Africa from 2008–2015 by a researcher or with the help of doctors and adherence counsellors. Recruitment procedures did not differ across groups. All participants who consented were screened for eligibility and childhood trauma exposure either in person at their clinic or telephonically. Those who met initial screening criteria subsequently underwent neuromedical, neuropsychiatric, neurocognitive, and neuroimaging assessments. Participants were reimbursed for travel costs to and from assessments.

Measures

Age, gender, marital status, ethnicity, years of education, and employment status were captured. A comprehensive history was obtained from, and a general physical examination conducted in, all patients. Virologic markers of disease progression (CD4 lymphocyte count and viral loads) were obtained from blood samples.

The Childhood Trauma Questionnaire Short Form (CTQ-SF) was used to retrospectively elucidate trauma exposure and to categorise HIV-positive and HIV-negative women into trauma (score of 41 or higher on the CTQ indicating moderate to extreme trauma) and non-trauma exposure groups (score of 25–40 on the CTQ). The CTQ-SF is a 28-item self-report inventory that provides valid screening for histories of abuse and neglect (up until the age of 18). It assesses five types of maltreatment including, emotional, physical, and sexual abuse, and emotional and physical neglect. Scores range from 25–125, with higher scores indicating higher levels of trauma exposure during childhood (Bernstein et al. 2003). The Cronbach alpha for the CTQ was excellent ($\alpha = 0.83$).

Neurocognitive function was measured with 10 tests that have been used commonly in HIV research and cover 7 ability domains (learning, delayed recall, processing speed, attention/working memory, executive function, verbal fluency, motor ability) (Heaton et al. 2010). Test instructions and stimuli were adapted as necessary to fit the cultural context of South Africa and translated into Afrikaans and isiXhosa using standard test adaptation techniques. Tests were administered in English, Afrikaans or isiXhosa according to the participants' self

report of best language. We have previously reported group differences using this test battery (Spies et al. 2012a).

Neuroimaging Data acquisition and analysis

Data were acquired using a MAGETOM Siemens Allegra syngo MR 3 Tesla scanner, with a four channel head coil. The following imaging parameters were used: TR = 2530ms; TE = 1.53ms; TI = 1100ms; flip angle = 7 degrees; voxel size = 1.3×1.0×1.3mm; field of view = 256mm. Region specific grey matter volumetric measurements were acquired through Freesurfer software which is publicly available and includes probabilistic volumetric segmentation for white matter and subcortical regions and cortical surface reconstruction and parcellation for neocortical measures (Dale et al. 1999; Desikan et al. 2006; Fischl et al. 1999; Fischl and Dale 2000; Fischl et al. 2002; Fischl et al. 2004). The software was used in the completely automated fashion and quality control was implemented throughout the process. The processing pipeline is computationally intensive and as such, we utilised custom batching scripts on the Centre for High Performance Computing (CHPC) in Cape Town. All data were analysed using SPSS for Windows (version 22) and Statistica (version 12). Descriptive statistics were computed for all demographic and clinical information. We used analysis of variance (ANOVA) to assess group differences in estimated intracranial vault, age, education and brain volumes. Estimated intracranial vault was added as a covariate to control for individual differences in head size. Analysis of covariance (ANCOVA) using age and estimated intracranial vault as covariates was subsequently undertaken. Pearson's correlations assessed for associations between brain volumes and neurocognitive function as well as for associations between estimated intracranial vault and trauma subtypes on the CTQ. Homogeneity of slopes ANCOVA was conducted with regional brain volumes as dependent variables, group and neurocognitive function as independent variables. The interaction variable (group by neurocognitive outcomes) tested the relationships (slopes) between brain volumes and neurocognitive outcomes across the groups. Estimated intracranial vault, age, and education were controlled for as covariates in the models. The significance level was two sided and set at 0.05 respectively. Post-hoc tests (Fisher's LSD and Bonferroni) were conducted to correct for multiple comparisons.

Results

Demographic characteristics

The mean age of participants was 30.3 years (minimum of 18 years, maximum of 50 years). The mean level of education was 10.3 years (minimum of 5 years, maximum of 14 years). The groups significantly differed in age and education, with HIV-negative participants younger and more educated than HIV-positive counterparts. The majority of the women were Black (86.3%), Xhosa speaking (85.6%), and right handed (84.2%). Further demographic details are presented in Table 1.

Clinical characteristics

Of the HIV-positive women, 30 (48.4%) reported being on antiretroviral treatment (ART). The mean CD4 lymphocyte count was 431.20 (minimum of 25 and a maximum of 1529). The mean viral load was 116172.08, ranging from lower than the detectable limit to

3200000 copies/ml. There were no significant group differences with respect to clinical variables such as CD4 lymphocyte count and viral load.

Childhood Trauma

Sixty one participants reported no childhood trauma. Among the HIV infected women ($n = 62$), a total of 32 reported histories of childhood trauma, one in the low to moderate range (41–55), 11 (17.7%) in the moderate to severe range (56–72) and 20 (32.3%) in the severe to extreme range (73–125). Among the HIV uninfected women ($n = 62$), a total of 31 reported histories of childhood trauma, 20 (32.3%) in the mild to moderate range (41–55), 7 (11.3%) in the moderate to severe range (56–72) and 4 (6.45%) in the severe to extreme range (73–125). The overall mean score on the CTQ was 49.34 with a minimum of 25 and a maximum of 116. The most commonly reported childhood trauma type was emotional abuse with a mean of 11.55, followed by emotional neglect ($M = 11.34$), physical neglect ($M = 9.76$), physical abuse ($M = 9.30$) and lastly, sexual abuse ($M = 7.38$). See table 4. Pearson's correlation assessed the association between intracranial vault volume (ICV) and CTQ sub-scale scores. ICV was significantly associated with childhood physical neglect ($r = 0.195$, $p = 0.030$).

Regional volumetric differences between groups

Brain volumes of the following regions were significantly different among the four groups as revealed by ANCOVA, using age and intracranial vault as covariates: right ACC ($p < 0.05$), bilateral hippocampi, corpus callosum, left and right caudate, and left and right putamen ($p < 0.01$). Mean brain volume was significantly smaller in HIV-positive women with childhood trauma, compared to the other three groups, for all the abovementioned regions. For the left hippocampus, mean brain volume was smallest in HIV-positive women with childhood trauma, followed by HIV-negative women with childhood trauma. Mean brain volume of the left amygdala and right amygdala was smallest in the dually affected group although group differences did not reach statistical significance ($p = 0.05$). See table 2.

Associations between regional brain volumes and neurocognition

There were no significant correlations between CD4 lymphocyte count and viral load and any of the neurocognitive variables. However, in the HIV-infected group CD4 lymphocyte count was significantly correlated with left frontal lobe volume ($p = 0.029$). There was no association between viral load and any of the brain regions of interest.

For the whole sample ($N = 124$), homogeneity of slopes ANCOVA controlling for age, education and intracranial vault revealed a number of significant group dependent associations between brain volumes and neurocognitive variables. Significant associations were evident between the left frontal lobe and Trails Making Test A ($p = 0.04$), right ACC and Grooved Pegboard Test ($p = 0.01$) and WAIS-III Digit Symbol ($p = 0.04$), left hippocampus and Hopkins Verbal Learning Test [HVLT] ($p = 0.03$) and Category Fluency Test ($p = 0.04$), corpus callosum and Controlled Oral Word Association Test [COWAT] ($p = 0.01$), WAIS-III Digit Symbol ($p = 0.01$) and WMS-III Spatial Span ($p = 0.01$), left amygdala and WAIS-III Digit Symbol ($p = 0.01$) and Colour Trails 2 ($p = 0.02$), right amygdala and COWAT ($p = 0.04$), left caudate and WAIS-III Digit Symbol ($p = 0.03$),

Stroop Colour Word Test ($p = 0.01$), and Paced Auditory Serial Addition Task [PASAT] ($p = 0.02$), respectively. Mean plots revealed that for all neurocognitive variables listed above, HIV-positive women with trauma performed more poorly (i.e. had lower mean scores or higher mean scores on tests where higher scores indicate poorer performance) than their HIV-positive counterparts without trauma and their HIV-negative counterparts with and without trauma (see table 3).

Discussion

The results of the present study demonstrated significant group volumetric differences for the right ACC, bilateral hippocampi, corpus callosum, left and right caudate, and left and right putamen. Mean volumes of the abovementioned regions were smallest in HIV-positive women with childhood trauma compared to all other groups. Mean volumes of the left and right amygdala were also smallest in HIV-positive women with childhood trauma compared to all other groups, although the probability value did not reach significance. The results of the present study suggest that in the dually affected group (HIV and childhood trauma), brain volumes were significantly smaller, compared to women singly affected by HIV. This finding supports our primary hypothesis that women with HIV and childhood trauma would have significantly reduced regional brain volumes (striatum, hippocampus, amygdala, corpus callosum, and anterior cingulate) compared to HIV-infected women without childhood trauma. This finding is consistent with previous studies. Research conducted in single risk groups suggest that both HIV and childhood trauma separately result in reduced volumes in regions such as the ACC, hippocampus, corpus callosum, amygdala, and caudate nuclei (Ances et al. 2012; Ances et al. 2012; Andersen et al. 2008; Andersen et al. 2008; Bremner et al. 2003; Bremner et al. 1997; Cohen et al. 2006; Cohen et al. 2006; Cohen et al. 2010a; Cohen et al. 2010a; Cohen et al. 2010b; Cohen et al. 2010b; De Bellis et al. 2000; Paul et al. 2008b; Seckfort et al. 2008; Stein et al. 1997; Teicher et al. 2004; Vythilingam et al. 2002). In contrast to the only available study by Clark and colleagues, who did not find any significant volumetric differences in regions of interest such as the ACC, hippocampus, caudate and putamen, we found evidence for smaller brain volumes in the right ACC, bilateral hippocampi, corpus callosum, left and right caudate, and left and right putamen among dually affected women. Although only a trend was evident, we also found that mean volumes of the left and right amygdala were smaller in HIV-positive women with ELS. In contrast, Clark and colleagues reported larger amygdala volumes in HIV-positive individuals with high levels of ELS in their study (Clark et al. 2012) which differed from our own with regards to HIV disease severity and the proportion of individuals on treatment. In the Clark et al. study (Clark et al. 2012) the majority of HIV-infected participants were on antiretroviral treatment and had higher CD4 lymphocyte counts suggestive of milder disease. The mode of transmission (homosexual or heterosexual) was not reported and the sample consisted of both males and females. The dually affected group (HIV+ with high ELS) was also smaller in size than in the present study. In the present study, HIV transmission was heterosexual, less than half of the sample was on antiretroviral therapy and the lower group mean CD4 lymphocyte count is suggestive of greater disease severity. The present study also included an all-female sample. The aforementioned differences between the two studies make comparisons difficult, thus, future studies investigating the volumetric effects of the

dual burden of HIV and ELS on the amygdala in particular are needed. In the present study, perhaps greater childhood trauma severity may have rendered the trend in amygdala differences between groups significant.

Results of the present study suggest that intracranial vault (ICV) was significantly associated with physical neglect on the CTQ. A significant positive correlation was found between ICV and childhood physical neglect. Moreover, although just shy of significance ($p = 0.05$), a positive correlation was also evident between ICV and childhood physical abuse. This finding is to some extent similar to a previous study although there are differences in the direction of the relationship (Fennema-Notestine et al. 2002). In that study, supratentorial cranial vault was negatively correlated with the severity of childhood physical abuse in a sample of female victims of IPV (Fennema-Notestine et al. 2002). Although there was no significant difference in ICV between the groups in the current study, a larger mean ICV was evident in both the dually affected group (HIV+ trauma) and the traumatised control group (HIV- trauma), compared with the HIV-positive and HIV-negative nontraumatised groups. In addition, the mean score for childhood physical neglect and abuse was highest in the dually affected group, followed by the HIV uninfected group with ELS. Thus, it may be possible that increased ICV in these women may be a neurodevelopmental epiphenomenon, perhaps as a result of increased levels of childhood physical abuse/neglect. However, causal inferences are not possible and this hypothesis needs to be examined in future work.

Associations between regional brain volumes and neurocognitive variables (grouped by domain) revealed significant associations between the left frontal lobe and speed of information processing, right ACC, motor skills and processing speed, left hippocampus, learning and language/verbal fluency, corpus callosum, language/verbal fluency, processing speed and attention/working memory, left amygdala, processing speed and abstraction/executive functioning, right amygdala and language/verbal fluency, left caudate, processing speed, abstraction/executive functioning and attention/working memory, respectively. Women singly affected by HIV performed more poorly on the majority of these tasks than their HIV-negative counterparts, except on three tasks measuring processing speed, attention/working memory and motor skills (WAIS-III Digit Symbol, Trail Making Test A, and Grooved Pegboard Test). For motor skills and attention/working memory (Trail Making Test A, and Grooved Pegboard Test), women dually affected (HIV-positive with trauma) performed the worst, followed by HIV-negative women with trauma. These findings provide evidence for the potential contributory role of childhood trauma to brain alterations and neurocognitive decline in HIV-infected individuals. Mean scores revealed that dually affected women (HIV-positive with childhood trauma) performed the worst across all three groups on all except one task of verbal fluency (COWAT), on which HIV-positive women without trauma had a slightly lower mean score. The finding that dually affected women performed more poorly on these tasks than all other groups provides support for our second hypothesis that among those dually affected, brain volumetric differences would be consistent with poorer neurocognitive functioning.

HIV is frequently associated with deficits in higher order brain function, including deficits in memory (learning), psychomotor speed, executive functions, and attention are frequent (Grant 2008). Studies have reported a link between neurocognitive impairment and

alterations in brain structure, function, and metabolite concentrations in HIV-infected individuals (Castelo et al. 2006; Cohen et al. 2010a; Harezlak et al. 2011; Melrose et al. 2008; Moore et al. 2006; Paul et al. 2008b). Similarly, studies have documented trauma related neurocognitive impairments in executive functions, memory, intellectual development, language, verbal learning, spatial working memory, and psychomotor speed (Choi et al. 2009; Clark et al. 2003; Jelinek et al. 2006; Lagarde, Doyon, Brunet 2010; Majer et al. 2010; Palmer et al. 1997). Larger amygdala volumes in HIV-infected individuals with high levels of ELS were recently reported to be associated with greater impairment on the WAIS-III Digit Symbol Test, a test of processing speed (Clark et al. 2012). In keeping with this, the results of the present study also revealed an association between the amygdala and the WAIS-III Digit Symbol Test, a test of speed of information processing. Although there was no significant difference between groups on amygdala volume (only a trend), mean amygdala volume was smallest in the dually affected group (HIV+ with trauma). Moreover, performance on the WAIS-III Digit Symbol Test was poorest in this group compared to all other groups. The findings of both Clark et al. and the present study therefore suggest that HIV and ELS interact to influence amygdala volume, which is associated with neurocognitive dysfunction in individuals infected with HIV. However, as previously alluded to, this hypothesis warrants further investigation given the discrepancies between these two studies and the paucity of research on this topic. In addition to the amygdala findings, the present study also found evidence for an association between the left frontal lobe, right ACC, corpus callosum and left caudate and processing speed. The dually affected group had lower scores (i.e. worse performance) on tests of processing speed compared to all other groups and this was significantly associated with the above-mentioned brain regions.

A limitation of the present study was the retrospective assessment of childhood trauma and its contribution to recall bias. A prospective study will better explain the effects of childhood trauma on the brain. Moreover, the CTQ-SF does not include any questions around whether the abuse reported was single isolated event(s) or repeated exposures. This would be important to elucidate going forward. There was a difference in antiretroviral therapy among groups with more women on antiretroviral treatment in the trauma group compared with infected women without trauma. The influence this difference in treatment status has on findings needs further exploration. Moreover, the length of antiretroviral treatment and time since HIV infection was also not taken into account and should be explored in follow-up studies. Stratification of groups by these variables would require large sample sizes. Despite these limitations, this is to our knowledge only the second study to provide evidence of a combined impact of HIV and childhood trauma on brain volumetrics and associated neurocognitive performance and the first to do so in an all-female cohort with more advanced disease.

Conclusion

The present study found evidence for smaller volumes in the right ACC, bilateral hippocampi, corpus callosum, left and right caudate, and left and right putamen in HIV-infected women with childhood trauma. Although causality cannot be inferred, findings also suggest that alterations in the left frontal lobe, right ACC, left hippocampus, corpus callosum, left and right amygdala, and left caudate may be associated with poorer

neurocognitive performance in the domains of processing speed, attention/working memory, abstraction/executive functions, motor skills, learning, and language/fluency with these effects more pronounced in women living with both HIV and childhood trauma. This highlights the potential contributory role of developmental trauma in brain morphological and neurocognitive aberrations in HIV-infected individuals. Future longitudinal studies that track the course of neurocognitive and brain volumetric changes are needed to provide more clarity.

Acknowledgments

This research was supported by the South African Research Chair in PTSD hosted by Stellenbosch University, funded by the DST and administered by NRF, and the Faculty of Medicine and Health Sciences, Stellenbosch University (Deputy Dean's strategic fund for postdoctoral fellows). Additional research support was provided by the HIV Neurobehavioral Research Center (HNRC; National Institutes of Health Grant P30-MH62512). We would like to thank the Centre for High Performance Computing, Council for Scientific and Industrial Research, Cape Town for storage, processing and technical support. Professor Martin Kidd from the department of Statistics and Actuarial Sciences of Stellenbosch University provided statistical assistance.

References

- Allers CT, Benjack KJ. Connections between childhood abuse and HIV infection. *Journal of Counseling & Development*. 1991; 70:309–313.
- Ances BM, Ortega M, Vaida F, Heaps J, Paul R. Independent effects of HIV, aging, and HAART on brain volumetric measures. *J Acquir Immune Defic Syndr*. 2012; 59:469–477. [PubMed: 22269799]
- Ances BM, Roc AC, Wang J, Korczykowski M, Okawa J, Stern J, Kim J, Wolf R, Lawler K, Kolson DL, Detre JA. Caudate blood flow and volume are reduced in HIV+ neurocognitively impaired patients. *Neurology*. 2006; 66:862–866. [PubMed: 16567703]
- Andersen SL, Tomada A, Vincow ES, Valente E, Polcari A, Teicher MH. Preliminary evidence for sensitive periods in the effect of childhood sexual abuse on regional brain development. *J Neuropsychiatry Clin Neurosci*. 2008; 20:292–301. [PubMed: 18806232]
- Andersson N, Cockcroft A, Shea B. Gender-based violence and HIV: Relevance for HIV prevention in hyperendemic countries of southern africa. *Aids*. 2008; 22:73–86.
- Antinori A, Arendt G, Becker JT, Brew BJ, Byrd DA, Cherner M, Clifford DB, Cinque P, Epstein LG, Goodkin K, et al. Updated research nosology for HIV-associated neurocognitive disorders. *Neurology*. 2007; 69:1789–1799. [PubMed: 17914061]
- Aylward EH, Henderer JD, McArthur JC, Brettschneider PD, Harris GJ, Barta PE, Pearlson GD. Reduced basal ganglia volume in HIV-1-associated dementia: Results from quantitative neuroimaging. *Neurology*. 1993; 43:2099–2104. [PubMed: 8413973]
- Becker JT, Maruca V, Kingsley LA, Sanders JM, Alger JR, Barker PB, Goodkin K, Martin E, Miller EN, Ragin A, et al. Factors affecting brain structure in men with HIV disease in the post-HAART era. *Neuroradiology*. 2012; 54:113–121. [PubMed: 21424708]
- Becker JT, Sanders J, Madsen SK, Ragin A, Kingsley L, Maruca V, Cohen B, Goodkin K, Martin E, Miller EN, et al. Subcortical brain atrophy persists even in HAART-regulated HIV disease. *Brain Imaging Behav*. 2011; 5:77–85. [PubMed: 21264551]
- Bernstein DP, Stein JA, Newcomb MD, Walker E, Pogge D, Ahluvalia T, Stokes J, Handelsman L, Medrano M, Desmond D, Zule W. Development and validation of a brief screening version of the childhood trauma questionnaire. *Child Abuse Negl*. 2003; 27:169–190. [PubMed: 12615092]
- Bremner JD, Vythilingam M, Vermetten E, Southwick SM, McGlashan T, Nazeer A, Khan S, Vaccarino LV, Soufer R, Garg PK. MRI and PET study of deficits in hippocampal structure and function in women with childhood sexual abuse and posttraumatic stress disorder. *Am J Psychiatry*. 2003; 160:924–932. [PubMed: 12727697]
- Bremner JD, Randall P, Vermetten E, Staib L, Bronen RA, Mazure C, Capelli S, McCarthy G, Innis RB, Charney DS. Magnetic resonance imaging-based measurement of hippocampal volume in

- posttraumatic stress disorder related to childhood physical and sexual abuse - a preliminary report. *Biol Psychiatry*. 1997; 41:23–32. [PubMed: 8988792]
- Cardenas VA, Meyerhoff DJ, Studholme C, Kornak J, Rothlind J, Lampiris H, Neuhaus J, Grant RM, Chao LL, Truran D, et al. Evidence for ongoing brain injury in human immunodeficiency virus-positive patients treated with antiretroviral therapy. *J Neurovirol*. 2009; 15:324–333. [PubMed: 19499454]
- Castelo JM, Sherman SJ, Courtney MG, Melrose RJ, Stern CE. Altered hippocampal-prefrontal activation in HIV patients during episodic memory encoding. *Neurology*. 2006; 66:1688–1695. [PubMed: 16769942]
- Centers for Disease Control and Prevention. [Accessed 07 Jun 2015] Revised Classification System for HIV Infection and Expanded Surveillance Case Definition for AIDS Among Adolescents and Adults. 1993. Available from: <http://www.cdc.gov/mmwr/preview/mmwrhtml/00018871.htm>
- Choi J, Jeong B, Rohan ML, Polcari AM, Teicher MH. Preliminary evidence for white matter tract abnormalities in young adults exposed to parental verbal abuse. *Biol Psychiatry*. 2009; 65:227–234. [PubMed: 18692174]
- Clark CR, McFarlane AC, Morris P, Weber DL, Sonkilla C, Shaw M, Marcina J, Tochon-Danguy HJ, Egan GF. Cerebral function in posttraumatic stress disorder during verbal working memory updating: A positron emission tomography study. *Biol Psychiatry*. 2003; 53:474–481. [PubMed: 12644352]
- Clark US, Cohen RA, Sweet LH, Gongvatana A, Devlin KN, Hana GN, Westbrook ML, Mulligan RC, Jerskey BA, White TL, et al. Effects of HIV and early life stress on amygdala morphometry and neurocognitive function. *J Int Neuropsychol Soc*. 2012; 18:657–668. [PubMed: 22621973]
- Cohen M, Deamant C, Barkan S, Richardson J, Young M, Holman S, Anastos K, Cohen J, Melnick S. Domestic violence and childhood sexual abuse in HIV-infected women and women at risk for HIV. *Am J Public Health*. 2000; 90:560–565. [PubMed: 10754970]
- Cohen RA, Grieve S, Hoth KF, Paul RH, Sweet L, Tate D, Gunstad J, Stroud L, McCaffery J, Hitsman B, et al. Early life stress and morphometry of the adult anterior cingulate cortex and caudate nuclei. *Biol Psychiatry*. 2006; 59:975–982. [PubMed: 16616722]
- Cohen RA, Harezlak J, Schifitto G, Hana G, Clark U, Gongvatana A, Paul R, Taylor M, Thompson P, Alger J, et al. Effects of nadir CD4 count and duration of human immunodeficiency virus infection on brain volumes in the highly active antiretroviral therapy era. *J Neurovirol*. 2010a; 16:25–32. [PubMed: 20113183]
- Cohen RA, Harezlak J, Gongvatana A, Buchthal S, Schifitto G, Clark U, Paul R, Taylor M, Thompson P, Tate D, et al. Cerebral metabolite abnormalities in human immunodeficiency virus are associated with cortical and subcortical volumes. *J Neurovirol*. 2010b; 16:435–444. [PubMed: 20961212]
- Dale AM, Fischl B, Sereno MI. Cortical surface-based analysis. I. segmentation and surface reconstruction. *Neuroimage*. 1999; 9:179–194. [PubMed: 9931268]
- De Bellis MD, Keshavan MS, Spencer S, Hall J. N-acetylaspartate concentration in the anterior cingulate of maltreated children and adolescents with PTSD. *Am J Psychiatry*. 2000; 157:1175–1177. [PubMed: 10873933]
- Desikan RS, Segonne F, Fischl B, Quinn BT, Dickerson BC, Blacker D, Buckner RL, Dale AM, Maguire RP, Hyman BT, et al. An automated labeling system for subdividing the human cerebral cortex on MRI scans into gyral based regions of interest. *Neuroimage*. 2006; 31:968–980. [PubMed: 16530430]
- Ellis RJ, Badiee J, Vaida F, Letendre S, Heaton RK, Clifford D, Collier AC, Gelman B, McArthur J, Morgello S, et al. CD4 nadir is a predictor of HIV neurocognitive impairment in the era of combination antiretroviral therapy. *Aids*. 2011; 25:1747–1751. [PubMed: 21750419]
- Fennema-Notestine C, Ellis RJ, Archibald SL, Jernigan TL, Letendre SL, Notestine RJ, Taylor MJ, Theilmann RJ, Julaton MD, Croteau DJ, et al. Increases in brain white matter abnormalities and subcortical gray matter are linked to CD4 recovery in HIV infection. *J Neurovirol*. 2013; 19:393–401. [PubMed: 23838849]

- Fennema-Notestine C, Stein MB, Kennedy CM, Archibald SL, Jernigan TL. Brain morphometry in female victims of intimate partner violence with and without posttraumatic stress disorder. *Biol Psychiatry*. 2002; 52:1089–1101. [PubMed: 12460692]
- Fischl B, Dale AM. Measuring the thickness of the human cerebral cortex from magnetic resonance images. *Proc Natl Acad Sci USA*. 2000; 97:11050–11055. [PubMed: 10984517]
- Fischl B, Sereno MI, Dale AM. Cortical surface-based analysis. II: Inflation, flattening, and a surface-based coordinate system. *Neuroimage*. 1999; 9:195–207. [PubMed: 9931269]
- Fischl B, van der Kouwe A, Destrieux C, Halgren E, Segonne F, Salat DH, Busa E, Seidman LJ, Goldstein J, Kennedy D, et al. Automatically parcellating the human cerebral cortex. *Cereb Cortex*. 2004; 14:11–22. [PubMed: 14654453]
- Fischl B, Salat DH, Busa E, Albert M, Dieterich M, Haselgrove C, van der Kouwe A, Killiany R, Kennedy D, Klaveness S, et al. Whole brain segmentation: Automated labeling of neuroanatomical structures in the human brain. *Neuron*. 2002; 33:341–355. [PubMed: 11832223]
- Grant I. Neurocognitive disturbances in HIV. *International Review of Psychiatry*. 2008; 20:33–47. [PubMed: 18240061]
- Harezlak J, Buchthal S, Taylor M, Schifitto G, Zhong J, Daar E, Alger J, Singer E, Campbell T, Yiannoutsos C, et al. Persistence of HIV-associated cognitive impairment, inflammation, and neuronal injury in era of highly active antiretroviral treatment. *Aids*. 2011; 25:625–633. [PubMed: 21297425]
- Heaton RK, Grant I, Butters N, White DA, Kirson D, Atkinson JH, McCutchan JA, Taylor MJ, Kelly MD, Ellis RJ. The HNRC 500-neuropsychology of HIV infection at different disease stages. HIV neurobehavioral research center. *J Int Neuropsychol Soc*. 1995; 1:231–251. [PubMed: 9375218]
- Heaton RK, Cysique LA, Jin H, Shi C, Yu X, Letendre S, Franklin DR Jr, Ake C, Vigil O, Atkinson JH, et al. Neurobehavioral effects of human immunodeficiency virus infection among former plasma donors in rural china. *J Neurovirol*. 2010; 16:185–188. [PubMed: 20443740]
- Holt JL, Kraft-Terry SD, Chang L. Neuroimaging studies of the aging HIV-1-infected brain. *J Neurovirol*. 2012; 18:291–302. [PubMed: 22653528]
- Jelinek L, Jacobsen D, Kellner M, Larbig F, Biesold KH, Barre K, Moritz S. Verbal and nonverbal memory functioning in posttraumatic stress disorder (PTSD). *J Clin Exp Neuropsychol*. 2006; 28:940–948. [PubMed: 16822734]
- Jernigan TL, Archibald S, Hesselink JR, Atkinson JH, Velin RA, McCutchan JA, Chandler J, Grant I. Magnetic resonance imaging morphometric analysis of cerebral volume loss in human immunodeficiency virus infection. the HNRC group. *Arch Neurol*. 1993; 50:250–255. [PubMed: 8442702]
- Jernigan TL, Archibald SL, Fennema-Notestine C, Taylor MJ, Theilmann RJ, Julaton MD, Notestine RJ, Wolfson T, Letendre SL, Ellis RJ, et al. Clinical factors related to brain structure in HIV: The CHARTER study. *J Neurovirol*. 2011; 17:248–257. [PubMed: 21544705]
- Jewkes R, Penn-Kekana L, Levin J, Ratsaka M, Schrieber M. Prevalence of emotional, physical and sexual abuse of women in three south african provinces. *South African Medical Journal*. 2001; 91:421–428. [PubMed: 11455808]
- Jewkes R, Levin J, Mbananga N, Bradshaw D. Rape of girls in south africa. *The Lancet*. 2002; 359:319–320.
- Kalichman SC and Simbayi LC. Sexual assault history and risks for sexually transmitted infections among women in an african township in cape town, south africa. *AIDS Care*. 2004; 16:681–689. [PubMed: 15370057]
- Kumar AM, Fernandez JB, Singer EJ, Commings D, Waldrop-Valverde D, Ownby RL, Kumar M. Human immunodeficiency virus type 1 in the central nervous system leads to decreased dopamine in different regions of postmortem human brains. *J Neurovirol*. 2009; 15:257–274. [PubMed: 19499455]
- Lagarde G, Doyon J, Brunet A. Memory and executive dysfunctions associated with acute posttraumatic stress disorder. *Psychiatry Res*. 2010; 177:144–149. [PubMed: 20381880]
- Letendre SL, Ellis RJ, Everall I, Ances B, Bharti A, McCutchan JA. Neurologic complications of HIV disease and their treatment. *Top HIV Med*. 2009; 17:46–56. [PubMed: 19401607]

- Majer M, Nater UM, Lin JM, Capuron L, Reeves WC. Association of childhood trauma with cognitive function in healthy adults: A pilot study. *BMC Neurol.* 2010; 10:61–70. [PubMed: 20630071]
- Malan-Muller S, Hemmings SM, Spies G, Kidd M, Fennema-Notestine C, Seedat S. Shorter telomere length - A potential susceptibility factor for HIV-associated neurocognitive impairments in south african women. *PLoS One.* 2013; 8(3):e58351. [PubMed: 23472184]
- Mehta MA, Golembo NI, Nosarti C, Colvert E, Mota A, Williams SC, Rutter M, Sonuga-Barke EJ. Amygdala, hippocampal and corpus callosum size following severe early institutional deprivation: The english and romanian adoptees study pilot. *J Child Psychol Psychiatry.* 2009; 50:943–951. [PubMed: 19457047]
- Melrose RJ, Tinaz S, Castelo JM, Courtney MG, Stern CE. Compromised fronto-striatal functioning in HIV: An fMRI investigation of semantic event sequencing. *Behav Brain Res.* 2008; 188:337–347. [PubMed: 18242723]
- Moore DJ, Masliah E, Rippeth JD, Gonzalez R, Carey CL, Cherner M, Ellis RJ, Achim CL, Marcotte TD, Heaton RK, et al. Cortical and subcortical neurodegeneration is associated with HIV neurocognitive impairment. *Aids.* 2006; 20:879–887. [PubMed: 16549972]
- Ortega M, Heaps JM, Joska J, Vaida F, Seedat S, Stein DJ, Paul R, Ances BM. HIV clades B and C are associated with reduced brain volumetrics. *J Neurovirol.* 2013; 19:479–487. [PubMed: 24078556]
- Palmer LK, Armsworth M, Swank PR, Bush G, Frantz C, Copley J. The neuropsychological sequelae of chronically psychologically traumatized children. *Archives of Clinical Neuropsychology.* 1997; 12:379–380.
- Paul R, Cohen R, Navia B, Tashima K. Relationships between cognition and structural neuroimaging findings in adults with human immunodeficiency virus type-1. *Neurosci Biobehav Rev.* 2002; 26:353–359. [PubMed: 12034135]
- Paul R, Henry L, Grieve SM, Guilmette TJ, Niaura R, Bryant R, Bruce S, Williams LM, Richard CC, Cohen RA, et al. The relationship between early life stress and microstructural integrity of the corpus callosum in a non-clinical population. *Neuropsychiatr Dis Treat.* 2008a; 4:193–201. [PubMed: 18728817]
- Paul RH, Ernst T, Brickman AM, Yiannoutsos CT, Tate DF, Cohen RA, Navia BA, ACTG 301 team, ACTG 700 team, HIV MRS Consortium. Relative sensitivity of magnetic resonance spectroscopy and quantitative magnetic resonance imaging to cognitive function among nondemented individuals infected with HIV. *J Int Neuropsychol Soc.* 2008b; 14:725–733. [PubMed: 18764968]
- Seckfort D, Paul R, Grieve S, Vandenberg B, Bryant R, Williams L, Gordon E. Early life stress on brain structure and function across the lifespan: A preliminary study. *Brain Imaging and Behavior.* 2008; 2:49–58.
- Seedat S, Videen JS, Kennedy CM, Stein MB. Single voxel proton magnetic resonance spectroscopy in women with and without intimate partner violence-related posttraumatic stress disorder. *Psychiatry Research: Neuroimaging.* 2005; 139:249–258. [PubMed: 16055312]
- Spies G, Fennema-Notestine C, Archibald SL, Cherner M, Seedat S. Neurocognitive deficits in HIV-infected women and victims of childhood trauma. *AIDS Care.* 2012a; 24:1126–1235. [PubMed: 22672200]
- Spies G, Afifi TO, Archibald SL, Fennema-Notestine C, Sareen J, Seedat S. Mental health outcomes in HIV and childhood maltreatment: A systematic review. *Syst Rev.* 2012b; 1:4053.
- Statistics South Africa. [Accessed 07 Jun 2015] Mid-year population estimates. 2014. Available from: <http://www.statssa.gov.za/publications/P0302/P03022014.pdf>
- Stein MB, Koverola C, Hanna C, Torchia MG, McClarty B. Hippocampal volume in women victimized by childhood sexual abuse. *Psychol Med.* 1997; 27:951–959. [PubMed: 9234472]
- Teicher MH, Dumont NL, Ito Y, Vaituzis C, Giedd JN, Andersen SL. Childhood neglect is associated with reduced corpus callosum area. *Biol Psychiatry.* 2004; 56:80–85. [PubMed: 15231439]
- Tottenham N, Hare TA, Quinn BT, McCarry TW, Nurse M, Gilhooly T, Millner A, Galvan A, Davidson MC, Eigsti IM, et al. Prolonged institutional rearing is associated with atypically large amygdala volume and difficulties in emotion regulation. *Dev Sci.* 2010; 13:46–61. [PubMed: 20121862]
- Towgood KJ, Pitkanen M, Kulasegaram R, Fradera A, Kumar A, Soni S, Sibtain NA, Reed L, Bradbeer C, Barker GJ, et al. Mapping the brain in younger and older asymptomatic HIV-1 men:

Frontal volume changes in the absence of other cortical or diffusion tensor abnormalities. *Cortex*. 2012; 48:230–241. [PubMed: 21481856]

Troeman ZC, Spies G, Cherner M, Archibald SL, Fennema-Notestine C, Theilmann RJ, Spottiswoode B, Stein DJ, Seedat S. Impact of childhood trauma on functionality and quality of life in HIV-infected women. *Health Qual Life Outcomes*. 2011; 9:7525.

Valcour V, Chalermchai T, Sailasuta N, Marovich M, Lerdlum S, Suttichom D, Suwanwela NC, Jagodzinski L, Michael N, Spudich S, et al. Central nervous system viral invasion and inflammation during acute HIV infection. *J Infect Dis*. 2012; 206:275–282. [PubMed: 22551810]

Vythilingam M, Heim C, Newport J, Miller AH. Childhood trauma associated with smaller hippocampal volume in women with major depression. *Am J Psychiatry*. 2002; 159:2072–2080. [PubMed: 12450959]

Walton G, Co SJ, Milloy MJ, Qi J, Kerr T, Wood E. High prevalence of childhood emotional, physical and sexual trauma among a canadian cohort of HIV-seropositive illicit drug users. *AIDS Care*. 2011; 23:714–721. [PubMed: 21390877]

Socio-demographic and clinical characteristics of HIV-positive women with and without childhood trauma and HIV-negative women with and without childhood trauma

Table 1

	Total sample (N = 124)	HIV+ trauma (n = 32)	HIV+ no trauma (n = 30)	HIV – trauma (n = 31)	HIV – no trauma (n = 31)	F	p
Age	30.27 (7.94)	34.90 (6.60)	31.06 (6.92)	29.22 (9.02)	25.77 (6.32)	8.557	0.000**
Ethnicity							
African	120	32	28	31	29		
Coloured	4	0	2	0	2		
Education						8.904	0.000**
Mean (SD)	10.29 (1.79)	9.03 (2.20)	10.53 (1.19)	10.64 (1.58)	11.00 (1.36)		
< grade 8	16	11	2	2	1		
> grade 8	108	21	28	29	30		
Handedness						1.420	0.240
Left	7	1	3	0	3		
Right	117	31	27	31	28		
Home language							
Afrikaans	4	0	2	0	2		
isiXhosa	119	32	28	30	29		
Other	1	0	0	1	0		
Household income							
< R40,000/\$3714	119	32	27	30	30		
> R40,000	5	0	3	1	1		
Marital status							
Single	90	26	22	20	22		
Cohabiting/Married	31	5	8	9	9		
Separated/Divorced	3	1	0	2	0		
Employed							
Yes	32	6	11	6	9		
No	92	26	19	25	22		

	Total sample (N = 124)	HIV + trauma (n = 32)	HIV + no trauma (n = 30)	HIV – trauma (n = 31)	HIV – no trauma (n = 31)	F	p
ARV treatment							
Yes	30	7	23	0	0		
No	94	9	23	31	31		
CD 4 count						0.020	0.887
Mean	431.20	435.91	426.00	N/A	N/A		
SD	267.75	295.155	239.01	N/A	N/A		
Minimum	25	25	124	N/A	N/A		
Maximum	1529	1529	1053	N/A	N/A		
Viral load						0.892	0.349
Mean	116172.08	169754.10	62590.07	N/A	N/A		
SD	439037.76	605528.30	139822.78	N/A	N/A		
Minimum	0	0	0	N/A	N/A		
Maximum	3200000	3200000	641937	N/A	N/A		

p < 0.01

F = Analysis of Variance (ANOVA); HIV + = HIV positive; HIV – = HIV negative; African = South Africans belonging to the Black racial group; Coloured = South Africans belonging to the mixed race group; isiXhosa = one of the official indigenous African languages spoken in South Africa; Afrikaans = a West Germanic language spoken natively in South Africa

Table 2

Cortical volume measurements for select brain regions obtained using Freesurfer ($N = 124$)

Brain Region (volume/mm ³)	HIV+ with trauma ($n = 32$) M(SD)	HIV- with trauma ($n = 31$) M(SD)	HIV+ without trauma ($n = 30$) M(SD)	HIV- without trauma ($n = 31$) M(SD)	p
Left frontal lobe	725.84 (174.27)	783.22 (163.14)	713.60 (126.20)	779.96 (141.11)	0.400
Right frontal lobe	988.09 (239.23)	974.19 (222.51)	912.43 (165.85)	939.77 (158.70)	0.384
Left ACC	1547.06 (410.37)	1866.45 (554.08)	1675.33 (470.46)	1596.90 (410.84)	0.076
Right ACC	1773.34 (421.83)	2020.83 (444.02)	2098.86 (557.52)	2018.03 (480.70)	0.044*
Left hippocampus	3369.50 (461.37)	3591.54 (375.08)	3829.46 (435.37)	3631.19 (362.81)	0.000**
Right hippocampus	3385.09 (426.98)	3615.00 (333.11)	3779.03 (525.48)	3551.41 (462.85)	0.004**
Corpus Callosum	2731.84 (516.66)	3215.48 (534.92)	3052.36 (581.53)	3097.41 (438.96)	0.009**
Left Amygdala	1266.43 (211.40)	1355.41 (210.58)	1390.33 (168.74)	1315.16 (196.04)	0.054
Right Amygdala	1400.90 (247.36)	1451.58 (243.44)	1431.10 (193.84)	1410.61 (228.58)	0.569
Left caudate	3228.37 (438.61)	3564.51 (467.31)	3715.60 (501.04)	3513.22 (457.39)	0.003**
Right caudate	3459.56 (433.37)	3766.74 (474.05)	3885.10 (509.06)	3659.38 (429.03)	0.008**
Left putamen	5156.31 (639.81)	5872.45 (570.78)	5550.63 (768.79)	5633.74 (708.07)	0.006**
Right putamen	4999.71 (501.91)	5621.90 (619.95)	5426.06 (587.93)	5379.03 (647.67)	0.008**

Note: HIV+ = HIV positive; HIV- = HIV negative; ACC = Anterior Cingulate Cortex

* $p < 0.05$;

** $p < 0.01$

Table 3

Mean raw scores for significantly associated neurocognitive variables categorised by group

Mean raw score	HIV– no trauma (n = 31)	HIV+ no trauma (n = 30)	HIV+ with trauma (n = 32)	HIV– with trauma (n = 31)
Trail Making A *	54.61	53.56	69.12	51.58
Grooved Pegboard (non-dominant hand) *	75.58	76.66	97.12	77.51
WAIS-III Digit Symbol	53.19	49.83	36.62	51.54
HVLT learning	24.29	22.83	21.62	23.22
Category Fluency (animals)	12.64	12.16	10.65	12.67
COWAT	23.90	21.63	22.00	22.77
WMS-III Spatial Span	11.90	11.56	10.00	11.35
Color Trails 2 *	116.25	117.90	135.12	112.16
Stroop (Color-Word)	34.00	30.16	23.71	30.96
PASAT	23.38	20.76	17.59	24.93

Asterisk denotes that higher scores indicate poorer performance.

HIV + = HIV positive; HIV – = HIV negative; HVLT = Hopkins Verbal Learning Test; WAIS-III = Wechsler Adult Intelligence Scale; WMS-III = Wechsler Memory Scale; COWAT = Controlled Oral Word Association Task (with letters FAS for English, LBS for Afrikaans and IBS for Xhosa); Stroop = Stroop Color and Word Test; PASAT = Paced Auditory Serial Addition Test

Table 4

Childhood Trauma Questionnaire (CTQ) scores ($N = 124$)

Group		Physical abuse	Sexual abuse	Emotional abuse	Emotional neglect	Physical neglect	CTQ Total
HIV – no trauma ($n = 31$)	Mean	5.22	5.00	6.87	7.48	6.61	31.19
	Min	5.00	5.00	5.00	5.00	5.00	25.00
	Max	8.00	5.00	15.00	13.00	12.00	39.00
HIV + no trauma ($n = 30$)	Mean	5.96	5.33	6.03	8.83	6.90	33.06
	Min	5.00	5.00	5.00	5.00	5.00	25.00
	Max	10.00	9.00	12.00	19.00	13.00	40.00
HIV + with trauma ($n = 32$)	Mean	15.65	10.68	17.43	16.90	15.31	76.00
	Min	5.00	5.00	9.00	8.00	5.00	54.00
	Max	44.00	25.00	25.00	25.00	24.00	110.00
HIV – with trauma ($n = 31$)	Mean	10.06	8.35	15.48	11.87	9.96	55.74
	Min	5.00	5.00	5.00	5.00	5.00	41.00
	Max	25.00	25.00	71.00	25.00	18.00	116.00

Min = minimum; Max = maximum; HIV + = HIV positive; HIV – = HIV negative