CORRESPONDENCE

Screening and Treatment in Retinopathy of Prematurity

by PD Dr. med. Andreas Stahl and Prof. Dr. med. Wolfgang Göpel
in issue 43/2015

On Pathogenesis

Retinopathy of prematurity is a chronic retinal hypoxia, caused by the presence of high concentrations of fetal red blood cells containing fetal hemoglobin in the first months of life. Visible signs of hypoxia are retinal neovascularization. Fetal blood has a higher oxygen affinity, which causes retinal hypoxia. Time for manifestation, disease period, frequency, severity, and regression are all associated with the presence of fetal red blood cells containing fetal hemoglobin. In full-term newborns, the switch to adult hemoglobin is mostly completed, precluding this disease from occurring.

Thus, the difference between a preterm and a full-term newborn is that the fetal erythrocytes are still present in higher concentrations (according to gestational age) in a preterm newborn, while they have been mostly replaced by adult erythrocytes in the full-term newborn.

Fetal red blood cells and fetal hemoglobin are of causal significance for the incidence of retinopathy of prematurity. Therefore, prevention should have priority over therapy.

DOI: 10.3238/arztebl.2016.0328a

REFERENCES

1. Hammerstein W: Pathophysiologie und Prophylaxe der Frühgeborenenretinopathie. *Klin Mbl Augenheilk* 1991; 199: 177–82.
2. Stahl A, Göpel W: Screening and treatment in retinopathy of prematurity. *Dtsch Arztebl Int* 2015; 112: 730–5.

Prof. Dr. med. Wolfgang Hammerstein
Kyritz
prof.hammerstein@t-online.de

Conflict of interest statement

The author declares that no conflict of interest exists.

In Reply:

We thank the author for commenting on the role that oxygen affinity of fetal hemoglobin could play in the development of retinopathy of prematurity. This aspect must certainly be considered—in addition to numerous other pathomechanistically relevant factors—in the assessment of retinopathy of prematurity.

In this context, we would like to mention a question that is widely discussed at the moment, about the effects of erythropoietin on the risk of retinopathy of prematurity. Two recently published Cochrane Reviews have described an increased risk of developing

retinopathy of prematurity following iatrogenic doses of erythropoietin (1, 2). In addition to the time point at which erythropoietin administration begins, the duration of treatment may be equally relevant—an aspect that was less focused on in the Cochrane Reviews. Work on animal models has shown that erythropoietin administration for retinopathy of prematurity during stages with active pathological angiogenesis can further increase preretinal proliferation, thereby worsening those stages. In contrast, erythropoietin administration that was limited to the very early stages of retinopathy even had a protective effect in the animal models (3). In clinical reality, however, administration of erythropoietin is often frequently repeated and is necessary for prolonged periods of time. Perhaps most relevant to the risk of retinopathy of prematurity might therefore be those administration courses in which erythropoietin is given over longer periods, including during the phase of retinal angiogenic activation (phase II of retinopathy of prematurity).

In general, the aim should be to obtain the best possible postnatal management, which has the main initial goal of ensuring survival of the preterm infant, yet at the same time obtaining the lowest possible rate of stages of retinopathy of prematurity that require treatment. For those stages of retinopathy of prematurity that cannot be adequately controlled despite optimal conditions of systemic factors, it is essential to have a timely detection of indication for an ophthalmological treatment and possibly a targeted therapeutic intervention.

DOI: 10.3238/arztebl.2016.0328b

REFERENCES

1. Aher SM, Ohlsson A: Late erythropoietin for preventing red blood cell transfusion in preterm and/or low birth weight infants. *Cochrane Database Syst Rev* 2014; 4: CD004868.
2. Ohlsson A, Aher SM: Early erythropoietin for preventing red blood cell transfusion in preterm and/or low birth weight infants. *Cochrane Database Syst Rev* 2014; 4: CD004863.
3. Chen J, Connor KM, Aderman CM, Smith LEH: Erythropoietin deficiency decreases vascular stability in mice. *J Clin Invest* 2008; 118: 526–33.
4. Stahl A, Göpel W: Screening and treatment in retinopathy of prematurity. *Dtsch Arztebl Int* 2015; 112: 730–5.

PD Dr. med. Andreas Stahl
Klinik für Augenheilkunde, Universitätsklinikum Freiburg
andreas.stahl@uniklinik-freiburg.de

Prof. Dr. med. Wolfgang Göpel
Neonatologie, Universitätsklinikum Schleswig-Holstein, Campus Lübeck

Conflict of interest statement

Prof. Göpel has received travel and accommodation reimbursement and well as honoraria for preparation of scientific meetings from Chiese, AbbVie, and Novartis.

PD Dr. Stahl has received consultant fees, travel and accommodation reimbursement, honoraria for preparation of scientific meetings, and financial support for research (third-party funds) from Novartis