Published in final edited form as:

Bio Protoc. 2015 August 5; 3(15): .

Pulmonary Myeloperoxidase Activity

Tammy Regena Ozment*

Department of Surgery, East Tennessee State University Quillen College of Medicine, Johnson City, USA

Abstract

Neutrophils are considered one of the first responders of the innate immune response. Their primary activities are to migrate to sites of infection by chemotaxis and trans-migration across the endothelium (Gaines *et al.*, 2005). Once at the site of infection, they phagocytize microbes and kill them. Critical to the neutrophil's ability to kill microbes are the multiple degradative enzymes contained within granules. The activity of these enzymes is non-specific, and therefore, neutrophils also contribute to tissue damage at the site of infection (Gaines and Berliner, 2005). Measurement of neutrophil infiltration into tissues is one way to gauge the severity of infection, inflammation, and tissue damage (Ayala *et al.*, 2002). Myeloperoxidase is found in the primary granules of neutrophils and is an effective measure of neutrophil infiltration into tissues (Gaines and Berliner, 2005).

Materials and Reagents

1. Fresh or snap-frozen tissues
2. Liquid nitrogen
3. MPO Fluoremetric Detection Kit (Assay Designs, catalog number: 907-029)
4. N-ethylmaleimide (Sigma-Aldrich, catalog number: E2068)
5. Hexadecyltrimethylammonium (Sigma-Aldrich, catalog number: 52366)
6. Deionized water
7. DMSO (Sigma-Aldrich, catalog number: D8418)
8. Assay buffer (see Recipes)
9. Hydrogen Peroxide (see Recipes)
10. Detection Reagent (see Recipes)
11. Reaction cocktail (see Recipes)

*For correspondence: ozmentsk@etsu.edu.

Equipment

1. 14 ml Polystyrene test tubes
2. 2 ml Microfuge tubes
3. Black-welled 96 well plates
4. Polytron homogenizer
5. Sonicator
6. Fluorescent plate reader
7. Refrigerated centrifuge

Procedure

1. Weigh out 50 mg tissue into polycarbonate test tubes containing 0.5-1 ml ice cold 1× assay buffer with 10 mM N-ethylmaleimide.
2. Homogenize to just disrupt the tissue by placing the tip of the homogenizer in the bottom of the tube and switching the machine on, agitating the tube to move the tip of the homogenizer throughout the liquid and then switching the machine off after approximately one sec. Repeat until the tissues are just disrupted, usually 9 more times for a total of 10 pulses.
3. Centrifuge at $500 \times g$ for 10 min at 4 °C.
4. Discard supernatant, add 500 µl ice cold 1× assay buffer containing 0.5% hexadecyltrimethylammonium to the pellet and transfer to a 2 ml microfuge tube.
5. Homogenize to lyse the cells by placing the tip of the homogenizer in the bottom of the tube and switching the machine on, agitating the tube to move the tip of the homogenizer throughout the liquid and then switching the machine off after approximately 5 sec. Repeat 9 times for a total of 10 pulses. Place on ice.
6. Sonicate to further lyse the cells at 50% power for 10 sec. Place on ice. Repeat 2 times for a total of 3 pulses.
7. Snap freeze in liquid nitrogen and thaw at room temperature by placing in liquid nitrogen, immediately removing from the nitrogen and leave the samples at room temperature until completely thawed. Repeat snap freeze and thaw once.
8. Store at −80 °C until assayed.
9. Prior to assay, serially dilute the included MPO standard in assay buffer.

10. To assay, add 50 μ l of sample or the MPO standard to the bottom of a black 96 well plate. Add 50 μ l Reaction Cocktail provided by the kit. All samples and standards should be run in duplicate.
11. Incubate at room temperature in the dark for 30 min.
12. Read the fluorescence at 550 nm excitation and 595 nm emission every 10 min until 60 min incubation.
13. Plot the standard concentration vs. the maximum relative fluorescence units to create a standard curve (Figure 1).
14. Determine the concentration of the unknowns from the standard curve.

Recipes

1. 1 $\times$ Assay buffer 4 ml MPO assay buffer concentrate 36 ml deionized water
2. Hydrogen Peroxide 22.7 μ l 3% hydrogen peroxide 977 μ l deionized water
3. Detection Reagent Supplied vial 500 μ l DMSO
4. Reaction cocktail 50 μ l detection reagent 5 μ l hydrogen peroxide 4.875 ml 1 $\times$ assay buffer

Acknowledgments

This protocol is adapted from Ozment *et al.* (2012).

References

1. Ayala A, Chung CS, Lomas JL, Song GY, Doughty LA, Gregory SH, Cioffi WG, LeBlanc BW, Reichner J, Simms HH, Grutkoski PS. Shock-induced neutrophil mediated priming for acute lung injury in mice: divergent effects of TLR-4 and TLR-4/FasL deficiency. *Am J Pathol.* 2002; 161(6): 2283–2294. [PubMed: 12466142]
2. Gaines P, Chi J, Berliner N. Heterogeneity of functional responses in differentiated myeloid cell lines reveals EPRO cells as a valid model of murine neutrophil functional activation. *J Leukoc Biol.* 2005; 77(5):669–679. [PubMed: 15673544]
3. Ozment TR, Ha T, Breuel KF, Ford TR, Ferguson DA, Kalbfleisch J, Schweitzer JB, Kelley JL, Li C, Williams DL. Scavenger receptor class a plays a central role in mediating mortality and the development of the pro-inflammatory phenotype in polymicrobial sepsis. *PLoS Pathog.* 2012; 8(10):e1002967. [PubMed: 23071440]

Figure 1.
Representative Standard Curve