Published in final edited form as:

Horm Cancer. 2014 December ; 5(6): 390–404. doi:10.1007/s12672-014-0195-9.

The G Protein-Coupled Estrogen Receptor-1, GPER-1, Promotes Fibrillogenesis via a Shc-Dependent Pathway Resulting in Anchorage-Independent Growth

Hilary T. Magruder,

Division of Hematology and Oncology, Rhode Island Hospital and the Warren Alpert Medical School of Brown University, Providence, RI 02903, USA

Jeffrey A. Quinn,

Division of Hematology and Oncology, Rhode Island Hospital and the Warren Alpert Medical School of Brown University, Providence, RI 02903, USA

Jean E. Schwartzbauer,

Department of Molecular Biology, Princeton University, Princeton, NJ 08544, USA

Jonathan Reichner,

Division of Surgical Research, Rhode Island Hospital and the Warren Alpert Medical School of Brown University, Providence, RI 02903, USA

Allan Huang, and

Division of Surgical Research, Rhode Island Hospital and the Warren Alpert Medical School of Brown University, Providence, RI 02903, USA

Edward J. Filardo

Division of Hematology and Oncology, Rhode Island Hospital and the Warren Alpert Medical School of Brown University, Providence, RI 02903, USA

Edward J. Filardo: ed@radixbiosolutions.com

Abstract

The G protein-coupled estrogen receptor-1, GPER-1, coordinates fibronectin (FN) matrix assembly and release of heparan-bound epidermal growth factor (HB-EGF). This mechanism of action results in the recruitment of FN-engaged integrin $\alpha 5 \beta 1$ to fibrillar adhesions and the formation of integrin $\alpha 5 \beta 1$ -Shc adaptor protein complexes. Here, we show that GPER-1 stimulation of murine 4 T1 or human SKBR3 breast cancer cells with 17β -estradiol ($E2\beta$) promotes the formation of focal adhesions and actin stress fibers and results in increased cellular adhesion and haptotaxis on FN, but not collagen. These actions are also induced by the xenoestrogen, bisphenol A, and the estrogen receptor (ER) antagonist, ICI 182, 780, but not the inactive stereoisomer, 17α -estradiol ($E2\alpha$). In addition, we show that GPER-1 stimulation of breast cancer cells allows for FN-dependent, anchorage-independent growth and FN fibril

Correspondence to: Edward J. Filardo, ed@radixbiosolutions.com.

Present Address: E. J. Filardo, Radix BioSolutions, Inc., Georgetown, TX 78626, USA

Conflict of Interest The authors declare that they have no conflict of interest.

formation in “hanging drop” assays, indicating that these GPER-1-mediated actions occur independently of adhesion to solid substrata. Stable expression of Shc mutant Y317F lacking its primary tyrosyl phosphorylation site disrupts E2 β -induced focal adhesion and actin stress fiber formation and abolishes E2 β -enhanced haptotaxis on FN and anchorage-dependent growth. Collectively, these data demonstrate that E2 β action via GPER-1 enhances cellular adhesivity and FN matrix assembly and allows for anchorage-independent growth, cellular events that may allow for cellular survival, and tumor progression.

Introduction

Fibronectin (FN) plays a major role in cellular adhesion, growth, and survival, and it is important for processes such as wound healing [1], vascular growth [2], and embryonic development [3]. On the contrary, altered expression of FN, or perturbations in the specific recognition of FN by integrin $\alpha 5 \beta 1$, has been associated with the development of cancer and fibrosis [4, 5]. FN is synthesized in a soluble form as a dimeric glycoprotein that is assembled into an insoluble fibrillar matrix in a complex, dynamic cell-mediated process that is initiated by its specific recognition by integrin $\alpha 5 \beta 1$ via individual Arg-Gly-Asp (RGD)-binding sites on each monomer thereby facilitating integrin clustering. Upon FN engagement, integrin $\alpha 5 \beta 1$ undergoes conformational alterations associated with increased receptor affinity [6]. FN-occupied integrin $\alpha 5 \beta 1$ is then recruited to sites of close cell matrix contact known as “focal adhesions” that are enriched in tyrosyl-phosphorylated proteins and actin stress fibers where robust anchorage to FN occurs. The local concentration of integrin-bound FN increases, allowing bound FN molecules to more readily interact with one another and form short FN fibrils between cells, thus beginning the process of fibrillogenesis. Conversion of soluble FN to insoluble fibrils proceeds when cryptic FN-binding sites are exposed along the length of bound FN by contractile forces that stretch FN by pulling on their FN-bound integrin receptors [7] and partially unfolding FN, unmasking cryptic FN-binding sites [8, 9], and allowing nearby FN molecules to associate. This FN-FN interaction enables the soluble, cell-associated fibrils to branch and stabilize into an insoluble FN matrix. Fragmentation of FN uncovers non-RGD-binding sites leading to enhanced integrin $\alpha 4 \beta 1$ adhesion and FN matrix contractility [10], illustrating the influence of matrix proteases on provisional FN matrix assembly.

A number of studies have shown that FN is critical to normal homeostasis of the mammary gland and is associated with the development of breast cancer. Namely, the addition of exogenous FN negatively impacts acinar differentiation in the mammary gland and creates a microenvironment conducive to the growth of mammary epithelia [5]. Integrin $\alpha 5 \beta 1$ and FN are prominently expressed in the mammary gland, and their basal expression is increased during active proliferation of mammary gland tissue in mice suggesting that this FN-integrin interaction may be required for hormone-dependent proliferation in the mammary gland [11]. In addition, transgenic mice expressing dominant-negative integrin $\beta 1$ show disrupted mammary gland development that is associated in a loss of AKT activation and Shc-dependent extracellular regulated kinase-1 and -2 (Erk-1/-2) activation [12]. Moreover, successful implantation of human mammary tumor xenografts in immunocompromised mice is facilitated by coadministration of exogenous FN, indicating a survival advantage for

tumor cells that interact with FN [13]. This observation is supported by studies that have shown that mammary adenocarcinoma cells are capable of converting soluble FN into fibrils [14] resulting in increased responsiveness to growth factors and enhanced anchorage-independent growth [15]. The survival of tumor cells under these imposed in vitro growth conditions is reflective of their capacity to assemble a provisional extracellular matrix [14] and to circumvent death signals promoted by mechanosensors that report reduced tensile forces [16]. As measured in a two-dimensional environment, ligation of integrin $\alpha 5 \beta 1$ to FN-coated substrata is sufficient to promote intracellular signals associated with cellular growth and survival, including activation of Src, focal adhesion kinase (FAK), B/AKT, and Erk-1/-2 [17–20]. Under circumstances where either integrin density or its binding sites on adhesive proteins are limiting, serum-derived factors have been shown to facilitate the recruitment of tyrosyl-phosphorylated proteins (FAK, Src, vinculin, and paxillin) into focal adhesion plaques and induce the formation of RhoA-dependent actin stress fibers [21]. However, matrix engagement by integrin is not sufficient to promote subsequent cell growth responses, and FN polymerization is a critical requirement for measurable adhesion-dependent growth on planar surfaces [22]. Regulatory roles for phosphatidylinositol 3-OH kinase, FAK, Src-like kinases, and phospho-paxillin in FN matrix assembly have been suggested [23, 24]. Studies evaluating FN assembly in a three-dimensional environment have shown that exogenous FN can facilitate fibrillogenesis [25] but that this cellular activity is not sufficient for anchorage-independent growth by mammary adenocarcinoma cells, as they also must become responsive to external growth factors [14, 15, 26].

Shc is a nonenzymatic adaptor protein that participates in kinase signaling cascades and generally functions as a signaling intermediary to determine growth factor responsiveness and extracellular matrix (ECM) engagement [27]. Shc protein is expressed as three isoforms (p66Shc, p52Shc, and p46Shc) that are synthesized as a result of differential ribosomal initiation start sites on the same genetic locus. Their shared carboxyl terminus encodes PTB and SH2 phosphotyrosine-binding domains that are separated by an intervening collagen homologous (CH) 1 domain. Shc proteins are recruited to tyrosine kinase receptors and nonreceptor tyrosine kinases via interactions between their PTB and SH2 binding domains and phosphotyrosines [27]. On the other hand, Shc proteins physically associate with integrin via an indirect linkage involving the SH3 domain of Src-like kinases and the conserved proline-rich collagen homologous (CH1) domain on Shc that lies between PTB and SH2 [28]. While all three Shc isoforms are recruited to focal adhesions and share a primary tyrosine phosphorylation site and common functional motifs that promote their interaction with other signaling effectors, different cell biological roles have been assigned for the long and short forms of Shc. p52Shc and p46Shc have been linked to signals that promote cellular survival and proliferation, primarily through their ability to couple to the Raf-Mek-Erk protein kinase signaling axis via Grb-2/Sos/Ras [27]. In contrast, p66Shc uncouples the Ras-to-Erk protein kinase cascade and is associated with RhoA-dependent anoikis [29] and thus, is best known as a proapoptotic protein. This observation is consistent with the fact that p66Shc is poorly expressed in hematopoietic lineage cells that are considered to be anchorage-independent and insensitive to substrate stiffness [30–33]. Similarly, lung carcinoma cell lines that lack p66Shc display aggressive metastatic behavior, anchorage-independent growth, and bypass anoikis [34]. Finally, patients whose breast or

colon cancers express higher ratios of total tyrosyl-phosphorylated Shc to p66Shc are linked to poor prognosis [35, 36], further suggesting that failure to regulate Shc is associated with more advanced cancer.

Tumors that arise from the mammary gland exhibit biological behaviors that are described dichotomously as either estrogen- or growth factor-dependent. This categorization is largely derived from analysis of known receptors for estrogen (ERs) and epidermal growth factor (EGFR)-related receptors in breast tumor biopsies and the fact that there is a strong inverse relationship between expression of ER and EGFR [37]. G protein-coupled estrogen receptor-1 (GPER-1)/GPR30 represents a newly appreciated estrogen receptor whose expression in primary breast tumors is directly linked to tumor size and metastasis [38–41], a relationship diametrically opposed to the one shared between ER and these same tumor progression variables. This observation suggests that GPER-1 plays a distinct role from ER in breast cancer biology and is consistent with the fact that GPER-1 and ER are structurally distinct receptors that promote estrogen-mediated signals measured with different metrics and kinetics. While ER bears structural homology shared by the members of the nuclear steroid hormone receptor superfamily and functions as a hormone-inducible transcription factor, GPER-1 belongs to the most broadly studied class of cell surface receptors, the G protein-coupled receptor (GPCR) superfamily. GPER-1 promotes rapid signals attributed to this receptor class, including stimulation of adenylyl cyclase [42] and EGFR activation via the release of membrane-tethered heparan-bound epidermal growth factor (HB-EGF) [43]. We have previously shown that EGFR transactivation by GPER-1 requires activation of the FN receptor, integrin $\alpha 5\beta 1$, in breast cancer cells, as measured by its recruitment to fibrillar adhesions, the conversion of soluble FN to a detergent-insoluble form, and the association of integrin $\alpha 5\beta 1$ with the signaling adaptor, Shc [44].

Here, the influence of this GPER-1-integrin $\alpha 5\beta 1$ -Shc-dependent signaling mechanism on breast cancer cell adhesion was further evaluated by measuring its influence on cancer cell cytoarchitecture, adhesion, and haptotactic responses on immobilized FN and by evaluating their role in fibrillogenesis and growth in anchorage-independent conditions.

Materials and Methods

Cell Culture

SKBR3 (ER α -, ER β -, GPER-1+) breast cancer cells were obtained from the American Type Culture Collection (ATTC) (Manassas, VA). SKBR3 variants expressing dominant-negative Shc (Shc317Y/F) and dominant-negative GPER (GPER 154) were generated as described previously [44]. ER-negative murine 4 T1 breast cancer cells were obtained from the ATCC. 4 T1 cells expressing GPER 154 were generated as described previously [44]. All cells were grown in phenol red-free (PRF) DMEM/Ham's F12 media (1:1) with 5 % fetal bovine serum and 25 μ g/ml gentamicin.

Growth Factors, Estrogens, Antiestrogens, and Matrix Proteins

Water-soluble 17 β -estradiol (E2 β), 17 α -estradiol (E2 α), and angiotensin II (ATII) were purchased from Sigma (St. Louis, MO). Bisphenol A (BPA) was a kind gift from the Hixon

Lab at Brown University (Providence, RI). ICI 182, 780 was purchased from Tocris Bioscience (Ellisville, MO). Bovine, human, and rat FN were purchased from EMD Millipore (Milford, MA).

Antibodies

mAB IC3 specific for rat FN was a kind gift from the Schwarzbauer Lab at Princeton University (NJ) and has been previously described [45]. Phosphotyrosine-specific mAB, 4G10, was purchased from Upstate Biotechnology, Inc. Rabbit polyclonal antibodies (AB1949) specific for the cytoplasmic tail of integrin $\alpha 5$ subunit protein was purchased from Chemicon. Inhibitory rat anti-mouse integrin $\alpha 5\beta 1$ monoclonal antibody, clone BMB4, was purchased from Millipore. Alexa fluor dye-conjugated secondary antibodies were purchased from Molecular Probes, Inc. (Eugene, OR)/Invitrogen.

Cellular Stimulation and Protein Extraction

Conditions for quiescence, cell stimulation, and protein extraction were discussed previously [43].

Immunofluorescence

Focal adhesions were visualized in 4 T1 and SKBR3 cells that were seeded onto glass coverslips in PRF-DMEM/F12 medium containing FN-reduced serum and allowed to adhere overnight at 37 C. The following day, serum was removed by washing in PRF-DMEM/F12. Cells were then cultured in the same media in the absence of serum for an additional 30 h. Serum-starved cells were fed 2- μ g/ml rat FN in the absence or presence of ligand (10 nM E2 α , 10 nM E2 β , 1 μ M ICI 182, 780, 10 nM BPA) for 2 h. Cells were then washed, fixed for 5 min in 4 % paraformaldehyde, permeabilized in 0.05 % Triton for 60 s, and blocked in 5 % BSA/PBS for 30 min. Cells were incubated with phosphotyrosine-specific 4G10 antibody diluted 1:500 in PRF-DMEM/F12 containing 5 % BSA for 60 min. Coverslips were washed in PRF-DMEM/F12, and cell-associated antibodies were detected using Alexa 594-conjugated anti-mouse IgG diluted 1:1,000 and delivered in PRF-DMEM/F12 containing 5 % BSA for 30 min. After staining, coverslips were washed and mounted on glass slides in Vectashield/4'6-diamidino-2-phenylindole (Vector Laboratories, Inc., Burlingame, CA). FN fibril formation was examined in SKBR3, SKBR3 GPER 154, and SKBR3 Shc317Y/F, and 4 T1 cells that were seeded onto glass coverslips in PRF-DMEM/F12 medium containing FN-reduced serum. Starved cells were fed rat plasma FN (25 μ g/ml) in PRF-DMEM/F12 medium in the presence of ligand (10 nM E2 α , 10 nM E2 β , 1 μ M ICI 182, 780, 10 nM BPA) for 18 h and then fixed and prepared for immunostaining as above. Fixed cells were stained with IC3 ascites diluted 1:1,000 and delivered in PBS containing 1 % BSA for 60 min. IC3 mAB was detected by staining with Alexa 594-conjugated anti-mouse IgG (1:1,000) and processed for microscopy. Integrin $\alpha 5\beta 1$ and stress fibers were visualized in SKBR3, SKBR3 GPER 154, and SKBR3 Shc317Y/F cells that were seeded onto glass coverslips in PRF-DMEM/F12 medium containing FN-reduced serum and allowed to adhere overnight at 37 C. After adhesion, serum was removed by washing 3 $\times$ with PRF-DMEM/F12, and the cells were then cultured in the same media in the absence of serum for an additional 30 h. Serum-starved cells were fed 2- μ g/ml rat FN in the absence or presence of E2 β (10 nM) for 2 h. Cells were then washed, fixed for 5 min in 4 % paraformaldehyde, permeabilized in

0.05 % Triton for 60 s, and blocked in 5 % BSA/PBS for 30 min. Cells were incubated with rabbit polyclonal antibodies (AB1949) specific for the cytoplasmic tail of integrin $\alpha 5$ subunit protein diluted 1:500 and TRITC-phalloidin diluted 1:500 in PRF-DMEM/F12 containing 5 % BSA for 60 min. Coverslips were washed in PRF-DMEM/F12, and cell-associated antibodies were detected using Alexa 594-conjugated anti-rabbit IgG diluted 1:1,000 and delivered in PRF-DMEM/F12 containing 5 % BSA for 30 min. After staining, coverslips were washed and mounted on glass slides in Vectashield/4',6-diamidino-2-phenylindole (Vector Laboratories, Inc., Burlingame, CA). All immunofluorescent images were visualized with a Nikon Eclipse 80i microscope (Nikon, Inc., Melville, NY) equipped with a Nikon Plan Fluor 100 $\times$ 0.5–1.3 Oil Iris with differential interference contrast and epifluorescent capabilities. Digital images were captured using a QImaging Retiga 2000R digital camera and Nikon imaging software (Elements Basic Research 3.0).

Adhesion Assay

Forty-eight-well plates were coated with 200 μ l of PRF-DMEM/F12 containing 2- μ g/ml human FN or 10- μ g/ml collagen overnight. Wells were blocked with 5 % BSA PRF-DMEM/F12 for 1 h. SKBR3, SKBR3 GPER 154, and SKBR3 Shc317Y/F cells were seeded in triplicate, left untreated or treated with 10 nM E2 β , and allowed to adhere for 2 h. Nonadherent cells were gently washed away with PRF-DMEM/F12. Adherent cells were fixed and stained with 0.4 % crystal violet and 4 % ethanol in water for 5 min, and then washed 2 $\times$ in large volumes of water. Crystal violet dye was extracted with 10 % acetic acid, and absorbance was measured spectrophotometrically at 550 nm. All data points were determined from triplicate assays and expressed as the mean $\pm$ standard deviation. Nonspecific adhesion was subtracted as determined from cells that were seeded in BSA in the absence of substratum.

Boyden Chamber Migration Assay

Haptotaxis assays were conducted using modified Boyden chambers consisting of a porous polycarbonate membrane (6.5- μ m thickness, 8- μ m pores; Transwells, CoStar corporation, Cambridge, MA) [46]. The lower surfaces of the Transwell membrane were coated by adding 500 μ l of serum free, PRF-DMEM/F12 containing 2- μ g/ml human FN, or 10- μ g/ml collagen to the lower reservoir overnight. The underneath surface of the membrane was then blocked in 5 % BSA in for 1 h. SKBR3, SKBR3 GPER 154, and SKBR3 Shc317Y/F (10⁵) cells were placed in the upper reservoirs of the Transwell in serum free, PRF-DMEM/F12 and left untreated or treated with 10 nM E2 β and allowed to migrate overnight at 37 C. Nonmigrated cells were removed from the upper surface of the membrane using a Q-tip, and cells remaining attached to the lower surface were fixed in ethanol and stained with 0.4 % crystal violet in sodium borate buffer, pH 9.2 for 5 min, and then washed 2 $\times$ in large volumes of water. Dye was eluted from the migrant cells using acetic acid and measured spectrophotometrically at 550 nm. Each data point was measured in triplicate and measured as the mean plus or minus the standard deviation. Nonspecific migration was subtracted as determined from cells that were seeded in BSA in the absence of substratum.

Anchorage-Independent Growth

SKBR3, SKBR3 GPER 154, SKBR3 Shc317Y/F, 4 T1 vector, and 4 T1 GPER 154 cells (1×10^4) were seeded into PRF DMEM-F12 media in 0.35 % agarose in the absence or presence of E2 β (10 nM) and 10 % fetal bovine serum which was FN-depleted using gelatin-conjugated sepharose as described by Pierschbacher et al. [47] and supplemented with exogenous FN (2 μ g/ml). Cells were grown for 10 days at 37 C in a humidified chamber at 5 % CO₂. In some assays, inhibitory rat anti-integrin mouse $\alpha 5\beta 1$ monoclonal antibody, clone BMB4 from Millipore, or control nonimmune rat antibodies were incorporated in the agar overlay. Cultures were weighed every 2 days, and evaporated water was replaced as needed. Images of colonies were captured at $\times 100$ magnification (Brightfield). Colonies of greater than 20 cells were enumerated by direct counting.

Anchorage-Independent Fibrillogenesis

Anchorage-independent fibrillogenesis was determined from “hanging drop” assays as previously described [48–50]. Briefly, 4 T1 vector or 4 T1 GPER 154 cells were placed in suspension in serum-free media supplemented with or without agonist (10 nM E2 α , 10 nM E2 β , 10 nM ATII, 1 μ M ICI 182, 780, RGD, RGE) and rhodamine-labeled bovine FN (30 μ g/ml). For each treatment, 10 $\times$ 15- μ l aliquots were distributed onto the underside of a 100-mm Petri dish lid in a humidified chamber. Hanging drop cultures were incubated for 18 h at 37 C. Cells were fixed, stained with DAPI, and transferred to a glass slide. Images were captured using a Nikon 80i inverted fluorescent microscope fitted with a Retiga color camera at $\times 100$ magnification. Multiple Z-axis sections were reconstructed into three-dimensional images using Nikon imaging software.

Results

Estradiol-Induced Mobilization of $\alpha 5\beta 1$ into Focal Adhesions, the Formation of Actin Stress Fibers, and FN Fibril Formation Require GPER-1 and the Shc Signaling Adapter Protein

Src-like kinases, integrin $\alpha 5\beta 1$, and Shc have been identified as integral components of a signaling pathway leading to E2 β -mediated transactivation of the EGFR and FN matrix assembly [43, 44]. In the latter study, we showed that stimulation of human SKBR3 cells with E2 β results in the recruitment of FN-occupied integrin $\alpha 5\beta 1$ into fibrillar adhesions at the cell periphery [44] but did not address the role of GPER-1 or Shc in the formation of focal adhesion plaques, presumed precursory adhesion structures which give rise to fibrillar adhesions. To gain knowledge as to whether GPER-1 action was necessary for E2 β -induced clustering of integrin $\alpha 5\beta 1$ into focal adhesions and the formation of actin stress fibers, these cellular activities were compared in vector control SKBR3 cells and in a derivative line of SKBR3 cells expressing a dominant-negative form of GPER-1 (GPER 154) (Fig. 1). In these experiments, SKBR3 vector control and SKBR3 GPER 154 cells were seeded onto coverslips, serum-starved, and stimulated with E2 β in the presence of exogenous FN for 2 h. Cells were fixed and stained with integrin $\alpha 5\beta 1$ -specific antibodies (green) and rhodamine-conjugated phalloidin to identify polymerized actin stress fibers (red). While vector control SKBR3 cells demonstrated prominent actin stress fibers that colocalized with integrin $\alpha 5\beta 1$ in focal adhesions, neither actin stress fibers nor integrin $\alpha 5\beta 1$ -enriched focal adhesions

were observed in unstimulated cells (Fig. 1a). Similarly, SKBR3 cells expressing a dominant-negative form of GPER-1 (GPER 154) that were stimulated with E2 β were also unable to form actin stress fibers or concentrate integrin $\alpha 5\beta 1$ into focal adhesions (Fig. 1a) suggesting that GPER-1 was required for the cellular activation of events that recruit integrin $\alpha 5\beta 1$ to focal adhesions and promote actin stress fiber formation. As previously demonstrated, GPER-1 stimulation with E2 β resulted in the formation of FN fibrils (red), while dominant-negative GPER-1 compromised FN fibril formation (Fig. 1a).

To establish the requirement of Shc in E2 β -induced integrin $\alpha 5\beta 1$ recruitment to focal adhesions and the formation of actin stress fibers and FN fibril formation, these cellular events were evaluated in SKBR3 cells expressing control vector or a mutant Shc protein lacking its primary tyrosyl phosphorylation site, Shc317Y/F (Fig. 1b). Focal adhesions and actin stress fibers were not detectable in quiescent, unstimulated control or Shc317Y/F cells. Following exposure to E2 β , integrin $\alpha 5\beta 1$ was recruited into prominent focal adhesions that coaligned with the termini of actin stress fibers, while SKBR3 Shc317Y/F cells showed an impaired ability with regards to these integrin activation events. Expression of Shc317Y/F in SKBR3 cells negatively affected E2 β -induced clustering of integrin $\alpha 5\beta 1$ into focal adhesions and showed less prominent focal adhesion plaques that appeared to be disordered with regards to their alignment with the termini of focal adhesions (Fig. 1b). Likewise, E2 β -induced FN fibril formation observed in control SKBR3 cells was prohibited in SKBR3 Shc317Y/F cells.

Collectively, these results indicate that Shc signaling following GPER-1 activation is required to promote the recruitment of integrin $\alpha 5\beta 1$ to focal adhesions and to induce actin stress fiber formation and consequent FN fibril formation. We have previously shown that Shc317Y/F accumulates relative to Shc wild-type protein on integrin $\alpha 5\beta 1$ and cells expressing this mutant Shc protein fail to form fibrillar adhesions [44]. Thus, our current observations may further suggest that the primary tyrosyl phosphorylation site on Shc is not required for entry of integrin $\alpha 5\beta 1$ into focal adhesions but that this site is required for its subsequent recruitment to fibrillar adhesions.

The Influence of the Xenoestrogen, Bisphenol A, or the ER Antagonist, ICI 182, 780, on the Recruitment of Tyrosyl-Phosphorylated Proteins to Focal Adhesions and the Production of Fibronectin Fibrils

Xenoestrogens and ER antagonists act as GPER-1 agonists [40, 43, 51, 52]. To determine whether these estrogenic steroids also influence GPER-1-dependent formation of focal adhesions and FN fibrils, these integrin activation events were assessed in SKBR3 cells expressing vector or GPER 154 protein (Fig. 2a). For these experiments, cells were made quiescent by seeding them onto glass coverslips in FN-reduced serum followed by serum deprivation. Quiescent cells were left untreated or exposed to E2 β ; its inactive stereoisomer (E2 α), the xenoestrogen, BPA, or the ER antagonist, ICI 182, 780 and focal adhesions were assessed by measuring the clustering of tyrosyl-phosphorylated proteins by immunostaining with the phosphotyrosine (pptyr)-specific monoclonal antibody, 4G10. Stimulation with E2 β , BPA, or ICI 182, 780 resulted in the recruitment of tyrosyl-phosphorylated proteins into focal adhesion plaques (Fig. 2a). Focal adhesions were similarly measured in murine 4

T1 breast cancer cells that were stimulated with the ER antagonist, ICI 182, 780, or the xenoestrogen, BPA (data not shown). Focal adhesions were neither measured in quiescent cells nor in cells that were stimulated with E2 α (Fig. 2a) nor EGF (data not shown). These results suggest that estrogen-stimulated enrichment of tyrosyl-phosphorylated proteins to focal adhesion plaques occurs independently of the ER and does not require EGF stimulation.

Previously, we have shown that stimulation of human SKBR3 cells with E2 β results in the recruitment of FN-occupied integrin $\alpha 5 \beta 1$ to the cell periphery into fibrillar adhesions, specialized adhesion structures at which FN fibrils form [44]. Here, we addressed the capacity of SKBR3 cells to form FN fibrils in response to stimulation with ICI 182, 780 or BPA. As shown in Fig. 2b, either ICI 182, 780 or BPA as well as E2 β resulted in the formation of FN fibrils that were detected at the periphery of SKBR3 cells. FN fibril formation was not observed in SKBR3 cells that were left untreated or stimulated with E2 α (Fig. 2b) or EGF (data not shown). Similar observations were measured in murine 4 T1 breast cancer cells. In both cell types, expression of GPER 154 protein prohibited FN fibril formation (Fig. 2b and data not shown), demonstrating that these estrogenic hormones are capable of promoting FN fibril formation (fibrillogenesis) in ER-negative breast cancer cells that are attached to planar surfaces coated with this ECM protein.

Stimulation of GPER-1 Selectively Enhances the Adhesivity of Human Breast Cancer Cells for FN-Coated Substrata by a Mechanism that Requires the Primary Tyrosyl Phosphorylation Site on Shc

In many instances, agonists that employ G protein-coupled receptors promote enhanced cellular adhesive interactions by modulating the affinity of integrins for their cognate ECM proteins and also by inducing the recruitment of integrins to focal adhesion plaques, a process referred to as “inside-out” integrin signaling [53, 54]. To examine the influence of GPER-1 stimulation by E2 β on the adhesion of SKBR3 vector control breast cancer cells for immobilized adhesive ligand, SKBR3 or SKBR3 GPER 154 cells were detached and exposed to E2 β or left untreated and seeded into polystyrene wells coated with various concentrations of FN or collagen I (COLL) (Fig. 3). Following a 2-h incubation time at 37 C, cells that were not firmly attached were gently washed away, adherent cells were fixed, cellular attachment was assessed by staining the remaining adherent cells with crystal violet, and attachment was measured as a function of eluted dye recovered from the resulting adherent cells. Unstimulated SKBR3 cells adhered to both FN- and COLL-coated substrata in a dose-dependent fashion with maximum cell adhesion measured at coating concentrations of 20 and 10 $\mu\text{g/ml}$ of FN and COLL, respectively (data not shown). A 2-fold increase ($p=0.034$) in the capacity of E2 β -stimulated versus unstimulated SKBR3 cells to adhere to wells coated with suboptimal concentrations of FN (2 $\mu\text{g/ml}$) was observed (Fig. 3a), which was associated with increased cellular spreading. In contrast, more modest differences in enhanced E2 β -mediated adhesion to COLL-coated substrata were measured (Fig. 3) with no discernible difference in cellular spreading (data not shown). E2 β -increased adhesivity was eliminated in SKBR3 cells expressing GPER 154 suggesting that E2 β promotes adhesion of SKBR3 breast cancer cells in a GPER-1-dependent manner. Similarly, Shc was tested for its involvement in GPER-1-enhanced adhesivity by comparing the

relative capacity of SKBR3 and SKBR3 Shc317Y/F cells to adhere to immobilized FN or COLL (Fig. 3b). There was no significant increase ($p=0.46$) in adhesion between the E2 β -stimulated and untreated SKBR3 Shc317Y/F cells, suggesting that Shc is also involved in the signaling events that lead to increased adhesion to FN.

Cellular adhesion to planar surfaces is greatly strengthened by cell spreading [55] and is often associated with increased cellular motility as measured in haptotactic responses on immobilized adhesive ligands. To examine whether GPER-1 and Shc promote increased migration on FN-coated substrata, SKBR3 vector control, SKBR3 GPER 154, or SKBR3 Shc317Y/F cells were seeded in the presence of 10 nM E2 β or left untreated) into the upper reservoirs of a modified Boyden chambers containing a porous polycarbonate membrane (10- μ m thickness, 8- μ m pore) whose undersurface was coated with adhesive ligand (2 μ g/ml FN or 10 μ g/ml COLL) (Fig. 4). Cell migration was measured by determining the number of cells that were capable of migrating from the upper reservoir across the membrane to its undersurface. On FN-coated membranes, SKBR3 vector control cells stimulated with E2 β showed a 6.5-fold increase in their capacity to migrate compared with unstimulated cells in this assay (Fig. 4). SKBR3 cells that were plated onto membranes that were coated with higher concentrations of FN (5–20 μ g/ml) did not show increased haptotaxis when stimulated with E2 β (data not shown), suggesting that GPER-1-enhanced migration was the product of increased recruitment of FN receptors to cellular adhesion sites (data not shown). E2 β -enhanced haptotaxis on FN was abrogated in SKBR3 GPER 154 cells demonstrating that this migratory response was dependent upon GPER-1 action (Fig. 4). Likewise, expression of Shc317Y/F also impeded E2 β -enhanced haptotaxis on FN (Fig. 4). SKBR3 cells migrated equally well on COLL-coated substrata independent of E2 β stimulation, and Shc317Y/F had no impact on COLL migration suggesting that GPER-1 signaling did not enhance migration on this ECM protein (data not shown).

GPER-1 Promotes Anchorage-Independent Growth by Promoting Fibrillogenesis in a Three-Dimensional Environment

Since the conversion of soluble FN into fibrils by mammary adenocarcinoma cells has been linked to enhanced anchorage-independent growth [14] and GPER-1 action results in fibrillogenesis on planar surfaces (Figs. 1 and 2) [44], the possibility that GPER-1-mediated fibrillogenesis is required for anchorage-independent growth was evaluated (Figs. 5 and 6). To address this hypothesis, human SKBR3 or murine 4 T1 breast cancer cells were seeded into semisolid media supplemented with FN-depleted fetal bovine serum in the presence of increasing amounts of exogenous FN. Under conditions of FN depletion, neither SKBR3 nor 4 T1 cells were able to form colonies whereas both cell types readily formed colonies in the presence of 10 % fetal bovine serum which had not been FN-depleted (data not shown and Fig. 6a). However, either cell line readily formed colonies in FN-reduced conditions with 2- μ g/ml exogenous FN (Figs. 5 and 6) provided that E2 β was also present. Neither cell line grew in the presence of exogenous E2 α and FN. Expression of GPER 154 in either cell background effectively prohibited E2 β -dependent, anchorage-independent growth (Figs. 5 and 6b), suggesting that these growth properties are GPER-1-dependent. The specificity of GPER 154 for inhibiting GPER-1 action in this assay was demonstrated by the observation that the substitution of E2 β for exogenous angiotensin II (ATII), which acts through its

dedicated cognate G protein-coupled receptor, could restore FN-dependent, anchorage-independent growth (data not shown and Fig. 6b). Expression of Shc317Y/F in the SKBR3 cell background had an inhibitory effect on colony formation suggesting that Shc is also required for E2 β -dependent growth in semisolid media (Fig. 5). Inclusion of inhibitory anti-integrin $\alpha 5\beta 1$ antibodies in semisolid media containing murine 4 T1 cells showed that integrin $\alpha 5\beta 1$ engagement is necessary for anchorage-independent growth under these conditions of limiting FN and exogenous E2 β (Fig. 6c). Collectively, these data indicate that GPER-1 stimulation and exogenous FN is required for estradiol-dependent, anchorage-independent growth.

To directly address the capacity of GPER-1 to promote fibrillogenesis in a three-dimensional environment, murine 4 T1 vector control or 4 T1 GPER 154 breast cancer cells were cultured in hanging drop assays in which rhodamine-labeled FN was incorporated [48–50]. As demonstrated in Fig. 7, 4 T1 cells that were cultured in suspension (in the absence of a substratum) overnight in the presence of E2 β were capable of forming FN fibrils in this anchorage-independent assay (Fig. 7a). Similarly, FN fibrils were also measured by 4 T1 cells that were stimulated with ATII. FN fibril formation was not measured in untreated or E2 α -treated cells. GPER 154 specifically inhibited fibrillogenesis in hanging drop cultures of 4 T1 cells that were stimulated by E2 β but had no effect on ATII-induced fibrils (Fig. 7a). The ER antagonist, ICI 182, 780, also induced the formation of FN fibrils in the hanging drop assay (Fig. 7b), and this cellular activity was inhibited by GPER 154 (data not shown). Moreover, hanging drop cultures of E2 β -stimulated 4 T1 breast cancer cells that were supplemented with soluble RGD peptide were unable to form FN fibrils; however, control RGE peptide did not have a negative effect on anchorage-independent fibrillogenesis (Fig. 7b). Taken together, the data in Figs. 5, 6 and 7 suggests that GPER-1 promotes anchorage-independent growth through its ability to synthesize FN fibrils.

Discussion

Evidence is provided here that estrogenic hormones act via GPER-1 to enhance integrin $\alpha 5\beta 1$ -dependent adhesion of breast cancer cells to FN via a signaling mechanism that requires tyrosyl phosphorylation of the Shc adaptor protein. Specifically, we show that GPER-1 stimulation promotes the following: (i) the formation of focal adhesions leading to the reorganization of actin stress fibers; (ii) enhanced cellular adhesivity and haptotaxis on immobilized FN; (iii) anchorage-independent FN fibril formation; and (iv) FN-dependent, anchorage-independent growth. We have previously reported that integrin $\alpha 5\beta 1$ and Shc are integral components of a signaling pathway leading to estrogen-mediated EGFR transactivation and FN matrix assembly on planar surfaces [44]. Collectively, these data support a model suggesting that GPER-1 coordinates two key cellular events required for the survival of breast cancer cells that escape the confines of glandular epithelia and invade the surrounding tissue parenchyma, namely, responsiveness to soluble growth factors and the capacity to form a provisional ECM (Fig. 8). Our findings are consistent with studies in mice that have shown a requirement for exogenous FN for efficient tumor cell implantation [56] and integrin $\beta 1$ and Shc for homeostasis of the mammary gland [12].

Estrogenic hormones regulate mammary gland homeostasis and, in certain instances, influence the cellular behavior of tumors that are derived from this tissue. However, the mechanism by which estrogen regulates FN-adhesive function has remained unclear. A direct influence of estrogen or the xenoestrogen, resveratrol, on cell behavior has been shown by studies measuring cytoarchitectural alterations in estrogen receptor-negative cells cultured in serum in response to hormone stimulation [57, 58]. Similar cytostructural changes, including enhanced actin stress fiber formation and the establishment of prominent focal adhesions, have been measured in long-term cultures of ER-positive MCF-7 breast cancer cells [59, 60] or endometrial cancer cells [57] stimulated with tamoxifen or the pure ER antagonist, ICI 182, 780, suggesting that alternative estrogen receptors may influence the interaction of these cancer cells derived from the female reproductive tract with their ECM. Here, we provide direct evidence that ICI 182, 780 or BPA act via the membrane estrogen receptor, GPER-1, to promote focal adhesion plaque formation, actin stress fiber assembly, and fibrillogenesis (Figs. 1 and 2). A result that is consistent with other reports that BPA acts via GPER-1 to induce rapid signaling effects [61] and gene transactivation [52] by breast cancer cells.

Modulation of integrin affinity for their adhesive ligands is commonly accomplished as the result of intracellular signals initiated by the interaction of external soluble mediators with GPCRs, a process referred to as inside-out integrin signaling [53]. For example, affinity upregulation of integrin $\alpha\text{IIb}\beta 3$ for its ligand fibrinogen, a key event in thrombus formation, occurs in response to stimulation of platelets with ADP, epinephrine, or thrombin, whose receptors are GPCRs [62]. Likewise, $\beta 1$ and $\beta 2$ integrins on leukocytes exhibit increased affinity for their adhesive proteins in response to a broad array of immunomodulatory substances that act through GPCRs, including, but not limited to, f-Leu-Met-Phe, chemokine, and complement cascade products [63]. Similarly, our findings here indicate that estrogen action via GPER-1 activates integrin $\alpha 5\beta 1$ resulting in its recruitment to focal adhesions (Figs. 1 and 2), increased adhesion and haptotaxis on planar surfaces coated with FN (Figs. 3 and 4), and FN matrix assembly in two- and three-dimensional environments (Figs. 1, 2, and 5, 6 and 7) by breast cancer cells. Since the experiments presented here do not directly address conformational alterations in the external domains of integrin $\alpha 5\beta 1$ associated with FN affinity, it is not possible to formally conclude whether GPER-1-enhanced cellular adhesivity occurs via inside-out signaling. However, our findings are consistent with prior observations that have shown that GPCR activation promotes allosteric changes within the ligand-binding domains of $\beta 1$, $\beta 2$, and $\beta 3$ integrins resulting in enhanced adhesive function [63–65]. Our findings suggest that enhanced integrin $\alpha 5\beta 1$ adhesive action of breast cancer cells for FN does not appear to be solely relegated to signaling by GPER-1, as angiotensin II stimulation of breast cancer cells also enhanced fibrillogenesis (Fig. 6).

The adaptor protein Shc is intrinsically involved in intracellular signaling events that determine growth factor responsiveness and bidirectional integrin signaling [27]. Our work presented here provides additional support for this idea by showing that GPER-1 mediates enhanced integrin $\alpha 5\beta 1$ -Shc-dependent adhesivity and further supports the concept that integrin $\alpha 5\beta 1$ and Shc form a signaling node that regulates FN matrix assembly and the release of membrane-tethered HB-EGF by SKBR3 breast cancer cells [44]. Further support for integrin $\alpha 5\beta 1$ and Shc as an integration point in growth factor responsiveness and ECM

signaling is demonstrated by the report that vascular endothelial cells coordinate FN-dependent adhesivity and vEGF receptor activation [66]. Similar to the work by Sweet and colleagues in vascular cells [66], we show that Shc is required for enhanced adhesion of breast cancer cells to FN-coated substrata, but not collagen (Fig. 3). This is consistent with our prior report that recruitment of Shc to integrin $\alpha 2\beta 1$, a primary collagen receptor for SKBR3 breast cancer cells, occurred in the absence of stimulation and was not further enhanced by GPER-1 activation [44]. In this context, the data presented here support the concept that Shc recruitment is associated with enhanced cellular adhesion to FN, but not collagen. However, this difference may simply reflect synchrony in Shc activation that results from plating unattached cells to their adhesive ligands and does not account for the fact that Shc-integrin recruitment is likely a dynamic process that is associated with assembly and disassembly of focal adhesion structures [67]. It is interesting that Wary et al. [68] further noted that interaction of integrin $\alpha 5\beta 1$ or $\alpha v\beta 3$ with FN resulted in cellular proliferation, while engagement of integrin $\alpha 2\beta 1$ or integrin $\alpha 6\beta 1$ with their adhesive ligands results in cell cycle withdrawal, suggesting that differential cellular responses to extracellular matrices of different compositions may depend on the ability of a class of integrins to activate Shc signaling [68].

Shc couples with integrin during matrix engagement [68–70] and localizes to focal adhesions in attached cells [71], adhesive events associated with increased contractility tension at integrin anchorage points [29, 72]. These observations suggest that Shc is able to regulate integrin affinity by its capacity to interpret adhesive interactions with the ECM that is, in turn, determined by its phosphorylation status and interactions with integrin and proteins that accumulate in focal adhesions. The physical association with integrin is determined by an indirect interaction between the SH3 domain of Src-like kinases and the conserved proline-rich CH1 domain encoded within Shc while its conserved PTB domain directs focal adhesion targeting [29]. p66Shc uniquely contains an additional collagen homology domain, CH2, at its N-terminus; however, simple mutational analysis studies indicate that CH2 does not influence targeting to integrin or focal adhesions. Consistent with results reported by others examining the requirements for Shc complex formation with other integrins [64, 65], the primary tyrosyl phosphorylation site on Shc (Tyr317 on p52Shc) is dispensable with regards to integrin association [44], although it serves as an SH2 binding site that promotes the recruitment of Grb2 and a number of proteins that concentrate in focal adhesions, including FAK and Src-like kinases [73, 74]; tyrosine phosphatases, PTPN12, SHP-2 [69]; and tyrosine-phosphorylated proteins such as paxillin, talin, and vinculin [75]. In fact, expression of Shc317Y/F did not block GPER-1-enhanced association of endogenous Shc isoforms with immunopurified integrin $\alpha 5\beta 1$, but rather resulted in its accumulation relative to the endogenous p52 and p46Shc isoforms [44], suggesting a possible role for protein tyrosine phosphatases (PTPs) for the release of Shc from integrin $\alpha 5\beta 1$. This hypothesis is consistent with the concept that PTPs are well known to regulate focal adhesion disassembly, cell spreading, and migration [76–78]. The concept that focal adhesion formation is a dynamic process may suggest that disassembly and reassembly of Shc-integrin complexes is necessary for the formation of more ordered focal adhesions and actin stress cables that may be aligned through SH2 binding proteins [67]. Interestingly, although expression of Shc317Y/F did block GPER-1-mediated FN fibril formation [44], it

did not completely impede GPER-1-enhanced recruitment of integrin $\alpha 5\beta 1$ into focal adhesions as integrin $\alpha 5\beta 1$ recruitment occurred in the context of mutant Shc, but resulted in the formation of poorly organized focal adhesions which were not properly aligned with actin stress fibers (Fig. 1). It is noteworthy that SKBR3 317Y/F cells express approximately equivalent amounts of p52Shc317Y/F relative to endogenous wild-type p52Shc and p46Shc isoforms [44]; yet, mutant Shc accumulates on integrin $\alpha 5\beta 1$ relative to wild-type Shc following GPER-1 stimulation. This Shc-mediated defect is also associated with a complete blockade of GPER-1-mediated FN matrix assembly and EGFR transactivation [44], cellular activities that lie downstream of integrin recruitment to focal adhesions. One plausible explanation for this observation is that both mutant and wild-type Shc are equally recruited to integrin $\alpha 5\beta 1$ during integrin clustering and the initial phases of focal adhesion formation but that the presence of Shc317Y/F inhibits its dissociation from integrin $\alpha 5\beta 1$, an event that may be linked to focal contact elongation and disassembly, events that may be required for subsequent centripetal movement of integrin $\alpha 5\beta 1$ to fibrillar adhesions and release of membrane-associated proHB-EGF. Anchorage-independent growth by tumor cells is the best-known predictor of experimental metastasis in mice and the best measure of resident stem cells in human tumors [14, 15]. In our study, we shown that estrogen action via GPER-1 leads to FN-dependent, anchorage-independent growth by human SKBR3 and murine 4 T1 breast cancer cells (Figs. 5 and 6). Prior work investigating the capacity of mammary adenocarcinoma cells to undergo anchorage-independent growth demonstrated a correlation between their capacity to convert soluble FN into fibrils and increased responsiveness to growth factor [14, 15]. As discussed above, the p66Shc isoform appears to play a unique role in sensing cell adhesion as p66Shc promotes anoikis via RhoA activation in detached cells [29]. These findings are consistent with studies which have shown that lung cancer cells lacking p66Shc display traits associated with advanced cancer, including anchorage-independent growth [34]. However, GPER-1-enhanced adhesivity to FN does not simply appear to be a result of preferential recruitment of the p52Shc or p46Shc isoforms to integrin $\alpha 5\beta 1$ as human MDA-MB-231 breast cancer cell lines that express GPER-1 and p66Shc remain competent with regards to their capacity to promote integrin $\alpha 5\beta 1$ -dependent EGFR transactivation and enhanced FN adhesivity and anchorage-independent growth [Quinn, Magruder, and Filardo, unpublished results]. Interestingly, a direct linear increase in tumor stem cell activity of human SKBR3 cells has been reported as measured by their capacity to form mammospheroids in low adhesive media following serial passage through immunocompromised mice treated with epirubicin [79]. This finding suggests that chemotherapeutic drugs indirectly provide an environment that favors FN matrix assembly and anchorage-independent growth. It is tempting to speculate that in this harsh environment in which GPER-1-positive breast cancer cells that were presented with estrogenic hormones may show enhanced tumor stem cell activity and anchorage-independent growth.

Taken together, the data presented here support the hypothesis that GPER-1 signals via integrin $\alpha 5\beta 1$ and Shc to coordinate growth factor responsiveness and anchorage-independent growth, events critical for breast cancer progression. GPER-1 expression in primary breast tumors directly varies with tumor size and metastasis, markers of advanced disease, a relationship that is diametrically opposed to that shared by ER and these same prognostic variables [38]. The more recent finding that GPER-1 expression in triple negative

breast cancer directly varies with disease progression underscores a potential role for this newly appreciated estrogen receptor in advanced disease [41]. Our work here, defining the cell biological influence of GPER-1 stimulation on FN matrix assembly and adhesion to this ECM protein, provides insight into the cellular mechanisms by which estrogens may promote the survival of metastatic breast cancer cells.

Acknowledgments

This work was supported, in part, by a Research Scholar Award from the American Cancer Society (RSG-02-194-01) and by National Institutes of Health Award CA119165-01A2. We would also like to thank the Hixon Lab at Brown University for their kind gift of Bisphenol A for our studies.

References

1. Stoffels JM, Zhao C, Baron W. Fibronectin in tissue regeneration: timely disassembly of the scaffold is necessary to complete the build. *Cell Mol Life Sci*. 2013; 70:4243–4253. [PubMed: 23756580]
2. Astrof S, Hynes RO. Fibronectins in vascular morphogenesis. *Angiol*. 2009; 12:165–175.
3. Francis SE, Goh KL, Hodivala-Dilke K, Bader BL, Stark M, Davidson D, Hynes RO. Central roles of $\alpha 5 \beta 1$ integrin and fibronectin in vascular development in mouse embryos and embryoid bodies. *Arterioscler Thromb Vasc Biol*. 2002; 22:927–933. [PubMed: 12067900]
4. Faraldo MM, Deugnier MA, Thiery JP, Glukhova MA. Growth defects induced by perturbation of $\beta 1$ -integrin function in the mammary gland epithelium result from a lack of MAPK activation via the Shc and Akt pathways. *EMBO Rep*. 2001; 2:431–437. [PubMed: 11375936]
5. Williams CM, Engler AJ, Slone RD, Galante LL, Schwarzbauer JE. Fibronectin expression modulates mammary epithelial cell proliferation during acinar differentiation. *Cancer Res*. 2008; 68:3185–3192. [PubMed: 18451144]
6. Faull RJ, Kovach NL, Harlan JM, Ginsberg MH. Affinity modulation of integrin $\alpha 5 \beta 1$: regulation of the functional response by soluble fibronectin. *J Cell Biol*. 1993; 121:155–162. [PubMed: 8458867]
7. Hocking DC, Sottile J, Langenbach KJ. Stimulation of integrin-mediated cell contractility by fibronectin polymerization. *J Biol Chem*. 2000; 275:10673–10682. [PubMed: 10744764]
8. Ingham KC, Brew SA, Huff S, Litvinovich SV. Cryptic self-association sites in type III modules of fibronectin. *J Biol Chem*. 1997; 272:1718–1724. [PubMed: 8999851]
9. Klotzsch E, Smith ML, Kubow KE, Muntwyler S, Little WC, Beyeler F, Gourdon D, Nelson BJ, Vogel V. Fibronectin forms the most extensible biological fibers displaying switchable force-exposed cryptic binding sites. *Proc Natl Acad Sci U S A*. 2009; 106:18267–18272. [PubMed: 19826086]
10. Valenick LV, Hsia HC, Schwarzbauer JE. Fibronectin fragmentation promotes $\alpha 4 \beta 1$ integrin-mediated contraction of a fibrin-fibronectin provisional matrix. *Exp Cell Res*. 2005; 309:48–55. [PubMed: 15992798]
11. Woodward TL, Mienaltowski AS, Modi RR, Bennett JM, Haslam SZ. Fibronectin and the $\alpha 5 \beta 1$ integrin are under developmental and ovarian steroid regulation in the normal mouse mammary gland. *Endocrinol*. 2001; 142:3214–3222.
12. Faraldo MM, Deugnier MA, Tlouzeau S, Thiery JP, Glukhova MA. Perturbation of $\beta 1$ -integrin function in involuting mammary gland results in premature dedifferentiation of secretory epithelial cells. *Mol Biol Cell*. 2002; 13(10):3521–3531. [PubMed: 12388754]
13. Price JE. Metastasis from human breast cancer cell lines. *Breast Cancer Res Treat*. 1996; 39:93–102. [PubMed: 8738609]
14. Saulnier R, Bhardwaj B, Klassen J, Leopold D, Rahimi N, Tremblay E, Mosher D, Elliott B. Fibronectin fibrils and growth factors stimulate anchorage-independent growth of murine mammary carcinoma. *Exp Cell Res*. 1996; 222:360–369. [PubMed: 8598224]
15. Qiao H, Saulnier R, Patrykat A, Rahimi N, Raptis L, Rossiter J, Tremblay E, Elliott B. Cooperative effect of hepatocyte growth factor and fibronectin in anchorage-independent survival

- of mammary carcinoma cells: requirement for phosphatidylinositol 3-kinase activity. *Cell Growth Differ.* 2000; 11:123–133. [PubMed: 10714768]
16. Discher DM, Noy D, Strazalka J, Ye S, Moser CC, Lear JD, Blasie JK, Dutton PL. Design of amphiphilic protein maquettes: controlling assembly, membrane insertion, and cofactor interactions. *Biochemistry.* 2005; 44(37):12329–12343. [PubMed: 16156646]
 17. Frisch SM, Francis H. Disruption of epithelial cell-matrix interactions induces apoptosis. *J Cell Biol.* 1994; 124(4):619–626. [PubMed: 8106557]
 18. Frisch SM, Vuori K, Ruoslahti E, Chan-Hui PY. Control of adhesion-dependent cell survival by focal adhesion kinase. *J Cell Biol.* 1996; 134:793–799. [PubMed: 8707856]
 19. Martin SW, Butcher AJ, Berrow NS, Richards MW, Paddon RE, Turner DJ, Dolphin AC, Sihra TS, Fitzgerald EM. Phosphorylation sites on calcium channel alpha 1 and beta subunits regulate ERK-dependent modulation of neuronal N-type calcium channels. *Calcium.* 2006; 39(3):275–292.
 20. Green JA, Berrier AL, Pankov R, Yamada KM. beta1 integrin cytoplasmic domain residues selectively modulate fibronectin matrix assembly and cell spreading through talin and Akt-1. *J Biol Chem.* 2009; 284:8148–8159. [PubMed: 19144637]
 21. Ridley AJ, Hall A. The small GTP-binding protein rho regulates the assembly of focal adhesions and actin stress fibers in response to growth factors. *Cell.* 1992; 70:389–399. [PubMed: 1643657]
 22. Sottile J, Hocking DC, Swiatek PJ. Fibronectin matrix assembly enhances adhesion-dependent cell growth. *J Cell Sci.* 1998; 111:2933–2943. [PubMed: 9730985]
 23. Wierzbicka-Patynowski I, Schwarzbauer JE. Regulatory role for SRC and phosphatidylinositol 3-kinase in initiation of fibronectin matrix assembly. *J Biol Chem.* 2002; 277:19703–19708. [PubMed: 11912200]
 24. Wierzbicka-Patynowski I, Mao Y, Schwarzbauer JE. Continuous requirement for pp 60-Src and phospho-paxillin during fibronectin matrix assembly by transformed cells. *J Cell Physiol.* 2007; 210:750–756. [PubMed: 17187346]
 25. Mao Y, Schwarzbauer JE. Stimulatory effects of a three-dimensional microenvironment on cell mediated fibronectin fibrillogenesis. *J Cell Sci.* 2005; 118:4427–4436. [PubMed: 16159961]
 26. Accornero P, Miretti S, Cucuzza LS, Martignani E, Baratta M. Epidermal growth factor and hepatocyte growth factor cooperate to enhance cell proliferation, scatter, and invasion in murine mammary epithelial cells. *J Mol Endocrinol.* 2010; 44:115–125. [PubMed: 19850646]
 27. Ravichandran KS. Signaling via Shc family adapter proteins. *Oncogene.* 2001; 20:6322–6330. [PubMed: 11607835]
 28. Wary KK, Mariotti A, Zurzolo C, Gianetti FG. A requirement of caveolin-1 and associated kinase Fyn in integrin signaling and anchorage-dependent cell growth. *Cell.* 1998; 94(5):625–634. [PubMed: 9741627]
 29. Ma Z, Myers DP, Wu RF, Nwariaku FE, Terada LS. p66Shc mediates anoikis through RhoA. *J Cell Biol.* 2007; 179:23–31. [PubMed: 17908916]
 30. Migliaccio E, Mele S, Salcini AE, Pelicci G, Lai KM, Superti-Furga G, Pawson T, Di Fiore PP, Lanfrancone L, Pelicci PG. Opposite effects of the p52shc/p46shc and p66shc splicing isoforms on the EGF receptor-MAP kinase-fos signalling pathway. *EMBO J.* 1997; 16:706–716. [PubMed: 9049300]
 31. Bonati A, Carlo-Stella C, Lunghi P, Albertini R, Pinelli S, Migliaccio E, Sammarelli G, Savoldo B, Tabilio A, Dall'Aglio PP, Pelicci PG. Selective expression and constitutive phosphorylation of SHC proteins [corrected] in the CD34+ fraction of chronic myelogenous leukemias. *Cancer Res.* 2000; 60:728–732. [PubMed: 10676660]
 32. Zhang L, Lorenz U, Ravichandran KS. Role of Shc in T-cell development and function. *Immunol Rev.* 2003; 191:183–195. [PubMed: 12614360]
 33. Giles AJ, Bender TP, Ravichandran KS. The adaptor protein Shc plays a key role during early B cell development. *J Immunol.* 2009; 183:5468–5476. [PubMed: 19828641]
 34. Ma Z, Liu Z, Wu RF, Terada LS. p66(Shc) restrains Ras hyperactivation and suppresses metastatic behavior. *Oncogene.* 2010; 29:5559–5567. [PubMed: 20676142]
 35. Frackelton AR Jr, Lu L, Davol PA, Bagdasaryan R, Hafer LJ, Sgroi DC. p66 Shc and tyrosine-phosphorylated Shc in primary breast tumors identify patients likely to relapse despite tamoxifen therapy. *Breast Cancer Res.* 2006; 8:R73. [PubMed: 17196107]

36. Grossman SR, Lyle S, Resnick MB, Sabo E, Lis RT, Rosinha E, Liu Q, Hsieh CC, Bhat G, Frackelton AR Jr, Hafer LJ. p66 Shc tumor levels show a strong prognostic correlation with disease outcome in stage IIA colon cancer. *Clin Cancer Res*. 2007; 13:5798–5804. [PubMed: 17908971]
37. Lee CS, Hall RE, Alexander IE, Koga M, Shine J, Sutherland RL. Inverse relationship between estrogen receptor and epidermal growth factor receptor mRNA levels in human breast cancer cell lines. *Growth Factors*. 1990; 3(2):97–103. [PubMed: 2206558]
38. Filardo EJ, Graeber CT, Quinn JA, Resnick MB, Giri D, DeLellis RA, Steinhoff MM, Sabo E. Distribution of GPR30, a seven-membrane-spanning estrogen receptor, in primary breast cancer and its association with clinicopathological determinants of tumor progression. *Clin Cancer Res*. 2006; 12:6359–6366. [PubMed: 17085646]
39. Arias-Pulido H, Royce M, Gong Y, Joste N, Lomo L, Lee SJ, Chaheer N, Verschraegen C, Lara J, Prossnitz ER, Cristofanilli M. GPR30 and estrogen receptor expression: new insights into hormone dependence of inflammatory breast cancer. *Breast Cancer Res Treat*. 2010; 123:51–58. [PubMed: 19902352]
40. Ignatov A, Ignatov T, Weissenborn C, Eggemann H, Bischoff J, Semczuk A, Roessner A, Costa SD, Kalinski T. G-protein-coupled estrogen receptor GPR30 and tamoxifen resistance in breast cancer. *Breast Cancer Res Treat*. 2011; 128:457–466. [PubMed: 21607586]
41. Steiman J, Peralta EA, Louis S, Kamel O. Biology of the estrogen receptor, GPR30, in triple negative breast cancer. *Am J Surg*. 2013; 206:698–703. [PubMed: 24011568]
42. Filardo EJ, Quinn JA, Frackelton AR, Bland KI. Estrogen action via the G protein-coupled receptor, GPR30: stimulation of adenylyl cyclase and cAMP-mediated attenuation of the epidermal growth factor receptor-to-MAPK signaling axis. *Mol Endocrinol*. 2002; 16:70–84. [PubMed: 11773440]
43. Filardo EJ, Quinn JA, Bland KI, Frackelton AR. Estrogen-induced activation of Erk-1 and Erk-2 requires the G-protein coupled receptor homolog, GPR30, and occurs via the epidermal growth factor receptor through release of HB-EGF. *Mol Endocrinol*. 2000; 14:1649–1660. [PubMed: 11043579]
44. Quinn JA, Graeber T, Frackelton R, Kim M, Schwarzbauer JE, Filardo EJ. Coordinate regulation of estrogen-mediated fibronectin matrix assembly and epidermal growth factor receptor transactivation by the G protein-coupled receptor, GPR30. *Mol Endocrinol*. 2009; 23:1052–1064. [PubMed: 19342448]
45. Sechler JL, Takada Y, Schwarzbauer JE. Altered rate of fibronectin matrix assembly by deletion of the first type III repeats. *J Cell Biol*. 1996; 134:573–583. [PubMed: 8707839]
46. Filardo EJ, Brooks PC, Deming SL, Damsky C, Cheresch DA. Requirement of the NPXY motif in the integrin $\beta 3$ subunit cytoplasmic tail for melanoma cell migration in vitro and in vivo. *J Cell Biol*. 1995; 130:441–450. [PubMed: 7542248]
47. Pierschbacher MD, Hayman EG, Ruoslahti E. Location of the cell-attachment site in fibronectin with monoclonal antibodies and proteolytic fragments of the molecule. *Cell*. 1981; 26(2):259–267. [PubMed: 6174240]
48. Foty R. A simple hanging drop cell culture protocol for generation of 3D spheroids. *J Vis Exp*. 2011; 6(51)
49. Bartosh TJ, Ylostalo JH. Preparation of anti-inflammatory mesenchymal stem/precursor cells (MSCs) through sphere formation using hanging-drop culture technique. *Curr Protoc Stem Cell Biol*. 2014; 6(28)
50. Archacka K, Pozzobon M, Repele A, Rossi CA, Campanella M, De Coppi P. Culturing muscle fibres in hanging drop: a novel approach to solve an old problem. *Biol Cell*. 2014; 106(2):72–82. [PubMed: 24405025]
51. Tsai C, Wu H, Lin C, Lin Y, Chao A, Wang T, Hsueh S, Lai C, Wang H. Estradiol and tamoxifen induce cell migration through GPR30 and activation of focal adhesion kinase (FAK) in endometrial cancers with low or without nuclear estrogen receptor α (ER α). *PLoS ONE*. 2013; 8(9):e72999. [PubMed: 24039841]
52. Pupo M, Pisano A, Lappano R, Santolla MF, De Francesco EM, Abonante S, Rosano C, Maggiolini M. Bisphenol A induces gene expression changes and proliferative effects through

- GPER in breast cancer cells and cancer-associated fibroblasts. *Environ Health Perspect.* 2012; 120(8):1177–1182.
53. Ginsberg MH, Partridge A, Shattil SJ. Integrin regulation. *Curr Opin Cell Biol.* 2005; 17(5):509–516. [PubMed: 16099636]
 54. Luo BH, Carman CV, Springer TA. Structural basis of integrin regulation and signaling. *Annu Rev Immunol.* 2007; 25:619–647. [PubMed: 17201681]
 55. O'Toole TE, Loftus JC, Plow EF, Glass AA, Harper JR, Ginsberg MH. Efficient surface expression of platelet GPIIb-IIIa requires both subunits. *Blood.* 1989; 74(1):14–18. [PubMed: 2752106]
 56. Price J. Metastasis from human breast cancer cell lines. *Breast Cancer Res Treat.* 1996; 39(1):93–102. [PubMed: 8738609]
 57. Acconcia F, Barnes CJ, Kumar R. Estrogen and tamoxifen induce cytoskeletal remodeling and migration in endometrial cells. *Endocrinol.* 2006; 147:1203–1212.
 58. Azios NG, Krishnamoorthy L, Harris M, Cubano LA, Cammer M, Dharmawardhane SF. Estrogen and resveratrol regulate Rac and Cdc42 signaling to the actin cytoskeleton of metastatic breast cancer cells. *Neoplasia.* 2007; 9(2):147–158. [PubMed: 17356711]
 59. Sapino A, Pietribiasi F, Bussolati G, Marchisio PC. Estrogen- and tamoxifen-induced rearrangement of cytoskeletal and adhesion structures in breast cancer MCF-7 cells. *Cancer Res.* 1986; 46:2526–2531. [PubMed: 2421880]
 60. Ehlers EM, Schubert C. Differences in morphology and cytoskeleton of MCF-7 and MX-1 cells after therapy with OH-tamoxifen and the pure estrogen antagonist ZM 182780. An immunofluorescence and scanning electron microscopic study. *Ann Anat.* 1999; 181:231–236. [PubMed: 10363104]
 61. Dong S, Terasaka S, Kiyama R. Bisphenol A induces a rapid activation of Erk1/2 through GPR30 in human breast cancer cells. *Environ Pollut.* 2011; 159(1):212–218. [PubMed: 20875696]
 62. Cossemans JMEM, Iserbyt BF, Deckmyn H, Heemskerk JMW. Multiple ways to switch platelet integrins on and off. *J Thromb Haemost.* 2008; 6(8):1253–1261. [PubMed: 18513212]
 63. Oberyshyn T, Conti C, Ross M, Oberyshyn A, Tober K, Rackoff A, Robertson F. $\beta 2$ integrin/ICAM-1 adhesion molecule interactions in cutaneous inflammation and tumor promotion. *Carcinogenesis.* 1998; 19(3):445–455. [PubMed: 9525279]
 64. Philips DR, Prasad KS, Manganello J, Bao M, Nannizzi-Alaimo L. Integrin tyrosine phosphorylation in platelet signaling. *Curr Opin Cell Biol.* 2001; 13(5):541–554. [PubMed: 11544021]
 65. Cowan KJ, Law DA, Phillips DR. Identification of Shc as the primary protein binding to the tyrosine-phosphorylated $\beta 3$ subunit of $\alpha \text{IIb}\beta 3$ during outside-in integrin platelet signaling. *J Biol Chem.* 2000; 275:36423–36429. [PubMed: 10964917]
 66. Sweet DT, Chen Z, Wiley DM, Bautch VL, Tzima E. The adaptor protein Shc integrates growth factor and ECM signaling during postnatal angiogenesis. *Blood.* 2012; 119:1946–1955. [PubMed: 22096252]
 67. Barberis L, Wary KK, Fiucci G, Liu F, Hirsch E, Brancaccio M, Altruda F, Tarone G, Giancotti FG. Distinct roles of adaptor protein Shc and focal adhesion kinase in integrin signaling to ERK. *J Biol Chem.* 2000; 275:36532–36540. [PubMed: 10976102]
 68. Wary K, Mainiero F, Isakoff SJ, Marcantonio EE, Giancotti FG. The adaptor protein Shc couples a class of integrins to the control of cell cycle progression. *Cell.* 1996; 87:733–743. [PubMed: 8929541]
 69. Habib T, Herrera R, Decker SJ. Activators of protein kinase C stimulate association of Shc and the PEST tyrosine phosphatase. *J Biol Chem.* 1994; 269(41):25243–25246. [PubMed: 7929214]
 70. Deshmukh L, Gorbatyuk V, Vinogradova O. Integrin $\beta 3$ phosphorylation dictates its complex with the Sch phosphotyrosine-binding (PTB) domain. *J Biol Chem.* 2010; 285:34875–34884. [PubMed: 20739287]
 71. Angers-Loustau A, Cote JF, Charest A, Dowbenko D, Spencer S, Lasky LA, Tremblay. Protein tyrosine phosphatase-PEST regulates focal adhesion disassembly, migration, and cytokinesis in fibroblasts. *J Cell Biol.* 1999; 144(5):1019–1031. [PubMed: 10085298]
 72. Debnath J. p66Shc and Ras: controlling anoikis from the inside-out. *Oncogene.* 2010; 29:5556–5558. [PubMed: 20711240]

73. Charest A, Wagner J, Jacob S, McGlade CJ, Tremblay ML. Phosphotyrosine-independent binding of Shc to the NPLH sequence of murine protein-tyrosine phosphatase-PEST. Evidence for extended phosphotyrosine binding/phosphotyrosine interaction domain recognition specificity. *J Biol Chem.* 1996; 271(14):8424–2429. [PubMed: 8626541]
74. Sachdev S, Bu Y, Gelman IH. Paxillin-Y118 phosphorylation contributes to the control of Src-induced anchorage-independent growth by FAK and adhesion. *BMC Cancer.* 2009; 9(12)
75. Petita B, Thiery JP. Focal adhesions: structure and dynamics. *Biol Cell.* 2000; 92:477–494. [PubMed: 11229600]
76. Arregui CO, Balsamo J, Lilien J. Impaired integrin-mediated adhesion and signaling in fibroblasts expressing a dominant-negative mutant PTP1B. *J Cell Biol.* 1998; 143(3):861–873. [PubMed: 9813103]
77. Yu DH, Qu CK, Henegariu O, Lu X, Feng GS. Protein-tyrosine phosphatase Shp-2 regulates cell spreading, migration, and focal adhesion. *J Biol Chem.* 1998; 273(33):21125–21131. [PubMed: 9694867]
78. Zheng Y, Lu Z. Regulation of tumor cell migration by protein tyrosine phosphatase (PTP)-proline-, glutamate-, serine-, and threonine-rich sequence (PEST). *Chin J Cancer.* 2013; 32(2):75–83. [PubMed: 23237212]
79. Yu F, Yao H, Zhu P, Zhang X, Pan Q, Gong C, Huang Y, Hu X, Su F, Lieberman J, Song E. Let-7 regulates self renewal and tumorigenicity of breast cancer cells. *Cell.* 2007; 131:1109–1123. [PubMed: 18083101]

Fig. 1.

E2 β -induced recruitment of integrin $\alpha 5\beta 1$ to focal adhesions and the formation of actin stress fibers and FN fibrils are GPER- and Shc-dependent. **a** (*top panel*) SKBR3 cells expressing vector or GPER 154 and **b** (*top panel*) SKBR3 cells expressing vector or Shc317Y/F were seeded onto coverslips, serum starved, and stimulated with E2 β (10 nM) and exogenous rat FN (2 μ g/ml). Cells were incubated for 2 h, fixed with 4 % paraformaldehyde, permeabilized with detergent, and stained with integrin $\alpha 5\beta 1$ -specific antibodies (*green*) and Alexa-594-phalloidin (*red*). Nuclei were stained with DAPI (*blue*). **a** (*bottom panel*) SKBR3 cells expressing vector or GPER 154 and **b** (*bottom panel*) SKBR3 cells expressing vector or Shc317Y/F were seeded onto coverslips, serum starved, and stimulated with E2 β (10 nM) and exogenous rat FN (25 μ g/ml). Cells were incubated for 18 h, fixed with 4 % paraformaldehyde, and stained with rat FN-specific mAB, IC3 (*red*). Nuclei were stained with DAPI (*blue*)

Fig. 2.

E2β stimulation of human ER-negative breast cancer cells induces the formation of focal adhesions and FN fibrils. **a** Human SKBR3 cells grown on glass coverslips were serum-starved and stimulated with either E2β (10 nM), E2α (10 nM), BPA (10 nM), or ICI 182,780 (1 μM), and incubated for 2 h in the presence of exogenous rat FN (2 μg/ml). Following incubation, cells were fixed in 4 % paraformaldehyde, permeabilized with detergent, and focal adhesions were detected using phosphotyrosine-specific antibodies (*red*). Nuclei were stained with DAPI (*blue*). **b** SKBR3 cells grown on glass coverslips were serum-starved and stimulated as described above. Cells were incubated for 18 h in the presence of exogenous rat FN (25 μg/ml), fixed in 4 % paraformaldehyde, and FN fibrils were detected using rat FN-specific mAb, IC3 (*red*). Nuclei were stained with DAPI (*blue*)

Fig. 3.

GPER-1 stimulation promotes SKBR3 cell adhesion onto FN-coated, but not collagen-coated, substrata in a Shc-dependent manner. **a** SKBR3 cells expressing vector or GPER 154 and **b** SKBR3 cells expressing vector or Shc317Y/F were seeded onto 48-well plates coated with FN (2 μg/ml) or collagen (10 μg/ml) and allowed to attach for 2 h at 37 C in the absence or presence of E2β (10 nM). After adhesion, unattached cells were removed by gentle washing, and the remaining adherent cells were fixed and stained with crystal violet. Excess crystal violet was washed away, and cell-associated crystal violet was extracted with 10 % acetic acid. Absorbance was measured at 550 nm. Each data point represents the mean±standard deviation of triplicate samples. Nonspecific adhesion as measured on BSA-coated wells has been subtracted

Fig. 4. GPER-1 stimulation enhances Shc-dependent haptotaxis of human SKBR3 breast cancer cells on FN-coated, but not collagen-coated, substrata. SKBR3 vector, GPER 154, or Shc317Y/F cells were left untreated or treated with E2β (10 nM) and seeded into the transwells of modified Boyden chambers containing a porous polycarbonate membrane (10-μm thickness, 8-μm pore) that were left untreated or coated with either FN (2 μg/ml) or collagen (10 μg/ml) and incubated overnight at 37 C. Nonmigrant cells were removed from the top of each chamber, and migrant cells on the lower surface of the membranes were fixed and stained with crystal violet. The number of migrated cells were counted and images were taken at×40 magnification

Fig. 5.

GPER-1 enhances FN-dependent, anchorage-independent growth of SKBR3 cells. SKBR3 vector, GPER 154, or Shc317Y/F cells were seeded into phenol red-free DMEM-F12 media containing 2 % FN-reduced serum in 0.35 % agarose in the absence or presence of E2β (10 nM) and supplemented with exogenous FN (2 μg/ml). Cells were grown for 10 days at 37 C in a humidified chamber. Cultures were weighed every 2 days, and water was replaced as needed. Images of colonies were captured at×10 magnification (*brightfield*). Examples shown above are representative views of multiple experiments

Fig. 6.

E2 β stimulation alters colony morphology of ER-negative mouse breast cancer cells grown in soft agar. **a** 4 T1 cells were seeded into phenol red-free DMEM-F12 media containing 2 % FN-reduced serum in 0.35 % agarose in the absence or presence of E2 β (10 nM) and supplemented with exogenous FN (2 μ g/ml). Cells were grown for 10 days at 37 C in a humidified chamber. Cultures were weighed every 2 days, and water was replaced as needed. Images of colonies were captured at $\times 10$ magnification (*brightfield*). Examples shown above are representative views of multiple experiments. **b** 4 T1 vector or 4 T1 GPER 154 cells were seeded into phenol red-free DMEM-F12 media containing 2 % serum in 0.35 % agarose and left untreated or stimulated with E2 β (10 nM) or ATII (10 nM) in the presence or absence of exogenous FN (2 μ g/ml). Growth conditions were the same as above. **c** 4 T1 cells were seeded into phenol red-free DMEM-F12 media containing 2 % serum in 0.35 % agarose and left untreated or stimulated with E2 β (10 nM), ICI 182, 780 (1 μ M), IgG and E2 β (10 nM), or anti-integrin $\alpha 5\beta 1$ and E2 β (10 nM) in the presence or absence of exogenous FN (2 μ g/ml). Growth conditions are the same as above

Fig. 7.

GP130 promotes FN fibril formation of mouse 4 T1 breast cancer cells cultured in hanging drops. **a** 4 T1 vector or GP130^{-/-} cells were placed in suspension in serum-free media supplemented with or without E2 α (10 nM), E2 β (10 nM), or angiotensin II (ATII) (10 nM), and rhodamine-labeled bovine FN (30 μ g/ml). For each treatment, 10 $\times$ 15- μ l aliquots were distributed onto the underside of a 100-mm Petri dish lid in a humidified chamber. Hanging drop cultures were incubated for 18 h at 37 C. Cells were fixed, stained with DAPI, and transferred to a glass slide. Images were captured using a Nikon 80i inverted fluorescent microscope fitted with a Retiga color camera at $\times$ 100 magnification. Multiple Z-axis sections were reconstructed into three-dimensional images using Nikon imaging software. Examples shown above are representative views of multiple experiments. **b** 4 T1 vector cells were placed in serum-free media supplemented with or without ICI 182, 780 (1 μ M), or E2 β (10 nM) with RGD or RGE, and rhodamine-labeled bovine FN (30 μ g/ml). Hanging drops were incubated and analyzed as described in **a**

Fig. 8.

Proposed mechanism of E2 β activation of integrin $\alpha 5\beta 1$ via GPER-1 signaling. E2 β binding to GPER-1 at the plasma membrane promotes the dissociation of heterotrimeric G proteins into an α subunit and a $\beta\gamma$ subunit on the cytoplasmic side of the plasma membrane. The $\beta\gamma$ subunit promotes activation of Shc and its recruitment to integrin $\alpha 5\beta 1$ and the formation of focal adhesions. Dissociation of Shc from integrin $\alpha 5\beta 1$ is required for its subsequent centripetal movement to fibrillar adhesions and the release of membrane-tethered proHB-EGF